The impact of antenatal balanced plate nutrition education for pregnant women on

birth weight: a cluster randomized controlled trial in rural Bangladesh

Authors:

Morseda Chowdhury[1,2], Camille Raynes-Greenow[2], Patrick J. Kelly[2], Ashraful Alam[2],

Kaosar Afsana[3], Michael J. Dibley[2]

Affiliations:

[1]Health Nutrition and Population Programme, BRAC, BRAC Centre, 75 Mohakhali, Dhaka

1212, Bangladesh

[2]The University of Sydney, Faculty of Medicine and Health, Sydney School of Public Health,

Camperdown, New South Wales 2006, Australia

[3]James P Grant School of Public Health (JPGSPH), BRAC University, Dhaka, Bangladesh

Corresponding author: Michael J Dibley, Room 328 Edward Ford Building (A27) Sydney

School of Public Health, the University of Sydney, NSW 2006, Australia.

Phone: +61 2 9351 520   Email: michael.dibley@sydney.edu.au

Short running head: Balanced plate nutrition education & birth weight

Abstract

Background: Low birth weight (LBW) is a global public health problem with the highest prevalence in South Asia. It is strongly associated with maternal under nutrition. In South Asia, intra household food distribution is inequitable with lower dietary adequacy in women, regardless of pregnancy. Evidence that nutrition education improves diet during pregnancy and reduces LBW is weak.

Objective: We aimed to assess the impact of a nutrition education intervention for pregnant women on infant birth weight in a low-income population.

Design: We conducted a parallel, two arm, cluster-randomized controlled trial in rural Bangladesh, with 36 clusters randomized to receive the intervention (n=445 participants) or standard care (n=448). Eligible participants were ≤12 weeks gestation. Intervention participants received education about eating balanced meals to meet daily dietary requirements with the right proportions of food groups (a balanced plate) and practical demonstration from their first trimester until delivery. Primary outcomes were mean birth weight and the incidence of LBW. We included all participants in analyses.

Results: The mean birth weight difference between the intervention and control groups was 127.5 g (95% confidence interval (CI): 11.1, 243.9; p=0.032), and the intervention reduced the risk of LBW by 57% (relative risk (RR): 0.43; 95% CI: 0.25, 0.75; p=0.003). Post hoc analyses showed a higher birth weight and a greater reduction in LBW amongst adolescent mothers (mean difference: 299.1 g; 95% CI: 101.6, 496.6; p=0.003 and RR: 0.28; 95% CI: 0.11 to 0.71; p=0.007). The mean number of food groups consumed was significantly higher from the third month of pregnancy onwards in the intervention compared to the control group.

41   **Conclusions:** A community-based balanced plate nutrition education intervention effectively

42   increased mean birth weight and reduced the proportion of infants with LBW in a rural

43   population of Bangladesh.

44   **Keywords:** Low birth weight, maternal nutrition education, food demonstrations, dietary

45   diversity, cluster RCT, Bangladesh

Introduction

Birth weight is a powerful predictor of infant survival.[1] Low birth weight, defined as less than 2500 grams (g), is a composite measure of restricted fetal growth and short duration of gestation, major determinants of infant mortality, childhood growth and development, and adulthood well-being.[1,2] The lower the birth weight, the higher the risk of mortality.[3,4] Globally, in 2010, there were an estimated 18 million LBW babies, of whom 59% were term, but small for gestational age and 41% were preterm.[5] An estimated 606,500 neonatal deaths occur annually in low and middle-income countries due to small for gestational age, which accounts for 22% of neonatal deaths.[6] South Asia has the highest prevalence of LBW globally (34%).[6] Bangladesh is ranked fourth in the global burden of LBW.[5]

LBW is strongly associated with maternal under-nutrition[7], which induces suboptimal placental growth, thereby influencing nutrient delivery, altering fetal hormones, and contributing to restricted development.[8] A World Health Organization study with 111,000 women reported that mothers in the lowest quartile of pre-pregnancy weight carried an elevated risk of IUGR of 2.55 (95% confidence interval (CI): 2.3, 2.7) and LBW of 2.38 (95% CI: 2.1, 2.5) compared to the upper quartile.[9] In India, the odds ratio for LBW was three times greater in severe energy-deficient low BMI groups than normal BMI. [10] Consistent with the mechanisms and epidemiological evidence is the high prevalence of LBW in rural Bangladeshi women who have a chronic shortfall of food energy throughout pregnancy.[11] In South Asia, intra-household food distribution is inequitable with lower energy share, animal source food, and dietary adequacy in women, regardless of pregnancy.[12]

There is little data on nutrition education's effects on birth weight in low or middle-income countries, although four meta-analyses have assessed impacts. The first in 1993 found nutrition education significantly increased the consumption of protein and vitamins, and birth

weight of infants by 300 g. However, the studies lacked statistical power or did not report

estimates of effect size.[13] The second meta-analysis in 2012 found a 105 g (95% CI: 18, 193

g) increase in mean birth weight with nutrition education and counseling in pregnancy. When

stratified by the study site, the effect failed to reach significance in low and middle-income

countries (152 g; 95% CI: -81, 384 g).[14] The third meta-analysis in 2014 reported that dietary

counselling significantly increased birth weight in three studies (149 g; p=0·039), but two had

a small sample size, and another had methodological limitations.[15] The fourth, a Cochrane

review,[16] identified two small trials that reported significantly higher birth weight in the

intervention group (mean difference 489.8 g, 95% CI: 427.9, 551.6 g). There are no reports of

trials of antenatal nutrition education in rural Bangladesh.

Thus, the evidence for nutrition education to improve diet and maternal and neonatal health

indicators, including birth weight, remains weak. Most evidence comes from trials with

design or analysis limitations[14] and high-income countries where counselling is by

professional dietitians or nutritionists.[15] There is a critical knowledge gap about antenatal

nutrition education's efficacy and its potential to reduce LBW in South Asia, including

Bangladesh. Filling this gap can potentially contribute to the World Health Assembly's

global goal of a 30% reduction in the LBW rate by 2025.[17.] Therefore, we investigated the

impact of antenatal nutrition education with demonstrations of a nutritionally energy-protein

balanced diet on infants' birth weight in a low-income population.

Subjects and Methods

Study design and participants

We conducted a two-arm, parallel, cluster randomized controlled trial (cRCT). The James P.

Grant School of Public Health; BRAC University Ethical Review Committee; Dhaka;

Bangladesh approved the study protocol (No 53, 25th May 2015).

We have described the study design.[18] In brief, the study was in the rural district of Sherpur,

which has a low human development index, a high percentage in the lowest economic

quintile (37·1%) compared to the national average (21·7%) [19] and 48% below the poverty

line.[20] The Food Security Atlas of Bangladesh classifies Sherpur as having a high/very high

household food insecurity level.[21] In 2016 in Sherpur the neonatal mortality (41/1000 live

births) and under-five mortality (59/1000 live births) were higher than the national average

(30 and 47/1000 respectively).[22]

Sherpur has five sub-districts: Jhenaigati (population 160,452), Nakla (189,685), Nalitabari

(251,361), Sherpur Sadar (497,179) and Sreebardi (259,648). Pregnant women, self-reporting

to a hospital or formal practitioner (doctor, nurse, or Family Welfare Visitor), receive

nutrition education as part of antenatal care (ANC). BRAC (a national NGO; formerly known

as Bangladesh Rural Advancement Committee) implements a community-based Maternal,

Neonatal, and Child Health (MNCH) program characterized by prospective pregnancy

surveillance and basic ANC led by community health workers (CHW) known as *Shasthya*

Kormi. One *Shasthya Kormi* serves an area of ~10,000 people, along with ten health

volunteers (*Shasthya Shebika*). The *Shasthya Kormi* are village women who have basic 10-

years schooling and have received training on health, nutrition, and sanitation issues for six

weeks, followed by bi-monthly refreshers.

In our study, a cluster was the catchment area of one *Shasthya Kormi,* which provided

logistical convenience and minimized intervention spills into the control group. *Shasthya*

Kormi, with at least six months' experience in MNCH, were eligible. Participants were

pregnant women 15-49 years, gestation 12 weeks or less, permanently residing in the area,

and not expecting to move from the usual residence until birth. We used the established

BRAC pregnancy detection system using monthly menstrual surveillance to identify new

pregnancies. Women with delayed menstruation for six weeks or more underwent a urine test.

We excluded pregnant women with self-reported chronic disease. All women excluded from

the study continued to receive standard care and counselling as in the control clusters. A

Shasthya Kormi visited each confirmed pregnant woman, screened for eligibility, and

obtained verbal informed consent. Recruitment took place from 1st October- 31st December

2016.

Randomization and masking

We used proportionate stratified sampling to randomly select 36 of 135 *Shasthya Kormi*

clusters (4, 5, 7, 13, and 7 from the five sub-districts). The clusters were dispersed across the

sub-districts and sufficiently separated to minimize the risk of contamination. We selected 36

as the maximum number of *Shasthya Kormi,* who could be overseen by a single BRAC

supervisor, and we randomized with 1:1 allocation. All eligible pregnant women served by

one *Shasthya Kormi* received the intervention or standard care as per the random assignment.

We could not mask families or the research team to group allocation; however, we blinded

researchers assessing primary and secondary outcomes to the study hypotheses. We also used

an objective outcome (birth weight), which is less prone to ascertainment bias.

Procedures

Intervention *Shasthya Kormi* received two days of nutrition training, including a balanced

diet and counselling techniques. They also practiced community counselling skills under

supervision and received one-day training on taking consent and data collection. Enrolled

participants were visited monthly to provide ANC that included standard nutrition advice. In

intervention clusters, balanced plate intervention replaced the standard nutrition advice using

a culturally appropriate diet for Bangladeshi pregnant women.[23] This contained seven-food

groups (cereal, lentil, animal protein, vegetables, fruits, milk, and oil), giving a daily 2500

kcal energy and included essential micro and macronutrients. The balanced plate was a

combination of foods in appropriate portion sizes to meet requirements. Before the trial, we

conducted focus groups to identify locally preferred foods and dietary restrictions. We pilot

tested the menu to assess community acceptability and the feasibility of incorporating it into

ANC services.

We delivered the intervention individually. Initially, participants received information on a

balanced diet and benefits for maternal and fetal health. Then, they learned to prepare a

"balanced plate" by demonstrating the right proportions of food groups using the prescribed

menu.[18] The community health worker displayed different combinations of food on the plate,

with appropriate alternatives, such as replacing meat (an expensive protein source) with

cheaper eggs or farm fishes. We emphasized foods commonly missed in the regular diet, such

as milk, fruits, and coloured vegetables. Women were encouraged to increase their frequency

of food intake to three meals and two snacks per day. To demonstrate measurements, we used

a bowl of 250 ml commonly found in most homes. The community health worker invited

family decision-makers to observe the demonstration and assist in making a balanced plate.

Delivering the intervention took 45 minutes: 10 for individual counselling, 10 for household

counselling, and 25 for demonstration. To help recall messages, we provided a pictorial food

chart and a written menu[18]. At the first visit, the community health worker delivered eight

nutrition messages with a practical demonstration. In subsequent visits, they emphasized

messages that were less familiar or difficult to remember. In the counselling, they excluded

messages that were easy to recall, such as drinking eight glasses of water or cooking with oil.

They also dropped the messages if it was clear the women practiced the behaviour. As the

number of messages decreased, the community health worker reduced the visit time to 25

minutes. The mean number of intervention visits was 3.4.

Home visits continued until birth or the end of the pregnancy; ANC from other sources was

not discouraged. Husbands were encouraged to re-allocate family budgets to purchase

nutritious food and mothers-in-law to re-distribute family food to increase their daughter-in-

law's share.

In the control group, pregnant women received the same frequency of home and standard

nutrition advice except for the balanced plate demonstrations. Standard advice included

recommendations to consume all food groups and take iron-folic acid and calcium

supplements.

Shasthya Kormi collected demographic, socio-economic, and reproductive data at enrolment

plus birth weight measurements within 72 hours of birth. Families notified birth or pregnancy

loss (miscarriage, abortion, and stillbirth) immediately to their nearby *Shasthya Shebika*.

One field supervisor oversaw the 36 *Shasthya Kormi,* recorded field activities, and conducted

follow-up interviews with 5% of participants to confirm visits and measure protocol

adherence.

Outcomes

Primary outcomes were the mean birth weight of live-born infants and the proportion of

newborns with birth weight less than 2500 g. The *Shasthya Kormi* had many years of

experience measuring birth weight, but we retrained them with their standard equipment, a 5

kg Salter spring balance scale with an accuracy of 100 g. They calibrated the scales with a

standard weight every week. We also checked the scales monthly during refresher training.

For mothers who birthed in health facilities, we recorded the birth weight measured at the

facility.

For secondary outcomes, the *Shasthya Kormi* used structured forms to record fetal loss

during pregnancy, birth outcome, delivery information, and neonatal or maternal death. They

also used semi-structured questionnaires to collect dietary data, with the food list adapted

from Food and Nutrition Technical Assistance II Project (FANTA-2).[24] These questionnaires

captured information about foods, beverages, and supplements consumed 24 hours before the

interview. From this data we formed ten food groups as follows: (1) grains, white roots and

tubers; (2) pulses (beans, peas and lentils); (3) nuts and seeds; (4) dairy; (5) meat, poultry and

fish; (6) eggs; (7) dark green leafy vegetables; (8) other vitamin A-rich fruits and vegetables;

(9) other vegetables; and (10) other fruits. *Shasthya Kormi* collected dietary data monthly,

and we calculated the mean number of food groups consumed to assess diet quality.[25] We

defined animal source protein as consumption of any combination of dairy, meat, poultry,

fish or egg food groups.

Statistical analysis

We required 720 pregnant women to achieve 80% power to detect a difference of 100 g in

birth weight between groups. We assumed a mean birth weight of 2531 g (standard deviation

(SD) 415 g) in the control group based on the findings from a similar trial in Bangladesh.[26]

We set a significance level of 0·05 (two-sided), and assumed an intra-cluster correlation

coefficient (ICC) of 0·03[27], and 36 clusters with an average of 10-15 live births per cluster

per month. To adjust for 10% pregnancy loss and 15% loss to follow-up based on similar

trials,[26,28], we inflated the required sample size to 900 pregnant women.

We analysed the data with Stata version 15 and conducted intention to treat analyses. We

assigned the intervention randomly at the cluster level, but assessed outcomes at the

individual level. We estimated the mean between-group difference in birth weight using

linear regression and the relative risk for LBW using binomial regression. For both, we

included the study group as a covariate and adjusted for cluster randomization using

Generalised Estimating Equations (GEE).[29] Similarly, we estimated the mean number of food

groups consumed using GEE linear regression with the treatment group as a covariate and

adjusting for baseline imbalances by including covariates.

We conducted post hoc subgroup analyses for age, education, family income, and place of

delivery. We added an interaction term to assess whether there was a difference in

intervention effect for any sub-group, using a significance level of 0·01.

Role of the funding source

The funders had no role in study design, participant selection, data collection, and data

analyses. The corresponding author had full access to the study's data and final responsibility

for the decision to publish.

Results

We identified 2,154 newly pregnant women from August 2015 to February 2016 in the

selected clusters. Of these women, 937 did not meet the eligibility criteria (603 had a

gestation of >12 weeks, 147 were not permanent residents, and 187 had planned to move out

of the study area for childbirth) and 324 declined to participate. We recruited 893 participants,

445 in 18 intervention, and 448 in 18 control clusters, and we lost 44 participants during

follow-up, 25 (5·6%) intervention, and 19 (4·2%) control. Birth weight was missing for 22

participants (Figure 1). There were 42 pregnancy losses (abortion or stillbirth), and six

neonatal deaths within 24 hours of birth.

There were no important differences in baseline characteristics between the groups (Table 1).

There were small differences for mother's age, mother's and father's education, and parity

but not enough to warrant an adjustment in analyses. Slightly more intervention participants

received four or more ANC visits, but for antenatal care from health centres, this trend

reversed (Table 1). When we combined both ANC data sources, the intervention participants

received slightly more ANC (intervention 61.7% vs. control 55.6%). There was a slightly

higher percentage of intervention participants delivered in a facility (20.7% vs. 16.6%) but

this variable reflects the small differences in ANC. Mean birth weight was higher in facility

deliveries (2,915 g, SD 560 g) than in-home births (2,769 g, SD: 410 g).

Supervisors attended 5% of balanced plate demonstrations to observe the appropriateness of

message delivery, the use of the menu and food chart, demonstration, problem-solving, and

counselling. There were approximately 6-7 of 8 messages delivered in 80% of the

participants. The time taken for the demonstrations gradually decreased with increasing

numbers of visits. Counselling family members were largely dependent on availability;

husbands were present at 15-20% of visits, other family members at >80% of visits.

We found a statistically significant difference in mean birth weight and LBW (Tables 2 and

3). The mean difference in birth weight between groups was 127·5 g (95% CI: 11.1, 243·9;

p=0·032) adjusted for clustering. The estimated relative risk (RR) of LBW was 57% lower in

the intervention group (RR: 0·43; 95% CI: 0·25, 0·75; p=0·003) adjusted for clustering.

We conducted subgroup analyses of mother's age, education, and family income (Tables 2

and 3) and found a significant intervention effect on birth weight in adolescents vs. non-

adolescents. Mean difference in birth weight between groups was greater in adolescents

(299·1 g; 95% CI: 101·6, 496·6; p=0·003; vs 95·6 g; 95% CI: -17·3, 208·4; p=0·097) (p-value

for interaction 0·009) as was reduction in LBW (RR: 0·28; 95% CI: 0·11, 0·71; p=0·007 vs

RR: 0·54; 95% CI: 0·29, 0·98; p=0·044), although this was not statistically significant (p-

value for interaction 0·22). There was no evidence of treatment modification by the mother's

education or family income (Tables 2 and 3).

To assess bias in ascertaining birth weight, we stratified by participants birthed in a facility

and whose babies were weighed by independent observers and those who birthed at home

and found no evidence of treatment modification (p=0.78).

After adjustment for ANC, the use of iron and folic acid, and clustering, the reduction in

mean birth weight and LBW was smaller in the intervention group (Supplemental Tables S1

and S2). The adjusted difference was 121·8 g (95% CI: 11.7, 231·9; p=0·030), which is 5.7 g

less than the unadjusted analysis (Table 2). The adjusted relative risk of LBW was 0·53 (95%

CI: 0·29, 0·98; p=0·042), which was smaller than the unadjusted relative risk in Table 3.

The mean number of food groups consumed increased in the intervention group and was

significantly higher from the third month of pregnancy. The largest difference was in the

eighth month of pregnancy when intervention participants consumed 5.6 food groups/day

(95% CI: 5.19, 6.14) compared to 3.8 (95% CI: 3.40, 4.12) in control participants (Fig 2).

Starting from the fifth month of pregnancy, the intervention group's mean number of food

groups exceeded five (the number for adequate dietary diversity[25]). The mean number of

animal-source protein foods consumed was significantly higher from the fourth month of

pregnancy (Figure 2).

Discussion

We demonstrated that antenatal nutrition education with practical demonstrations of

preparing a balanced diet increases infant birth weight and reduces LBW in rural Bangladesh.

There was a small birth weight reduction after adjusting for differences in antenatal care and

iron/folic acid use. The effect of the intervention was greater in adolescent mothers. The

findings provide strong evidence of the effectiveness of a sustainable intervention suitable for

implementation in community-based health care systems. They demonstrate the potential

impact of nutrition education to improve intrahousehold food distribution as a tool to reduce

low birth weight in Bangladesh and across South Asia.

The process evaluation of the trial [30] provided insights into why the intervention was so

impactful. It identified key elements in the intervention strategy, such as practical

demonstration of portion sizes; addressing local food perceptions; demystifying animal-

source foods; engaging husbands and mothers-in-law; leveraging women's social networks;

and harnessing community health workers' social role, that played a crucial role in achieving

the desired adherence. In particular, the approach of involving the mothers-in-law and

husbands changed their attitudes towards a fairer allocation of foods to pregnant women. The

husbands adhered to the messages and purchased more and nutritious foods for their wives by

adjusting the shopping budget and even working extra hours.

Comparison with other studies

Our intervention appears more effective than interventions using micronutrient or food

supplements. A large cluster RCT (22,405 pregnancies) in Bangladesh assessed the impact of

antenatal micronutrient supplementation and found a 54 g (95% CI: 41, 66) increase in birth

weight and a 12% reduction (RR: 0.88; 95% CI: 0·85, 0·91) in LBW.[26] Another cluster RCT

of lipid-based micronutrient supplementation showed a 41 g birth weight increment without a

significant effect on LBW.[28]

Providing micronutrient supplements even with additional food energy is unlikely to address

the fundamental problems of insufficient food intake in pregnancy related to food taboos and

inequitable intra-household food distribution. The earlier MINIMat RCT (3267 singleton

newborns)[31], which tested early and late starts to food supplements with micronutrients in

pregnancy, found no significant main-effects on birth weight. However, the combination of

multiple micronutrients and early food supplementation did improve infant survival. The lack

of a control group without food supplements might explain the lack of birth weight response.

In contrast, an observational study in Bangladesh reported a 118 g increase in birth weight

with daily 608 kcal food supplements combined with iron-folic acid sustained for more than

four months.[32] Although this evidence is weak, it supports using food-based solutions to

address nutrition problems.

Adolescent pregnancy has an increased risk of adverse birth outcomes, including LBW.[33] If

severely undernourished, the growing pregnant adolescent and foetus compete for nutrients.[34]

A recent study from rural Bangladesh found a cessation of linear growth, weight loss, and fat

and lean body mass depletion in pregnant adolescents.[34] This greater nutritional susceptibility

helps explain the greater effect of the balanced plate education in this group.

The balanced-plate intervention significantly increased maternal dietary diversity (Fig 2).

Change in dietary patterns is the likely cause of the higher birth weight in the intervention

group. A recent cohort study[35] followed 374 women from their first antenatal care visit to

assess associations between dietary diversity during pregnancy and LBW in rural Ethiopia.

The LBW risk doubled with a dietary diversity score of less than four food groups (ARR:

2.06; 95% CI: 1.03, 4.11). A facility-based case-control study[36] from rural Ethiopia, which

examined iron, folic acid and nutrition counselling, snack consumption, dietary diversity, and

maternal undernutrition, also reported similar findings.

There was a significant six-fold increase in the adjusted odds of LBW with an inadequate

minimum dietary diversity score (AOR: 6.65; 95% CI, 2.31, 19.16). A cross-sectional study[37]

from Ghana also reported a protective effect of higher dietary diversity scores for LBW

(AOR: 0.10; 95% CI: 0.04, 0.13 per standard deviation change score, p=<0.0001). This

evidence supports our findings that improved and more diverse dietary intake leads to a

reduction in LBW.

Participants in the intervention group consumed more animal-sourced protein foods (Fig 2),

reaching a peak at eight months of pregnancy of two animal-sourced foods/day. A large

cross-sectional study of pregnant women in Shaanxi province, northwestern China, reported

an average weekly consumption of five animal-sourced foods.[38] As the frequency of animal-

sourced food increased by one time/week, the average birth weight increased by 3.24 g (95%

CI: 1.09, 5.39). Unlike increasing animal-sourced foods in the diet, the consumption of

excess protein supplements (>20% of energy as protein) can harm fetal growth.[39]

Despite earlier evidence of a positive impact of nutrition education on pregnancy outcomes,

the delivery method has received little attention. It is particularly important in low literate and

poor self-efficacy populations, such as in rural Bangladesh, where women require dietary

counselling to be accessible, feasible, and sanctioned by the community. A review of over

300 studies found that nutrition education is more likely to be effective when focusing on

behaviour and action than knowledge alone. And it is also likely to be more effective if

grounded in theory.[40]   Fishbein and Yzer note that "any given behaviour is most likely to

occur; if one has a strong intention to perform the behaviour; if a person has the necessary

skills and abilities required to perform the behaviour; and if no environmental constraints are

preventing the behavioural performance." [41] To positively impact dietary behaviour, we

designed our intervention to embrace all aspects of this theory. We adopted an innovative

counselling technique, including a practical demonstration of a balanced diet and engaged

family decision-makers. Similar nutrition counselling intervention was found effective in

increasing age-specific complementary feeding practices in rural Bangladesh.[42]

In Bangladesh, the main source of nutrition education for pregnant women is through ANC,

but the contact time is insufficient to affect nutrition change.[43]   Thus, WHO. Proposes

upskilling CHWs in delivering dietary advice is an appropriate approach to complement

health services.[44]   In our study, we integrated the balanced plate intervention into BRAC's

existing ANC package delivered by CHWs. The same staff in a recent intervention

demonstrated their successful counselling on breastfeeding and complementary feeding.[45]

Previously, we also found that CHWs can positively influence family decision-makers to

increase pregnant women's iron-folic acid consumption.[46]

Strengths and limitations of this study

The evidence presented from this trial is robust. Key strengths are the cluster, randomized

controlled trial design with sufficient statistical power to detect clinically important changes

in outcomes. The delivery of the trial treatments through an established home-based antenatal

care service led to a balance between the treatment groups in the level of antenatal contacts

with community health workers. The unblinded intervention was counterbalanced by

objective outcomes and blinding participants and data collectors to the aims and hypotheses.

The trial had a very high follow-up (93%). We used intention-to-treat analyses and adjusted

for unbalanced baseline factors and clustering. Our intervention, based on well-designed

qualitative formative research, was logistically simple and culturally appropriate. The key to

our success was conducting the trial within existing ANC frameworks. The delivery strategy

resonates with maternal and child nutrition recommendations to achieve adequate coverage of

nutrition-specific interventions by reaching populations in need.[47] However, we acknowledge

some limitations. First, the *Shasthya Kormi*, who delivered the intervention, also collected

birth weight data. Measuring birth weight and providing nutrition counselling are routine

tasks for these staff. By blinding the SK to study outcome, we would expect non-differential

bias. Also, we found no difference in the effects between women who had birth weight

measured by hospital staff versus those who birthed at home with birth weight measured by

the SK. Secondly, using catchment areas as clusters might introduce contamination, with the

possibility of *Shasthya Kormi* exchanging information. However, this would have biased

results towards a null effect. To prevent this, we trained intervention and control group

Shasthya Kormi separately at regular refresher sessions. Thirdly, we could not quantify food

intake and intra-household food distribution, which might have provided further evidence of

pathways to the impact we observed. Finally, we excluded over 30% of pregnant women

identified mainly because they were already in the second or third trimester when we

detected their pregnancy. There is no information about the characteristics of these excluded

women and thus no way to compare them to the women included in the trial. This limits our

assessments of the external validity of the trial findings.

Conclusions and policy implications

Our results indicate that a balanced plate nutrition education during pregnancy increases birth

weight and reduces LBW. Our trial offers a pragmatic food-based strategy for resource-poor

settings. The CHW-led intervention is simple, feasible, and easy to implement in a country

with a shortage of human health resources. The intervention is also scalable through existing

public and private community health infrastructure.

References

1.   Black RE, Allen LH, Bhutta ZA, et al. Maternal and child undernutrition: global and
398   regional exposures and health consequences. *Lancet.* 2008;371:243-260.
399   2.   Cesar G Victora, Linda Adair, Caroline Fall, et al. Maternal and child undernutrition:
400   consequences for adult health and human capital. *Lancet.* 2008;371:340-357.
401   3.   Wilcox AJ, Russell IT. Birthweight and Perinatal Mortality: II. On Weight-Specific
402   Mortality. *International Journal of Epidemiology.* 1983;12(3):319-325.
403   4.   Watkins WJ, Kotecha SJ, Kotecha S. All-cause mortality of low birthweight infants in
404   infancy, childhood, and adolescence: population study of England and Wales. *PLoS*
405   *medicine.* 2016;13(5):e1002018.
406   5.   Lee AC, Katz J, Blencowe H, et al. National and regional estimates of term and
407   preterm babies born small for gestational age in 138 low-income and middle-income
408   countries in 2010. *Lancet Glob Health* 2013;1:e26-36.
409   6.   Lee AC, Kozuki N, Cousens S, et al. Estimates of burden and consequences of infants
410   born small for gestational age in low and middle income countries with
411   INTERGROWTH-21st standard: analysis of CHERG datasets. *BMJ.* 2017;358(j3677).
412   7.   Han Z, Mulla S, Beyene J, Liao G, McDonald SD. Maternal underweight and the risk
413   of preterm birth and low birth weight: a systematic review and meta-analyses.
414   *International journal of epidemiology.* 2010;40(1):65-101.
415   8.   Triunfo S, Lanzone A. Impact of maternal under nutrition on obstetric outcomes.
416   *Journal of endocrinological investigation.* 2015;38(1):31-38.
417   9.   World Health Organization. *Maternal anthropometry and pregnancy outcomes: a*
418   *WHO collaborative study.* World Health Organization; 1995.
419   10.   Naidu A, Rao N. Body mass index: a measure of the nutritional status in Indian
420   populations. *European journal of clinical nutrition.* 1994;48:S131-140.
421   11.   Islam MZ, Akhtaruzzaman M, Lamberg-Allardt C. Nutritional status of women in
422   Bangladesh: comparison of energy intake and nutritional status of a low income rural
423   group with a high income urban group. *Asia Pacific Journal of Clinical Nutrition.*
424   2004;13(1):61-68.
425   12.   Harris-Fry H, Shrestha N, Costello A, Saville NM. Determinants of intra-household
426   food allocation between adults in South Asia – a systematic review. *International*
427   *Journal for Equity in Health.* 2017;16(107).
428   13.   Boyd NR, Jr., Windsor RA. A meta-evaluation of nutrition education intervention
429   research among pregnant women. *Health education quarterly.* 1993;20(3):327-345.
430   14.   Girard AW, Olude O. Nutrition education and counselling provided during
431   pregnancy: effects on maternal, neonatal and child health outcomes. *Paediatric and*
432   *Perinatal Epidemiology.* 2012;26(s1):191-204.
433   15.   Gresham E, Collins CE, Mishra GD, Byles JE, Hure AJ. Diet quality before or during
434   pregnancy and the relationship with pregnancy and birth outcomes: the Australian
435   Longitudinal Study on Women's Health. *Public Health Nutrition.* 2016;19(16):2975–
436   2983.
437   16.   Ota E, Hori H, Mori R, Tobe-Gai R, Farrar D. Antenatal dietary education and
438   supplementation to increase energy and protein intake. *Cochrane Database of*
439   *Systematic Reviews.* 2015(6).
440   17.   Lopes KdS, Ota E, Shakya P, et al. Effects of nutrition interventions during pregnancy
441   on low birth weight: an overview of systematic reviews. *BMJ Global Health.*
442   2017;2(e000389).

443   18.   Chowdhury M, Raynes-Greenow C, Alam A, Dibley M. Making a balanced plate for
444   pregnant women to improve birthweight of infants: a study protocol for a cluster
445   randomised controlled trial in rural Bangladesh. *BMJ Open.* 2017;7(e015393).

446   19.   Sen B, Ali Z. Spatial Inequality in Social Progress in Bangladesh. *The Bangladesh*
447   *Development Studies.* 2009;XXXII(2).

448   20.   *Poverty Maps of Bangladesh 2010.* World Bank Bangladesh, Bangladesh Bureau of
449   Statistics, and World Food Programme2011.

450   21.   *The Food Security Atlas of Bangladesh: Towards a poverty and hunger free*
451   *Bangladesh.* Planning Commission, Bangladesh Bureau of Statistics and UN World
452   Food Programme. 2004.

453   22.   National Institute of Population Research and Training (NIPORT), International
454   Centre for Diarrhoeal Disease Research Bangladesh (icddrb), MEASURE Evaluation.
455   *Bangladesh District Level Socio-demographic and Health Care Utilization Indicators.*
456   Dhaka, Bangladesh, and Chapel Hill, NC, USA: NIPORT, icddr,b, and MEASURE
457   Evaluation;2019.

458   23.   Nahar Q, Choudhury S, Faruque MO, Sultana SSS, Siddiquee MA. *Desirable Dietary*
459   *Pattern for Bangladesh.* National Food Policy Capacity Strengthening
460   Programme;2013.

461   24.   Arimond M, Torheim LE, Wiesmann D, Joseph M, Carriquiry A. Dietary diversity as
462   a measure of the micronutrient adequacy of women's diets: results from rural
463   Bangladesh site. *Washington, DC: Food and Nutrition Technical Assistance II*
464   *Project (FANTA-2), FHI.* 2009;360.

465   25.   FAO and FHI. Minimum dietary diversity for women: a guide for measurement.
466   *Rome: FAO.* 2016.

467   26.   West Jr PK, Shamim AA, Mehra S, et al. Effect of Maternal Multiple Micronutrient
468   vs Iron–Folic Acid Supplementation on Infant Mortality and Adverse Birth Outcomes
469   in Rural Bangladesh: The JiVitA-3 Randomized Tria. *JAMA.* 2014;312(24):2649-
470   2658.

471   27.   Zeng L, Cheng Y, Dang S, et al. Impact of micronutrient supplementation during
472   pregnancy on birth weight, duration of gestation, and perinatal mortality in rural
473   western China: double blind cluster randomised controlled trial. *BMJ.*
474   2008;337(7680):1211 - 1215.

475   28.   Mridha MK, Matias SL, Chaparro CM, et al. Lipid-based nutrient supplements for
476   pregnant women reduce newborn stunting in a cluster-randomized controlled
477   effectiveness trial in Bangladesh. *American Society for Nutrition.* 2016;103:236-249.

478   29.   Rabe-Hesketh S, Skrondal A. *Multilevel and longitudinal modeling using Stata.*
479   STATA press; 2008.

480   30.   Alam A, Chowdhury M, Dibley MJ, Raynes-Greenow C. How can we improve the
481   consumption of a nutritionally balanced maternal diet in rural bangladesh? The key
482   elements of the "balanced plate" intervention. *International Journal of Environmental*
483   *Research and Public Health.* 2020;17(17):6289.

484   31.   Persson LÅ, Arifeen S, Ekström E-C, Rasmussen KM, Frongillo EA, Team MS.
485   Effects of prenatal micronutrient and early food supplementation on maternal
486   hemoglobin, birth weight, and infant mortality among children in Bangladesh: the
487   MINIMat randomized trial. *Jama.* 2012;307(19):2050-2059.

488   32.   Shaheen R, Francisco Ad, Arifeen SE, Ekström E-C, Persson LÅ. Effect of prenatal
489   food supplementation on birth weight: an observational study from Bangladesh1–. *Am*
490   *J Clin Nutr.* 2006;83:1355–1361.

33. King JC. The risk of maternal nutritional depletion and poor outcomes increases in
early or closely spaced pregnancies. *The Journal of nutrition.* 2003;133(5):1732S-
1736S.
34. Rah JH, Christian P, Shamim AA, Arju UT, Labrique AB, Rashid M. Pregnancy and
lactation hinder growth and nutritional status of adolescent girls in rural Bangladesh.
The Journal of nutrition. 2008;138(8):1505-1511.
35. Zerfu TA, Umeta M, Baye K. Dietary diversity during pregnancy is associated with
reduced risk of maternal anemia, preterm delivery, and low birth weight in a
prospective cohort study in rural Ethiopia. *The American journal of clinical nutrition.*
2016;103(6):1482-1488.
36. Girma S, Fikadu T, Agdew E, et al. Factors associated with low birthweight among
newborns delivered at public health facilities of Nekemte town, West Ethiopia: a case
control study. *BMC pregnancy and childbirth.* 2019;19(1):220.
37. Abubakari A, Jahn A. Maternal dietary patterns and practices and birth weight in
Northern Ghana. *PloS one.* 2016;11(9):e0162285.
38. Yan H, Dang S, Mi B, et al. Study on relationship between mother's animal sourced
food intake during pregnancy and neonate birth weight. *Zhonghua liu xing bing xue
za zhi= Zhonghua liuxingbingxue zazhi.* 2017;38(5):615-620.
39. Liberato SC, Singh G, Mulholland K. Effects of protein energy supplementation
during pregnancy on fetal growth: a review of the literature focusing on contextual
factors. *Food & nutrition research.* 2013;57(1):20499.
40. Contento IR. Nutrition education: linking research, theory, and practice. *Asia Pac J
Clin Nutr.* 2008;17(1):176-179.
41. Fishbein M, Yzer MC. Using Theory to Design Effective Health Behavior
Interventions. *Communication Theory.* 2003;13(2):164-183.
42. Sanghvi T, Haque R, Roy S, et al. Achieving behaviour change at scale: Alive &
Thrive's infant and young child feeding programme in Bangladesh. *Maternal & Child
Nutrition.* 2016;12(Suppl. 1):141-154.
43. Ahmed SM, Hossain MA, Chowdhury AMR, Bhuiya AU. The health workforce crisis
in Bangladesh: shortage, inappropriate skill-mix and inequitable distribution. *Human
Resources for Health.* 2011;9(3).
44. Lehmann U, Sanders D. *Community health workers: what do we know about them?
The state of the evidence on programmes, activities, costs and impact on health
outcomes of using community health workers.* Geneva: World Health
Organization;2007.
45. Alive & Thrive. *Alive & Thrive's approach and results in Bangladesh: Rapid
improvements in infant and young child feeding practices resulting from a large-scale
program.* Dhaka, Bangladesh: Alive & Thrive;2015.
46. Alam A, Rasheed S, Khan NU, et al. How can formative research inform the design
of an iron-folic acid supplementation intervention starting in first trimester of
pregnancy in Bangladesh? *BMC Public Health.* 2015;15(374).
47. Bhutta ZA, Das JK, Rizvi A, et al. Evidence-based interventions for improvement of
maternal and child nutrition: what can be done and at what cost? *The Lancet.*
2013;382:452-477.

Acknowledgments

This study is part of research towards a Ph.D. at The University of Sydney funded by
Endeavour Postgraduate Scholarship (Ph.D.); Department of Education and Training,
Australian Government. The study received partial support from the James P. Grant School
of Public Health, BRAC University, Bangladesh, and the Sydney Medical School, The
University of Sydney, Australia. We thank the *Shasthya Kormi* and the *Shasthya Shebika* of
BRAC, and the pregnant women and their families in Sherpur for their time in this study.

Funding:

The study was funded by James P. Grant School of Public Health; Bangladesh, and Sydney
Medical School Foundation; Australia

Author contributions:

MC conceived the overall study, developed the study design and data collection methods, and
conducted the data analysis, interpreted data, and wrote the first draft of the manuscript. MJD
and CRG provided critical input regarding study design, sample size calculation, and
outcome evaluation and the statistical analysis plan. KA provided practical guidance in
formulating the implementation strategy. AA provided crucial input on formative research
and process evaluation design. PK provided technical support for data analyses. CRG and
MC obtained funds for the intervention. CRG provided critical input to all drafts and
approved the final version for submission. MJD, KA, PK and AA critically reviewed the
manuscript and approved the final version for submission. MC, MJD, CRG, PK, KA and AA
agreed to be accountable for all investigations necessary to resolve questions related to
accuracy or integrity of all or any part of the work.

Competing interests:

None to declare

Data and materials availability:

The data described in this manuscript, the code book, and the analytic code will be made

available upon request pending approval by the investigators of an application to use the data.

Supplemental Information

S1 Table: The mean birth weight and the adjusted mean difference in birth weight of

balanced -plate intervention compared to usual programs.

S2 Table: Percentage of low birth weight and adjusted relative risk of balanced-plate

intervention compared to usual programs.

 Table 1: Cluster, baseline maternal and household, and antenatal care and delivery

 characteristics by study arm

	Intervention (n=382)	Control (n=397)
Cluster characteristics		
Number of clusters	18	18
Mean (SD) population in clusters in 2016	9775 (2007)	9639 (1763)
Mean (SD) number of households in clusters	2315 (288)	2321 (419)
Mean (SD) number of women of reproductive age	2639 (542)	2631 (481)
Mean (SD) number of pregnant women enrolled per clusters	24.7 (1.0)	24.9 (0.5)
Baseline maternal and household characteristics		
Mother's age (years):		
Mean (SD)	23·0 (4·6)	24·0 (4.3)
15-19	80 (21·7)	64 (17.0)
20-34	280 (76·1)	307 (81·4)
>34	8 (2·2)	6 (1·6)
Missing	14	20
Mother's education;		
No education	82 (22·0)	89 (23·1)
Primary	250 (67·2)	242 (62·7)
Secondary	27 (7·3)	35 (9·1)
Higher	13 (3·5)	20 (5·2)
Missing	10	11
Monthly family income (SD) in Bangladesh Taka	7308 (4303)	7213 (4438)
Less than 5000	87 (23.9)	79 (21.2)
5000 to less than 10,000	190 (52.2)	209 (56.2)
10,000 or more	87 (23.9)	84 (22.6)
Missing	18	25
Father's education:		
No education	130 (35·0)	151 (39.7)
Primary	188 (50·7)	169 (44·5)
Secondary	34 (9·2)	39 (10.3)
Higher	19 (5·1)	21 (5·5)
Missing	11	17
Marital Status		
Married	372 (97.4)	386 (97.3)
Not married	10 (2.6)	11 (2.7)
Parity:		

	Intervention (n=382)	Control (n=397)
Mean (SD)	1·8 (1·2)	1·9 (1·3)
0	36 (10·3)	48 (12·6)
1	134 (38·5)	115 (30·3)
≥2	178 (51·2)	217 (57·1)
Missing	34	17
Previous pregnancy loss		
Yes	67 (19·3)	87 (22·9)
No	281 (80·8)	293 (77·1)
Missing	34	17
Gestation at enrolment (weeks) Mean (SD)	10·1 (3·2)	9·6 (3·2)
Missing	27	2
Antenatal care and delivery characteristics		
Iron Folic Acid Supplements		
Mean number consumed (SD)	45.6 (61.8)	29.0 (46.8)
Any use	117 (30.6)	70 (17.6)
None	265 (69.4)	327 (83.4)
Antenatal care visit by SK		
Mean number of visits (SD)	3.8 (2.6)	3.6 (2.2)
<4 visits	158 (41.1)	198 (49.9)
≥4 visits	224 (58.6)	199 (50.1)
Antenatal care in health centers		
Mean number of visits (SD)	0.7 (1.1)	1.0 (1.4)
No Visit	227 (59.4)	246 (51.9)
<4 visits	143 (37.4)	169 (42.6)
≥4 visits	12 (3.1)	22 (5.5)
Place of delivery		
Facility	79 (20.7)	66 (16.6)
Home	280 (73.3)	323 (81.4)
Unknown	23 (6.0)	8 (2.0)
Mode of delivery		
Normal	331 (86.7)	361 (90.9)
C-Section	33 (8.6)	34 (8.6)
Unknown	18 (4.7)	2 (0.5)
Time of birth weight measurement		
Within 24h	266 (69.6)	290 (73.1)
24 to 48h	75 (19.6)	61 (15.4)
48 to 72h	13 (3.4)	19 (4.8)
Unknown	28 (7.3)	27 (6.8)

Numbers are frequencies (and column percentage) unless stated otherwise

Table 2: The mean birth weight and the mean difference in birth weight of balanced-plate intervention compared to usual programs.

	Number of participants		Birth weight (g)				Mean difference			
	Intervention	Control	Intervention		Control		Estimate	(95%CI)	P-value	P-value for interaction
			Mean (SD)		Mean (SD)					
Overall	382	397	2861.0	(444.2)	2736.8	(423.2)	127.5	(11.1, 243.9)	0.032	
Mother's age[1] (years)										0.009
<20	80	64	2916.3	(471.1)	2637.5	(481.0)	299.1	(101.6, 496.6)	0.003	
>=20	288	313	2847.6	(443.4)	2762.6	(401.1)	95.6	(-17.3, 208.4)	0.097	
Mother's education[2]										0.330
None	82	89	2869.5	(460.6)	2693.3	(339.7)	175.1	(29.9, 320.4)	0.018	
Primary	250	242	2839.2	(443.9)	2761.6	(394.4)	92.9	(-28.0, 213.7)	0.132	
Secondary or higher	40	55	2982.5	(435.5)	2741.8	(588.0)	226.1	(-33.5, 485.8)	0.088	
Family income[3]										0.236
Low	87	79	2819.5	(397.9)	2663.3	(420.4)	166.8	(10.7, 322.8)	0.036	
Medium	190	209	2806.8	(421.7)	2750.7	(390.8)	51.8	(-70.0, 173.6)	0.404	
High	87	84	2995.4	(514.5)	2791.7	(433.2)	183.3	(-40.9, 407.4)	0.109	

573 Birth weight was analyzed using linear regression. We adjusted all models for clusters using generalized estimating equation (GEE) models, assuming an exchangeable correlation structure and applied a sandwich

574 estimator to standard errors. [1] n=745 due to 34 missing maternal age values. [2] n=758 due to 21 missing maternal education values. [3] n=736 due to 43 missing family income values.

577 **Table 3: Percentage of low birth weight and relative risk of balanced-plate intervention compared to usual programs.**

	Number of participants		Low birth weight (<2,500g)				Relative risk			
	Intervention	Control	Intervention n (%)		Control n (%)		Estimate	(95%CI)	P-value	P-value for interaction
Overall	382	397	37	(10)	87	(22)	0.43	(0.25, 0.75)	0.003	
Mother's age[1] (years)										0.217
<20	80	64	6	(8)	17	(27)	0.28	(0.11, 0.71)	0.007	
>=20	288	313	31	(11)	62	(20)	0.54	(0.29, 0.98)	0.044	
Mother's education[2]										0.240
None	82	89	11	(13)	22	(25)	0.54	(0.28, 1.05)	0.069	
Primary	250	242	25	(10)	47	(19)	0.50	(0.26, 0.93)	0.029	
Secondary or higher	40	55	1	(3)	12	(22)	0.11	(0.02, 0.8)	0.029	
Family income[3]										0.441
Low	87	79	9	(10)	25	(32)	0.31	(0.13, 0.76)	0.010	
Medium	190	209	23	(12)	41	(20)	0.60	(0.32, 1.14)	0.117	
High	87	84	5	(6)	12	(14)	0.45	(0.13, 1.53)	0.199	

578 Low birth weight was analyzed using binomial regression with a log link. We adjusted all models for clusters using generalized estimating equation (GEE) models, assuming an

579 exchangeable correlation structure and applied sandwich estimator to standard errors. [1] n=745 due to 34 missing maternal age values. [2] n=758 due to 21 missing maternal

580 education values. [3] n=736 due to 43 missing family income values

581   **Figure 1: Trial flow chart**

582
583

585 **Figure 2: The mean number and 95% confidence intervals* of food groups (A) and**

586 **animal-source protein foods (B) consumed by women per day by treatment group.**

588 * The estimated the mean number of food groups consumed and 95% confidence intervals were calculated using

589 linear regression with the study arm as a covariate using Generalised Estimating Equations (GEE) assuming an

590 exchangeable correlation structure and applying the sandwich estimator to standard errors.