The impact of antenatal balanced plate nutrition education for pregnant women on

birth weight: a cluster randomised controlled trial in rural Bangladesh

Authors:

Morseda Chowdhury[1, 2], Camille Raynes-Greenow[2], Patrick J. Kelly[2], Ashraful Alam[2],

Kaosar Afsana[3], Sk Masum Billah[2,4], Michael J. Dibley[2]

Affiliations:

[1] Health Nutrition and Population Programme, BRAC, BRAC Centre, 75 Mohakhali, Dhaka

1212, Bangladesh

[2] The University of Sydney, Faculty of Medicine and Health, Sydney School of Public Health,

Camperdown, New South Wales 2006, Australia

[3] James P Grant School of Public Health, (JPGSPH), BRAC University, Dhaka, Bangladesh

[4] Maternal and Child Health Division, International Centre for Diarrhoeal Disease Research,

Bangladesh (icddr,b), Dhaka 1212, Bangladesh

Corresponding author: Michael J Dibley, Room 328 Edward Ford Building (A27) Sydney

School of Public Health, the University of Sydney, NSW 2006, Australia.

Phone: +61 2 9351 520   Email: michael.dibley@sydney.edu.au

Short running head: Balanced plate nutrition education & birth weight

Keywords: Low birth weight, maternal nutrition education, food demonstrations, dietary

diversity, cluster RCT, Bangladesh

Trial registration: The Australian New Zealand Clinical Trials Registry number

ACTRN12616000080426

Abstract

Low birth weight (LBW) is a global public health problem with the highest prevalence in South Asia. It is strongly associated with maternal undernutrition. In South Asia, intra-household food distribution is inequitable, with lower dietary adequacy in women. Evidence that nutrition education improves diet during pregnancy and reduces LBW is weak. We assessed the impact of nutrition education for pregnant women on birth weight in rural Bangladesh. We conducted a parallel, two-arm, cluster-randomised controlled trial, with 36 clusters allocated equally to intervention (n=445) or standard care (n=448). From their first trimester until delivery, intervention participants received education about eating balanced meals to meet daily dietary requirements with diverse food groups. The primary outcome of mean birth weight was 127.5 g higher in the intervention compared to control women, and the intervention reduced the risk of LBW by 57%. Post hoc analyses showed a significantly higher birth weight and a greater reduction in LBW amongst adolescent mothers. The mean number of food groups consumed was significantly higher in the intervention from the third month of pregnancy than in the control. A community-based balanced plate nutrition education intervention effectively increased mean birth weight and reduced LBW, and improved dietary diversity in rural Bangladeshi women.

Keywords: Low birth weight, maternal diet, dietary diversity, antenatal, nutrition education

Introduction

Birth weight is a powerful predictor of infant survival.[1] Low birth weight, defined as less than 2500 grams (g), is a composite measure of restricted fetal growth and short duration of gestation, major determinants of infant mortality, childhood growth and development, and adulthood well-being.[1,2] The lower the birth weight, the higher the risk of mortality.[3,4] Globally, in 2010, there were an estimated 18 million LBW babies, of whom 59% were term but small for gestational age, and 41% were preterm.[5] An estimated 606,500 neonatal deaths occur annually in low and middle-income countries due to small for gestational age, which accounts for 22% of neonatal deaths.[6] South Asia has the highest prevalence of LBW globally (34%).[6] Bangladesh is ranked fourth in the global burden of LBW.[5]

LBW is strongly associated with maternal under-nutrition[7], which induces suboptimal placental growth, influencing nutrient delivery, altering fetal hormones, and contributing to restricted development.[8] A World Health Organization study with 111,000 women reported that mothers in the lowest quartile of pre-pregnancy weight carried an elevated risk of IUGR of 2.55 (95% confidence interval (CI): 2.3, 2.7) and LBW of 2.38 (95% CI: 2.1, 2.5) compared to the upper quartile.[9] In India, the odds ratio for LBW was three times greater in severe energy-deficient low BMI groups than in normal BMI. [10] Consistent with the mechanisms and epidemiological evidence is the high prevalence of LBW in rural Bangladeshi women who have a chronic shortfall of food energy throughout pregnancy.[11] In South Asia, intra-household food distribution is inequitable with lower energy share, animal source food, and dietary adequacy in women, regardless of pregnancy.[12]

There is little data on nutrition education's effects on birth weight in low or middle-income countries, although four meta-analyses have assessed impacts. The first, in 1993, found that nutrition education significantly increased the consumption of protein and vitamins and the

birth weight of infants by 300 g. However, the studies lacked statistical power or did not

report effect size estimates.[13] The second meta-analysis in 2012 found a 105 g (95% CI: 18,

193 g) increase in mean birth weight with nutrition education and counselling in pregnancy.

When stratified by the study site, the effect failed to reach significance in low and middle-

income countries (152 g; 95% CI: -81, 384 g).[14] The third meta-analysis in 2014 reported that

dietary counselling significantly increased birth weight in three studies (149 g; p=0·039), but

two had a small sample size, and another had methodological limitations.[15] The fourth, a

Cochrane review,[16] identified two small trials that reported significantly higher birth weight

in the intervention group (mean difference 489.8 g, 95% CI: 427.9, 551.6 g). There are no

reports of trials of antenatal nutrition education in rural Bangladesh.

Thus, the evidence for nutrition education to improve diet and maternal and neonatal health

indicators, including birth weight, remains weak. Most evidence comes from trials with

design or analysis limitations[14] and high-income countries where counselling is by

professional dietitians or nutritionists.[15] There is a critical knowledge gap about antenatal

nutrition education's efficacy and potential to reduce LBW in South Asia, including

Bangladesh. Filling this gap can potentially contribute to the World Health Assembly's global

goal of a 30% reduction in the LBW rate by 2025.[17.] Therefore, we investigated the impact of

antenatal nutrition education with demonstrations of a nutritionally energy-protein balanced

diet on infants' birth weight in a low-income population.

85   Subjects and Methods

86   Study design and participants

We conducted a two-arm, parallel, cluster randomised controlled trial (cRCT). The James P.

Grant School of Public Health; the BRAC University Ethical Review Committee, Dhaka,

Bangladesh, approved the study protocol (No 53, 25th May 2015).

We have described the study design.[18] In brief, the study was in the rural district of Sherpur,

which has a low human development index, a high percentage in the lowest economic

quintile (37·1%) compared to the national average (21·7%) [19] and 48% below the poverty

line.[20] The Food Security Atlas of Bangladesh classifies Sherpur as having a high/very high

household food insecurity level.[21] In 2016 in Sherpur the neonatal mortality (41/1000 live

births) and under-five mortality (59/1000 live births) were higher than the national average

(30 and 47/1000 respectively).[22]

Sherpur has five sub-districts: Jhenaigati (population 160,452), Nakla (189,685), Nalitabari

(251,361), Sherpur Sadar (497,179) and Sreebardi (259,648). Pregnant women, self-reporting

to a hospital or formal practitioner (doctor, nurse, or Family Welfare Visitor), receive

nutrition education as part of antenatal care (ANC). BRAC (a national NGO; formerly known

as Bangladesh Rural Advancement Committee) implements a community-based Maternal,

Neonatal, and Child Health (MNCH) program characterised by prospective pregnancy

surveillance and basic ANC led by community health workers (CHW) known as *Shasthya*

Kormi. One *Shasthya Kormi* serves an area of ~10,000 people, along with ten health

volunteers (*Shasthya Shebika*). The *Shasthya Kormi* are village women who have basic 10-

years schooling and have received training on health, nutrition, and sanitation issues for six

weeks, followed by bi-monthly refreshers.

In our study, a cluster was the catchment area of one *Shasthya Kormi,* which provided

logistical convenience and minimised intervention spills into the control group. *Shasthya*

Kormi, with at least six months of experience in MNCH, were eligible. Participants were

pregnant women 15-49 years, gestation 12 weeks or less, permanently residing in the area,

and not expecting to move from the usual residence until birth. We used the established

BRAC pregnancy detection system for monthly menstrual surveillance to identify new

pregnancies. Women with delayed menstruation for six weeks or more underwent a urine test.

We excluded pregnant women with self-reported chronic disease. All women excluded from

the study continued to receive standard care and counselling as in the control clusters. A

Shasthya Kormi visited each confirmed pregnant woman, screened for eligibility, and

obtained verbal informed consent. Recruitment took place from 1st October- 31st December

2016.

Randomisation and masking

We used proportionate stratified sampling to randomly select 36 of 135 *Shasthya Kormi*

clusters (4, 5, 7, 13, and 7 from the five sub-districts). The clusters were dispersed across the

sub-districts and sufficiently separated to minimise the risk of contamination. We selected 36

as the maximum number of *Shasthya Kormi,* whom a single BRAC supervisor could oversee,

and we randomised with 1:1 allocation. All eligible pregnant women served by one *Shasthya*

Kormi received the intervention or standard care per the random assignment.

We could not mask families or the research team to group allocation; however, we blinded

researchers assessing primary and secondary outcomes to the study hypotheses. We also used

an objective outcome (birth weight) less prone to ascertainment bias.

Procedures

Intervention *Shasthya Kormi* received two days of nutrition training, including a balanced

diet and counselling techniques. They also practised community counselling skills under

supervision and received one-day training on taking consent and data collection. Enrolled

participants were visited monthly to provide ANC that included standard nutrition advice. In

intervention clusters, balanced plate intervention replaced the standard nutrition advice using

a culturally appropriate diet for Bangladeshi pregnant women.[23] This contained seven-food

groups (cereal, lentils, animal protein, vegetables, fruits, milk, and oil), giving a daily 2500

kcal energy and essential micro and macronutrients. The balanced plate was a combination of

foods in appropriate portion sizes to meet requirements. Before the trial, we conducted focus

groups to identify locally preferred foods and dietary restrictions. We pilot-tested the menu to

assess community acceptability and the feasibility of incorporating it into ANC services.

We delivered the intervention individually. Initially, participants received information on a

balanced diet and the maternal and fetal health benefits. Then, they learned to prepare a

"balanced plate" by demonstrating the right proportions of food groups using the prescribed

menu.[18] The community health worker displayed different food combinations on the plate

with appropriate alternatives, such as replacing meat (an expensive protein source) with

cheaper eggs or farm fish. We emphasised foods commonly missed in the regular diet, such

as milk, fruits, and coloured vegetables. Women were encouraged to increase their frequency

of food intake to three meals and two snacks per day. We used a bowl of 250 ml, commonly

found in most homes, to demonstrate measurements. The community health worker invited

family decision-makers to observe the demonstration and assist in making a balanced plate.

Delivering the intervention took 45 minutes: 10 for individual counselling, 10 for household

counselling, and 25 for demonstration. To help recall messages, we provided a pictorial food

chart and a written menu[18]. The community health worker delivered eight nutrition messages

during the first visit with a practical demonstration. In subsequent visits, they emphasised

messages that were less familiar or difficult to remember. They excluded easily recalled

messages in the counselling, such as drinking eight glasses of water or cooking with oil. They

also dropped the messages if it was clear the women practised the behaviour. As the number

of messages decreased, the community health worker reduced the visit time to 25 minutes.

The mean number of intervention visits was 3.4.

Home visits continued until birth or the end of the pregnancy; ANC from other sources was

not discouraged. Husbands were encouraged to reallocate family budgets to purchase

nutritious food and mothers-in-law to re-distribute family food to increase their daughter-in-

law's share.

In the control group, pregnant women received the same frequency of home and standard

nutrition advice except for the balanced plate demonstrations. Standard advice recommended

consuming all food groups and taking iron-folic acid and calcium supplements.

Shasthya Kormi collected demographic, socio-economic, and reproductive data at enrolment

plus birth weight measurements within 72 hours of birth. Families immediately notified birth

or pregnancy loss (miscarriage, abortion, and stillbirth) to their nearby *Shasthya Shebika*.

One field supervisor oversaw the 36 *Shasthya Kormi,* recorded field activities, and conducted

follow-up interviews with 5% of participants to confirm visits and measure protocol

adherence.

Outcomes

Primary outcomes were the mean birth weight of live-born infants and the proportion of

newborns with a birth weight less than 2500 g. The *Shasthya Kormi* had many years of

experience measuring birth weight, but we retrained them with their standard equipment, a 5

kg Salter spring balance scale with an accuracy of 100 g. They calibrated the scales with a

standard weight every week. We also checked the scales monthly during refresher training.

For mothers birthed in health facilities, we recorded the birth weight measured at the facility.

For secondary outcomes, the *Shasthya Kormi* used structured forms to record fetal loss

during pregnancy, birth outcome, delivery information, and neonatal or maternal death. They

also used monthly semi-structured questionnaires to collect dietary data, with the food list

adapted from Food and Nutrition Technical Assistance II Project (FANTA-2).[24] These

questionnaires captured information about foods, beverages, and supplements consumed 24

hours before the interview. From this data, we formed seven food groups as follows: (1)

grains, white roots and tubers; (2) pulses (beans, peas and lentils), nuts and seeds; (3) dairy;

(4) meat, poultry and fish; (5) eggs; (6) Vitamin A-rich fruits and vegetables; (7) other fruit

and vegetables. We calculated the mean number of food groups consumed to assess diet

quality[25] and compared the mean number of food groups between the treatment groups and

the relative differences for each food group. We defined animal source protein as

consumption of any combination of dairy, meat, poultry, fish or egg food groups.

Statistical analysis

We required 720 pregnant women to achieve 80% power to detect a difference of 100 g in

birth weight between groups. We assumed a mean birth weight of 2531 g (standard deviation

(SD) 415 g) in the control group based on the findings from a similar trial in Bangladesh.[26]

We set a significance level of 0·05 (two-sided) and assumed an intra-cluster correlation

coefficient (ICC) of 0·03[27] and 36 clusters with an average of 10-15 live births per cluster per

month. To adjust for 10% pregnancy loss and 15% loss to follow-up based on similar

trials,[26,28], we inflated the required sample size to 900 pregnant women.

We analysed the data with Stata version 17 and conducted intention-to-treat analyses. We

assigned the intervention randomly at the cluster level but assessed outcomes at the

individual level. We estimated the mean between-group difference in birth weight using

linear regression and the relative risk for LBW using binomial regression. We included the

study group as a covariate and adjusted for cluster randomisation using Generalised

Estimating Equations (GEE).[29] We estimated the intervention effect on maternal dietary

diversity and consuming individual food groups in the last 24 hours by multilevel linear and

Poisson mixed models, respectively. The linear mixed model estimated the difference in the

mean number of food groups consumed (out of 7), and the Poisson model estimated the

relative risk of consuming individual food groups between intervention and control arms. In

each model, we included study arms (intervention =1 and control =0) and gestational months

(2-9) as fixed effects with an interaction term between them. We also included two random

effect variables for adjusting the cluster RCT design and the repeated measurements at 2-9

gestational months. After each model, we used the post-estimation *lincom* command to

estimate the intervention effect at each gestational month.

We conducted post hoc subgroup analyses for the modifying effects of age, education, family

income, and place of delivery on the primary outcome, low birth weight. We added an

interaction term to assess whether there was a difference in intervention effect for any sub-

group, using a significance level of 0·01.

Role of the funding source

221 The funders had no role in study design, participant selection, data collection, and analyses.

222 The corresponding author had full access to the study's data and final responsibility for the

223 decision to publish.

Results

We identified 2,154 newly pregnant women from August 2015 to February 2016 in the

selected clusters. Of these women, 937 did not meet the eligibility criteria (603 had a

gestation of >12 weeks, 147 were not permanent residents, and 187 had planned to move out

of the study area for childbirth), and 324 declined to participate. We recruited 893

participants, 445 in 18 intervention clusters and 448 in 18 control clusters, and we lost 44

participants during follow-up, 25 (5·6%) intervention, and 19 (4·2%) control. Birth weight

was missing for 22 participants (Figure 1). There were 42 pregnancy losses (abortion or

stillbirth) and six neonatal deaths within 24 hours of birth.

There were no important differences in baseline characteristics between the groups (Table 1).

There were small differences in the mother's age, mother's and father's education, and parity

but not enough to warrant an adjustment in analyses. Slightly more intervention participants

received four or more ANC visits, but this trend reversed for antenatal care from health

centres (Table 1). When we combined both ANC data sources, the intervention participants

received slightly more ANC (intervention 61.7% vs. control 55.6%). A somewhat higher

percentage of intervention participants delivered in a facility (20.7% vs. 16.6%), but this

variable reflects the small differences in ANC. Mean birth weight was higher in facility

deliveries (2,915 g, SD 560 g) than in-home births (2,769 g, SD: 410 g).

Supervisors attended 5% of balanced plate demonstrations to observe the appropriateness of

message delivery, the use of the menu and food chart, demonstration, problem-solving, and

counselling. Approximately 6-7 of 8 messages were delivered to 80% of the participants. The

time taken for the demonstrations gradually decreased with increasing numbers of visits.

Counselling family members largely depended on availability; husbands were present at 15-

20% of visits, and other family members at >80% of visits.

We found a statistically significant difference in mean birth weight and LBW (Tables 2 and

3). The mean difference in birth weight between groups was 127·5 g (95% CI: 11.1, 243·9;

p=0·032) adjusted for clustering. The estimated relative risk (RR) of LBW was 57% lower in

the intervention group (RR: 0·43; 95% CI: 0·25, 0·75; p=0·003) adjusted for clustering.

We conducted subgroup analyses of the mother's age, education, and family income (Tables

2 and 3) and found a significant intervention effect on birth weight in adolescents vs. non-

adolescents. Mean difference in birth weight between groups was greater in adolescents

(299·1 g; 95% CI: 101·6, 496·6; p=0·003; vs 95·6 g; 95% CI: -17·3, 208·4; p=0·097) (p-value

for interaction 0·009) as was reduction in LBW (RR: 0·28; 95% CI: 0·11, 0·71; p=0·007 vs

RR: 0·54; 95% CI: 0·29, 0·98; p=0·044), although this was not statistically significant (p-

value for interaction 0·22). There was no evidence of treatment modification by the mother's

education or family income (Tables 2 and 3).

To assess bias in ascertaining birth weight, we stratified by participants birthed in a facility

whose babies were weighed by independent observers and those birthed at home and found

no evidence of treatment modification (p=0.78).

After adjustment for ANC, using iron and folic acid, and clustering, the reduction in mean

birth weight and LBW was smaller in the intervention group (Supplemental Tables S1 and

S2). The adjusted difference was 121·8 g (95% CI: 11.7, 231·9; p=0·030), which is 5.7 g less

than the unadjusted analysis (Table 2). The adjusted relative risk of LBW was 0·53 (95% CI:

0·29, 0·98; p=0·042), which was smaller than the unadjusted relative risk in Table 3.

The mean number of food groups consumed increased in the intervention group and was

significantly higher from the third month of pregnancy (Table 4, Fig 2). The largest

difference was in the eighth month of pregnancy (Table 4, Fig 2). The mean number of

animal-source protein foods consumed was significantly higher from the fourth month of

pregnancy. At the fourth gestational month the intervention participants consumed 1.5 animal

source foods / day (95% CI: 1.38, 1.64) compared to 1.1 (95% CI: 1.02, 1.25) in control

participants. The largest difference was in the eighth gestational month when intervention

participants consumed 1.9 animal source foods/day (95% CI: 1.68, 2.15) compared to 1.1

(95% CI: 1.02, 1.24) in control participants.

The relative consumption rate of beans, legumes and nuts, dairy food and other fruits and

vegetables was significantly higher in the intervention than in the comparison group after the

third or fourth gestational month (Table 4). The largest relative increases were for eggs, dairy

foods and other fruits and vegetables after the fourth or fifth gestational month (Table 4).

Discussion

We demonstrated that antenatal nutrition education with practical demonstrations of

preparing a balanced diet increases infant birth weight and reduces LBW in rural Bangladesh.

After adjustment for ANC, using iron and folic acid, and clustering, there was a higher mean

birth weight in the intervention group of public health significance (121·8 g) than in the

control group. The effect of the intervention was greater in adolescent mothers. The findings

provide strong evidence of the effectiveness of a sustainable intervention suitable for

implementation in community-based health care systems.   They demonstrate the potential

impact of nutrition education to improve intrahousehold food distribution as a tool to reduce

low birth weight in Bangladesh and across South Asia.

The process evaluation of the trial[30] provided insights into why the intervention was so

impactful. It identified key elements in the intervention strategy that was crucial in achieving

the desired adherence. These include the practical demonstration of portion sizes, addressing

local food perceptions, demystifying animal-source foods, engaging husbands and mothers-

in-law, leveraging women's social networks, and harnessing community health workers'

social roles. In particular, involving the mothers-in-law and husbands changed their attitudes

towards a fairer allocation of foods to pregnant women. The husbands adhered to the

messages and purchased more nutritious foods for their wives by adjusting the shopping

budget and even working extra hours.

Comparison with other studies

Our intervention appears more effective than interventions using micronutrients or food

supplements. A large cluster RCT (22,405 pregnancies) in Bangladesh assessed the impact of

antenatal micronutrient supplementation and found a 54 g (95% CI: 41, 66) increase in birth

weight and a 12% reduction (RR: 0.88; 95% CI: 0·85, 0·91) in LBW.[26] Another cluster RCT

of lipid-based micronutrient supplementation showed a 41 g birth weight increment without a

significant effect on LBW.[28]

Providing micronutrient supplements, even with additional food energy, is unlikely to address

the fundamental problems of insufficient food intake in pregnancy related to food taboos and

inequitable intra-household food distribution. The earlier MINIMat RCT (3267 singleton

newborns)[31], which tested early and late starts to food supplements with micronutrients in

pregnancy, found no significant main effects on birth weight. However, combining multiple

micronutrients and early food supplementation improved infant survival. The lack of a

control group without food supplements might explain the lack of birth weight response. In

contrast, an observational study in Bangladesh reported a 118 g increase in birth weight with

daily 608 kcal food supplements combined with iron-folic acid sustained for more than four

months.[32] Although this evidence is weak, it supports using food-based solutions to address

nutrition problems.

Adolescent pregnancy has an increased risk of adverse birth outcomes, including LBW.[33] If

severely undernourished, the growing pregnant adolescent and foetus compete for nutrients.[34]

A recent study from rural Bangladesh found a cessation of linear growth, weight loss, and fat

and lean body mass depletion in pregnant adolescents.[34] This greater nutritional susceptibility

helps explain the greater effect of balanced plate education in this group.

The balanced-plate intervention significantly increased maternal dietary diversity (Table 4,

Fig 2). Change in dietary patterns is the likely cause of the higher birth weight in the

intervention group. A recent cohort study[35] followed 374 women from their first antenatal

care visit to assess associations between dietary diversity during pregnancy and LBW in rural

Ethiopia. The LBW risk doubled with a dietary diversity score of less than four food groups

(ARR: 2.06; 95% CI: 1.03, 4.11). A facility-based case-control study[36] from rural Ethiopia,

which examined iron, folic acid and nutrition counselling, snack consumption, dietary

diversity, and maternal undernutrition, also reported similar findings.

There was a significant six-fold increase in the adjusted odds of LBW with an inadequate

minimum dietary diversity score (AOR: 6.65; 95% CI, 2.31, 19.16). A cross-sectional study[37]

from Ghana also reported a protective effect of higher dietary diversity scores for LBW

(AOR: 0.10; 95% CI: 0.04, 0.13 per standard deviation change score, p=<0.0001). This

evidence supports our findings that improved and more diverse dietary intake leads to a

reduction in LBW.

Participants in the intervention group consumed more animal-sourced protein foods, reaching

a peak at eight months of gestation of approximately two animal-sourced foods/day. A large

cross-sectional study of pregnant women in Shaanxi province, northwestern China, reported

an average weekly consumption of five animal-sourced foods.[38] As the frequency of animal-

sourced food increased by one time/week, the average birth weight increased by 3.24 g (95%

CI: 1.09, 5.39). Unlike increasing animal-sourced foods in the diet, consuming excess protein

supplements (>20% of energy as protein) can harm fetal growth.[39]

Despite earlier evidence of the positive impact of nutrition education on pregnancy outcomes,

the delivery method has received little attention. It is particularly important in low-literate

and poor self-efficacy populations, such as rural Bangladesh, where women require dietary

counselling to be accessible, feasible, and sanctioned by the community. A review of over

300 studies found that nutrition education is more likely to be effective when focusing on

behaviour and action than knowledge alone. And it is also likely to be more effective if

grounded in theory.[40]   Fishbein and Yzer note that "any given behaviour is most likely to

occur; if one has a strong intention to perform the behaviour; if a person has the necessary

skills and abilities required to perform the behaviour; and if no environmental constraints are

preventing the behavioural performance." [41] To positively impact dietary behaviour, we

designed our intervention to embrace all aspects of this theory. We adopted an innovative

counselling technique, including a practical demonstration of a balanced diet and engaged

family decision-makers. Similar nutrition counselling intervention was found effective in

increasing age-specific complementary feeding practices in rural Bangladesh.[42]

In Bangladesh, the main source of nutrition education for pregnant women is through ANC,

but the contact time is insufficient to affect nutrition change.[43] Thus, WHO. Proposes that

upskilling CHWs in delivering dietary advice is an appropriate approach to complement

health services.[44] In our study, we integrated the balanced plate intervention into BRAC's

existing ANC package delivered by CHWs. The same staff in a recent intervention

demonstrated their successful counselling on breastfeeding and complementary feeding.[45]

Previously, we also found that CHWs can influence family decision-makers to increase

pregnant women's iron-folic acid consumption.[46]

Strengths and limitations of this study

The evidence presented from this trial is robust. Key strengths are the cluster, randomised

controlled trial design with sufficient statistical power to detect clinically important changes

in outcomes. The delivery of the trial treatments through an established home-based antenatal

care service led to a balance between the treatment groups in the level of antenatal contact

with community health workers. The unblinded intervention was counterbalanced by

objective outcomes and blinding participants and data collectors to the aims and hypotheses.

The trial had a very high follow-up (93%), and we used intention-to-treat analyses and

adjusted for unbalanced baseline factors and clustering. Our intervention was logistically

simple and culturally appropriate based on well-designed qualitative formative research. The

key to our success was conducting the trial within existing ANC frameworks. The delivery

strategy resonates with maternal and child nutrition recommendations to achieve adequate

coverage of nutrition-specific interventions by reaching needy populations.[47] However, we

acknowledge some limitations. First, *Shasthya Kormi*, who delivered the intervention,

collected birth weight data. Measuring birth weight and providing nutrition counselling are

routine tasks for these staff. By blinding the SK to the study outcome, we would expect non-

differential bias. Also, we found no difference in the effects between women who had birth

weight measured by hospital staff versus those who were birthed at home with birth weight

measured by the SK. Secondly, using catchment areas as clusters might introduce

contamination, with the possibility of *Shasthya Kormi* exchanging information. However, this

would have biased results towards a null effect. To prevent this, we trained intervention and

control group *Shasthya Kormi* separately at regular refresher sessions. Thirdly, we could not

quantify food intake and intra-household food distribution, which might have provided

further evidence of pathways to the observed impact. Finally, we excluded over 30% of

pregnant women who identified because they were already in the second or third trimester

when we detected their pregnancy. There is no information about the characteristics of these

excluded women, and thus no way to compare them to those included in the trial. This lack of

data limits our assessments of the external validity of the trial findings.

Conclusions and policy implications

Our results indicate that a balanced plate nutrition education during pregnancy increases birth

weight and reduces LBW. Our trial offers a pragmatic food-based strategy for resource-poor

environments, especially where there is gender-biased intrahousehold food allocation. The

CHW-led intervention is simple, feasible, and easy to implement in settings with an existing

community nutrition outreach program. The intervention can potentially be adapted in

Bangladesh for implementation through the existing community health infrastructure.

References

1.   Black RE, Allen LH, Bhutta ZA, et al. Maternal and child undernutrition: global and
regional exposures and health consequences. *Lancet.* 2008;371:243-260.
2.   Cesar G Victora, Linda Adair, Caroline Fall, et al. Maternal and child undernutrition:
consequences for adult health and human capital. *Lancet.* 2008;371:340-357.
3.   Wilcox AJ, Russell IT. Birthweight and Perinatal Mortality: II. On Weight-Specific
Mortality. *International Journal of Epidemiology.* 1983;12(3):319-325.
4.   Watkins WJ, Kotecha SJ, Kotecha S. All-cause mortality of low birth weight infants
in infancy, childhood, and adolescence: a population study of England and Wales.
PLoS medicine. 2016;13(5):e1002018.
5.   Lee AC, Katz J, Blencowe H, et al. National and regional estimates of term and
preterm babies born small for gestational age in 138 low-income and middle-income
countries in 2010. *Lancet Glob Health* 2013;1:e26-36.
6.   Lee AC, Kozuki N, Cousens S, et al. Estimates of burden and consequences of infants
born small for gestational age in low and middle income countries with
INTERGROWTH-21st standard: analysis of CHERG datasets. *BMJ.* 2017;358(j3677).
7.   Han Z, Mulla S, Beyene J, Liao G, McDonald SD. Maternal underweight and the risk
of preterm birth and low birth weight: a systematic review and meta-analyses.
International journal of epidemiology. 2010;40(1):65-101.
8.   Triunfo S, Lanzone A. Impact of maternal under nutrition on obstetric outcomes.
Journal of endocrinological investigation. 2015;38(1):31-38.
9.   World Health Organization. *Maternal anthropometry and pregnancy outcomes: a
WHO collaborative study.* World Health Organization; 1995.
10.   Naidu A, Rao N. Body mass index: a measure of the nutritional status in Indian
populations. *European journal of clinical nutrition.* 1994;48:S131-140.
11.   Islam MZ, Akhtaruzzaman M, Lamberg-Allardt C. Nutritional status of women in
Bangladesh: comparison of energy intake and nutritional status of a low income rural
group with a high income urban group. *Asia Pacific Journal of Clinical Nutrition.*
2004;13(1):61-68.
12.   Harris-Fry H, Shrestha N, Costello A, Saville NM. Determinants of intra-household
food allocation between adults in South Asia – a systematic review. *International
Journal for Equity in Health.* 2017;16(107).
13.   Boyd NR, Jr., Windsor RA. A meta-evaluation of nutrition education intervention
research among pregnant women. *Health education quarterly.* 1993;20(3):327-345.
14.   Girard AW, Olude O. Nutrition education and counselling provided during
pregnancy: effects on maternal, neonatal and child health outcomes. *Paediatric and
Perinatal Epidemiology.* 2012;26(s1):191-204.
15.   Gresham E, Collins CE, Mishra GD, Byles JE, Hure AJ. Diet quality before or during
pregnancy and the relationship with pregnancy and birth outcomes: the Australian
Longitudinal Study on Women's Health. *Public Health Nutrition.* 2016;19(16):2975–
2983.
16.   Ota E, Hori H, Mori R, Tobe-Gai R, Farrar D. Antenatal dietary education and
supplementation to increase energy and protein intake. *Cochrane Database of
Systematic Reviews.* 2015(6).
17.   Lopes KdS, Ota E, Shakya P, et al. Effects of nutrition interventions during pregnancy
on low birth weight: an overview of systematic reviews. *BMJ Global Health.*
2017;2(e000389).

18.   Chowdhury M, Raynes-Greenow C, Alam A, Dibley M. Making a balanced plate for
450   pregnant women to improve birthweight of infants: a study protocol for a cluster
451   randomised controlled trial in rural Bangladesh. *BMJ Open.* 2017;7(e015393).
452   19.   Sen B, Ali Z. Spatial Inequality in Social Progress in Bangladesh. *The Bangladesh*
453   *Development Studies.* 2009;XXXII(2).
454   20.   *Poverty Maps of Bangladesh 2010.* World Bank Bangladesh, Bangladesh Bureau of
455   Statistics, and World Food Programme2011.
456   21.   *The Food Security Atlas of Bangladesh: Towards a poverty and hunger free*
457   *Bangladesh.* Planning Commission, Bangladesh Bureau of Statistics and UN World
458   Food Programme. 2004.
459   22.   National Institute of Population Research and Training (NIPORT), International
460   Centre for Diarrhoeal Disease Research Bangladesh (icddrb), MEASURE Evaluation.
461   *Bangladesh District Level Socio-demographic and Health Care Utilization Indicators.*
462   Dhaka, Bangladesh, and Chapel Hill, NC, USA: NIPORT, icddr,b, and MEASURE
463   Evaluation;2019.
464   23.   Nahar Q, Choudhury S, Faruque MO, Sultana SSS, Siddiquee MA. *Desirable Dietary*
465   *Pattern for Bangladesh.* National Food Policy Capacity Strengthening
466   Programme;2013.
467   24.   Arimond M, Torheim LE, Wiesmann D, Joseph M, Carriquiry A. Dietary diversity as
468   a measure of the micronutrient adequacy of women's diets: results from rural
469   Bangladesh site. *Washington, DC: Food and Nutrition Technical Assistance II*
470   *Project (FANTA-2), FHI.* 2009;360.
471   25.   FAO and FHI. Minimum dietary diversity for women: a guide for measurement.
472   *Rome: FAO.* 2016.
473   26.   West Jr PK, Shamim AA, Mehra S, et al. Effect of Maternal Multiple Micronutrient
474   vs Iron–Folic Acid Supplementation on Infant Mortality and Adverse Birth Outcomes
475   in Rural Bangladesh: The JiVitA-3 Randomized Tria. *JAMA.* 2014;312(24):2649-
476   2658.
477   27.   Zeng L, Cheng Y, Dang S, et al. Impact of micronutrient supplementation during
478   pregnancy on birth weight, duration of gestation, and perinatal mortality in rural
479   western China: double blind cluster randomised controlled trial. *BMJ.*
480   2008;337(7680):1211 - 1215.
481   28.   Mridha MK, Matias SL, Chaparro CM, et al. Lipid-based nutrient supplements for
482   pregnant women reduce newborn stunting in a cluster-randomised controlled
483   effectiveness trial in Bangladesh. *American Society for Nutrition.* 2016;103:236-249.
484   29.   Rabe-Hesketh S, Skrondal A. *Multilevel and longitudinal modeling using Stata.*
485   STATA press; 2008.
486   30.   Alam A, Chowdhury M, Dibley MJ, Raynes-Greenow C. How can we improve the
487   consumption of a nutritionally balanced maternal diet in rural bangladesh? The key
488   elements of the "balanced plate" intervention. *International Journal of Environmental*
489   *Research and Public Health.* 2020;17(17):6289.
490   31.   Persson LÅ, Arifeen S, Ekström E-C, Rasmussen KM, Frongillo EA, Team MS.
491   Effects of prenatal micronutrient and early food supplementation on maternal
492   hemoglobin, birth weight, and infant mortality among children in Bangladesh: the
493   MINIMat randomised trial. *Jama.* 2012;307(19):2050-2059.
494   32.   Shaheen R, Francisco Ad, Arifeen SE, Ekström E-C, Persson LÅ. Effect of prenatal
495   food supplementation on birth weight: an observational study from Bangladesh1–.*Am*
496   *J Clin Nutr.* 2006;83:1355–1361.

33. King JC. The risk of maternal nutritional depletion and poor outcomes increases in
early or closely spaced pregnancies. *The Journal of nutrition.* 2003;133(5):1732S-
1736S.
34. Rah JH, Christian P, Shamim AA, Arju UT, Labrique AB, Rashid M. Pregnancy and
lactation hinder growth and nutritional status of adolescent girls in rural Bangladesh.
The Journal of nutrition. 2008;138(8):1505-1511.
35. Zerfu TA, Umeta M, Baye K. Dietary diversity during pregnancy is associated with
reduced risk of maternal anemia, preterm delivery, and low birth weight in a
prospective cohort study in rural Ethiopia. *The American journal of clinical nutrition.*
2016;103(6):1482-1488.
36. Girma S, Fikadu T, Agdew E, et al. Factors associated with low birthweight among
newborns delivered at public health facilities of Nekemte town, West Ethiopia: a case
control study. *BMC pregnancy and childbirth.* 2019;19(1):220.
37. Abubakari A, Jahn A. Maternal dietary patterns and practices and birth weight in
Northern Ghana. *PloS one.* 2016;11(9):e0162285.
38. Yan H, Dang S, Mi B, et al. Study on relationship between mother's animal sourced
food intake during pregnancy and neonate birth weight. *Zhonghua liu xing bing xue
za zhi= Zhonghua liuxingbingxue zazhi.* 2017;38(5):615-620.
39. Liberato SC, Singh G, Mulholland K. Effects of protein energy supplementation
during pregnancy on fetal growth: a review of the literature focusing on contextual
factors. *Food & nutrition research.* 2013;57(1):20499.
40. Contento IR. Nutrition education: linking research, theory, and practice. *Asia Pac J
Clin Nutr.* 2008;17(1):176-179.
41. Fishbein M, Yzer MC. Using Theory to Design Effective Health Behavior
Interventions. *Communication Theory.* 2003;13(2):164-183.
42. Sanghvi T, Haque R, Roy S, et al. Achieving behaviour change at scale: Alive &
Thrive's infant and young child feeding programme in Bangladesh. *Maternal & Child
Nutrition.* 2016;12(Suppl. 1):141-154.
43. Ahmed SM, Hossain MA, Chowdhury AMR, Bhuiya AU. The health workforce crisis
in Bangladesh: shortage, inappropriate skill-mix and inequitable distribution. *Human
Resources for Health.* 2011;9(3).
44. Lehmann U, Sanders D. *Community health workers: what do we know about them?
The state of the evidence on programmes, activities, costs and impact on health
outcomes of using community health workers.* Geneva: World Health
Organization;2007.
45. Alive & Thrive. *Alive & Thrive's approach and results in Bangladesh: Rapid
improvements in infant and young child feeding practices resulting from a large-scale
program.* Dhaka, Bangladesh: Alive & Thrive;2015.
46. Alam A, Rasheed S, Khan NU, et al. How can formative research inform the design
of an iron-folic acid supplementation intervention starting in first trimester of
pregnancy in Bangladesh? *BMC Public Health.* 2015;15(374).
47. Bhutta ZA, Das JK, Rizvi A, et al. Evidence-based interventions for improvement of
maternal and child nutrition: what can be done and at what cost? *The Lancet.*
2013;382:452-477.

Acknowledgments

This study is part of research towards a PhD at The University of Sydney funded by

Endeavour Postgraduate Scholarship (PhD); Department of Education and Training,

Australian Government. The study received partial support from the James P. Grant School

of Public Health, BRAC University, Bangladesh, and the Sydney Medical School, The

University of Sydney, Australia. We thank the *Shasthya Kormi,* the *Shasthya Shebika* of

BRAC, and the pregnant women and their families in Sherpur for their time in this study.

Funding:

James P. Grant School of Public Health; Bangladesh, and Sydney Medical School

Foundation; Australia

Author contributions:

MC conceived the overall study, developed the study design and data collection methods, and

conducted the data analysis, interpreted data, and wrote the first draft of the manuscript. MJD

and CRG provided critical input regarding study design, sample size calculation, outcome

evaluation and the statistical analysis plan. KA provided practical guidance in formulating the

implementation strategy. AA provided crucial input on formative research and process

evaluation design. PK provided technical support for data analyses, and SMB conducted the

dietary analyses. CRG and MC obtained funds for the intervention. CRG, MJD, KA, PK, AA

and SMB critically reviewed the manuscript and approved the final version for submission.

MC, MJD, CRG, PK, KA and AA agreed to be accountable for all investigations necessary to

resolve questions related to the accuracy or integrity of any part of the work.

Competing interests:

None to declare

Data and materials availability:

The data described in this manuscript, the code book, and the analytic code will be made

available upon request, pending approval by the investigators of an application to use the data.

Supplemental Information

S1 Table: The mean birth weight and the adjusted mean difference in birth weight of

balanced -plate intervention compared to usual programs.

S2 Table: Percentage of low birth weight and adjusted relative risk of balanced-plate

intervention compared to usual programs.

Table 1: Cluster, baseline maternal and household, and antenatal care and delivery

characteristics by study arm

	Intervention (n=382)	Control (n=397)
Cluster characteristics		
Number of clusters	18	18
Mean (SD) population in clusters in 2016	9775 (2007)	9639 (1763)
Mean (SD) number of households in clusters	2315 (288)	2321 (419)
Mean (SD) number of women of reproductive age	2639 (542)	2631 (481)
Mean (SD) number of pregnant women enrolled per clusters	24.7 (1.0)	24.9 (0.5)
Baseline maternal and household characteristics		
Mother's age (years):		
Mean (SD)	23·0 (4·6)	24·0 (4.3)
15-19	80 (21·7)	64 (17.0)
20-34	280 (76·1)	307 (81·4)
>34	8 (2·2)	6 (1·6)
Missing	14	20
Mother's education;		
No education	82 (22·0)	89 (23·1)
Primary	250 (67·2)	242 (62·7)
Secondary	27 (7·3)	35 (9·1)
Higher	13 (3·5)	20 (5·2)
Missing	10	11
Monthly family income (SD) in Bangladesh Taka	7308 (4303)	7213 (4438)
Less than 5000	87 (23.9)	79 (21.2)
5000 to less than 10,000	190 (52.2)	209 (56.2)
10,000 or more	87 (23.9)	84 (22.6)
Missing	18	25
Father's education:		
No education	130 (35·0)	151 (39·7)
Primary	188 (50·7)	169 (44·5)
Secondary	34 (9·2)	39 (10.3)
Higher	19 (5·1)	21 (5·5)
Missing	11	17
Marital Status		
Married	372 (97.4)	386 (97.3)
Not married	10 (2.6)	11 (2.7)
Parity:		

	Intervention (n=382)	Control (n=397)
Mean (SD)	1·8 (1·2)	1·9 (1·3)
0	36 (10·3)	48 (12·6)
1	134 (38·5)	115 (30·3)
≥2	178 (51·2)	217 (57·1)
Missing	34	17
Previous pregnancy loss		
Yes	67 (19·3)	87 (22·9)
No	281 (80·8)	293 (77·1)
Missing	34	17
Gestation at enrolment (weeks) Mean (SD)	10·1 (3·2)	9·6 (3·2)
Missing	27	2
Antenatal care and delivery characteristics		
Iron Folic Acid Supplements		
Mean number consumed (SD)	45.6 (61.8)	29.0 (46.8)
Any use	117 (30.6)	70 (17.6)
None	265 (69.4)	327 (83.4)
Antenatal care visit by SK		
Mean number of visits (SD)	3.8 (2.6)	3.6 (2.2)
<4 visits	158 (41.1)	198 (49.9)
≥4 visits	224 (58.6)	199 (50.1)
Antenatal care in health centres		
Mean number of visits (SD)	0.7 (1.1)	1.0 (1.4)
No Visit	227 (59.4)	246 (51.9)
<4 visits	143 (37.4)	169 (42.6)
≥4 visits	12 (3.1)	22 (5.5)
Place of delivery		
Facility	79 (20.7)	66 (16.6)
Home	280 (73.3)	323 (81.4)
Unknown	23 (6.0)	8 (2.0)
Mode of delivery		
Normal	331 (86.7)	361 (90.9)
C-Section	33 (8.6)	34 (8.6)
Unknown	18 (4.7)	2 (0.5)
Time of birth weight measurement		
Within 24h	266 (69.6)	290 (73.1)
24 to 48h	75 (19.6)	61 (15.4)
48 to 72h	13 (3.4)	19 (4.8)
Unknown	28 (7.3)	27 (6.8)

Numbers are frequencies (and column percentage) unless stated otherwise

Table 2: The mean birth weight and the mean difference in birth weight of balanced-plate intervention compared to usual programs.

	Number of participants		Birth weight (g)				Mean difference			
	Intervention	Control	Intervention		Control		Estimate	(95%CI)	P-value	P-value for interaction
			Mean (SD)		Mean (SD)					
Overall	382	397	2861.0	(444.2)	2736.8	(423.2)	127.5	(11.1, 243.9)	0.032	
Mother's age[1] (years)										0.009
<20	80	64	2916.3	(471.1)	2637.5	(481.0)	299.1	(101.6, 496.6)	0.003	
>=20	288	313	2847.6	(443.4)	2762.6	(401.1)	95.6	(-17.3, 208.4)	0.097	
Mother's education[2]										0.330
None	82	89	2869.5	(460.6)	2693.3	(339.7)	175.1	(29.9, 320.4)	0.018	
Primary	250	242	2839.2	(443.9)	2761.6	(394.4)	92.9	(-28.0, 213.7)	0.132	
Secondary or higher	40	55	2982.5	(435.5)	2741.8	(588.0)	226.1	(-33.5, 485.8)	0.088	
Family income[3]										0.236
Low	87	79	2819.5	(397.9)	2663.3	(420.4)	166.8	(10.7, 322.8)	0.036	
Medium	190	209	2806.8	(421.7)	2750.7	(390.8)	51.8	(-70.0, 173.6)	0.404	
High	87	84	2995.4	(514.5)	2791.7	(433.2)	183.3	(-40.9, 407.4)	0.109	

 Birth weight was analysed using linear regression. We adjusted all models for clusters using generalised estimating equation (GEE) models, assuming an exchangeable correlation structure and applied a sandwich

 estimator to standard errors. [1] n=745 due to 34 missing maternal age values. [2] n=758 due to 21 missing maternal education values. [3] n=736 due to 43 missing family income values.

582   **Table 3: Percentage of low birth weight and relative risk of balanced-plate intervention compared to usual programs.**

| | Number of participants | | Low birth weight (<2,500g) | | | | Relative risk | | | |
	Intervention	Control	Intervention n (%)		Control n (%)		Estimate	(95%CI)	P-value	P-value for interaction
Overall	382	397	37	(10)	87	(22)	0.43	(0.25, 0.75)	0.003	
Mother's age[1] (years)										0.217
<20	80	64	6	(8)	17	(27)	0.28	(0.11, 0.71)	0.007	
>=20	288	313	31	(11)	62	(20)	0.54	(0.29, 0.98)	0.044	
Mother's education[2]										0.240
None	82	89	11	(13)	22	(25)	0.54	(0.28, 1.05)	0.069	
Primary	250	242	25	(10)	47	(19)	0.50	(0.26, 0.93)	0.029	
Secondary or higher	40	55	1	(3)	12	(22)	0.11	(0.02, 0.8)	0.029	
Family income[3]										0.441
Low	87	79	9	(10)	25	(32)	0.31	(0.13, 0.76)	0.010	
Medium	190	209	23	(12)	41	(20)	0.60	(0.32, 1.14)	0.117	
High	87	84	5	(6)	12	(14)	0.45	(0.13, 1.53)	0.199	

583   Low birth weight was analysed using binomial regression with a log link. We adjusted all models for clusters using generalised estimating equation (GEE) models, assuming an exchangeable

584   correlation structure and applied a sandwich estimator to standard errors. [1] n=745 due to 34 missing maternal age values. [2] n=758 due to 21 missing maternal education values. [3] n=736 due to

585   43 missing family income values

587 **Table 4: Effect of balanced plate nutrition education intervention on maternal dietary diversity and individual food groups**

Indicators		Months of pregnancy							
		2	3	4	5	6	7	8	9
N	Inter.	131	367	364	349	339	302	257	159
	Cont.	167	362	369	366	345	318	272	178
Dietary Diversity	Mean Diff. (95%CI)	0.19 (-0.11, 0.49)	0.47 **(0.20, 0.73)**	0.63 **(0.36, 0.90)**	0.80 **(0.53, 1.07)**	0.98 **(0.71, 1.25)**	1.04 **(0.77, 1.31)**	1.14 **(0.86, 1.42)**	0.96 **(0.67, 1.25)**
Starchy food	RR (95% CI)	1.16 (0.95, 1.41)	1.01 (0.99, 1.03)	1.03 (0.99, 1.08)	1.02 (0.98, 1.07)	1.02 (0.99, 1.06)	1.03 (0.97, 1.08)	1.05 (0.97, 1.13)	1.09 (0.92, 1.30)
Beans, legumes, nuts	RR (95% CI)	1.19 (0.82, 1.75)	**1.37 (1.06, 1.76)**	**1.40 (1.11, 1.77)**	**1.38 (1.07, 1.79)**	**1.50 (1.17, 1.93)**	**1.66 (1.29, 2.14)**	**1.86 (1.37, 2.52)**	**1.81 (1.11, 2.93)**
Dairy food	RR (95% CI)	**2.31 (1.09, 4.91)**	1.44 (0.78, 2.66)	1.69 (0.82, 3.51)	**2.36 (1.10, 5.08)**	**2.81 (1.35, 5.85)**	**2.97 (1.40, 6.32)**	**2.92 (1.54, 5.54)**	**2.34 (1.21, 4.55)**
Fish and meat	RR (95% CI)	1.12 (0.9, 1.39)	1.00 (0.97, 1.04)	1.07 (0.99, 1.14)	1.05 (0.97, 1.13)	1.06 (1.00, 1.13)	**1.09 (1.01, 1.17)**	1.06 (0.97, 1.15)	1.14 (0.95, 1.36)
Eggs	RR (95% CI)	1.48 (0.72, 3.04)	1.97 (1.00, 3.88)	**2.51 (1.28, 4.94)**	**2.92 (1.59, 5.36)**	**3.59 (2.00, 6.41)**	**3.59 (1.91, 6.77)**	**3.82 (2.17, 6.73)**	**4.65 (2.18, 9.88)**
Vitamin A rich foods	RR (95% CI)	1.17 (0.7, 1.96)	1.12 (0.72, 1.75)	1.15 (0.87, 1.52)	1.20 (0.92, 1.57)	1.26 (0.98, 1.62)	1.14 (0.87, 1.49)	1.26 (0.95, 1.67)	1.37 (0.97, 1.94)
Other fruits & vegetables	RR (95% CI)	1.92 (0.80, 4.63)	**2.59 (1.24, 5.43)**	**2.38 (1.19, 4.75)**	**2.87 (1.34, 6.14)**	**2.88 (1.32, 6.25)**	**3.64 (1.71, 7.73)**	**5.34 (2.54, 11.19)**	**4.55 (2.04, 10.16)**

588 Inter.= Intervention, Cont.= Control, Mean Diff.= Mean Difference, CI= Confidence Interval, RR= Relative Risk, 95% CI presented in bold font is statistically significant (p-
589 value <0.05)

590   **Figure 1: Trial flow chart**

591
592

Figure 2: The mean number and 95% confidence intervals* of food groups consumed

by women per day by treatment group.

* The estimated the mean number of food groups consumed in the last 24 hours and 95% confidence intervals

were calculated using multilevel linear mixed models with study arms (intervention =1 and control =0) and

gestational months (2-9) as fixed effects with an interaction term between them and two random effect variables

for adjusting the cluster RCT design and the repeated measurements.