Echinocandin heteroresistance causes prophylaxis failure and facilitates breakthrough *Candida parapsilosis* infection

Authors: Bing Zhai [1,2,3,*,#], Chen Liao [4,*], Siddharth Jaggavarapu [5,6,7,8,*], Yuanyuan Tang [1,*], Thierry Rolling [2,3], Yating Ning [9,10], Tianshu Sun [10,11], Sean A. Bergin [12], Mergim Gjonbalaj [2,3], Edwin Miranda [13], N. Esther Babady [2,12], Oliver Bader [14], Ying Taur [2,3,15], Geraldine Butler [12], Li Zhang [9,10], Joao B. Xavier [4], David S. Weiss [5,6,7,8,#], and Tobias M. Hohl [2,3,15,#]

Institutions: [1] CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China

[2] Infectious Disease Service, Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

[3] Immunology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

[4] Computational and Systems Biology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[5] Emory Antibiotic Resistance Center, Atlanta, GA, USA.

[6] Department of Microbiology and Immunology, Emory University, Atlanta, GA, USA.

[7] Emory Vaccine Center, Atlanta, GA, USA.

[8] Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, GA, USA.

[9] Department of Laboratory Medicine, State Key Laboratory of Complex Severe and Rare

Diseases, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences,

Beijing, China.

[10] Beijing Key Laboratory for Mechanisms Research and Precision Diagnosis of Invasive Fungal

Diseases, Beijing, China.

[11] Medical Research Centre, State Key Laboratory of Complex Severe and Rare Diseases, Peking

Union Medical College Hospital, Chinese Academy of Medical Science, Beijing, China

[12] School of Biomolecular and Biomedical Science, Conway Institute, University College

Dublin, Belfield, Dublin 4, Ireland.

[13] Clinical Microbiology Service, Department of Laboratory Medicine, Memorial Sloan

Kettering Cancer Center, New York, NY, USA.

[14] Institute for Medical Microbiology and Virology, University Medical Center Göttingen, 37075

Göttingen, Germany.

[15] Department of Medicine, Weill Cornell Medical College, New York, NY, USA.

[*] : Contributed equally.

[#] : Address correspondence to: hohlt@mskcc.org, david.weiss@emory.edu or to

bing.zhai@siat.ac.cn

Abstract

Breakthrough infections of patients on antimicrobial prophylaxis represent a significant

and often unexplained cause of morbidity. Here, we reveal that in high-risk patients on

micafungin prophylaxis heteroresistance – the presence of a phenotypically unstable, low

frequency subpopulation of resistant cells (~1 in 10,000) – underlies breakthrough bloodstream

infections by *Candida parapsilosis* misclassified as susceptible by standard antimicrobial

susceptibility testing. By analyzing 219 clinical *C. parapsilosis* isolates from North America,

Europe, and Asia, we demonstrate widespread micafungin heteroresistance. To facilitate

detection of micafungin heteroresistance, we constructed a predictive machine learning

framework that classifies isolates as heteroresistant or susceptible by a maximum of ten genomic

features. Our results connect heteroresistance to unexplained prophylaxis failure and demonstrate

a proof-of-principle diagnostic approach with the potential to inform clinical decisions.

Introduction

Antimicrobial resistance leads to over one million annual deaths worldwide and is a

severe public health problem [1]. Heteroresistance is a form of antimicrobial resistance in which a

minority subpopulation (sometimes as rare as one in one million cells) of resistant cells co-exists

with a majority population of susceptible cells [2-6]. The minimum inhibitory concentration (MIC)

of the resistant cells for a given antimicrobial drug is typically much higher than that of the

susceptible cells within the same isolate [4]. Therefore, treatment with an antimicrobial drug kills

susceptible cells but does not kill rare resistant cells, which survive and replicate at

concentrations lower than the MIC of the resistant subpopulation. Due to the low frequency of

resistant cells, heteroresistance may be undetected by standard antimicrobial susceptibility

testing (AST) which typically relies on assays that detect the outgrowth of these cells after 24-48

hours of incubation.

Heteroresistance has been demonstrated to mediate treatment failure in murine infection

models [7-10], but it has been challenging to determine its role in antimicrobial failure in human

infections [11,12]. Among other confounding variables, many serious infections are treated with

multiple antibiotics, which prevents directly linking treatment failure and resistance to a single

drug. In contrast, prophylaxis typically relies on administering a single antimicrobial drug, thus

providing a simplified scenario to study the impact of drug-resistant phenotypes on antimicrobial

failure.

In patients undergoing allogeneic hematopoietic cell transplantation (allo-HCT),

randomized prospective clinical trials have established antifungal prophylaxis as the standard of

care to prevent fungal infections and to reduce all-cause mortality [13-16]. Among the three major

classes of antifungal drugs, azoles and echinocandins have been approved by the FDA for

prophylaxis [17]. Echinocandin drugs (i.e., micafungin, caspofungin, and anidulafungin) inhibit β-

1,3-glucan synthase, which is encoded by the *FKS1* gene and has no mammalian homologs [18].

Since echinocandins have a favorable toxicity profile, they are commonly used for antifungal

prophylaxis in high-risk clinical scenarios.

Candida parapsilosis is one of the most common *Candida* species isolated among

patients with invasive candidiasis [19]. Life-threatening *C. parapsilosis* infections caused by

fluconazole-resistant isolates in COVID-19 patients have been documented worldwide [20-23]. *C.*

parapsilosis can survive in the gastrointestinal tract as a commensal organism and translocate

into the bloodstream during microbial dysbiosis and mucosal injury [24]. Compared to other

Candida species, *C. parapsilosis* has higher echinocandin MICs due to a natural P660A

polymorphism in the target Fks1 protein [25]. While classical resistance to echinocandins is rarely

observed in this organism [26,27], echinocandin prophylaxis or exposure has been linked to an

increased frequency of *C. parapsilosis* infections in different medical facilities [28,29], with an

unclear cause of drug failure.

Here, we describe heteroresistance to micafungin in *Candida parapsilosis* isolates

obtained from fecal and blood samples of allo-HCT recipients. Importantly, our results show that

patients with intestinal colonization of heteroresistant *C. parapsilosis* isolates have a

significantly higher chance of developing breakthrough infections during micafungin

prophylaxis, a significant advance in our understanding of the clinical relevance of

heteroresistance in prophylaxis failure. Moreover, current antifungal susceptibility tests

misclassified heteroresistant *C. parapsilosis* isolates as susceptible, highlighting an urgent need

to develop novel diagnostic tools to identify heteroresistance and to inform clinical decision-

making. To advance the rapid diagnosis of the heteroresistant phenotype, we employed a large

collection of 219 clinical isolates from three continents. We then developed a machine learning

framework to predict micafungin heteroresistance from genomic variations. This proof-of-

concept approach to identify micafungin heteroresistance in *C. parapsilosis* could help optimize

antifungal drug selection, thereby avoiding prophylaxis failure, and improving patient care.

Results

Candida parapsilosis is a frequent cause of micafungin breakthrough bloodstream infections.

To analyze breakthrough fungal bloodstream infections (BSIs) during allo-HCT, we performed a cohort study that included 952 patients who underwent a total of 983 allo-HCTs from January 2016 to December 2020 at Memorial Sloan Kettering Cancer Center (MSKCC) (Extended Data Fig. 1). In the majority of allo-HCTs (n = 966), patients received antifungal prophylaxis with intravenous micafungin (100-150 mg/day) starting seven to ten days prior to allo-HCT for a median of 17 days, prior to switching to triazole-based prophylaxis per institutional practices [30]. We identified a total of 25 micafungin breakthrough BSIs that occurred within 30 days post transplantation (Extended Data Fig. 1), with a clear predominance of *Candida parapsilosis* as the etiologic agent in these infections (12 out of 25 cases, Fig. 1a).

Fig. 1. Micafungin heteroresistance in *C. parapsilosis* bloodstream isolates. (a) Micafungin

breakthrough bloodstream infections (BSIs) in MSKCC allo-HCT recipients between day 0 and

30 post transplantation (between Jan. 2016 and Dec. 2020). The "others" category includes four

episodes of breakthrough BSI by *Candida kefyr*, *Trichosporon asahii*, *Capronia munkii*, and

Saccharomyces cerevisiae. **(b)** Micafungin E-test results of two representative *C. parapsilosis*

isolates, MSK65 (heteroresistant) and MSK804 (susceptible). **(c – h)** Population-level responses

of MSK65 and MSK804 to micafungin. **(c, f)** Micafungin population analysis profile (PAP) of

isolates **(c)** MSK65 and **(f)** MSK804. The % survival was calculated compared to growth on

drug-free agar. **(d, g)** CFU dynamics of total and resistant subpopulations of **(d)** MSK65 and **(g)**

MSK804 in YPD broth containing 2 µg/mL micafungin. Cultures were plated at indicated time

points for enumeration of total (black, circles) and resistant (colored, squares) cells. **(e, h)**

Frequency of micafungin-resistant subpopulation of isolates **(e)** MSK65 and **(h)** MSK804 during

continuous passage. The frequency was calculated after a 24-hour culture in drug-free YPD broth

(Pre-treatment), followed by passaging in YPD broth with 2 µg/mL micafungin (In micafungin)

for 48 hours, and a second passage in drug-free YPD broth for 72 hours (Subculture). Bar height:

mean; Error bar: standard deviation (n = 3). (i) Graphical illustration of the dynamics of the

resistant subpopulation in a micafungin heteroresistant isolate in the presence (+ mica) or

absence of micafungin (-mica). Blue cells are susceptible, orange cells are resistant (unstable),

and grey cells are those killed by micafungin.

Micafungin heteroresistance is prevalent in isolates from *C. parapsilosis* breakthrough

infections.

We were able to retrieve 12 *C. parapsilosis* bloodstream isolates from 10 allo-HCT

recipients on micafungin prophylaxis between day 0 and 30 post transplantation (Fig. 1a). In

addition, we retrieved three *C. parapsilosis* isolates from three patients who received micafungin

prophylaxis after day 30 of transplantation (Extended Data Fig. 1). Notably, all 15 isolates were

classified susceptible to micafungin by broth microdilution (Table 1) and E-test (i.e., Epsilometer

test; Fig. 1b) assays. However, neither of these conventional ASTs had sufficient sensitivity to

determine if there was a low frequency subpopulation of replicating cells resistant to micafungin,

i.e., heteroresistance. Thus, we re-analyzed all isolates by the population analysis profile (PAP)

assay, the gold standard for detecting heteroresistance [4] (see Methods). In PAP assays, 10^6 fungal

cells were plated on solid agar plates that contained micafungin at indicated concentrations. In

this study, we defined an isolate as heteroresistant to micafungin when it contained resistant cells

with a frequency between 0.0001% (i.e., limit of detection) and 50% across three consecutive

serially diluted micafungin concentrations that included 4 µg/ml (2x breakpoint concentration,

consistent with the European Committee on Antimicrobial Susceptibility Testing guidelines [31],

Supplementary Table 1).

As an example, isolate MSK65 was identified as heteroresistant since it harbored a minor

subpopulation of micafungin-resistant cells that survived at concentrations up to 32 µg/ml (Fig.

1c). The resistant cells rapidly replicated and became dominant during micafungin exposure (Fig.

1d), indicating that they were not persister cells which cannot replicate exponentially in the

presence of an antimicrobial drug [32]. However, the frequency of resistant cells reverted to the

baseline level during subsequent growth in the absence of micafungin, indicating that they were

not classical resistant cells (Fig. 1e). In contrast, MSK804 was entirely killed by micafungin

exposure without the emergence of a resistant subpopulation (Fig. 1f-h). This isolate was

classified as micafungin-susceptible. We confirmed that the heteroresistant phenotypes of blood

isolates were consistent across three PAP replicates (Extended Data Fig. 2a and 2b).

Using PAP assays, we found that 11 out of the 15 blood isolates were heteroresistant,

with the frequency of resistant subpopulations ranging from 0.008% to 0.8% (Table 1). We

further determined the MICs for both the susceptible (MIC-S) and resistant (MIC-R)

subpopulations in these 11 heteroresistant isolates (see Methods for definition of MIC-S and

MIC-R). Due to the predominance of susceptible cells, the clinical MIC values obtained from

broth microdilution aligned well with the respective MIC-S values (Table 1). However, the MIC-

R values of these isolates ranged from 32 to 64 µg/ml, indicating an increase of 32- to 64-fold

when compared to their corresponding MIC-S values. In addition to micafungin, these isolates

were generally, but not uniformly, heteroresistant to two other echinocandin drugs, caspofungin

and anidulafungin (Extended Data Fig. 2c and 2d). Together, these findings demonstrate the

phenomenon of heteroresistance (HR) to echinocandins in *C. parapsilosis* clinical isolates.

Isolate	Clinical MIC	Clinical designation	PAP	MIC-S	MIC-R	Fold diff.	Resistance freq.
MSK65	1	S	HR	1	64	64	0.033%
MSK67	1	S	HR	1	64	64	0.033%
MSK478	1	S	HR	1	64	64	0.12%
MSK485	1	S	HR	1	64	64	0.075%
MSK794	2	S	HR	1	64	64	0.667%
MSK795	2	S	HR	1	64	64	0.1%
MSK802	1	S	HR	1	64	64	0.8%
MSK804	1	S	S	1	n.d.	n.a.	n.a.
MSK811	1	S	S	1	n.d.	n.a.	n.a.
MSK812	2	S	HR	2	64	32	0.375%
MSK1191	1	S	S	1	n.d.	n.a.	n.a.
MSK2384	2	S	HR	1	64	64	0.014%
MSK2386	1	S	S	1	n.d.	n.a.	n.a.
MSK2387	1	S	HR	1	32	32	0.04%
MSK2390	0.5	S	HR	1	32	32	0.008%

Table 1. Micafungin susceptibility profiles of 15 bloodstream *C. parapsilosis* isolates from

micafungin breakthrough infections. Clinical minimum inhibitory concentration (MIC) data

and susceptibility phenotype were determined by the YeastOne assay. The MIC-S (MIC of the

susceptible subpopulation) and MIC-R (MIC of the resistant subpopulation) values were

determined by the population analysis profile (PAP) curves (see Methods for details). S,

susceptible; HR, heteroresistant; n.d. – not detected; n.a. – not available. The fold difference

(Fold diff.) represents the fold change between MIC-S and MIC-R. The resistance frequency

(Resistance freq.) indicates the proportion of resistant cells in each isolate that survived at the

micafungin breakpoint concentration (2 ug/ml [31]). Isolates MSK65 and MSK67 were recovered

from different blood cultures of patient 1 in Fig. 2c, and isolates MSK485 (HR) and MSK1191

(S) were isolated from different blood cultures of patient 3 in Fig. 2c.

Intestinal colonization by a micafungin heteroresistant strain increases the risk of

breakthrough bloodstream infection during prophylaxis.

A previous study showed that the intestinal tract is a major reservoir for *C. parapsilosis*

which can translocate across the intestinal mucosa into the bloodstream and cause BSI in allo-

HCT recipients [24]. To investigate whether micafungin-heteroresistant blood isolates originated

from the intestinal tract, we recovered 85 fecal *C. parapsilosis* isolates from 30 patients (5 of

whom developed BSI and were investigated in Fig. 1a). Among these isolates, 43 (50.6%)

originated from 14 of the patients and were classified as heteroresistant by PAP (Supplementary

Table 2). The remaining 16 of the 30 patients were colonized only by susceptible *C. parapsilosis*

isolates, and none of these patients developed breakthrough BSI. Among the 14 patients who

harbored heteroresistant isolates in their intestinal tract, five patients developed a breakthrough

C. parapsilosis BSI (Fig. 2a). These data establish an association (P = 0.014; two-sided Fisher's

Exact Test) between intestinal colonization with micafungin-heteroresistant *C. parapsilosis*

isolates and ensuing breakthrough BSI (Fig. 2b).

To confirm the gut-to-blood transmission of micafungin-heteroresistant *C. parapsilosis*,

we sequenced the genomes of fecal and blood *C. parapsilosis* isolates from the five patients with

breakthrough BSIs. In each patient, the bloodstream isolate was identical, defined as >99.999%

population average nucleotide identity (popANI) [33], to an intestinal isolate collected from the

feces prior to BSI, with both isolates having the same heteroresistant or susceptible phenotype

(Fig. 2c).

Interestingly, two patients (Patient 2 and 3) had a mixture of heteroresistant and

susceptible isolates in their feces before BSI (Fig. 2c). Longitudinal analysis revealed that in

Patient 2, the heteroresistant isolates replaced the susceptible isolates, leading to an ensuing BSI

by a heteroresistant isolate. In Patient 3, both susceptible and heteroresistant isolates translocated

to the bloodstream independently, resulting in two separate episodes of BSI (Fig. 2c). These

conclusions remained unchanged using an even more stringent popANI cutoff of 99.9999%.

Taken together, our analysis demonstrates that micafungin-heteroresistant *C. parapsilosis*

isolates can colonize the intestine and can subsequently translocate to the bloodstream, thereby

presenting a risk factor for the development of breakthrough fungal BSI.

In Patient 2, four of five genomes of susceptible isolates and all genomes of

heteroresistant isolates were identical within each category, as defined by the popANI metric.

Importantly, the susceptible isolates were distinct from the heteroresistant isolates at the same

popANI cutoff (Fig. 2c). Similarly, in Patient 3, two separate clusters of susceptible and

heteroresistant isolates were observed. These findings reinforce the notion that the

heteroresistance phenotype did not emerge from susceptible isolates during the micafungin

prophylaxis period in the patients that we examined. To test this further, we examined 16

patients from whom we isolated at least two identical fecal *C. parapsilosis* isolates at various

time points. Micafungin susceptibility phenotypes remained unchanged among these identical

isolates in 15 out of 16 patients (Extended Data Fig. 3). In one patient (patient 47), the phenotype

of two identical isolates changed from susceptible to heteroresistance, due to a two-fold change

in the MIC of the resistant subpopulation in the later isolate.

Fig. 2. Micafungin heteroresistance and risk of breakthrough *C. parapsilosis* BSIs in allo-HCT recipients. (a) A flow chart showing steps involved in determining the number of MSKCC allo-HCT recipients (2016 – 2020) with *C. parapsilosis* intestinal colonization and breakthrough BSIs. **(b)** Fisher's Exact test (two-sided) on patients with micafungin-heteroresistant *C. parapsilosis* intestinal colonization and breakthrough BSI. **(c)** Comparative genomic analysis between fecal and blood *C. parapsilosis* isolates. Longitudinal medical data are shown to the left for five allo-HCT recipients with *C. parapsilosis* breakthrough BSI, including antifungal drug administration (first row), the time and drug susceptibility of bloodstream isolate(s) (second row), the time of fecal sample collection and the drug susceptibility of corresponding fecal isolate (third row). Pairwise popANI (population average nucleotide identity) between blood (star) and fecal (circle) *C. parapsilosis* isolates were shown in a heatmap to the right of the

medical data. At a cutoff of 99.999%, red colors indicate "identical" isolates, and blue colors

indicate "distinct" isolates. The grey dashed line on the diagonal indicate self-comparisons

between the same isolates (i.e., popANI = 100%). Patient 1 and Patient 2 in this figure were

Patient 2 and Patient 3, respectively, in our previous study [24].

Association between micafungin heteroresistance phenotype and genomic variations in

clinical isolates from three continents.

To expand the scope of the study, we included 118 *C. parapsilosis* clinical isolates from

MSKCC (see Methods) and supplemented the collection with an additional 18 isolates from

other hospitals in the United States, 73 isolates from 32 different hospitals from 20 provinces in

China, 9 isolates from Germany (Göttingen), and one isolate from France (Nantes), yielding a

total of 219 isolates (Supplementary Table 2). Using the PAP method, we found the presence of

micafungin-heteroresistant *C. parapsilosis* isolates in hospitals from all three continents (68/136,

11/73, 7/10 micafungin-heteroresistant isolates from the North America, Asia, and Europe,

respectively). This finding highlights the global prevalence of this emerging and clinically

significant phenotype.

To understand the genomic underpinnings of micafungin heteroresistance, we sequenced

the genomes of these 219 isolates and profiled the genomic variations (GATK, see Methods).

Chromosome-level variations such as aneuploidy have been reported in antifungal

heteroresistance in *C. parapsilosis* and other human fungal pathogens[34-36]. We identified 12 out

of the 219 isolates (5.5%, eight heteroresistant and four susceptible isolates) with different

patterns of trisomy (seven, four, and one isolates have an extra copy of chromosome VI, III, and

V, respectively, see Extended Data Fig. 4). Although 8 out of 12 isolates with aneuploidy were

micafungin-heteroresistant, the study may lack power to confirm or reject a correlation (two-

sided Fisher's exact test, P = 0.066). An even larger collection of *C. parapsilosis* isolates would

help determine further a relationship between aneuploidy and micafungin heteroresistance.

Notably, most heteroresistant isolates (78 of 86) did not exhibit aneuploidy.

We next characterized copy number variations (CNVs), which have been linked with

azole tolerance [37], across the 219 *C. parapsilosis* genomes. We refined a previous algorithm on

CNV characterization [38] and defined the copy number of each ORF by two features (see

Methods and Extended Data Fig. 5 for illustrative examples): the first feature describes whether

an ORF presents a full or a partial amplification or deletion (CNV_categorical or CNV_cat), and

the second feature quantifies the copy number of amplified or deleted regions (CNV_quantitative

or CNV_quant). With this approach, we identified 3,317 ORFs that contain amplified or deleted

regions in at least one isolate in this dataset (Supplementary Table 3). Among the 2,774 ORFs

with amplifications, only 53 show overlaps with inverted repeats of at least 100 bp, suggesting

that these CNVs were unlikely to be technical artifacts (Supplementary Table 4).

We identified 43,531 single nucleotide variants (SNVs) and 3,722 small

insertion/deletions (INDELs) among the 219 isolates (Supplementary Table 5). These variations

served as the basis for constructing a phylogenetic tree (see Methods; Fig. 3). Notably, the

majority of heteroresistant isolates cluster within one clade, establishing a robust correlation

between the heteroresistance phenotype and phylogeny (Moran's I = 0.30, P < 0.0001).

Fig. 3. Phylogenetic tree of 219 *C. parapsilosis* **isolates collected from three continents.** The

outer circle indicates the micafungin susceptibility phenotype of each isolate.

289   We next sought to identify variations that are associated with micafungin heteroresistance

among the profiled SNVs, INDELs, and CNVs. We first encoded these variations as a mixture of

continuous (CNV_quant) and categorical (SNVs, INDELs, CNV_cat) features and then merged

redundant features if they correlated perfectly with each other across all 219 genomes (Methods).

This resulted in a non-redundant set of 6,806 features (Supplementary Table 6). Using logistic

regression (see Methods), we identified 453 features that were significantly associated with the

heteroresistance phenotype after multi-test correction (Fig. 4a, Supplementary Table 7). We

confirmed that the top five features with both positive and negative associations did not overlap

with the inverted repeats identified above (Supplementary Table 4). The most significant feature

(Feature 4376) was a homozygous alternative allele at the 5'UTR of CPAR2_206130 ($P_{adj.}$ =

2.4e-15). The *Saccharomyces cerevisiae* ortholog of this gene encodes a microtubule-associated

protein that plays a role in fungal cell wall organization [39] (Fig. 4b). The second most significant

feature (Feature 2154) was the full deletion of CPAR2_600440 (P = 2.9e-15), which encodes a

putative glycosylphosphatidylinositol (GPI)-anchored protein involved in cell wall integrity and

cell-cell interactions [40] (Fig. 4b). Both features remained the top-ranking ones when we

examined the associations by LASSO (Least Absolute Shrinkage and Selection Operator; see

Methods and Supplementary Table 7).

All 453 features with a significant correlation with heteroresistance also correlated with

phylogeny (Moran's I test, $P_{adj.} < 0.05$). Among these features, Feature 100 (a homozygous

alternative allele within CPAR2_109680, $P_{adj.} = 5.7e-10$) was the only one found in an

orthologue of the gene encoding the primary drug target. However, this correlation could be due

to phylogeny (Extended Data Fig. 6a). We applied a phylogenetic LASSO model to re-evaluate

the associations of genomic features with heteroresistance while controlling for their

phylogenetic dependences (see Methods). This approach yielded 40 features with non-zero

coefficients, where a homozygous alternative variant (Feature 4470) in CPAR2_206800 had the

largest positive coefficient (Supplementary Table 8). Although this feature did not completely

overlap with the heteroresistant isolates outside the major cluster, it successfully identified the

heteroresistance phenotype of three isolates within a phylogenetic region that is dominated by

susceptible isolates (Extended Data Fig. 6b). Taken together, the phenotype of micafungin

heteroresistance in *C. parapsilosis* is associated with genomic variations and these associations

are strongly, but not exclusively, coupled with phylogeny.

Fig. 4. Top _C. parapsilosis_ genomic features associated with micafungin heteroresistance.

(a) Volcano plot of 6,806 _C. parapsilosis_ genomic features. **(b)** The top two features positively

and negatively associated with micafungin heteroresistance and the annotations of the open

reading frames that contain the four features. Distribution of the features were plotted in

alignment with the phylogenetic tree. Chromosome locations: Feature 4376: Chr 2; Feature

2154: Chr 6; Feature 3654: Chr 8; Feature 2520: Chr 1. SNV, single nucleotide variation; CNV,

copy number variation.

Machine-learning-based diagnosis of micafungin heteroresistance

Given the clinical significance of micafungin heteroresistance, a rapid diagnosis with high accuracy is critical for the rapid administration of effective antifungal drugs. However, current laboratory approaches (i.e., PAP assay) require incubation times up to several days with high material and labor demands. To facilitate the diagnostic process, we developed a machine learning framework (Extended Data Fig. 7) that predicts the micafungin heteroresistance phenotype from the profiled genomic features found in the 219 clinical isolates. Given that multiple isolates from a single patient may share identical genomes (i.e., popANI ≥ 99.999%), separating these isolates into the training and test sets could lead to data leakage and compromise the model evaluation. To address this issue, we removed the duplicates of identical isolates collected from each patient (see Methods, Extended Data Fig. 8), resulting in a total of 160 non-redundant isolates for further analysis. The machine learning framework includes feature standardization, feature selection, hyperparameter optimization, and classification.

In the machine learning framework, we assessed three feature selection methods (logistic regression, LASSO, a neural-network-based approach ENNS [41]) along with three tree-based classifiers: Decision Tree (DT), Random Forest (RF), and Extreme Gradient Boosting (XGBoost). Our approach limited each feature selection method to choose a maximum of ten features to mitigate the risk of overfitting and, more importantly, to reduce the time and effort required to measure the selected features in the laboratory (see Discussion). All model hyperparameters, including the number of features selected, were optimized through a grid search in a 5-fold cross-validation setting. The 160 non-redundant isolates (Supplementary Table 9) were divided into a training set (80%) and a test set (20%), the process of which was repeated 50 times. On average, the combination of logistic regression and XGBoost consistently

outperformed the other eight combinations in F1 score (Fig. 5a) and other evaluation metrics

(precision, recall, and accuracy, Extended Data Fig. 9a). This combination achieved a median

precision of 0.78, a recall of 0.70, an F1 score of 0.74, and an accuracy of 0.84. This strategy can

be further simplified by developing SNV-only models, which have equivalent performance to the

models developed with SNVs, INDELs and CNVs (Fig. 5a). To confirm that the optimal model

captures more than spurious correlations, we randomly permuted the heteroresistance labels to

simulate the shuffled data and retrained the model. As expected, the resulting median F1 score

never exceeded 0.25 across three repetitions (Fig. 5b). This striking difference strongly indicated

that our model effectively captured meaningful genomic features of micafungin heteroresistance.

Given the strong association between heteroresistance and phylogeny (Fig. 3), we

developed an alternative phylogeny-based predictive approach (see Methods). This approach

placed genome sequences of the test isolates into a reference phylogenetic tree created by the

training isolates and conducted hidden-state prediction to infer their micafungin heteroresistance

labels. Interestingly, the best machine learning model achieved a slightly higher median F1 score

(0.74) than the phylogeny-based approach (0.71, Fig. 5b). Although the difference in

performance did not attain statistical significance (paired *t*-test, P = 0.457), both methods

substantially outperformed a popANI-based method (Fig. 5b), which simply assigned a test

isolate the same heteroresistance label as its most similar (i.e., highest popANI) isolate in the

training set. These findings indicate that the features chosen by the machine learning model

exploit the phylogenetic signal of heteroresistance and expand on it to achieve predictive power.

The most commonly misclassified isolates, including false positive and false negative predictions

(Extended Data Fig. 9b), showed micafungin susceptibility phenotypes inconsistent with their

phylogenetic neighbors. Collectively, our proof-of-concept machine learning framework

demonstrated its effectiveness in predicting the micafungin heteroresistant phenotype based on a

small set of genomic features. This innovative approach could facilitate the identification of

antimicrobial heteroresistance in clinical settings.

Fig. 5: Evaluation of machine learning model performance in predicting the micafungin

heteroresistance phenotype. F1 score is an evaluation metric that combines precision and

recall. Unless specified otherwise, all models selected features from a combined pool of SNV

(single nucleotide variant), INDEL (insertion/deletion), and CNV (copy number variation)

features. In all boxplots, each dot represents a single train-test split, which was randomly

repeated 50 times. **(a)** Machine learning models constructed by choosing among three feature

selection algorithms (LOGIT, Logistic regression; LASSO, Least Absolute Shrinkage and

Selection Operator; ENNS, Ensemble Neural Network Selection) and three classifiers (DT,

Decision Tree; RF, Random Forest; XGBoost, Extreme Gradient Boosting). **(b)** Comparison of

the best-performing combination (LOGIT+XGBoost) with alterative models. R1, R2, and R3

represent three repeats of retrained LOGIT+XGBoost models using randomly shuffled

heteroresistance data. The ANI (average nucleotide identity)-based model quantified genomic

similarity between isolates by popANI and assigned a test isolate the same heteroresistance label

of its genomically closest isolate in the training set. Phylogeny-based model placed test isolates

into a reconstructed phylogeny of training isolates and then performed hidden-state prediction.

Two-sided paired-sample *t*-test: exact P values are provided.

Discussion

This study shows that micafungin heteroresistance promotes breakthrough *C.*

parapsilosis infections in allo-HCT patients on micafungin prophylaxis. While both

heteroresistant and susceptible *C. parapsilosis* isolates could colonize the intestinal tract,

micafungin heteroresistant isolates were enriched in patients who developed breakthrough BSI.

Factors such as host immunity or gut microbiota composition may also contribute to fungal

translocation from the gut to the bloodstream. Nonetheless, our results show intestinal

colonization by a micafungin heteroresistant strain increases the risk of developing breakthrough

BSI. Importantly, we did not observe any instances of classical, stable resistance to micafungin

in this study. These findings highlight heteroresistance, rather than classical resistance, as a

major cause of breakthrough *C. parapsilosis* infections during echinocandin prophylaxis.

Identifying heteroresistance is challenging in the clinic because the resistant

subpopulation reverts to a very low frequency after removing the drug (i.e., the initial blood

culture process and the ensuing subculture of fungal cells for drug susceptibility testing).

Standard clinical ASTs fail to detect micafungin heteroresistance. PAP assay – the gold standard

for heteroresistance in research laboratories – is laborious and can take days to weeks to identify

slow-growing resistant cells. Isolate phylogeny, which we found to be a robust predictor of

micafungin heteroresistance in *C. parapsilosis*, requires whole genome sequencing which is also

time-consuming and computationally involved. In contrast, the machine learning models

introduced in this study include at most ten genomic features, which can be rapidly measured by

regular Sanger sequencing for SNVs and INDELs, quantitative PCR for CNVs. This approach

can bypass the time needed to culture fungal isolates during AST testing and is more precise than

phylogeny-based prediction (Fig. 5). Due to the increasing prevalence of antimicrobial

resistance, this proof-of-concept approach may advance current diagnostics and support rapid,

personalized clinical decisions in therapy and infection control. A future research direction is to

harness this strategy to detect intestinal colonization by heteroresistant *C. parapsilosis* using

metagenomic data.

Predicting micafungin heteroresistant phenotype can be improved in future studies by

using a larger, more diversified collection of *C. parapsilosis* isolates combined with their

micafungin susceptibility phenotypes. Classical resistance or tolerance to echinocandin drugs has

been attributed to chromosomal mutations in the *FKS* or *MSH2* genes in *Candida* species [25,42].

Here, with the exception of Feature 100 (SNP in CPAR2_109680, Extended Data Fig. 6a), we

did not identify variations in *FKS* or *MSH2* genes associated with micafungin heteroresistance.

In fact, insights from azole tolerance studies have indicated that, unlike conventional resistance

which often arises from mutations of the direct drug target, tolerance or heteroresistance may

hinge on changes in stress response pathways [2]. Elucidation of the biological functions of these

genes and the implications of the identified variations require further investigation, and could be

guided by future models trained on larger datasets. Future epidemiologic studies should also

clarify the potential spread of heteroresistant strains within hospital or healthcare settings.

Finally, we examined the presence of echinocandin heteroresistance in another medically

relevant fungal pathogen, *Candida auris*. *C. auris* are highly drug-resistant pathogenic fungi that

have rapidly spread across the world since their identification in 2009. More than 99% of the *C.*

auris isolates in the United States are resistant to fluconazole, 50-60% are resistant to

amphotericin B, and less than 10% are resistant to echinocandins [43-45]. Thus, echinocandin drugs

are the first-line empiric therapy for *C. auris* infections, as recommended by the CDC [46]. We

identified micafungin heteroresistance among *C. auris* isolates from the CDC/FDA AR Bank

using PAP analysis (Extended Data Fig. 10). Interestingly, recent reports describe echinocandin

treatment failures caused by susceptible *C. auris* as classified by standard AST, along with

emergence of resistance while the patients were being treated with echinocandins [43,44]. It is

worthwhile to determine if some of these isolates are heteroresistant to echinocandins, which

may link heteroresistance to these reported treatment failures. Such potential treatment failures,

together with the prophylaxis failures that we observed in allo-HCT recipients, would highlight

further the urgent need for rapid detection of echinocandin heteroresistance in clinical settings.

In summary, our study has uncovered the existence of heteroresistance to echinocandin

drugs in *Candida* species and has linked this phenotype to prophylaxis failures in allo-HCT

recipients. The proof-of-concept machine learning framework formulated here lays the

foundation for future development of rapid and precise diagnostic methods for heteroresistance.

Methods:

Inclusion criteria for allo-HCT recipients and *Candida parapsilosis* isolates at MSKCC

Patients that underwent allo-HCT at Memorial Sloan Kettering Cancer Center (MSKCC)

between Jan. 1st, 2016 to Dec. 31st, 2020 were examined for micafungin breakthrough fungal

bloodstream infections. In this study, we defined micafungin breakthrough bloodstream fungal

infections as a positive fungal blood culture if a patient received a minimum of one day of

micafungin prophylaxis prior to the blood culture positivity, or if micafungin prophylaxis was

discontinued within three days before the blood culture positivity. 15 isolates were retrieved

from documented breakthrough BSI in these patients: 12 isolates from 10 patients who

developed BSI between transplant day 0 to 30 and three isolates from three patients who

developed BSI after day 30. The medical record of non-allo-HCT patients with *Candida*

parapsilosis bloodstream infections from the same study period (2016 to 2020) were also

reviewed (Supplementary Table 2). A total of 14 bloodstream *C. parapsilosis* isolates from these

non-allo-HCT patients who developed *C. parapsilosis* BSI based on the criterion of having a

positive blood culture were obtained from clinical micro lab at MSKCC.

Fecal biospecimens from allo-HCT recipients were drawn from a prospective fecal

collection biobank, which has been described previously [47] (MSKCC IRB protocol 16-834).

Fecal samples from a subset of 205 patients (collected from -10 to 30 day of transplantation) had

been cultured for viable fungal isolates as previously described [24]. Among them, a total of 30

patients were confirmed with intestinal colonization by *C. parapsilosis* from fecal culture. For

the five patients with intestinal colonization of *C. parapsilosis* before BSI, at least one isolate

from each fecal sample was randomly picked for further analyses. For the remaining 25 patients

with intestinal colonization but no fungemia, one to two *C. parapsilosis* isolates from each *C.*

parapsilosis culture positive fecal samples were randomly picked. Detailed fecal isolate – patient

information data are listed in Supplementary Table 2. Fecal biospecimens from eight healthy

donors [48] were also examined for the *C. parapsilosis* culture positivity. Two fecal samples were

identified as positive and two *C. parapsilosis* isolates from each sample were randomly picked.

Taken together, there were a total of 89 fecal *C. parapsilosis* isolates from MSKCC included in

this study. Detailed information of patient demographics, antimicrobial exposure, bloodstream

infections, and fecal isolates are available in Supplementary Table 2.

C. parapsilosis **isolates from non-MSKCC hospitals**

Among the 18 *C. parapsilosis* isolates from United States other than MSKCC, nine

isolates were acquired from David Andes's lab at University of Wisconsin-Madison. The other

nine isolates were acquired from the CDC/FDA AR Bank. 73 Asian isolates were acquired from

China Hospital Invasive Fungal Surveillance Net (CHIF-NET). These isolates were collected

from 32 hospitals from 20 different provinces in China. Ten European isolates include one from

Nantes, France and nine from Göttingen, Germany. The detailed information of the isolates are

available in Supplementary Table 2.

Classification of micafungin heteroresistance

In this study, an isolate was classified as heteroresistant if the frequency of resistant cells

is at least 0.0001%, but remains lower than 50%, across all three consecutive serial-diluted

concentrations which include 4 µg/ml (2x breakpoint concentration, EUCAST [31]). The possible

set of three consecutive concentrations could be 1, 2, and 4 µg/ml, or 2, 4, and 8 µg/ml, or 4, 8,

and 16 µg/ml. Additionally, we determined the MIC-S (MIC of the susceptible subpopulation)

and MIC-R (MIC of the resistant subpopulation) from PAP curves. MIC-S was defined as the

minimum concentration at which survival subpopulation drops below 50%. On the other hand,

MIC-R was defined as the minimum concentration at which the survival subpopulation falls

below the detection limit (0.0001%) at this concentration and all other concentrations beyond it.

Population analysis profile assay (PAP)

YPD agar (Sigma) plates were made with each antifungal drug at the indicated

concentrations. The stocks of micafungin (Biovision), caspofungin (Biovision) and

anidulafungin (Biovision) were made by dissolving each drug in molecular biology grade water

(Corning). Overnight cultures of all the isolates were grown in 2 mL of YPD broth (RPI) in 8 mL

round-bottom tubes (Corning) at 35°C with 200 rpm shaking. Ten-fold serial dilutions of the

overnight cultures were made in PBS in 96-well plates (Corning). 5 µL of each dilution, from

undiluted to 10^{-6}, were spotted on the agar plates. Plates were incubated for 48 to 72 hours at

35°C before CFUs were enumerated. To calculate the percentage survival, CFUs from the plates

containing antifungals were divided by those from the plates without antifungals.

Time-kill assays

2 mL of YPD broth supplemented with 2 µg/mL micafungin for *C. parapsilosis* or 0.5

µg/mL micafungin for *C. auris*, were inoculated with 10^6 CFU from an overnight culture. These

YPD cultures with micafungin were incubated at 35°C with shaking at 200 rpm for up to 72

hours. At each indicated time points, 50 µL aliquots were acquired from the cultures and CFUs

were counted by plating serial dilutions on plates with and without micafungin at the same

concentrations as the respective broth cultures.

Antifungal resistance stability assay

For antifungal treatments, *C. parapsilosis* cultures were sub-cultured (1:100) from a drug-free overnight culture into YPD with 2 µg/mL micafungin and grown at 35°C with 200 rpm shaking for 48 hours. Micafungin-free subcultures were performed by diluting the micafungin culture 1:1,000 into fresh YPD media and the cultures were grown for 72 hours. Cultures were diluted in PBS and plated on YPD-agar plates with and without micafungin to determine the proportion of resistant *C. parapsilosis* cells.

E-tests

Micafungin E-test strips were obtained from Biomerieux (Marcy-l'Étoile, France). The E-tests were performed according to the manufacturer's instructions. Briefly, 5 colonies from an overnight YPD agar plate were dissolved into sterile PBS. The OD was adjusted to 0.5 McFarland standard. The solution was applied to RPMI 1640 plates with MOPS and 2% Glucose (Remel) with a cotton swab. The E-test strip was laid on the plate and the images were captured after 48-hour incubation at 35°C.

Average nucleotide identity (ANI) and identity between isolates

ANI is a quantitative metric for comparing the genomic similarity between two microbial genomes. We calculated the pairwise population ANI (popANI) values for all 219 *C. parapsilosis* isolates by the inStrain package [33] (version 1.6.3). A recommended threshold of 99.999% was used to determine whether two isolates are identical or distinct. However, pairwise comparisons may pose challenges in identifying identical isolates when more than two isolates are compared. For example, the identity between isolate A and C cannot be determined when

isolate A and B are identical (popANI ≥ 99.999%) and isolate B and C are identical (popANI ≥

99.999%), but isolate A and C are not identical (popANI < 99.999%).

To address this issue, we developed a computational algorithm to extract the largest

clusters in which the popANI value between any two isolates is at least 99.999% and no

additional isolate has popANI ≥ 99.999% with all isolates in the cluster. Within each such

cluster, all isolates are identical to each other. To identify these clusters, we first conducted

hierarchical clustering of the binarized popANI matrix at the 99.999% threshold. Using the

"fcluster" function in the Python SciPy package (version 1.10.1), we systematically reduced the

maximum allowed inter-cluster distance until the presence of a cluster where all pairwise

popANI values among its isolates are at least 99.999%. If multiple such clusters exist, we

selected the largest cluster and assigned it a cluster ID. Subsequently, isolates in this cluster were

removed from the popANI matrix, and hierarchical clustering was repeated on the remaining

isolates. We iterated these steps until no clusters containing more than one isolate could be

identified. The remaining isolates were each treated an autonomous cluster. By dividing all

isolates into clusters, we define that any two isolates are identical if they belong to the same

cluster; otherwise, they are distinct.

Genomic variation calling, filtering, and annotation

Reads were mapped against *C. parapsilosis* CDC317 reference genome [49] using BWA-

mem (version 0.7.17-r1188) with parameters -M -R 'RG', where 'RG' represents read group

information generated for each isolate. Duplicate reads were marked using the GATK (version

4.1.9.0) MarkDuplicates tool as per GATK best practices. SNVs and INDELs were identified

using the HaplotypeCaller tool from GATK. The resulted GVCF files were combined using the

CombineGVCFs tool and jointly genotyped using the GenotypeGVCFs tool from GATK.

The raw SNVs and INDEls were hard-filtered using the following criteria. Filter settings for

SNVs include mapping quality (< 40), variant quality score (< 30), quality by depth (< 2), strand

odds ratio (> 3), Fisher strand bias (> 60), mapping quality rank-sum test (< -12.5) and read

position rank-sum test (< -8). Filter settings for INDELS include quality by depth (< 2), variant

quality score (< 30), Fisher strand bias (> 200) and read position rank-sum test (< -20). Very low

coverage regions with a read depth < 10 and a genotype quality < 20 were set to missing on a

per-sample basis. Monomorphic (no alternative alleles were called for any of the samples) and

multiallelic (all sites at which only alternative alleles are called) variants, clusters of three or

more SNPs called in a 10-base pair window, and variants in repetitive regions identified by

TandemRepeatFinder[50] (version 4.09.1) were removed. In addition, we removed SNVs that are

within 10 bp of INDELs and allowed only one variant within a cluster of INDELs separated by 5

or fewer base pairs to pass. The remaining variants surviving the filtering process were exported

in table format using the GATK VariantsToTable tool. SNVs and INDEls were annotated using

the SnpEff tool[51] (version 5.0).

Characterization of copy number variation (CNV)

We characterized CNV for each open reading frame (ORF) across eight nuclear

chromosomes. The distribution of read depth within each ORF was computed using BEDTools

(version 2.31.0)[52]. This distribution was then transformed into a copy number distribution by

dividing the read depth by the average read depth across all diploid chromosomes, which was

then multiplied by two. Amplified or deleted regions within an ORF can be identified by

analyzing the number and position of peaks within the distribution. A normal diploid ORF

should display a single peak at the position of two copies, whereas amplifications or deletions

result in shifts in peak positions or the emergence of extra peaks. All peaks were identified from

smooth Gaussian kernels fitted to the distribution. For each peak, the percentage contribution to

the overall distribution was calculated by dividing the area under the peak (boundaries

determined as midpoints between this peak and nearest peaks) by the length of the ORF. Small

peaks that contributed less than 20% were deemed data noise and filtered out. The remaining

peak contributions were then renormalized.

Each CNV was characterized by two variables: a categorical one indicating the type of

CNV (normal, full or partial amplification, and full or partial deletion), and a continuous one

quantifying the copy number of the amplified or deleted regions. When only one peak remains,

the CNV type was categorized as "normal" if the peak centers at two copies, "full amplification"

if it centers at a copy number greater than two, or "full deletion" if it centers at a copy number

less than two. In cases where two peaks remain and one of them centers at two copies, the CNV

type was categorized as "partial amplification" if the other peak centers at a copy number greater

than two, or "partial deletion" if it centers at a copy number less than two. If more than two

peaks remain and one of them centers at two copies, the amplified or deleted region was

determined by the peak with the largest percent contribution among all peaks whose copy

number is not two. If two or more peaks remain but none of them center at two copies, both

variables were determined based on the location of the largest peak. In the extreme scenario

where all peak contributions were below 20% before filtering, these unfiltered small peaks were

grouped. The quantitative copy number of this grouped peak was calculated as a weighted

average of the copy numbers of these unfiltered peaks.

Identification of *C. parapsilosis* isolates with aneuploidy

Aneuploidy was identified through the presence of multiple peaks in the distribution of average ORF (open reading frame) coverage across the entire genome. Among the 219 isolates, 15 exhibited two peaks and none exhibited more than two. In isolates with two peaks, the center of the major peak corresponded to the average read depth mapped to normal diploid chromosomes, while the minor peak represented chromosome(s) with potential aneuploidy. The copy number of chromosomes with potential aneuploidy was calculated as the ratio of the minor peak's read depth to the major peak's, which was then multiplied by two. We identified 12 isolates with aneuploidy that met the criteria of the copy number ≥ 2.5 or ≤ 1.5.

Identification of inverted repeats

An inverted repeat consists of two arms of similar DNA sequences with one single stranded sequence followed downstream by its reverse complement. The inverted repeats in the *C. parapsilosis* reference genome CDC317 were identified using Inverted Repeats Finder (version 3.08). The input parameters include Match = 2, Mismatch = 3, Delta = 5, PM = 80, PI = 10, Minscore = 20, MaxLength = 100000, MaxLoop = 1000000.

Feature encoding and merging

SNVs, INDELs, and CNV types are categorical variables and were encoded by one-hot encoding. Specifically, each SNV or INDEL was encoded by two dummy variables: a binary variable indicating whether this variant is a heterozygous allele and a binary variable indicating whether this variant is a homozygous alternative allele. Each CNV type variable was encoded by four dummy variables, each of which indicates whether the type is "full amplification", "partial

amplification", "full deletion", or "partial deletion". No transformation was needed for the

quantitative CNV variable.

The total number of variables, including both dummy variables created for SNVs,

INDELs, CNV types and unaltered quantitative CNV variables, is 59,881. These variables were

termed "raw features". Additionally, we merged "raw features" that shared identical values

across the 219 isolates. These variants that perfectly correlate with each other across multiple

genomic loci suggest linkage equilibrium. A total of 6,806 merged features were identified.

Phylogenetic tree construction

The filtered VCF file containing genomic variants served as input for the RRHS tool[53]

(version, 1.0.0.2). This tool generated haploid FASTA sequences of equal length for each isolate

(parameter -n 1 generates one sequence per isolate). Subsequently, the phylogenetic tree was

constructed from these sequences using IQ-TREE 2 (version 2.0.7) [54]. IQ-TREE 2 was executed

with the following parameters: -T AUTO (automatically selected threads), -bb 1000 (ultrafast

bootstrap approximation [55] with 1000 replicates), -m MFP+ASC (use ModelFinder to determine

the best fit substitution model with ascertainment bias correction), and -alrt 1000 (1000 bootstrap

replicates for SH-like approximate likelihood ration test).

Logistic regression

We used the "Logit()" function implemented in the Python statsmodels package (version

0.14.0) to perform univariate logistic regression, i.e., the association with heteroresistance was

assessed for each feature individually. Raw P values were adjusted through the Bonferroni

correction. A significance threshold of 0.05 was applied to the adjusted P values.

LASSO and phylogenetic LASSO

The conventional LASSO (Least Absolute Shrinkage and Selection Operator) method

determines the coefficient vector β that associates genomic features (matrix X) with

heteroresistance label (vector Y) by minimizing $||Y - X\beta||_2^2/2N + \alpha||\beta||_1$, where α is a constant

that multiplies the regularization term, and $||\cdot||_1$ and $||\cdot||_2$ are L1 and L2 norm respectively. To

run LASSO analysis, we called the function "LassoCV" implemented in the Python's scikit-learn

library (version 1.2.2) with CV = 5 (5-fold cross validation) and n_alphas = 1,000 (number of α

along the regularization path).

The phylogenetic LASSO model is an extension of conventional LASSO by controlling

for variance explained by phylogeny. The technical details of this approach have been described

in the Supplementary Information of our previous study for bacterial genomics[56]. Briefly, the

matrix X (i.e., genomic features) and vector Y (i.e., heteroresistance label) need to be

transformed before applying conventional LASSO: $X_{new} \to (T'T)^{-1}T'X$, $Y_{new} \to (T'T)^{-1}T'Y$,

where matrix T is obtained from the phylogenetic variance-covariance matrix G by Cholesky

decomposition such as $TT' = G$. The phylogenetic variance-covariance matrix was computed

using the "vcv.phylo" function of the ape library[57] (version 5.7.1).

Machine-learning-based heteroresistance prediction

All machine learning models were developed using data from a training set and

subsequently assessed against a test set. As shown above, patient may harbor multiple identical

fecal and blood isolates due to repeated sampling and gut-to-blood translocation. If these

identical isolates were distributed into both the training and test sets, model performance may be

inflated as a result of data leakage. To avoid this potential risk, we systematically excluded

duplicate isolates sharing identical genomic sequences (see the section "Average nucleotide

identity (ANI) and identity between isolates" for the definition of identical isolates) within each

patient. This exclusion process was performed for isolates whose patient information is available.

The Random Forest (RF) and Extreme Gradient Boosting (XGBoost) models were built

using a fixed number of 1,000 trees. Other hyperparameters were tuned to maximize the F1 score

through grid search in a 5-fold cross-validation framework. The Decision Tree (DT) and RF

models shared three parameters, including the number of features chosen by an upstream feature

selection algorithm (n_features = 2, 4, 6, 8, 10), the number of features considered when

determining the optimal split (max_features = square root (n_features) and n_features), and the

maximum depth of each tree (max_depth = 2, 4, 6). The RF models had an extra tunable

parameter—the maximum percentage of samples each tree was trained on (max_samples = 0.5,

0.75, 1.0). In the XGBoost models, two unique parameters in addition to n_features and

max_depth were optimized: the percentage of features used for building each tree

(colsample_bytree = 0.5, 0.75, 1.0) and the learning rate controlling the step size at which model

weights were updated (learning_rate = 0.03, 0.1, 0.3).

The implementation of logistic regression and LASSO in Python has been described

above. In the case of logistic regression-based feature selection, features were ranked based on

their P values (from smallest to largest) and the top "n_features" were selected. When LASSO

was employed for feature selection, non-zero features were ranked based on their absolute values

(from largest to smallest). In situations where the coefficients of all features are zero, we

randomly selected "n_features" features from the entire set of features. When the number of non-

zero features was fewer than "n_features", all non-zero features were selected. The ENNS

algorithm has been implemented previously [41], see code at https://github.com/KaixuYang/ENNS.

The DT and RF models were constructed using the DecisionTreeClassifier and

RandomForestClassifier functions from Python's scikit-learn package (version 1.2.2),

respectively. The XGBoost models were developed using the XGBClassifier function from the

Python's XGBoost package (version 1.7.6). We used four metrics, including precision, recall, F1

score (= 2*precision*recall / (precision + recall)), and accuracy, to evaluate the performance of

machine learning models.

Phylogeny-based heteroresistance prediction

We used the PICRUSt2[58] pipeline (version 2.5.2) to predict heteroresistance phenotype.

For each train-test split, we began by performing sequence placement using the "place_seqs.py"

script. This step involved placing haploid sequences from test isolates into a phylogenetic tree

created from training isolates. The custom input files provided to the script "place_seqs.py"

included aligned haploid sequences of training isolates (generated by RRHS[53], version 1.0.0.2), a

phylogenetic tree of the training isolates and a model file specifying optimal tree parameters

(generated by RAxML-NG[59], version 1.2.0), as well as a hidden Markov model of aligned

training sequences (generated using hmmbuild from hmmer.org, version 3.1b2). The output of

the "place_seqs.py" script is a tree with placed test isolates. With this tree, we next ran hidden-

state prediction using the "hsp.py" script to infer their heteroresistance phenotype.

Data visualization

The data obtained from population analysis profile (PAP), time-kill, and antifungal

susceptibility assays were graphed by GraphPad prism (San Diego, version 9). All other data

were generated and visualized using Python's matplotlib package (version 3.7.1). All figures

underwent further editing in Adobe Illustrator (version 2021). Box plots, swarm plots, and

heatmaps were generated using the corresponding functions within Python's seaborn package

(version 0.12.2). In all boxplots, the central line represents the median, the box limits denote the

first and third quartiles respectively, and the whiskers extend to the smallest and largest values or

at most to 1.5× the interquartile range, whichever is smaller.

Other computational tools

In general, Python version 3.8.5 and R version 4.3.1 were used in this study for computational

analyses. Moran's I test for phylogenetic correlation was performed using "Moran.I" function

from the "ape" package (version 5.7.1). Phylogenetic trees were plotted and annotated using the

package "ggtree" (version 3.8.0). Fisher's exact test and paired-sample t-test were performed

using "fisher_exact" and "ttest_rel" from the SciPy (version 1.10.1) stats module respectively.

Data Availability Statement

The data that support the findings of this study are available from the corresponding authors

upon request. All sequencing data generated in this study will be deposited in the Sequence Read

Archive upon acceptance.

Code Availability Statement

Codes of this study are available at:

https://github.com/liaochen1988/EchinocandinHeteroresistance.git

Acknowledgements

We thank E. Pamer and K. Wolfe for discussions, and E. Zuniga for sequencing

assistance. We thank D. Andes for providing clinical *C. parapsilosis* isolates. This work was

supported by the National Key Research and Development Program of China 2021YFA0911300

(B.Z.), National Institutes of Health (NIH) grant nos. R01 AI093808 (T.M.H.), R21 AI105617

(T.M.H.), R21 AI156157 (T.M.H.), U19 AI158080 (D.S.W.), R01s AI141883 and AI148661

(D.S.W), U01 AI124275 (J.B.X.), R01 AI137269 (J.B.X. and Y.T.), an Investigator in the

Pathogenesis of Infectious Diseases Award from the Burroughs Wellcome Fund (T.M.H. and

D.S.W.), the Ludwig Center for Cancer Immunotherapy (T.M.H.), the Susan and Peter Solomon

Divisional Genomics Program (T.M.H.), Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) grant no. RO-5328/1-2 (T.R.), Science Foundation Ireland grant nos.

19/FFP/6668 (G.B.), 18/CRT/6214 (G.B.), and all authors from Memorial Sloan Kettering

Cancer Center were supported by NIH P30 CA008748.

Author contributions

The study was conceived by B.Z., C.L., S.J., D.S.W. and T.M.H.. MSKCC Clinical data

were collected and analyzed by Y.T.(MSKCC) and B.Z.. *C. parapsilosis* isolates from MSKCC

were collected and processed by B.Z., T.R., M.G., E.M., and N.E.B.. Isolates from China were

maintained and processed by Y.T., T.S., and L. Z.. Isolates from Europe were collected and

processed by S.B., G.B., and O. B.. Antifungal phenotypic assays were performed by S.J. and

Y.N.. Whole genome sequencing analyses were done by C.L. and Y.T.(SIAT). Association

analyses and machine learning modeling were done by C.L., Y.T.(SIAT), and J.B.X.. The

manuscript was written by B.Z., C.L., S.J., D.S.W. and T.M.H.. All co-authors reviewed and

edited the manuscript.

Extended Data Figures

Extended Data Fig. 1. Micafungin breakthrough infections in allo-HCT recipients in

MSKCC from Jan. 2016 to Dec. 2020. The dashed-line boxes indicate the patients or fungal

isolates analyzed in respective figure panels or tables.

Extended Data Fig. 2. Antifungal susceptibility of *C. parapsilosis* blood isolates to

echinocandin drugs. (a, b) Reproducibility of the PAP method in determining micafungin

heteroresistance: **a**, three repeats of PAP assays of the 15 *C. parapsilosis* blood isolates in Table

1. **b**, micafungin susceptibility phenotypes (susceptible or heteroresistant) classified by the PAP

curves in panel **a**. **(c, d)** Antifungal susceptibility of *C. parapsilosis* isolates to **(c)** caspofungin

and **(d)** anidulafungin. Isolates MSK65, MSK794, MSK795, and MSK2384 are heteroresistant to

micafungin (colored in orange), while the isolates MSK804, MSK811, and MSK2386 are

susceptible to micafungin (colored in blue). Among these isolates, the phenotypes of caspofungin

and anidulafungin heteroresistance were generally consistent with their micafungin

heteroresistance phenotype, with exceptions of MSK795 (heteroresistant to micafungin and

caspofungin, susceptible to anidulafungin) and MSK804 (susceptible to micafungin and

anidulafungin, heteroresistant to caspofungin). **(e)** Etest results of four additional *C. parapsilosis*

blood isolates. MSK67 and MSK795 are micafungin-heteroresistant (orange font), MSK811 and

MSK1191 are micafungin susceptible (blue font). Notably, both heteroresistant isolate

(MSK795) and susceptible isolate (MSK811) could have clean inhibition zone around the Etest

strip. Again, these observations showed the challenge of using Etest to differentiate micafungin-

heteroresistant and micafungin-susceptible isolates.

Extended Data Fig. 3. Tracking the stability of micafungin susceptibility phenotypes of

identical *C. parapsilosis* fecal isolates within patients. Each filled circle represents a fecal

isolate and all isolates shown in each row are identical. To simplify presentation, identical

isolates collected on the same day have been consolidated into a single, filled circle. The

susceptible and heteroresistant isolates obtained from Patient 2 were displayed at two separate

rows.

Extended Data Fig. 4. Twelve *C. parapsilosis* isolates with aneuploidy. The mean copy

number of consecutive fixed-size 5kb windows over the entire genome were plotted for the

isolates with aneuploidy. The mean copy number was computed as the mean read depth across

each 5kb window divided by that averaged over normal diploid chromosomes. Chr:

chromosome, S: susceptible, HR: heteroresistant.

Extended Data Fig. 5. Different types of copy number variations (CNVs). To illustrate our

approach for characterizing CNVs, we have provided five examples of open reading frames

(ORFs) with distinct CNV types. In each panel, the light gray curve represents distribution of

read depth of each position within an ORF. The solid red line fits a Gaussian kernel to this

distribution. The vertical dashed red line indicates the peak center of the fitted distribution. The

percentage contribution of each peak to the entire distribution is provided in the panel. The

feature values of these ORFs used in following analyses (e.g., statistical association, machine

learning) are shown in the table at the bottom right. Each ORF is described by a categorical

feature (CNV_cat) and a quantitative feature (CNV_quant). CNV_cat specifies the CNV types

(normal diploid ORF, partial amplification, full amplification, full deletion, partial deletion),

while CNV_quant specifies the copy number of the amplified or deleted regions (the value of

CNV_quant is 2 in the absence of amplification and deletion).

Extended Data Fig. 6. Distributions of Feature 100 (a) and 4470 (b) across the phylogenetic

tree. Both features are single nucleotide variants. The biological functions of the open reading

frames to which they belong are shown. *Saccharomyces cerevisiae*, *S.c.*

834 **Extended Data Fig. 7. Schematic diagram of machine learning framework in predicting**

835 **micafungin susceptibility phenotype.** LASSO: Least Absolute Shrinkage and Selection

836 Operator; ENNS: Ensemble Neural Network Selection; XGBoost: Extreme Gradient Boosting.

Extended Data Fig. 8. Hierarchical clustering of pairwise popANI (population average nucleotide identity) values across all 219 *C. parapsilosis* isolates. Each row or column represents an individual isolate. A gray box indicates a popANI value of at least 99.999% between the respective row and column isolates, while a white box indicates a value below this cutoff. Using the cutoff, we identified 91 clusters of varying sizes (*inset*, red circle; see Methods). For each cluster that contains at least two isolates, any pair of isolates within this cluster has a popANI value of at least 99.999%, indicating that these isolates are all identical to each other. The *inset* curve also shows the impact of the popANI threshold on the number of identified clusters. The detailed pairwise popANI values are available in Supplementary Table 9.

Extended Data Fig. 9. Performance of machine learning models. (a) Precision, recall, and

accuracy scores of machine learning models constructed with different combinations of feature

selection algorithms (LOGIT, LASSO, ENNS) and classifiers (DT, RF, XGBoost). In all

boxplots, each dot represents a single train-test split, which was randomly repeated 50 times. The

models were ranked based on their median scores in the ascending order. LOGIT+XGBoost

always achieved the highest median score. LOGIT, Logistic regression; LASSO, Least Absolute

Shrinkage and Selection Operator; ENNS, Ensemble Neural Network Selection; DT, Decision

Tree; RF, Random Forest; XGBoost, Extreme Gradient Boosting. **(b)** Misclassification (both

false-positive and negative predictions) percentage for all isolates misclassified at least once

among 50 random train-test splits. The number within parenthesis following each isolate

identifier indicates how many times the isolate was included in the test set.

Extended Data Fig. 10. Micafungin heteroresistance in *Candida auris* clinical isolates. *C.*

auris isolate **(a – d)** 386 and **(e – h)** 381 were obtained from CDC/FDA AR Bank. **(a, e)**

Micafungin E-test results for *C. auris* isolates **(a)** 386 and **(e)** 381. **(b, f)** Micafungin population

analysis profiles for *C. auris* isolates **(b)** 386 and **(f)** 381. Percent survival was calculated

compared to growth on drug-free agar. **(c, g)** CFU dynamics of total and resistant subpopulations

of *C. auris* isolates **(c)** 386 and **(g)** 381 under micafungin pressure (0.5 µg/mL). Cultures were

plated at indicated time points for enumeration of total (black, circles) and resistant (colored,

squares) cells. **(d, h)**, Frequency of micafungin-resistant subpopulation of *C. auris* isolates **(d)**

386 and **(h)** 381 during continuous passage. The frequency was calculated after a 24-hour growth

period in YPD broth (Pre-treatment), followed by a 48-hour passage in YPD broth with 0.5

µg/mL micafungin (In micafungin), and subsequently, a passage in YPD broth without

micafungin for 24 hours (Subculture). Bar height: mean; error bars: standard deviation (n=3).

Supplementary information:

Supplementary Table 1. Population analysis profile (PAP) data of all 219 *C. parapsilosis*

isolates in this study.

Supplementary Table 2. Information of patients and *C. parapsilosis* isolates in this study.

Supplementary Table 3. Copy number variations (CNVs) of open reading frames (ORFs)

identified from 219 *C. parapsilosis* isolates.

Supplementary Table 4. ORFs with amplification that overlap with inverted repeats of at least

100bp.

Supplementary Table 5. Single nucleotide variants (SNVs) and insertion/deletions (INDELs)

identified from the 219 *C. parapsilosis* isolates.

Supplementary Table 6. Mapping between raw and merged features. Raw features of single

nucleotide variants (SNVs), insertion/deletions (INDELs), and copy number variations (CNVs)

were merged if they held identical values across all 219 isolates. The identifiers of raw SNV and

INDEL features follow the pattern: "Chromosome ID:Position:SNP (or INDEL):Reference

allele:Alternative allele=X", where X=1, 2 denotes heterozygous and homozygous alternative

alleles respectively. For categorial CNVs, the raw feature identifier follows the pattern

"CNV_cat__[open reading frame identifier]=X", where X = 1, 2, 3, 4 represent full

amplification, partial amplification, full deletion, and partial deletion respectively. The identifiers

of raw quantitative CNV features has the format "CNV_quant__[open reading frame identifier]".

Supplementary Table 7. Logistic regression analysis of the correlation between genomic

features and micafungin heteroresistance. The logistic regression coefficients, along with their

lower and upper confidence intervals (CIs) and adjusted P values ($P_{adj.}$), are provided. Here, the

genomic features refer to the merged features in Supplementary Table 5. Only features with

adjusted P values < 0.05 are included in the table.

Supplementary Table 8. LASSO (Least Absolute Shrinkage and Selection Operator) and

phylogenetic LASSO analyses of the correlation between genomic features and micafungin

heteroresistance. Features with non-zero coefficients in at least one analysis were included in the

table.

Supplementary Table 9. Clustering of *C. parapsilosis* isolates based on popANI (population

average nucleotide identity). In each cluster, all isolates are identical to each other (i.e., all

pairwise popANI values ≥ 99.999%). The column "Used" indicates which isolates have been

included in the development of machine learning models.

Reference:

Antimicrobial Resistance, C. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* **399**, 629-655, doi:10.1016/S0140-6736(21)02724-0 (2022).

Berman, J. & Krysan, D. J. Drug resistance and tolerance in fungi. *Nat Rev Microbiol* **18**, 319-331, doi:10.1038/s41579-019-0322-2 (2020).

Andersson, D. I., Nicoloff, H. & Hjort, K. Mechanisms and clinical relevance of bacterial heteroresistance. *Nat Rev Microbiol* **17**, 479-496, doi:10.1038/s41579-019-0218-1 (2019).

El-Halfawy, O. M. & Valvano, M. A. Antimicrobial heteroresistance: an emerging field in need of clarity. *Clin Microbiol Rev* **28**, 191-207, doi:10.1128/CMR.00058-14 (2015).

Band, V. I. & Weiss, D. S. Heteroresistance: A cause of unexplained antibiotic treatment failure? *PLoS Pathog* **15**, e1007726, doi:10.1371/journal.ppat.1007726 (2019).

Mondon, P. *et al.* Heteroresistance to fluconazole and voriconazole in Cryptococcus neoformans. *Antimicrob Agents Chemother* **43**, 1856-1861, doi:10.1128/AAC.43.8.1856 (1999).

Sionov, E., Chang, Y. C., Garraffo, H. M. & Kwon-Chung, K. J. Heteroresistance to fluconazole in Cryptococcus neoformans is intrinsic and associated with virulence. *Antimicrob Agents Chemother* **53**, 2804-2815, doi:10.1128/AAC.00295-09 (2009).

Ben-Ami, R. *et al.* Heteroresistance to Fluconazole Is a Continuously Distributed Phenotype among Candida glabrata Clinical Strains Associated with In Vivo Persistence. *mBio* **7**, doi:10.1128/mBio.00655-16 (2016).

Band, V. I. *et al.* Antibiotic failure mediated by a resistant subpopulation in Enterobacter cloacae. *Nat Microbiol* **1**, 16053, doi:10.1038/nmicrobiol.2016.53 (2016).

Xiong, Y. *et al.* Impact of Carbapenem Heteroresistance Among Multidrug-Resistant ESBL/AmpC-Producing Klebsiella pneumoniae Clinical Isolates on Antibiotic Treatment in Experimentally Infected Mice. *Infect Drug Resist* **14**, 5639-5650, doi:10.2147/IDR.S340652 (2021).

Bert, F. *et al.* Prevalence, molecular epidemiology, and clinical significance of heterogeneous glycopeptide-intermediate Staphylococcus aureus in liver transplant recipients. *J Clin Microbiol* **41**, 5147-5152, doi:10.1128/JCM.41.11.5147-5152.2003 (2003).

Sieradzki, K., Roberts, R. B., Serur, D., Hargrave, J. & Tomasz, A. Heterogeneously vancomycin-resistant Staphylococcus epidermidis strain causing recurrent peritonitis in a dialysis patient during vancomycin therapy. *J Clin Microbiol* **37**, 39-44, doi:10.1128/JCM.37.1.39-44.1999 (1999).

Fisher, B. T. *et al.* Effect of Caspofungin vs Fluconazole Prophylaxis on Invasive Fungal Disease Among Children and Young Adults With Acute Myeloid Leukemia: A Randomized Clinical Trial. *JAMA* **322**, 1673-1681, doi:10.1001/jama.2019.15702 (2019).

Walsh, T. J. *et al.* Caspofungin versus liposomal amphotericin B for empirical antifungal therapy in patients with persistent fever and neutropenia. *N Engl J Med* **351**, 1391-1402, doi:10.1056/NEJMoa040446 (2004).

Slavin, M. A. *et al.* Efficacy and safety of fluconazole prophylaxis for fungal infections after marrow transplantation--a prospective, randomized, double-blind study. *J Infect Dis* **171**, 1545-1552, doi:10.1093/infdis/171.6.1545 (1995).

16   Goodman, J. L. *et al.* A controlled trial of fluconazole to prevent fungal infections in
patients undergoing bone marrow transplantation. *N Engl J Med* **326**, 845-851,
doi:10.1056/NEJM199203263261301 (1992).
17   Epstein, D. J., Seo, S. K., Brown, J. M. & Papanicolaou, G. A. Echinocandin prophylaxis
in patients undergoing haematopoietic cell transplantation and other treatments for
haematological malignancies. *J Antimicrob Chemother* **73**, i60-i72,
doi:10.1093/jac/dkx450 (2018).
18   Douglas, C. M. *et al.* Identification of the FKS1 gene of Candida albicans as the essential
target of 1,3-beta-D-glucan synthase inhibitors. *Antimicrob Agents Chemother* **41**, 2471-
2479, doi:10.1128/AAC.41.11.2471 (1997).
19   Toth, R. *et al.* Candida parapsilosis: from Genes to the Bedside. *Clin Microbiol Rev* **32**,
doi:10.1128/CMR.00111-18 (2019).
20   Daneshnia, F. *et al.* Worldwide emergence of fluconazole-resistant Candida parapsilosis:
current framework and future research roadmap. *Lancet Microbe* **4**, e470-e480,
doi:10.1016/S2666-5247(23)00067-8 (2023).
21   Trevijano-Contador, N. *et al.* Global Emergence of Resistance to Fluconazole and
Voriconazole in Candida parapsilosis in Tertiary Hospitals in Spain During the COVID-
19 Pandemic. *Open Forum Infect Dis* **9**, ofac605, doi:10.1093/ofid/ofac605 (2022).
22   Thomaz, D. Y. *et al.* A Brazilian Inter-Hospital Candidemia Outbreak Caused by
Fluconazole-Resistant Candida parapsilosis in the COVID-19 Era. *J Fungi (Basel)* **8**,
doi:10.3390/jof8020100 (2022).
23   Fekkar, A. *et al.* Hospital outbreak of fluconazole-resistant Candida parapsilosis:
arguments for clonal transmission and long-term persistence. *Antimicrob Agents*
Chemother **95**, doi:10.1128/AAC.02036-20 (2023).
24   Zhai, B. *et al.* High-resolution mycobiota analysis reveals dynamic intestinal
translocation preceding invasive candidiasis. *Nat Med* **26**, 59-64, doi:10.1038/s41591-
019-0709-7 (2020).
25   Garcia-Effron, G., Katiyar, S. K., Park, S., Edlind, T. D. & Perlin, D. S. A naturally
occurring proline-to-alanine amino acid change in Fks1p in Candida parapsilosis,
Candida orthopsilosis, and Candida metapsilosis accounts for reduced echinocandin
susceptibility. *Antimicrob Agents Chemother* **52**, 2305-2312, doi:10.1128/AAC.00262-08
(2008).
26   Pfaller, M. A., Moet, G. J., Messer, S. A., Jones, R. N. & Castanheira, M. Geographic
variations in species distribution and echinocandin and azole antifungal resistance rates
among Candida bloodstream infection isolates: report from the SENTRY Antimicrobial
Surveillance Program (2008 to 2009). *J Clin Microbiol* **49**, 396-399,
doi:10.1128/JCM.01398-10 (2011).
27   Ning, Y. *et al.* Decreased echinocandin susceptibility in Candida parapsilosis causing
candidemia and emergence of a pan-echinocandin resistant case in China. *Emerg*
Microbes Infect **12**, 2153086, doi:10.1080/22221751.2022.2153086 (2023).
28   Wang, E. *et al.* The ever-evolving landscape of candidaemia in patients with acute
leukaemia: non-susceptibility to caspofungin and multidrug resistance are associated with
increased mortality. *J Antimicrob Chemother* **70**, 2362-2368, doi:10.1093/jac/dkv087
(2015).

29   Lortholary, O. *et al.* Recent exposure to caspofungin or fluconazole influences the
epidemiology of candidemia: a prospective multicenter study involving 2,441 patients.
Antimicrob Agents Chemother **55**, 532-538, doi:10.1128/AAC.01128-10 (2011).
30   Rolling, T. *et al.* Haematopoietic cell transplantation outcomes are linked to intestinal
mycobiota dynamics and an expansion of Candida parapsilosis complex species. *Nat*
Microbiol **6**, 1505-1515, doi:10.1038/s41564-021-00989-7 (2021).
31   Giske, C. G., Turnidge, J., Canton, R., Kahlmeter, G. & Committee, E. S. Update from
the European Committee on Antimicrobial Susceptibility Testing (EUCAST). *J Clin*
Microbiol **60**, e0027621, doi:10.1128/JCM.00276-21 (2022).
32   Lewis, K. Persister cells, dormancy and infectious disease. *Nature Reviews Microbiology*
5, 48-56, doi:10.1038/nrmicro1557 (2007).
33   Olm, M. R. *et al.* inStrain profiles population microdiversity from metagenomic data and
sensitively detects shared microbial strains. *Nat Biotechnol* **39**, 727-736,
doi:10.1038/s41587-020-00797-0 (2021).
34   Sionov, E., Chang, Y. C. & Kwon-Chung, K. J. Azole heteroresistance in Cryptococcus
neoformans: emergence of resistant clones with chromosomal disomy in the mouse brain
during fluconazole treatment. *Antimicrob Agents Chemother* **57**, 5127-5130,
doi:10.1128/AAC.00694-13 (2013).
35   Stone, N. R. *et al.* Dynamic ploidy changes drive fluconazole resistance in human
cryptococcal meningitis. *J Clin Invest* **129**, 999-1014, doi:10.1172/JCI124516 (2019).
36   Yang, F. *et al.* Aneuploidy Underlies Tolerance and Cross-Tolerance to Drugs in Candida
parapsilosis. *Microbiol Spectr* **9**, e0050821, doi:10.1128/Spectrum.00508-21 (2021).
37   Todd, R. T. & Selmecki, A. Expandable and reversible copy number amplification drives
rapid adaptation to antifungal drugs. *Elife* **9**, e58349 (2020).
38   Zhai, B. *et al.* High-resolution mycobiota analysis reveals dynamic intestinal
translocation preceding invasive candidiasis. *Nature medicine* **26**, 59-64 (2020).
39   Lai, M. H., Silverman, S. J., Gaughran, J. P. & Kirsch, D. R. Multiple copies of PBS2,
MHP1 or LRE1 produce glucanase resistance and other cell wall effects in
Saccharomyces cerevisiae. *Yeast* **13**, 199-213, doi:10.1002/(SICI)1097-
0061(19970315)13:3<199::AID-YEA76>3.0.CO;2-Z (1997).
40   Oh, S.-H. *et al.* Agglutinin-like sequence (ALS) genes in the Candida parapsilosis species
complex: blurring the boundaries between gene families that encode cell-wall proteins.
Frontiers in microbiology **10**, 781 (2019).
41   Yang, K. & Maiti, T. ENNS: Variable Selection, Regression, Classification and Deep
Neural Network for High-Dimensional Data. *arXiv preprint arXiv:2107.03430* (2021).
42   Healey, K. R. *et al.* Prevalent mutator genotype identified in fungal pathogen Candida
glabrata promotes multi-drug resistance. *Nat Commun* **7**, 11128,
doi:10.1038/ncomms11128 (2016).
43   Ostrowsky, B. *et al.* Candida auris Isolates Resistant to Three Classes of Antifungal
Medications - New York, 2019. *MMWR Morb Mortal Wkly Rep* **69**, 6-9,
doi:10.15585/mmwr.mm6901a2 (2020).
44   Park, J. Y., Bradley, N., Brooks, S., Burney, S. & Wassner, C. Management of Patients
with Candida auris Fungemia at Community Hospital, Brooklyn, New York, USA, 2016-
2018(1). *Emerg Infect Dis* **25**, 601-602, doi:10.3201/eid2503.180927 (2019).

45   Kilburn, S. *et al.* Antifungal Resistance Trends of Candida auris Clinical Isolates, New
York-New Jersey, 2016-2020. *Antimicrob Agents Chemother*, aac0224221,
doi:10.1128/aac.02242-21 (2022).
46   CDC. *Recommendations for treatment of Candida auris infections*,
<https://www.cdc.gov/fungal/candida-auris/c-auris-treatment.html> (2021).
47   Taur, Y. *et al.* Reconstitution of the gut microbiota of antibiotic-treated patients by
autologous fecal microbiota transplant. *Sci Transl Med* **10**,
doi:10.1126/scitranslmed.aap9489 (2018).
48   Kim, S. G. *et al.* Microbiota-derived lantibiotic restores resistance against vancomycin-
resistant Enterococcus. *Nature* **572**, 665-669, doi:10.1038/s41586-019-1501-z (2019).
49   Butler, G. *et al.* Evolution of pathogenicity and sexual reproduction in eight Candida
genomes. *Nature* **459**, 657-662, doi:10.1038/nature08064 (2009).
50   Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids
Res* **27**, 573-580, doi:10.1093/nar/27.2.573 (1999).
51   Cingolani, P. *et al.* A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster
strain w1118; iso-2; iso-3. *Fly (Austin)* **6**, 80-92, doi:10.4161/fly.19695 (2012).
52   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033
(2010).
53   Lischer, H. E., Excoffier, L. & Heckel, G. Ignoring heterozygous sites biases
phylogenomic estimates of divergence times: implications for the evolutionary history of
microtus voles. *Mol Biol Evol* **31**, 817-831, doi:10.1093/molbev/mst271 (2014).
54   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534,
doi:10.1093/molbev/msaa015 (2020).
55   Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. *Mol Biol Evol* **35**, 518-522,
doi:10.1093/molbev/msx281 (2018).
56   Yan, J. *et al.* Bow-tie signaling in c-di-GMP: Machine learning in a simple biochemical
network. *PLoS Comput Biol* **13**, e1005677, doi:10.1371/journal.pcbi.1005677 (2017).
57   Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of Phylogenetics and Evolution in
R language. *Bioinformatics* **20**, 289-290, doi:10.1093/bioinformatics/btg412 (2004).
58   Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nature
biotechnology* **38**, 685-688 (2020).
59   Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast,
scalable and user-friendly tool for maximum likelihood phylogenetic inference.
Bioinformatics **35**, 4453-4455 (2019).
1087