

One Million and Counting: Estimates of Deaths in the United States from Ancestral SARS-CoV-2 and Variants

Jo Walker^{1,2,*}, Nathan D. Grubaugh^{1,2,3}, Gregg Gonsalves^{1,2}, Virginia Pitzer^{1,2}, Zain Rizvi⁴

¹ Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, Connecticut, USA

² Public Health Modeling Unit, Yale School of Public Health, New Haven, Connecticut, USA

³ Department of Ecology and Evolutionary Biology, Yale University, New Haven, Connecticut, USA

⁴ Public Citizen, Washington D.C., USA

*Corresponding author: can be reached at jo.walker@yale.edu

Abstract

Background: Over one million COVID-19 deaths have been recorded in the United States. Sustained global SARS-CoV-2 transmission has led to the emergence of new variants with increased transmissibility, virulence, and/or immune evasion. The specific burden of mortality from each variant over the course of the U.S. COVID-19 epidemic remains unclear.

Methods: We constructed an epidemiologic model using data reported by the CDC on COVID-19 mortality and circulating variant proportions to estimate the number of recorded COVID-19 deaths attributable to each SARS-CoV-2 variant in the U.S. We conducted sensitivity analysis to account for parameter uncertainty.

Findings: Of the 1,003,419 COVID-19 deaths recorded as of May 12, 2022, we estimate that 460,124 (46%) were attributable to WHO-designated variants. By U.S. Census Region, the South recorded the most variant deaths per capita (median estimate 158 per 100,000), while the Northeast recorded the fewest (111 per 100,000). Over 40 percent of national COVID-19 deaths were estimated to be caused by the combination of Alpha (median estimate 39,548 deaths), Delta (273,801), and Omicron (117,560).

Interpretation: SARS-CoV-2 variants that have emerged around the world have imposed a significant mortality burden in the U.S. In addition to national public health strategies, greater efforts are needed to lower the risk of new variants emerging, including through global COVID-19 vaccination, treatment, and outbreak mitigation.

Word Count

3660 (excluding abstract, tables, figures, references, acknowledgements, etc)

Introduction

Over one million deaths have been recorded in the United States from coronavirus disease 2019 (COVID-19). Since severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in Wuhan, China in late 2019, the virus has continued to evolve, with multiple new variants characterized by increased transmissibility, virulence, and/or immune evasion.

Variants reported around the world have repeatedly changed the trajectory of the U.S. epidemic. As of May 12, 2022, the World Health Organization (WHO) has named five variants of concern (VOC) that were originally detected in four different continents [**Table 1**]. All five variants of concern were more transmissible than the original virus, and spread to the U.S.¹⁻³ Delta (lineage B.1.617.2) was particularly severe from a clinical standpoint, and drove a surge of hospitalizations across the U.S in the summer and fall of 2021.⁴ While Omicron (lineage B.1.1.529) appears to cause less severe disease than Delta on a per-infection basis—likely due to a combination of lower intrinsic severity and the high observed concentration of cases among the previously vaccinated or infected⁵—its elevated transmissibility, including among the seropositive, resulted in the largest wave of SARS-CoV-2 transmission in the U.S. to date.^{6,7} Vaccines developed against the non-variant SARS-CoV-2 are less effective at protecting against infection with Delta and Omicron, but remain highly effective against severe disease.⁸⁻¹⁰

Table 1. SARS-CoV-2 variants of concern designated by the World Health Organization.^{11,12}

Pango Lineage	WHO Designation	First Detection Location	Earliest Documented Samples	Earliest U.S. Report	Downgrade in Status
B.1.1.7	Alpha	United Kingdom	Sept. 2020	Dec. 2020	Previous variant of concern: Mar. 2022
B.1.351	Beta	South Africa	May 2020	Jan. 2021	Previous variant of concern: Mar. 2022
P.1	Gamma	Brazil	Nov. 2020	Jan. 2021	Previous variant of concern: Mar. 2022
B.1.617.2	Delta	India	Oct. 2020	Mar. 2021	Not downgraded as of May 2022
B.1.1.529	Omicron	Botswana and South Africa	Nov. 2021	Dec. 2021	Not downgraded as of May 2022

In addition to variants of concern, the WHO and U.S. Centers for Disease Control and Prevention (CDC) have classified a number of additional SARS-CoV-2 lineages as variants of interest (VOI) or under monitoring (VUM).¹³ The variants have emerged in the context of sustained global transmission. More than 500 million COVID-19 cases have been reported globally, reflecting only a fraction of total infections, and giving SARS-CoV-2 many opportunities to mutate and evolve.¹⁴ One third of the global population has still not received a vaccine dose, limiting the benefits of vaccination on reducing transmission and viral replication at a population level.¹⁵

Recent studies have described clinical outcomes associated with SARS-CoV-2 variants, and analyzed factors associated with COVID-19 mortality in the U.S. including race and geography.^{16,17} The aim of our study is to estimate the burden of deaths associated with each SARS-CoV-2 variant in the U.S. We construct a model using data reported by the CDC on COVID-19 deaths by jurisdiction and circulating variant proportions.

Methods

To estimate the number of recorded COVID-19 deaths attributable to each variant, we back-distributed deaths to estimate the number of ultimately fatal COVID-19 cases testing positive on each day in each jurisdiction, and compared these counts to proportions of variants among sequenced cases in the same time period and location, adjusted for differences in disease severity between variants.

Data

Our analysis relies on provisional counts of COVID-19 deaths from the CDC's National Center for Health Statistics (NCHS).¹⁸ This dataset is based on death certificate records and reports the number of deaths with confirmed or presumed COVID-19 by jurisdiction (the 50 states, New York City, District of Columbia, and Puerto Rico) and time period (week, month, year, and total) of occurrence. All data were accessed May 12, 2022.

The NCHS data are subject to change as new death certificate records are received and processed by the CDC. While the majority of COVID-19 deaths are reported within 10 days of the date of death, state reporting timelines vary significantly.¹⁹ Counts from recent weeks may be incomplete. However, because adjustments are almost always upwards, the NCHS provisional COVID-19 deaths can be considered a lower bound of the number of recorded COVID-19 deaths that have occurred.

NCHS death counts are also displayed by the week of occurrence, rather than the day of occurrence. Since the delay distributions that we applied to match deaths to the appropriate variant proportions (see below) require analysis at the level of days, we imputed daily deaths within each week by sampling from a uniform distribution. Finally, for each jurisdiction, the NCHS suppresses death counts between 1 and 9 for privacy reasons. As a result, the specific week is not known for 3,628 of the 1,003,419 COVID-19 deaths recorded in this dataset (0.4% of deaths overall, 0.002% to 39.4% of deaths in each jurisdiction). In this analysis, we do not assign these deaths to a week or otherwise attribute them to individual variants, meaning that our variant-specific death estimates should be treated as lower bounds.

Variant proportions were retrieved either directly²⁰ from the CDC, or based on code and data made publicly available²¹ by the CDC. We considered 26 mutually-exclusive categories that deaths could be assigned to: the 5 current and former variants of concern (Alpha, Beta, Gamma, Delta, and Omicron), 19 other variants and lineages which have been recognized by the WHO and detected in the U.S. (A.23.1, A.27, A.28, B.1.1.318, B.1.1.519, B.1.214.2, B.1.617.3, B.1.620, B.1.640, C.16, C.36, Epsilon, Eta, Iota, Kappa, Lambda, Mu, Theta, or Zeta), a composite group of the early non-VOI/VOC/VBM virus lineages, which we will refer to as “non-variant”, and “unknown” (deaths for which the week of occurrence is suppressed).⁹

CDC genomic surveillance data was available from January 3rd, 2021. For the purpose of this analysis, we assumed that no variants were present in the U.S. prior to this date. In reality, there is evidence for circulation of the Alpha^{2,22,23} Epsilon²⁴, and Iota²² variants in the U.S. in late 2020, albeit at low frequencies: thus, estimates of the mortality burden for these variants should be considered conservative. For the period between January 3rd, 2021, and January 1st, 2022, we used publicly available¹⁹ code and data underpinning the CDC’s variant proportion analysis to generate weighted variant proportions for each jurisdiction and U.S. Health and Human Services (HHS) region in non-overlapping weekly intervals. If no data on sequenced cases was available for a week in any given jurisdiction, we substituted variant proportions from the wider HHS region.

For the period after January 1st, 2022, the code and data described above was not publicly available. Instead, we used variant proportions retrieved directly from the CDC for national and HHS-regional levels in non-overlapping two-week intervals.¹⁸ We assumed that all states in a region had the same variant distribution during any given 2-week period. This simplifying assumption likely only had a marginal impact on our results, since the overwhelming majority of cases sampled in this period were attributed to Omicron.

Analysis

Given the delays between symptom onset, testing, and death, we applied lags to infer the likely time of sample collection for new deaths. For our primary analysis, we assumed that sample collection for testing always took place either 0, 1, 2, or 3 days after symptom onset, with an associated probability of 25% for each of these 4 possible lag times. This is a simplifying assumption on our part. We also assumed the lag from symptom onset to death, $lag_{onset\ to\ death}$, follows a lognormal distribution with a mean and standard deviation of 20.2 and 11.6 days, respectively (mean and standard deviation of the $\log(lag) = 2.863$ and 0.534 respectively), truncated at 50 days. This corresponds to the best-fitting distribution for right-censored onset-to-death observations in Linton et al.²⁵

Before applying variant proportions to COVID-19 deaths, we adjusted variant proportions among cases for variant-specific differences in disease severity. For instance, during periods where Omicron and Delta co-circulated, we expect Delta to be overrepresented among deaths relative to its share of cases (most of which will not be fatal), since Delta infections are more likely to result in severe disease than Omicron infections (whether due to higher intrinsic severity, a greater concentration of cases in unvaccinated individuals, or a combination).⁵

We derived a variant i 's proportion of fatal cases, $p_{i, fatal}$ from its share of all cases $p_{i, all}$ and the relative risk of death RR_i among confirmed cases infected with variant i compared to non-variant virus in a given time period t as

$$p_{i, fatal, t} = \frac{\text{fatal cases caused by variant } i \text{ in period } t}{\text{fatal cases caused by all variants in period } t} = \frac{\text{Cases}_{all\ variants, t} * p_{i, all, t} * CFR_{non-variant} * RR_i}{\sum_{Variant = v}^{26} \text{Cases}_{all\ variants, t} * p_{v, all, t} * CFR_{non-variant} * RR_v} = \frac{p_{i, all, t} * RR_i}{\sum_{Variant = v}^{26} p_{v, all, t} * RR_v}$$

To parameterize RR_i , we derived data on the relative severity of variants from a European Centre for Disease Prevention and Control (ECDC) review which was last updated in early April 2022.²⁶ This review found no evidence for differences in severity (relative to the dominant variant(s) at the time of emergence) for the B.1.617.3, Epsilon, Eta, Iota, Kappa, Lambda, Mu, Theta, or Zeta variants. Evidence of differential severity was reported for the Alpha, Beta, Gamma, Delta, and Omicron variants [Table 2].

Table 2. Estimates of the relative risk of death for different SARS-CoV-2 Variants of Concern

SARS-CoV-2 Variant	Study	Effect Estimate	Notes
Alpha	Lin et al. ²⁷	1.41 (95% CI: 1.34 - 1.49) hazard ratio of death vs non-variant SARS-CoV-2 with fixed effects, 1.37 (95% CI: 1.15 - 1.60) with random effects	Meta-analysis of 12 studies
Beta	Lin et al. ²⁷	1.50 (95% CI: 1.26 - 1.74) hazard ratio of death vs non-variant SARS-CoV-2 with either fixed or random effects	Meta-analysis of 2 studies
Gamma	Lin et al. ²⁷	1.22 (95% CI: 1.02 - 1.42) hazard ratio of death vs non-variant SARS-CoV-2 with fixed effects, 1.06 (95% CI: 0.17 - 1.96) with random effects	Meta-analysis of 2 studies
Delta	Fisman et al. ²⁸	2.33 (95% CI: 1.54 - 3.31) adjusted odds ratio of death vs non-variant SARS-CoV-2	Retrospective cohort of 212,326 COVID-19 cases in Ontario between January and June 2021
Omicron	Lewnard et al. ²⁹	0.21 (95% CI: 0.10 - 0.42) hazard ratio of death for Omicron vs Delta cases	Prospective cohort comparing 42,515 Delta cases and 228,594 Omicron cases in Southern California
	Ulloa et al. ³⁰	0.12 (95% CI: 0.04 - 0.37) hazard ratio of death for Omicron vs Delta cases	Retrospective cohort. In this study, 0.03% (12 of 37,296) of Omicron cases died, compared to 0.5% (133 of 24,432) of Delta cases. Delta and

			Omicron cases were similar in terms of age (median = 33 and 30 years respectively) and sex (49.3% vs 50.1% female, respectively), but not vaccination status: 87.2% of Omicron cases had received at least one vaccine dose, compared to only 55.4% of Delta cases.
--	--	--	---

Based on the estimates above, we assumed that RR_i (the relative risk of death among confirmed cases caused by variant i vs non-variant in any given period of time), is 1.4 for Alpha, 1.5 for Beta, 1.2 for Gamma, 2.3 for Delta, 0.5 for Omicron ($\sim 0.21 * RR_{Delta}$), and 1 for the non-variant and all other variants in the primary analysis.

As a sensitivity analysis, we varied the assumed relative severity of Omicron from $RR_{Omicron} = 0.15$, the approximate ratio of observed Omicron and Delta case fatality rates in Ulloa et al. ($\frac{12/37296}{133/24432} = 0.059$) times $RR_{Delta} = 2.3$, to $RR_{Omicron} = 1$, which corresponds approximately to the upper 95% bound of the Omicron vs Delta hazard ratio of death estimated by Lewnard et al., 0.42, times $RR_{Delta} = 2.3$.^{29,30} The results from these sensitivity analyses provide upper and lower limits for the plausible number of recorded COVID-19 deaths caused by the Omicron variant.

Finally, after deriving appropriate severity-adjusted variant proportion values $p_{i, fatal}$ for each jurisdiction and time period, we estimated the number of recorded COVID-19 deaths caused by each variant i as

$$TotalDeaths_{Variant=i} = \sum_{t=1}^N FatalCases(sample\ collection\ day = t) * p_{i, fatal, Biweek(t)}$$

for each jurisdiction, where $Biweek(t)$ species the non-overlapping two-week period containing day t . We reported national, regional (based on the four U.S. Census Regions.³¹ For this analysis, we included Puerto Rico with states in the *South* region), and jurisdiction-specific estimates. Median estimates and 95% credible intervals were derived by running the model described above 250 times. Within each model run, we performed bootstrap resampling of the variant proportions at each location and time period before adjusting for severity, and resampled the day of death within the NCHS-specified week, the onset-to-death delay, and the onset-to-testing delay for each death in the dataset.

Results

National estimates

Of the 1,003,419 (299 per 100,000 people) recorded COVID-19 deaths in our dataset at the national level (the 50 states, the District of Columbia, and Puerto Rico), our median estimate is that 539,667 (95% credible interval: 538,928 - 540,466 deaths, 54% of all COVID-19 deaths, 161 deaths per 100,000) were caused by non-variant SARS-CoV-2, while 460,124 (95% CI: 459,325 - 460,863 deaths, 46% of deaths, 137 per 100,000) were caused by a WHO-classified variant [Table 3]. Of the variants that have had the greatest impact on the COVID-19 epidemic in the United States, we estimate that the Alpha variant caused a median of 39,548 recorded deaths (95% CI: 38,890 - 40,133 deaths, 4% of deaths, 12 per 100,000) across model runs, the Delta variant caused a median of 273,801 recorded deaths (95% CI: 273,483 - 274,168 deaths, 27% of deaths, 82 per 100,000), and the Omicron variant has so far caused a median of 117,560 recorded deaths (95% CI: 117,252 - 117,835 deaths, 12% of deaths, 35 per 100,000).

In sensitivity analyses where we increased or decreased the assumed severity of Omicron relative to other variants, estimates of recorded deaths caused by Delta ranged from medians of 265,228 deaths (95% CI: 264,898 - 265,600 deaths, 26% of deaths, 79 per 100,000) to 291,946 (95% CI: 291,623 - 292,319 deaths, 29% of deaths, 87 per 100,000) across model runs, while median estimates of recorded deaths caused by Omicron ranged from 99,023 (95% CI: 98,727 - 99,352 deaths, 10% of deaths, 30 per 100,000) to 126,327 (95% CI: 126,022 - 126,582 deaths, 13% of deaths, 38 per 100,000).

Of the COVID-19 deaths recorded by NCHS, 3,628 recorded deaths (0.3%, 1 per 100,000) could not be attributed to a variant because the week of death was not given at the state-level in the dataset. National estimates of the burden of recorded COVID-19 deaths attributable to each variant are given in Table 3.

Regional estimates

The Northeast, South, and Midwest recorded the most COVID-19 deaths per capita (328, 317, and 306 per 100,000 respectively), while the West recorded the fewest (244 per 100,000) [Figure 1A]. However, our estimates indicate the Northeast experienced the fewest COVID-19 *variant* deaths per capita (median 111 per 100,000, 95% credible interval: 111 - 112). As a result, variants accounted for 34% of all COVID-19 deaths in the Northeast, compared to nearly half of deaths in each of the other three census regions (Midwest: 46%, South: 50%, West: 49%) [Tables 4-5].

Non-Variant: The median estimated burden of recorded deaths caused by non-variant SARS-CoV-2 was highest in the Northeast with 215 COVID-19 deaths per 100,000 (123,103 deaths, 66% of all COVID-19 deaths in this region), and lowest in the West with 124 deaths per 100,000 (97,549 deaths, 51% of all COVID-19 deaths in this region) [Figure 1B, Tables 4-5]. New York City experienced the highest median incidence of deaths from non-variant

SARS-CoV-2 (315 deaths per 100,000, or 26,665 deaths overall, 76% of all COVID-19 deaths in NYC) [**Supplementary Tables 1-2**].

Alpha: After its emergence in the winter of 2020/21, the median estimated burden of deaths caused by the Alpha variant was highest in the South with 15 recorded COVID-19 deaths per 100,000 (19,143 deaths, 5% of all COVID-19 deaths in this region) and lowest in the West with 7 recorded COVID-19 deaths per 100,000 (5,339 deaths overall, 3% of all COVID-19 deaths in this region) [**Figure 1D, Tables 4-5**]. Michigan experienced the highest median incidence of Alpha deaths across all jurisdictions (31 deaths per 100,000, or 3,095 deaths overall, 10% of all COVID-19 deaths in Michigan) [**Supplementary Tables 1-2**].

Delta: The median estimated burden of deaths caused by the Delta variant was highest in the South with 97 recorded COVID-19 deaths per 100,000 (125,925 deaths, 30% of all COVID-19 deaths in this region), followed by the Midwest with 92 recorded COVID-19 deaths per 100,000 (63,049 deaths, 30% of all COVID-19 deaths in this region). The Northeast experienced the fewest deaths per capita with 57 recorded COVID-19 deaths per 100,000 (32,414 deaths, 17% of all COVID-19 deaths in this region) [**Figure 1E, Tables 4-5**]. West Virginia experienced the highest median incidence of Delta deaths (154 per 100,000, or 2,740 deaths overall, 40% of all COVID-19 deaths in West Virginia) across all jurisdictions [**Supplementary Tables 1-2**].

Omicron: So far, the median estimated burden of deaths caused by the Omicron variant has been highest in the South with 41 recorded COVID-19 deaths per 100,000 (52,912 deaths, 13% of all COVID-19 deaths in this region) and lowest in the West with 29 recorded COVID-19 deaths per 100,000 (22,738 deaths, 12% of all COVID-19 deaths in this region, respectively) [**Figure 1F, Tables 4-5**]. Mississippi has experienced the highest incidence of Omicron deaths to date (60 per 100,000, or 1,773 deaths overall, 13% of all COVID-19 deaths in Mississippi) across all jurisdictions [**Supplementary Tables 1-2**].

The specific number and share of Omicron and Delta COVID-19 deaths varied in sensitivity analyses where we assumed Omicron was more or less severe (compared to our primary analysis), but the overall regional patterns described above persisted [**Supplementary Tables 3-4**].

Other: The median estimated incidence of deaths from other variants (not Alpha, Delta, or Omicron) was higher in the West (16 COVID-19 deaths per 100,000) and Northeast (10 COVID-19 deaths per 100,000) than the Midwest and the South (5 and 6 recorded COVID-19 deaths per 100,000, respectively) [**Figure 1G, Tables 4-5**]. In addition to Alpha, the Epsilon and Iota variants accounted for a significant share of cases in the West and Northeast prior to the emergence of Delta.

Discussion

Sustained global transmission of SARS-CoV-2 has fueled the emergence of variants that have caused ~460,000 deaths in the U.S. Alpha, Delta and Omicron—which were first detected in Europe, Asia, and Africa—have led to ~430,000 deaths so far. The other variants, including those that were first detected in the U.S., accounted for an additional ~30,000 deaths.

More than 40 percent of national COVID-19 deaths were caused by SARS-CoV-2 variants first detected outside the United States. The phrase “no one is safe until everyone is safe” has been repeatedly invoked as a clarion call during the pandemic. Our analysis provides support for this claim by documenting the significant impact of SARS-CoV-2 variants, which have emerged in the context of uncontrolled transmission both locally and around the world, on U.S. mortality.

Our analysis also suggests the potential benefit of rapidly implementing strategies to reduce the impact of novel variants after the onset of a viral epidemic. We estimate that Delta and Omicron together have caused over 390,000 deaths in the U.S., despite the fact both variants were designated as variants of concern after multiple COVID-19 vaccines had received emergency authorization. Although causality cannot be established definitively, inequalities in global COVID-19 vaccine production and distribution, along with uncontrolled spread of the virus, may have contributed to viral evolution and circulation. For example, Omicron was detected in November 2021 by scientists in Botswana and South Africa, where less than a quarter of the population was fully vaccinated.³²

Robust global vaccination and treatment campaigns may continue to play an important role in reducing viral replication. A leading hypothesis holds that variants of concern have emerged from immunocompromised individuals with persistent infections that provide more opportunity for viral selection and evolution.³³⁻³⁵ While vaccine effectiveness is reduced against Delta and especially Omicron, booster doses provide modest, albeit waning, protection against infection and appear to reduce transmission risk by lowering infectious viral load.³⁶⁻³⁹ When breakthrough cases do occur, vaccinated persons also clear infections faster than unvaccinated persons.⁴⁰ Second-generation vaccines in development could yield significant benefit if they offer greater protection against transmission. In addition, monoclonal antibodies have received emergency authorization for preexposure prophylaxis, and emerging evidence suggests oral antiviral treatments lower SARS-CoV-2 viral load, and clear infections faster.⁴¹⁻⁴³

Finally, our analysis reveals the impact of the variants across the country. All regions faced a substantial burden of deaths from the variants. The Northeast recorded the lowest, but still substantial, number of variant deaths, both overall (63,542) and per-capita (111 per 100,000). Some regions were disproportionately impacted, likely reflecting differences in vaccination coverage, prior immunity⁴⁴, use of non-pharmaceutical interventions, demographics and social vulnerability. For example, the South recorded the most variant deaths per capita (median

estimate 158 per 100,000) and overall (205,729). It also experienced the highest deaths per capita in each of the Alpha, Delta, and Omicron waves (15, 97 and 41 per 100,000 respectively). Of the 18 states and territories included in the region, 10 had fully vaccinated less than 60% of their population as of early May 2022, according to the CDC.⁴⁵

Our analysis is limited by the quality of public data. NCHS data are considered provisional because of the lag between the date of death and the processing of death certificates by the CDC. Nonetheless, a potential lag would lead to underestimates of deaths caused by variants (due to undercounting of deaths in more recent weeks when Omicron is dominant). In general, reported deaths represent an undercount of total COVID-19 deaths, given limitations in testing and data.⁴⁶ In addition, there is considerable uncertainty associated with some state variant proportions due to insufficient sequencing. Our model also uses variant proportion figures based on clinical testing which may lag behind actual variant circulation. For example, wastewater surveillance suggests that Omicron was circulating in the country before the first case was reported.⁴⁷ This would also lead to underestimates of deaths caused by variants. Finally, we do not model disease transmission dynamics so our estimates should not be interpreted as the number of deaths that would have been prevented if novel variants did not emerge. Even in the absence of new variants, the non-variant SARS-CoV-2 would have continued to spread, resulting in additional deaths.

SARS-CoV-2 variants detected around the world have imposed a significant mortality burden in the U.S. In addition to national public health strategies, greater efforts are needed to reduce the risk of new variants emerging globally. However, U.S. government officials have warned that they are running out of COVID-19 funds.⁴⁸ Congressional lawmakers recently excluded \$5 billion in funding for global treatment and vaccination campaigns from a new spending proposal, and there has been limited investment in next-generation vaccines that may offer greater protection against transmission.^{49,50} Our analysis shows the significant impact of SARS-CoV-2 variants on U.S. public health. In doing so, it highlights the potential risk posed to Americans by new variants—a risk that is only amplified with low rates of global COVID-19 vaccination, and low availability of COVID-19 diagnostics, treatments, and prophylaxis.

Acknowledgements

The authors are grateful to Zena Ahmed for research assistance, to Dr. Daniel Weinberger for providing constructive feedback on the analysis, and to Drs. Prabasaj Paul and Molly Steele for assistance in identifying relevant data sources.

Author Contributions

Concept and design: JW, ZR

Statistical analysis: JW

Interpretation of data and findings: JW, NG, GG, VP, ZR

Drafting and revision of the manuscript: JW, NG, GG, VP, ZR

Conflicts of Interest

N.D.G. is a consultant for Tempus Labs and the National Basketball Association for work related to COVID-19 but is outside the submitted work. V.P. discloses reimbursement from Merck and Pfizer for travel to Scientific Input Engagements unrelated to COVID-19 and membership on the WHO Immunization and Vaccine-related Implementation Research Advisory Committee (IVIR-AC).

Data and Code Availability

Datasets and R code needed to replicate this analysis are available online at https://github.com/joewalker127/US_Deaths_COVID_Variants

References

- 1: Campbell F, Archer B, Laurenson-Schafer H, et al. Increased transmissibility and global spread of SARS-CoV-2 variants of concern as at June 2021. *Eurosurveillance*. 2021;26(24). doi:10.2807/1560-7917.es.2021.26.24.2100509
- 2: Alpert T et al. Early introductions and transmission of SARS-CoV-2 variant B.1.1.7 in the United States. *Cell*. 2021;184(10). doi: 10.1016/j.cell.2021.03.061
- 3: Earnest R et al. Comparative transmissibility of SARS-CoV-2 variants Delta and Alpha in New England, USA. *Cell Reports Medicine*. 2022;3(4). doi: 10.1016/j.xcrm.2022.100583
- 4: Taylor C., Whitaker M., Anglin O. et al. COVID-19–Associated Hospitalizations Among Adults During SARS-CoV-2 Delta and Omicron Variant Predominance, by Race/Ethnicity and Vaccination Status — COVID-NET, 14 States, July 2021–January 2022. *Morbidity and Mortality Weekly Report*. 2022;71(12):466-473. doi: 10.15585/mmwr.mm7112e2
- 5: Bhattacharyya R, Hanage W. Challenges in Inferring Intrinsic Severity of the SARS-CoV-2 Omicron Variant. *New England Journal of Medicine*. 2022;386:e14. doi: 10.1056/NEJMp2119682
- 6: Talent A, Hodcroft EB, Robertson DL. The Evolution and Biology of SARS-CoV-2 Variants. *Cold Spring Harbor Perspectives in Medicine*. 2022;12(3). doi: 10.1101/cshperspect.a041390
- 7: Clarke K et al. Seroprevalence of Infection-Induced SARS-CoV-2 Antibodies — United States, September 2021–February 2022. *Morbidity and Mortality Weekly Report*. 2022;71(17):606-608. doi: 10.15585/mmwr.mm7117e3
- 8: Andrews N, Stowe J, Kirsebom F, et al. Covid-19 vaccine effectiveness against the Omicron (b.1.1.529) variant. *New England Journal of Medicine*. 2022. doi: 10.1056/nejmoa2119451
- 9: Collie S, Moultrie H, Becker LG, Gray G. Effectiveness of BNT162b2 Vaccine against Omicron Variant in South Africa. *New England Journal of Medicine*. 2021. doi: 10.1056/NEJMc2119270


- 10: Sheikh A, McMenamin J, Taylor B, Robertson C. SARS-CoV-2 Delta VOC in Scotland: demographics, risk of hospital admission, and vaccine effectiveness. *Lancet*. 2021; 397(10293). Doi: 10.1016/S0140-6736(21)01358-1
- 11: Tracking SARS-CoV-2 variants. World Health Organization. Published April 12, 2022. <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
- 12: O'Toole Á, Scher E, Jackson B, et al. Pango Lineages: Latest epidemiological lineages of SARS-CoV-2. Cov-lineages. <https://cov-lineages.org/>
- 13: Centers for Disease Control and Prevention. SARS-CoV-2 Variant Classifications and Definitions. Published April 26, 2022. <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-classifications.html>
- 14: WHO Coronavirus (COVID-19) dashboard. World Health Organization. Published April 20, 2022. <https://covid19.who.int/>
- 15: Ritchie H, Mathieu E, Rodés-Guirao L, et al. Coronavirus (COVID-19) vaccinations. Our World in Data. Published March 5, 2020. <https://ourworldindata.org/covid-vaccinations>
- 16: Bialek S, Bowen V, Chow N, et al. Geographic differences in covid-19 cases, deaths, and incidence — United States, February 12–April 7, 2020. *Morbidity and Mortality Weekly Report*. 2020;69(15):465-471. doi: 10.15585/mmwr.mm6915e4
- 17: Bassett MT, Chen JT, Krieger N. Variation in racial/ethnic disparities in covid-19 mortality by age in the United States: A cross-sectional study. *PLOS Medicine*. 2020;17(10). doi: 10.1371/journal.pmed.1003402
- 18: Centers for Disease Control and Prevention. Provisional Death Counts for Coronavirus Disease 2019 (COVID-19). Centers for Disease Control and Prevention: National Center for Health Statistics. Published May 16, 2022. <https://www.cdc.gov/nchs/nvss/vsrr/covid19/index.htm>
- 19: Centers for Disease Control and Prevention. Technical notes: Provisional death counts for Coronavirus Disease (COVID-19). Centers for Disease Control and Prevention: National Center for Health Statistics. Published April 29, 2022. https://www.cdc.gov/nchs/nvss/vsrr/covid19/tech_notes.htm
- 20: Centers for Disease Control and Prevention. COVID Data Tracker: Variant Proportions. Published March 28, 2020. <https://covid.cdc.gov/covid-data-tracker/#variant-proportions>
- 21: Centers for Disease Control and Prevention. Genomic Surveillance of SARS-CoV-2 Circulating in the United States. https://github.com/CDCgov/SARS-CoV-2_Genomic_Surveillance
- 22: Petrone M et al. Combining genomic and epidemiological data to compare the transmissibility of SARS-CoV-2 variants Alpha and Iota. *Communications Biology*. 2022;5. doi: 10.1038/s42003-022-03347-3
- 23: Washington N et al. Emergence and rapid transmission of SARS-CoV-2 B.1.1.7 in the United States. *Cell*. 2021;184(10). doi: 10.1016/j.cell.2021.03.052
- 24: Deng X et al. Transmission, infectivity, and neutralization of a spike L452R SARS-CoV-2 variant. *Cell*. 2021;184(13). doi: 10.1016/j.cell.2021.04.025

- 25: Linton N, Kobayashi T, Yang Y, et al. Incubation period and other epidemiological characteristics of 2019 novel coronavirus infections with right truncation: A statistical analysis of publicly available Case Data. *Journal of Clinical Medicine*. 2020;9(2):538. doi: 10.3390/jcm9020538
- 26: SARS-COV-2 variants of concern as of 21 April 2022. European Centre for Disease Prevention and Control. Published April 22, 2022. <https://www.ecdc.europa.eu/en/covid-19/variants-concern>
- 27: Lin L, Liu Y, Tang X, He D. The disease severity and clinical outcomes of the SARS-COV-2 variants of concern. *Frontiers in Public Health*. 2021;9. doi: 10.3389/fpubh.2021.775224
- 28: Fisman DN, Tuite AR. Evaluation of the relative virulence of novel SARS-COV-2 variants: A retrospective cohort study in Ontario, Canada. *Canadian Medical Association Journal*. 2021;193(42). doi: 10.1503/cmaj.211248
- 29: Lewnard JA, Hong VX, Patel MM, Kahn R, Lipsitch M, Tartof SY. Clinical outcomes among patients infected with Omicron (B.1.1.529) SARS-COV-2 variant in Southern California. 2022. doi: 10.1101/2022.01.11.22269045
- 30: Ulloa AC, Buchan SA, Daneman N, Brown KA. Estimates of SARS-COV-2 omicron variant severity in Ontario, Canada. *JAMA*. 2022;327(13):1286. doi: 10.1001/jama.2022.2274
- 31: U.S. Census Bureau. Census regions and divisions of the United States. https://www2.census.gov/geo/pdfs/maps-data/maps/reference/us_regdiv.pdf
- 32: Speech made by CEPI CEO, Dr. Richard Hatchett, on the emergence of Omicron variant at today's World Health Assembly Special Session. CEPI. Published November 29, 2021. https://cepi.net/news_cepi/speech-made-by-cepi-ceo-dr-richard-hatchett-on-the-emergence-of-omicron-variant-at-todays-world-health-assembly-special-session/
- 33: Corey L, Beyrer C, Cohen MS, Michael NL, Bedford T, Rolland M. SARS-COV-2 variants in patients with immunosuppression. *New England Journal of Medicine*. 2021;385(6):562-566. doi:10.1056/nejmsb2104756
- 34: Burki T. The origin of SARS-COV-2 variants of concern. *The Lancet Infectious Diseases*. 2022;22(2):174-175. doi: 10.1016/s1473-3099(22)00015-9
- 35: Otto SP, Day T, Arino J, et al. The origins and potential future of SARS-COV-2 variants of concern in the evolving COVID-19 pandemic. *Current Biology*. 2021;31(14). doi:10.1016/j.cub.2021.06.049
- 36: Accorsi EK, Britton A, Fleming-Dutra KE, et al. Association between 3 doses of mRNA COVID-19 vaccine and symptomatic infection caused by the SARS-COV-2 omicron and Delta variants. *JAMA*. 2022;327(7):639. doi: 10.1001/jama.2022.0470
- 37: Puhach O, Adea K, Hulo N, et al. Infectious viral load in unvaccinated and vaccinated individuals infected with ancestral, delta or Omicron SARS-CoV-2. *Nature Medicine*. 2022. doi: 10.1038/s41591-022-01816-0
- 38: Lyngse FP, Mortensen LH, Denwood MJ, et al. SARS-COV-2 omicron VOC transmission in Danish households. *MedRxiv*. 2021. doi: 10.1101/2021.12.27.21268278

- 39: Prunas O, Warren JL, Crawford FW, et al. Vaccination with BNT162B2 reduces transmission of SARS-COV-2 to household contacts in Israel. *Science*. 2022;375(6585):1151-1154. doi: 10.1126/science.abl4292
- 40: Kissler SM, Fauver JR, Mack C, et al. Viral dynamics of SARS-COV-2 variants in vaccinated and unvaccinated persons. *New England Journal of Medicine*. 2021;385(26):2489-2491. doi: 10.1056/nejmc2102507
- 41: Levin MJ, Ustianowski A, De Wit S, et al. Intramuscular AZD7442 (Tixagevimab–CILGAVIMAB) for prevention of covid-19. *New England Journal of Medicine*. 2022. doi: 10.1056/nejmoa2116620
- 42: Fischer WA, Eron JJ, Holman W, et al. A phase 2a clinical trial of molnupiravir in patients with COVID-19 shows accelerated SARS-COV-2 RNA clearance and elimination of infectious virus. *Science Translational Medicine*. 2021;14(628). doi: 10.1126/scitranslmed.abl7430
- 43: Hammond J, Leister-Tebbe H, Gardner A, et al. Oral Nirmatrelvir for high-risk, nonhospitalized adults with covid-19. *New England Journal of Medicine*. 2022;386(15):1397-1408. doi: 10.1056/nejmoa2118542
- 44: Klassen F et al. Population immunity to pre-Omicron and Omicron SARS-CoV-2 variants in US states and counties through December 1, 2021. *MedRxiv*. 2022. doi: 10.1101/2021.12.23.21268272
- 45: Centers for Disease Control and Prevention. Covid Data tracker: COVID-19 Vaccinations in the United States. Published March 28, 2020.
https://covid.cdc.gov/covid-data-tracker/#vaccinations_vacc-total-admin-rate-total
- 46: Iuliano AD, Chang HH, Patel NN, et al. Estimating under-recognized COVID-19 deaths, United States, March 2020-May 2021 using an excess mortality modeling approach. *The Lancet Regional Health - Americas*. 2021;1:100019. doi: 10.1016/j.lana.2021.100019
- 47: Kirby AE, Welsh RM, Marsh ZA, et al.; New York City Department of Environmental Protection. Notes from the field: early evidence of the SARS-CoV-2 B.1.1.529 (Omicron) variant in community wastewater—United States, November–December 2021. *MMWR Morb Mortal Wkly Rep* 2022;71:103–5. <https://doi.org/10.15585/mmwr.mm7103a5>
- 48: Written Statement of Administrator Samantha Power, House Committee on Appropriations Subcommittee on State, Foreign Operations, and Related Programs. Published May 11, 2022.
<http://docs.house.gov/meetings/AP/AP04/20220511/114711/HHRG-117-AP04-Wstate-PowerA-20220511.pdf>
- 49: Diamond D, Rouben R. 'victory for the virus': Senators cut global aid from \$10B covid deal. *The Washington Post*. Published April 4, 2022.
- 50: Johnson CY. The next leap in coronavirus vaccine development could be a nasal spray. *The Washington Post*. Published April 11, 2022.
<https://www.washingtonpost.com/health/2022/04/10/nasal-covid-vaccine/>

Figures

Figure 1: Estimated COVID-19 Deaths per Capita by Variant and U.S. Census Region



Tables

Table 3: Estimated Number of Recorded COVID-19 Deaths in the U.S. Attributable to Each Variant

Variant	Attributable Recorded COVID-19 Deaths: Median Estimate (95% Credible Interval)	Attributable Recorded COVID-19 Deaths per 100,000: Median Estimate (95% Credible Interval)	Share of All Recorded COVID-19 Deaths: Median Estimate (95% Credible Interval)
Non-Variant	539,667 (538,928 - 540,466)	161 (161 - 161)	54% (54% - 54%)
All Variants	460,124 (459,325 - 460,863)	137 (137 - 138)	46% (46% - 46%)
Delta	273,801 (273,483 - 274,168)	82 (82 - 82)	27% (27% - 27%)
Omicron	117,560 (117,252 - 117,835)	35 (35 - 35)	12% (12% - 12%)
Alpha	39,548 (38,890 - 40,133)	12 (12 - 12)	4% (4% - 4%)
Epsilon	14,879 (14,230 - 15,528)	4 (4 - 5)	1% (1% - 2%)
Iota	5,614 (5,394 - 5,810)	2 (2 - 2)	1% (1% - 1%)
B.1.1.519	3,405 (3,114 - 3,715)	1 (1 - 1)	0% (0% - 0%)

Gamma	2,950 (2,878 - 3,039)	1 (1 - 1)	0% (0% - 0%)
B.1.621	557 (527 - 596)	0 (0 - 0)	0% (0% - 0%)
Beta	534 (478 - 585)	0 (0 - 0)	0% (0% - 0%)
P.2	460 (341 - 601)	0 (0 - 0)	0% (0% - 0%)
B.1.525	277 (204 - 354)	0 (0 - 0)	0% (0% - 0%)
B.1.1.318	173 (151 - 196)	0 (0 - 0)	0% (0% - 0%)
Lambda	170 (146 - 193)	0 (0 - 0)	0% (0% - 0%)
C.36	65 (45 - 81)	0 (0 - 0)	0% (0% - 0%)
B.1.214.2	34 (24 - 49)	0 (0 - 0)	0% (0% - 0%)
B.1.617.1	31 (24 - 40)	0 (0 - 0)	0% (0% - 0%)
C.16	24 (0 - 75)	0 (0 - 0)	0% (0% - 0%)
A.23.1	16 (8 - 28)	0 (0 - 0)	0% (0% - 0%)
A.28	6 (0 - 18)	0 (0 - 0)	0% (0% - 0%)

B.1.620	2 (1 - 4)	0 (0 - 0)	0% (0% - 0%)
Theta	1 (0 - 3)	0 (0 - 0)	0% (0% - 0%)
A.27	1 (0 - 4)	0 (0 - 0)	0% (0% - 0%)
B.1.617.3	1 (0 - 1)	0 (0 - 0)	0% (0% - 0%)
B.1.640	0 (0 - 1)	0 (0 - 0)	0% (0% - 0%)
Unknown (suppressed week)	3,628	1	0%
Total	1,003,419	299	100%

Table 4: Estimated COVID-19 Deaths per Capita in U.S. by Census Region and Variants, Primary Analysis

Census Region	Median Estimated Non-Variant COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Delta COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Omicron COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Alpha COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated COVID-19 Deaths per 100,000 from Other Variants (95% CI, % of Total)	Median Estimated COVID-19 Deaths per 100,000 from All Variants (95% CI, % of Total)	Recorded COVID-19 Deaths per 100,000: Unknown Variant (% of Total)	Total Recorded COVID-19 Deaths per 100,000
National	161 (161 - 161, 54%)	82 (82 - 82, 27%)	35 (35 - 35, 12%)	12 (12 - 12, 4%)	9 (8 - 9, 3%)	137 (137 - 138, 46%)	1 (0%)	299
Northeast (Connecticut, Maine, Massachusetts, New Hampshire, New York, Pennsylvania, Rhode Island, and Vermont)	215 (215 - 216, 66%)	57 (56 - 57, 17%)	33 (33 - 33, 10%)	12 (12 - 12, 4%)	10 (9 - 10, 3%)	111 (111 - 112, 34%)	2 (1%)	328

South (Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, Puerto Rico, South Carolina, Tennessee, Texas, Virginia, and West Virginia)	158 (158 - 159, 50%)	97 (96 - 97, 30%)	41 (40 - 41, 13%)	15 (14 - 15, 5%)	6 (6 - 6, 2%)	158 (157 - 158, 50%)	1 (0%)	317
Midwest (Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin)	163 (163 - 163, 53%)	92 (91 - 92, 30%)	34 (33 - 34, 11%)	12 (12 - 12, 4%)	5 (5 - 5, 2%)	142 (142 - 143, 46%)	1 (0%)	306
West (Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming)	124 (123 - 125, 51%)	67 (66 - 67, 27%)	29 (29 - 29, 12%)	7 (6 - 7, 3%)	16 (15 - 16, 7%)	118 (118 - 119, 49%)	1 (1%)	244

Table 5: Estimated COVID-19 Deaths in U.S. by Census Region and Variants, Primary Analysis

Census Region	Median Estimated Non-Variant COVID-19 Deaths (95% CI, % of Total)	Median Estimated Delta COVID-19 Deaths (95% CI, % of Total)	Median Estimated Omicron COVID-19 Deaths (95% CI, % of Total)	Median Estimated Alpha COVID-19 Deaths (95% CI, % of Total)	Median Estimated COVID-19 Deaths from Other Variants (95% CI, % of Total)	Median Estimated COVID-19 Deaths from All Variants (95% CI, % of Total)	Recorded COVID-19 Deaths: Unknown Variant (% of Total)	Total Recorded COVID-19 Deaths
National	539,667 (538,928 - 540,466, 54%)	273,801 (273,483 - 274,168, 27%)	117,560 (117,252 - 117,835, 12%)	39,548 (38,890 - 40,133, 4%)	29,208 (28,460 - 29,926, 3%)	460,124 (459,325 - 460,863, 46%)	3,628 (0%)	1,003,419
Northeast (Connecticut, Maine, Massachusetts, New Hampshire, New York, Pennsylvania, Rhode Island, and Vermont)	123,103 (122,827 - 123,377, 66%)	32,414 (32,250 - 32,565, 17%)	18,743 (18,585 - 18,898, 10%)	6,895 (6,659 - 7,085, 4%)	5,492 (5,272 - 5,765, 3%)	63,542 (63,268 - 63,818, 34%)	991 (1%)	187,636

South (Alabama, Arkansas, Delaware, District of Columbia, Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, Puerto Rico, South Carolina, Tennessee, Texas, Virginia, and West Virginia)	206,752 (206,211 - 207,325, 50%)	125,925 (125,671 - 126,154, 30%)	52,912 (52,711 - 53,124, 13%)	19,143 (18,715 - 19,521, 5%)	7,754 (7,310 - 8,162, 2%)	205,729 (205,156 - 206,270, 50%)	828 (0%)	413,309
Midwest (Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin)	112,246 (111,992 - 112,480, 53%)	63,049 (62,910 - 63,222, 30%)	23,174 (23,015 - 23,324, 11%)	8,176 (8,031 - 8,316, 4%)	3,458 (3,245 - 3,678, 2%)	97,856 (97,622 - 98,110, 46%)	671 (0%)	210,773
West (Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming)	97,549 (97,134 - 98,042, 51%)	52,409 (52,222 - 52,573, 27%)	22,738 (22,628 - 22,831, 12%)	5,339 (4,969 - 5,674, 3%)	12,502 (12,002 - 12,971, 7%)	93,014 (92,521 - 93,429, 49%)	1,138 (1%)	191,701

Supplementary Table 1: Estimated COVID-19 Deaths per Capita in the U.S. by Jurisdiction and Variant, Primary Analysis

Jurisdiction	Median Estimated Non-Variant COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Delta COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Omicron COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated Alpha COVID-19 Deaths per 100,000 (95% CI, % of Total)	Median Estimated COVID-19 Deaths per 100,000 from Other Variants (95% CI, % of Total)	Median Estimated COVID-19 Deaths per 100,000 from All Variants (95% CI, % of Total)	Recorded COVID-19 Deaths per 100,000: Unknown Variant (% of Total)	Total Recorded COVID-19 Deaths per 100,000
Alabama	206 (205 - 207, 54%)	112 (111 - 113, 29%)	50 (49 - 51, 13%)	11 (10 - 12, 3%)	4 (3 - 5, 1%)	177 (176 - 178, 46%)	0 (0%)	384
Alaska	27 (26 - 27, 15%)	97 (96 - 98, 56%)	16 (16 - 18, 9%)	1 (1 - 1, 1%)	3 (3 - 3, 2%)	118 (118 - 118, 68%)	30 (17%)	174
Arizona	201 (200 - 203, 54%)	112 (111 - 112, 30%)	40 (39 - 40, 11%)	6 (6 - 7, 2%)	16 (15 - 18, 4%)	174 (173 - 175, 46%)	0 (0%)	375

Arkansas	194 (193 - 195, 53%)	115 (114 - 117, 31%)	48 (47 - 49, 13%)	8 (7 - 9, 2%)	3 (2 - 5, 1%)	174 (173 - 175, 47%)	1 (0%)	369
California	137 (136 - 138, 57%)	46 (46 - 47, 19%)	28 (28 - 28, 12%)	7 (6 - 8, 3%)	23 (22 - 24, 10%)	105 (104 - 106, 43%)	0 (0%)	242
Colorado	107 (105 - 108, 47%)	79 (79 - 80, 35%)	27 (27 - 28, 12%)	9 (9 - 9, 4%)	6 (5 - 7, 2%)	121 (120 - 123, 53%)	0 (0%)	228
Connecticut	216 (215 - 218, 71%)	41 (40 - 41, 13%)	29 (28 - 30, 9%)	6 (6 - 7, 2%)	9 (7 - 10, 3%)	85 (83 - 86, 28%)	3 (1%)	304
Delaware	153 (152 - 155, 53%)	70 (69 - 72, 24%)	38 (36 - 39, 13%)	6 (5 - 7, 2%)	5 (3 - 6, 2%)	119 (117 - 120, 41%)	19 (6%)	291

District of Columbia	187 (185 - 189, 63%)	26 (25 - 28, 9%)	33 (31 - 34, 11%)	10 (8 - 12, 3%)	7 (5 - 10, 2%)	76 (74 - 78, 25%)	36 (12%)	298
Florida	134 (133 - 135, 42%)	116 (115 - 116, 36%)	39 (38 - 39, 12%)	25 (24 - 26, 8%)	9 (8 - 9, 3%)	188 (187 - 189, 58%)	0 (0%)	322
Georgia	150 (148 - 152, 50%)	92 (92 - 93, 31%)	38 (37 - 38, 13%)	16 (14 - 18, 5%)	4 (3 - 5, 1%)	150 (148 - 152, 50%)	0 (0%)	300
Hawaii	21 (21 - 22, 22%)	38 (38 - 39, 40%)	15 (15 - 16, 16%)	0 (0 - 1, 0%)	1 (0 - 1, 1%)	55 (55 - 55, 57%)	20 (20%)	96
Idaho	100 (99 - 102, 38%)	117 (116 - 118, 45%)	26 (26 - 27, 10%)	8 (7 - 9, 3%)	6 (5 - 8, 2%)	158 (156 - 159, 60%)	5 (2%)	262

Illinois	162 (161 - 162, 60%)	61 (60 - 62, 23%)	31 (31 - 32, 12%)	9 (9 - 10, 3%)	8 (7 - 8, 3%)	109 (109 - 110, 40%)	0 (0%)	271
Indiana	188 (187 - 189, 54%)	109 (108 - 109, 31%)	37 (36 - 38, 11%)	10 (10 - 11, 3%)	5 (4 - 5, 1%)	161 (160 - 161, 46%)	0 (0%)	349
Iowa	182 (180 - 185, 61%)	80 (80 - 81, 27%)	25 (24 - 26, 8%)	6 (5 - 6, 2%)	5 (2 - 7, 2%)	116 (113 - 119, 39%)	1 (0%)	300
Kansas	162 (161 - 163, 53%)	97 (96 - 98, 32%)	34 (33 - 35, 11%)	5 (4 - 5, 2%)	5 (4 - 6, 2%)	141 (140 - 142, 46%)	2 (1%)	304
Kentucky	157 (155 - 158, 43%)	137 (135 - 138, 37%)	57 (56 - 59, 16%)	12 (11 - 13, 3%)	5 (4 - 7, 1%)	212 (210 - 213, 57%)	0 (0%)	369

Louisiana	196 (195 - 197, 57%)	90 (89 - 91, 26%)	37 (36 - 38, 11%)	13 (12 - 14, 4%)	8 (7 - 9, 2%)	147 (146 - 148, 43%)	0 (0%)	344
Maine	55 (54 - 55, 30%)	76 (76 - 77, 42%)	29 (28 - 30, 16%)	4 (3 - 4, 2%)	3 (3 - 4, 2%)	112 (112 - 113, 62%)	14 (8%)	181
Maryland	150 (149 - 152, 60%)	49 (48 - 50, 19%)	32 (31 - 33, 13%)	15 (14 - 16, 6%)	5 (5 - 6, 2%)	101 (100 - 102, 40%)	0 (0%)	251
Massachusetts	186 (185 - 186, 71%)	40 (40 - 40, 15%)	25 (25 - 26, 10%)	6 (6 - 7, 2%)	4 (4 - 5, 2%)	76 (75 - 76, 29%)	0 (0%)	262
Michigan	151 (151 - 152, 48%)	99 (98 - 100, 31%)	32 (31 - 32, 10%)	31 (30 - 32, 10%)	5 (4 - 6, 2%)	167 (166 - 168, 52%)	0 (0%)	318

Minnesota	124 (124 - 124, 55%)	65 (65 - 66, 29%)	22 (21 - 22, 10%)	10 (10 - 10, 4%)	3 (3 - 3, 1%)	100 (100 - 100, 45%)	0 (0%)	224
Mississippi	247 (245 - 249, 55%)	120 (119 - 122, 27%)	60 (59 - 62, 13%)	14 (12 - 15, 3%)	5 (3 - 7, 1%)	199 (197 - 202, 45%)	0 (0%)	447
Missouri	171 (170 - 172, 51%)	110 (109 - 111, 33%)	38 (38 - 39, 11%)	9 (8 - 9, 3%)	5 (4 - 6, 1%)	162 (161 - 163, 49%)	0 (0%)	333
Montana	137 (135 - 138, 43%)	133 (131 - 134, 42%)	30 (29 - 31, 9%)	3 (3 - 4, 1%)	4 (3 - 5, 1%)	170 (169 - 172, 53%)	12 (4%)	319
Nebraska	140 (139 - 141, 56%)	73 (72 - 74, 29%)	26 (25 - 27, 10%)	5 (5 - 6, 2%)	3 (2 - 3, 1%)	107 (106 - 108, 43%)	4 (2%)	251

Nevada	162 (158 - 165, 46%)	116 (115 - 117, 33%)	46 (46 - 47, 13%)	10 (9 - 10, 3%)	16 (13 - 20, 5%)	188 (185 - 192, 54%)	1 (0%)	351
New Hampshire	85 (84 - 85, 48%)	56 (55 - 57, 31%)	20 (19 - 21, 11%)	2 (1 - 2, 1%)	1 (1 - 2, 1%)	79 (78 - 79, 44%)	14 (8%)	178
New Jersey	241 (240 - 242, 70%)	42 (41 - 43, 12%)	35 (34 - 35, 10%)	15 (14 - 17, 4%)	12 (11 - 13, 3%)	104 (103 - 105, 30%)	0 (0%)	345
New Mexico	187 (186 - 188, 51%)	124 (123 - 126, 34%)	39 (38 - 39, 11%)	7 (7 - 8, 2%)	7 (6 - 8, 2%)	177 (176 - 178, 48%)	3 (1%)	367
New York State (non-NYC)	202 (201 - 203, 64%)	57 (56 - 58, 18%)	35 (35 - 36, 11%)	11 (11 - 12, 4%)	11 (11 - 12, 4%)	115 (114 - 116, 36%)	0 (0%)	318

New York City	315 (314 - 316, 76%)	34 (33 - 34, 8%)	32 (31 - 32, 8%)	16 (15 - 17, 4%)	17 (16 - 18, 4%)	99 (97 - 100, 24%)	0 (0%)	413
North Carolina	138 (137 - 139, 51%)	83 (82 - 83, 31%)	34 (33 - 34, 12%)	10 (9 - 11, 4%)	6 (5 - 6, 2%)	132 (131 - 133, 49%)	0 (0%)	270
North Dakota	199 (198 - 200, 56%)	93 (91 - 95, 26%)	28 (27 - 31, 8%)	1 (1 - 1, 0%)	4 (3 - 5, 1%)	127 (126 - 128, 36%)	31 (9%)	357
Ohio	182 (181 - 183, 49%)	123 (122 - 124, 33%)	47 (46 - 48, 13%)	11 (11 - 12, 3%)	5 (4 - 6, 1%)	187 (185 - 188, 51%)	0 (0%)	369
Oklahoma	208 (206 - 210, 52%)	125 (123 - 127, 31%)	56 (55 - 58, 14%)	8 (7 - 9, 2%)	3 (2 - 5, 1%)	193 (191 - 194, 48%)	1 (0%)	402

Oregon	53 (52 - 53, 31%)	74 (73 - 74, 44%)	27 (27 - 28, 16%)	7 (6 - 7, 4%)	8 (7 - 9, 5%)	116 (115 - 117, 69%)	1 (0%)	170
Pennsylvania	199 (198 - 200, 56%)	96 (95 - 97, 27%)	38 (38 - 39, 11%)	15 (15 - 16, 4%)	7 (7 - 8, 2%)	157 (156 - 158, 44%)	0 (0%)	356
Puerto Rico	61 (60 - 62, 48%)	25 (24 - 26, 20%)	24 (23 - 24, 19%)	10 (9 - 10, 7%)	4 (3 - 6, 3%)	63 (62 - 64, 49%)	4 (3%)	129
Rhode Island	228 (227 - 229, 69%)	47 (46 - 48, 14%)	26 (24 - 27, 8%)	3 (3 - 4, 1%)	6 (5 - 7, 2%)	82 (81 - 83, 25%)	19 (6%)	329
South Carolina	178 (177 - 179, 52%)	105 (105 - 106, 31%)	45 (44 - 45, 13%)	10 (10 - 11, 3%)	6 (5 - 8, 2%)	167 (166 - 168, 48%)	0 (0%)	345

South Dakota	214 (213 - 215, 62%)	77 (76 - 78, 22%)	28 (27 - 29, 8%)	4 (3 - 4, 1%)	1 (1 - 2, 0%)	109 (109 - 110, 32%)	20 (6%)	344
Tennessee	174 (172 - 175, 46%)	126 (125 - 127, 34%)	55 (54 - 56, 15%)	17 (17 - 18, 5%)	4 (3 - 5, 1%)	202 (201 - 204, 54%)	0 (0%)	376
Texas	171 (170 - 173, 53%)	92 (91 - 92, 28%)	38 (38 - 38, 12%)	14 (13 - 15, 4%)	7 (6 - 8, 2%)	151 (149 - 152, 47%)	0 (0%)	322
Utah	68 (68 - 69, 44%)	59 (59 - 60, 39%)	17 (17 - 17, 11%)	3 (3 - 3, 2%)	4 (3 - 5, 3%)	83 (83 - 84, 54%)	2 (1%)	154
Vermont	16 (16 - 16, 17%)	27 (26 - 28, 27%)	16 (15 - 17, 16%)	0 (0 - 0, 0%)	0 (0 - 0, 0%)	42 (42 - 42, 44%)	38 (39%)	97

Virginia	117 (116 - 117, 52%)	61 (60 - 62, 27%)	33 (32 - 33, 15%)	9 (9 - 9, 4%)	4 (4 - 5, 2%)	107 (107 - 108, 48%)	0 (0%)	224
Washington	63 (62 - 64, 41%)	57 (56 - 57, 36%)	24 (24 - 24, 15%)	6 (5 - 6, 4%)	7 (6 - 7, 4%)	93 (92 - 94, 59%)	0 (0%)	156
West Virginia	150 (149 - 151, 39%)	154 (152 - 155, 40%)	58 (57 - 60, 15%)	13 (12 - 14, 3%)	4 (3 - 5, 1%)	229 (228 - 230, 60%)	6 (1%)	384
Wisconsin	134 (133 - 134, 54%)	80 (79 - 80, 32%)	27 (26 - 27, 11%)	5 (5 - 5, 2%)	4 (3 - 4, 1%)	115 (114 - 116, 46%)	0 (0%)	249
Wyoming	91 (90 - 92, 33%)	130 (128 - 132, 47%)	24 (23 - 26, 9%)	0 (0 - 1, 0%)	3 (2 - 4, 1%)	157 (156 - 158, 56%)	31 (11%)	280

Supplementary Table 2: Estimated COVID-19 Deaths in the U.S. by Jurisdiction and Variant, Primary Analysis

Jurisdiction	Median Estimated Non-Variant COVID-19 Deaths (95% CI, % of Total)	Median Estimated Delta COVID-19 Deaths (95% CI, % of Total)	Median Estimated Omicron COVID-19 Deaths (95% CI, % of Total)	Median Estimated Alpha COVID-19 Deaths (95% CI, % of Total)	Median Estimated COVID-19 Deaths from Other Variants (95% CI, % of Total)	Median Estimated COVID-19 Deaths from All Variants (95% CI, % of Total)	Recorded COVID-19 Deaths: Unknown Variant (% of Total)	Total Recorded COVID-19 Deaths
Alabama	10,402 (10,347 - 10,449, 54%)	5,660 (5,610 - 5,715, 29%)	2,534 (2,492 - 2,579, 13%)	547 (500 - 604, 3%)	181 (140 - 242, 1%)	8,925 (8,878 - 8,980, 46%)	10 (0%)	19,337
Alaska	195 (194 - 196, 15%)	714 (704 - 721, 56%)	120 (114 - 130, 9%)	8 (5 - 11, 1%)	21 (18 - 24, 2%)	863 (862 - 864, 68%)	217 (17%)	1,275
Arizona	14,659 (14,543 - 14,754, 54%)	8,115 (8,065 - 8,157, 30%)	2,899 (2,861 - 2,946, 11%)	460 (430 - 487, 2%)	1,176 (1,079 - 1,296, 4%)	12,647 (12,552 - 12,763, 46%)	15 (0%)	27,321

Arkansas	5,880 (5,842 - 5,909, 53%)	3,488 (3,436 - 3,526, 31%)	1,440 (1,408 - 1,483, 13%)	228 (202 - 258, 2%)	100 (73 - 142, 1%)	5,256 (5,227 - 5,294, 47%)	23 (0%)	11,159
California	53,817 (53,410 - 54,230, 57%)	18,219 (18,075 - 18,373, 19%)	11,085 (11,025 - 11,142, 12%)	2,866 (2,503 - 3,197, 3%)	9,127 (8,747 - 9,594, 10%)	41,300 (40,887 - 41,707, 43%)	13 (0%)	95,130
Colorado	6,199 (6,128 - 6,261, 47%)	4,598 (4,564 - 4,632, 35%)	1,591 (1,560 - 1,625, 12%)	533 (519 - 549, 4%)	329 (271 - 400, 2%)	7,052 (6,990 - 7,123, 53%)	14 (0%)	13,265
Connecticut	7,793 (7,736 - 7,849, 71%)	1,466 (1,433 - 1,491, 13%)	1,036 (1,011 - 1,065, 9%)	234 (220 - 250, 2%)	315 (262 - 378, 3%)	3,052 (2,996 - 3,109, 28%)	103 (1%)	10,948
Delaware	1,539 (1,522 - 1,554, 53%)	705 (690 - 720, 24%)	379 (365 - 394, 13%)	59 (50 - 69, 2%)	46 (31 - 61, 2%)	1,190 (1,175 - 1,207, 41%)	189 (6%)	2,918

District of Columbia	1,252 (1,239 - 1,268, 63%)	177 (165 - 190, 9%)	219 (206 - 231, 11%)	68 (55 - 80, 3%)	45 (31 - 64, 2%)	509 (493 - 522, 25%)	238 (12%)	1,999
Florida	29,179 (28,976 - 29,399, 42%)	25,174 (25,093 - 25,251, 36%)	8,428 (8,371 - 8,482, 12%)	5,422 (5,230 - 5,626, 8%)	1,919 (1,837 - 2,030, 3%)	40,941 (40,721 - 41,144, 58%)	4 (0%)	70,124
Georgia	16,226 (16,016 - 16,401, 50%)	9,966 (9,911 - 10,014, 31%)	4,061 (4,017 - 4,110, 13%)	1,705 (1,542 - 1,905, 5%)	455 (365 - 546, 1%)	16,194 (16,019 - 16,404, 50%)	15 (0%)	32,435
Hawaii	309 (304 - 314, 22%)	554 (546 - 562, 40%)	221 (214 - 230, 16%)	6 (4 - 8, 0%)	12 (7 - 17, 1%)	793 (788 - 798, 57%)	284 (20%)	1,386
Idaho	1,899 (1,874 - 1,930, 38%)	2,227 (2,212 - 2,241, 45%)	500 (489 - 511, 10%)	152 (137 - 169, 3%)	119 (92 - 147, 2%)	2,998 (2,967 - 3,023, 60%)	92 (2%)	4,989

Illinois	20,499 (20,401 - 20,578, 60%)	7,732 (7,656 - 7,796, 23%)	3,952 (3,890 - 4,017, 12%)	1,189 (1,149 - 1,237, 3%)	963 (882 - 1,054, 3%)	13,837 (13,758 - 13,935, 40%)	1 (0%)	34,337
Indiana	12,789 (12,731 - 12,846, 54%)	7,390 (7,327 - 7,439, 31%)	2,518 (2,469 - 2,572, 11%)	704 (660 - 754, 3%)	312 (273 - 365, 1%)	10,927 (10,870 - 10,985, 46%)	18 (0%)	23,734
Iowa	5,818 (5,736 - 5,903, 61%)	2,564 (2,541 - 2,585, 27%)	798 (781 - 817, 8%)	186 (171 - 201, 2%)	157 (71 - 238, 2%)	3,704 (3,619 - 3,786, 39%)	42 (0%)	9,564
Kansas	4,754 (4,722 - 4,778, 53%)	2,849 (2,823 - 2,874, 32%)	1,002 (979 - 1,025, 11%)	134 (120 - 149, 2%)	144 (120 - 177, 2%)	4,128 (4,104 - 4,160, 46%)	50 (1%)	8,932
Kentucky	7,075 (7,005 - 7,142, 43%)	6,180 (6,108 - 6,245, 37%)	2,591 (2,533 - 2,656, 16%)	550 (504 - 599, 3%)	221 (167 - 299, 1%)	9,546 (9,479 - 9,616, 57%)	20 (0%)	16,641

Louisiana	9,076 (9,030 - 9,126, 57%)	4,151 (4,107 - 4,198, 26%)	1,714 (1,683 - 1,745, 11%)	605 (565 - 648, 4%)	348 (303 - 396, 2%)	6,819 (6,769 - 6,865, 43%)	18 (0%)	15,913
Maine	748 (742 - 756, 30%)	1,048 (1,037 - 1,059, 42%)	395 (384 - 406, 16%)	50 (44 - 57, 2%)	46 (41 - 52, 2%)	1,540 (1,532 - 1,546, 62%)	192 (8%)	2,480
Maryland	9,277 (9,192 - 9,346, 60%)	3,022 (2,975 - 3,075, 19%)	1,957 (1,905 - 2,009, 13%)	899 (844 - 986, 6%)	331 (284 - 369, 2%)	6,209 (6,140 - 6,294, 40%)	12 (0%)	15,498
Massachusetts	12,966 (12,917 - 13,009, 71%)	2,788 (2,759 - 2,813, 15%)	1,767 (1,743 - 1,793, 10%)	432 (399 - 466, 2%)	296 (266 - 326, 2%)	5,281 (5,238 - 5,330, 29%)	25 (0%)	18,272
Michigan	15,227 (15,142 - 15,322, 48%)	9,961 (9,897 - 10,019, 31%)	3,191 (3,136 - 3,240, 10%)	3,095 (3,012 - 3,179, 10%)	512 (446 - 587, 2%)	16,761 (16,666 - 16,846, 52%)	4 (0%)	31,992

Minnesota	7,072 (7,053 - 7,089, 55%)	3,718 (3,694 - 3,741, 29%)	1,247 (1,225 - 1,271, 10%)	565 (551 - 577, 4%)	166 (153 - 180, 1%)	5,696 (5,679 - 5,715, 45%)	1 (0%)	12,769
Mississippi	7,290 (7,226 - 7,348, 55%)	3,551 (3,502 - 3,605, 27%)	1,773 (1,728 - 1,831, 13%)	404 (360 - 443, 3%)	156 (95 - 217, 1%)	5,884 (5,826 - 5,948, 45%)	10 (0%)	13,184
Missouri	10,560 (10,466 - 10,626, 51%)	6,804 (6,753 - 6,856, 33%)	2,356 (2,315 - 2,402, 11%)	528 (491 - 569, 3%)	285 (217 - 370, 1%)	9,974 (9,908 - 10,068, 49%)	10 (0%)	20,544
Montana	1,511 (1,495 - 1,523, 43%)	1,464 (1,449 - 1,482, 42%)	333 (315 - 347, 9%)	37 (29 - 45, 1%)	44 (32 - 60, 1%)	1,878 (1,866 - 1,894, 53%)	137 (4%)	3,526
Nebraska	2,751 (2,731 - 2,768, 56%)	1,436 (1,421 - 1,451, 29%)	512 (497 - 527, 10%)	102 (94 - 110, 2%)	50 (33 - 69, 1%)	2,099 (2,082 - 2,119, 43%)	87 (2%)	4,937

Nevada	5,082 (4,971 - 5,176, 46%)	3,641 (3,615 - 3,670, 33%)	1,460 (1,435 - 1,483, 13%)	305 (276 - 328, 3%)	513 (403 - 624, 5%)	5,918 (5,824 - 6,029, 54%)	23 (0%)	11,023
New Hampshire	1,176 (1,167 - 1,184, 48%)	775 (760 - 789, 31%)	276 (261 - 289, 11%)	24 (21 - 31, 1%)	18 (13 - 26, 1%)	1,093 (1,085 - 1,102, 44%)	201 (8%)	2,470
New Jersey	22,335 (22,202 - 22,450, 70%)	3,884 (3,823 - 3,943, 12%)	3,210 (3,148 - 3,266, 10%)	1,436 (1,332 - 1,529, 4%)	1,106 (1,033 - 1,212, 3%)	9,635 (9,520 - 9,768, 30%)	3 (0%)	31,973
New Mexico	3,958 (3,935 - 3,981, 51%)	2,634 (2,612 - 2,656, 34%)	815 (798 - 835, 11%)	153 (140 - 168, 2%)	147 (124 - 171, 2%)	3,751 (3,728 - 3,774, 48%)	53 (1%)	7,762
New York State (non-NYC)	22,990 (22,865 - 23,110, 64%)	6,484 (6,413 - 6,552, 18%)	4,012 (3,953 - 4,090, 11%)	1,305 (1,215 - 1,404, 4%)	1,300 (1,203 - 1,397, 4%)	13,102 (12,982 - 13,227, 36%)	9 (0%)	36,101

New York City	26,665 (26,571 - 26,786, 76%)	2,849 (2,795 - 2,904, 8%)	2,686 (2,637 - 2,740, 8%)	1,386 (1,301 - 1,470, 4%)	1,425 (1,339 - 1,517, 4%)	8,346 (8,225 - 8,440, 24%)	2 (0%)	35,013
North Carolina	14,590 (14,484 - 14,683, 51%)	8,722 (8,676 - 8,765, 31%)	3,540 (3,502 - 3,581, 12%)	1,074 (996 - 1,169, 4%)	582 (525 - 666, 2%)	13,924 (13,831 - 14,030, 49%)	4 (0%)	28,518
North Dakota	1,542 (1,533 - 1,550, 56%)	724 (704 - 739, 26%)	221 (207 - 240, 8%)	8 (5 - 11, 0%)	33 (26 - 42, 1%)	986 (978 - 995, 36%)	237 (9%)	2,765
Ohio	21,444 (21,277 - 21,606, 49%)	14,499 (14,391 - 14,623, 33%)	5,535 (5,420 - 5,647, 13%)	1,346 (1,275 - 1,429, 3%)	607 (473 - 747, 1%)	21,985 (21,823 - 22,152, 51%)	11 (0%)	43,440
Oklahoma	8,299 (8,231 - 8,365, 52%)	4,993 (4,921 - 5,059, 31%)	2,241 (2,184 - 2,299, 14%)	318 (275 - 373, 2%)	129 (91 - 185, 1%)	7,686 (7,620 - 7,754, 48%)	24 (0%)	16,009

Oregon	2,237 (2,201 - 2,269, 31%)	3,132 (3,103 - 3,156, 44%)	1,154 (1,132 - 1,177, 16%)	292 (269 - 311, 4%)	354 (318 - 395, 5%)	4,932 (4,900 - 4,968, 69%)	29 (0%)	7,198
Pennsylvania	25,834 (25,713 - 25,929, 56%)	12,438 (12,334 - 12,527, 27%)	4,971 (4,889 - 5,070, 11%)	1,978 (1,913 - 2,069, 4%)	918 (846 - 1,007, 2%)	20,312 (20,217 - 20,433, 44%)	5 (0%)	46,151
Puerto Rico	1,998 (1,956 - 2,037, 48%)	819 (797 - 835, 20%)	778 (762 - 799, 19%)	310 (290 - 330, 7%)	146 (110 - 186, 3%)	2,052 (2,013 - 2,094, 49%)	144 (3%)	4,194
Rhode Island	2,499 (2,485 - 2,513, 69%)	518 (505 - 531, 14%)	280 (266 - 292, 8%)	37 (33 - 42, 1%)	64 (50 - 77, 2%)	898 (884 - 912, 25%)	204 (6%)	3,601
South Carolina	9,236 (9,169 - 9,293, 52%)	5,475 (5,430 - 5,512, 31%)	2,323 (2,289 - 2,361, 13%)	543 (494 - 590, 3%)	329 (266 - 396, 2%)	8,665 (8,608 - 8,732, 48%)	4 (0%)	17,905

South Dakota	1,917 (1,911 - 1,924, 62%)	689 (679 - 701, 22%)	250 (238 - 261, 8%)	31 (26 - 37, 1%)	10 (6 - 13, 0%)	980 (973 - 986, 32%)	182 (6%)	3,079
Tennessee	12,117 (12,004 - 12,187, 46%)	8,791 (8,723 - 8,853, 34%)	3,818 (3,763 - 3,875, 15%)	1,217 (1,163 - 1,261, 5%)	278 (215 - 377, 1%)	14,102 (14,032 - 14,215, 54%)	3 (0%)	26,222
Texas	50,572 (50,151 - 50,960, 53%)	27,045 (26,966 - 27,122, 28%)	11,246 (11,178 - 11,320, 12%)	4,159 (3,918 - 4,395, 4%)	2,003 (1,690 - 2,369, 2%)	44,448 (44,060 - 44,869, 47%)	2 (0%)	95,022
Utah	2,278 (2,254 - 2,299, 44%)	1,975 (1,955 - 1,993, 39%)	568 (555 - 584, 11%)	103 (89 - 116, 2%)	129 (105 - 152, 3%)	2,775 (2,754 - 2,799, 54%)	72 (1%)	5,125
Vermont	106 (106 - 106, 17%)	171 (165 - 178, 27%)	103 (96 - 108, 16%)	0 (0 - 0, 0%)	0 (0 - 0, 0%)	274 (274 - 274, 44%)	247 (39%)	627

Virginia	10,102 (10,055 - 10,149, 52%)	5,270 (5,216 - 5,326, 27%)	2,831 (2,777 - 2,881, 15%)	790 (759 - 820, 4%)	381 (342 - 425, 2%)	9,270 (9,223 - 9,317, 48%)	8 (0%)	19,380
Washington	4,899 (4,829 - 4,958, 41%)	4,386 (4,352 - 4,416, 36%)	1,850 (1,824 - 1,882, 15%)	429 (406 - 454, 4%)	504 (448 - 576, 4%)	7,173 (7,114 - 7,243, 59%)	11 (0%)	12,083
West Virginia	2,672 (2,654 - 2,691, 39%)	2,740 (2,709 - 2,770, 40%)	1,036 (1,011 - 1,068, 15%)	230 (216 - 244, 3%)	72 (57 - 87, 1%)	4,079 (4,060 - 4,097, 60%)	100 (1%)	6,851
Wisconsin	7,877 (7,828 - 7,913, 54%)	4,694 (4,664 - 4,723, 32%)	1,584 (1,558 - 1,611, 11%)	287 (271 - 305, 2%)	210 (171 - 257, 1%)	6,775 (6,739 - 6,824, 46%)	28 (0%)	14,680
Wyoming	529 (523 - 535, 33%)	754 (743 - 763, 47%)	138 (131 - 150, 9%)	1 (0 - 4, 0%)	17 (11 - 23, 1%)	911 (905 - 917, 56%)	178 (11%)	1,618

Supplementary Table 3: Estimated Omicron and Delta COVID-19 Deaths in U.S. by Census Region, Sensitivity Analysis Assuming Higher or Lower Omicron Severity

Census Region	Analysis	Median Estimated Delta COVID-19 Deaths (95% CI)	Median Estimated Omicron COVID-19 Deaths (95% CI)	Median Estimated Delta COVID-19 Deaths per 100k (95% CI)	Median Estimated Omicron COVID-19 Deaths per 100k (95% CI)	Median Estimated % of COVID-19 Deaths from Delta (95% CI)	Median Estimated % of COVID-19 Deaths from Omicron (95% CI)
National	Higher Omicron Severity	265,228 (264,898 - 265,600)	126,327 (126,022 - 126,582)	79 (79 - 79)	38 (38 - 38)	26% (26% - 26%)	13% (13% - 13%)
	Lower Omicron Severity	291,946 (291,623 - 292,319)	99,023 (98,727 - 99,352)	87 (87 - 87)	30 (29 - 30)	29% (29% - 29%)	10% (10% - 10%)
Northeast (Connecticut, Maine, Massachusetts, New Hampshire, New York, Pennsylvania, Rhode Island, and Vermont)	Higher Omicron Severity	30,601 (30,458 - 30,746)	20,593 (20,452 - 20,728)	54 (53 - 54)	36 (36 - 36)	16% (16% - 16%)	11% (11% - 11%)
	Lower Omicron Severity	36,099 (35,925 - 36,249)	14,984 (14,847 - 15,140)	63 (63 - 63)	26 (26 - 26)	19% (19% - 19%)	8% (8% - 8%)
South (Alabama, Arkansas, Delaware, District of Columbia,	Higher Omicron Severity	123,151 (122,921 - 123,393)	55,766 (55,572 - 55,964)	94 (94 - 95)	43 (43 - 43)	30% (30% - 30%)	13% (13% - 14%)

Florida, Georgia, Kentucky, Louisiana, Maryland, Mississippi, North Carolina, Oklahoma, Puerto Rico, South Carolina, Tennessee, Texas, Virginia, and West Virginia)	Lower Omicron Severity	132,324 (132,051 - 132,547)	46,294 (46,108 - 46,525)	101 (101 - 102)	35 (35 - 36)	32% (32% - 32%)	11% (11% - 11%)
Midwest (Illinois, Indiana, Iowa, Kansas, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, South Dakota, and Wisconsin)	Higher Omicron Severity	60,446 (60,290 - 60,618)	25,823 (25,671 - 25,979)	88 (88 - 88)	38 (37 - 38)	29% (29% - 29%)	12% (12% - 12%)
	Lower Omicron Severity	68,011 (67,874 - 68,154)	18,140 (18,022 - 18,261)	99 (99 - 99)	26 (26 - 27)	32% (32% - 32%)	9% (9% - 9%)
West (Alaska, Arizona, California, Colorado, Hawaii, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington, and Wyoming)	Higher Omicron Severity	51,022 (50,841 - 51,195)	24,139 (24,039 - 24,249)	65 (65 - 65)	31 (31 - 31)	27% (27% - 27%)	13% (13% - 13%)
	Lower Omicron Severity	55,528 (55,338 - 55,679)	19,597 (19,498 - 19,683)	71 (70 - 71)	25 (25 - 25)	29% (29% - 29%)	10% (10% - 10%)

Supplementary Table 4: Estimated Omicron and Delta COVID-19 Deaths in U.S. by Jurisdiction, Sensitivity Analysis Assuming Higher or Lower Omicron Severity

Jurisdiction	Analysis	Median Estimated Delta COVID-19 Deaths (95% CI)	Median Estimated Omicron COVID-19 Deaths (95% CI)	Median Estimated Delta COVID-19 Deaths per 100k (95% CI)	Median Estimated Omicron COVID-19 Deaths per 100k (95% CI)	Median Estimated % of COVID-19 Deaths from Delta (95% CI)	Median Estimated % of COVID-19 Deaths from Omicron (95% CI)
Alabama	Higher Omicron Severity	5,572 (5,528 - 5,624)	2,624 (2,585 - 2,664)	111 (110 - 112)	52 (51 - 53)	29% (29% - 29%)	14% (13% - 14%)
Alabama	Lower Omicron Severity	5,893 (5,826 - 5,954)	2,295 (2,242 - 2,357)	117 (116 - 118)	46 (44 - 47)	30% (30% - 31%)	12% (12% - 12%)
Alaska	Higher Omicron Severity	707 (699 - 715)	127 (120 - 134)	96 (95 - 98)	17 (16 - 18)	55% (55% - 56%)	10% (9% - 11%)
Alaska	Lower Omicron Severity	730 (717 - 737)	104 (98 - 117)	100 (98 - 101)	14 (13 - 16)	57% (56% - 58%)	8% (8% - 9%)

Arizona	Higher Omicron Severity	7,922 (7,870 - 7,965)	3,094 (3,052 - 3,146)	109 (108 - 109)	43 (42 - 43)	29% (29% - 29%)	11% (11% - 12%)
Arizona	Lower Omicron Severity	8,483 (8,440 - 8,517)	2,530 (2,499 - 2,562)	117 (116 - 117)	35 (34 - 35)	31% (31% - 31%)	9% (9% - 9%)
Arkansas	Higher Omicron Severity	3,412 (3,360 - 3,453)	1,516 (1,482 - 1,559)	113 (111 - 114)	50 (49 - 52)	31% (30% - 31%)	14% (13% - 14%)
Arkansas	Lower Omicron Severity	3,647 (3,604 - 3,676)	1,281 (1,258 - 1,314)	121 (119 - 121)	42 (42 - 43)	33% (32% - 33%)	11% (11% - 12%)
California	Higher Omicron Severity	17,572 (17,435 - 17,726)	11,739 (11,685 - 11,807)	45 (44 - 45)	30 (30 - 30)	18% (18% - 19%)	12% (12% - 12%)
California	Lower Omicron Severity	19,749 (19,605 - 19,905)	9,545 (9,481 - 9,593)	50 (50 - 51)	24 (24 - 24)	21% (21% - 21%)	10% (10% - 10%)

Colorado	Higher Omicron Severity	4,481 (4,451 - 4,512)	1,708 (1,676 - 1,735)	77 (77 - 78)	29 (29 - 30)	34% (34% - 34%)	13% (13% - 13%)
Colorado	Lower Omicron Severity	4,844 (4,808 - 4,879)	1,342 (1,313 - 1,376)	83 (83 - 84)	23 (23 - 24)	37% (36% - 37%)	10% (10% - 10%)
Connecticut	Higher Omicron Severity	1,361 (1,330 - 1,386)	1,143 (1,117 - 1,172)	38 (37 - 38)	32 (31 - 33)	12% (12% - 13%)	10% (10% - 11%)
Connecticut	Lower Omicron Severity	1,691 (1,664 - 1,716)	806 (784 - 834)	47 (46 - 48)	22 (22 - 23)	15% (15% - 16%)	7% (7% - 8%)
Delaware	Higher Omicron Severity	671 (655 - 684)	413 (400 - 428)	67 (65 - 68)	41 (40 - 43)	23% (22% - 23%)	14% (14% - 15%)
Delaware	Lower Omicron Severity	780 (762 - 795)	303 (290 - 321)	78 (76 - 79)	30 (29 - 32)	27% (26% - 27%)	10% (10% - 11%)

District of Columbia	Higher Omicron Severity	159 (149 - 170)	236 (224 - 247)	24 (22 - 25)	35 (33 - 37)	8% (7% - 9%)	12% (11% - 12%)
District of Columbia	Lower Omicron Severity	222 (209 - 238)	173 (158 - 187)	33 (31 - 36)	26 (24 - 28)	11% (10% - 12%)	9% (8% - 9%)
Florida	Higher Omicron Severity	24,814 (24,734 - 24,888)	8,806 (8,760 - 8,852)	114 (114 - 114)	40 (40 - 41)	35% (35% - 35%)	13% (12% - 13%)
Florida	Lower Omicron Severity	26,101 (26,007 - 26,203)	7,449 (7,371 - 7,520)	120 (119 - 120)	34 (34 - 35)	37% (37% - 37%)	11% (11% - 11%)
Georgia	Higher Omicron Severity	9,754 (9,706 - 9,801)	4,281 (4,246 - 4,322)	90 (90 - 91)	40 (39 - 40)	30% (30% - 30%)	13% (13% - 13%)
Georgia	Lower Omicron Severity	10,481 (10,419 - 10,540)	3,520 (3,465 - 3,579)	97 (96 - 98)	33 (32 - 33)	32% (32% - 32%)	11% (11% - 11%)

Hawaii	Higher Omicron Severity	543 (535 - 550)	233 (227 - 240)	38 (37 - 38)	16 (16 - 17)	39% (39% - 40%)	17% (16% - 17%)
Hawaii	Lower Omicron Severity	586 (574 - 596)	188 (179 - 199)	41 (40 - 41)	13 (12 - 14)	42% (41% - 43%)	14% (13% - 14%)
Idaho	Higher Omicron Severity	2,199 (2,184 - 2,213)	528 (517 - 540)	116 (115 - 116)	28 (27 - 28)	44% (44% - 44%)	11% (10% - 11%)
Idaho	Lower Omicron Severity	2,292 (2,279 - 2,305)	434 (425 - 444)	121 (120 - 121)	23 (22 - 23)	46% (46% - 46%)	9% (9% - 9%)
Illinois	Higher Omicron Severity	7,228 (7,159 - 7,288)	4,470 (4,409 - 4,532)	57 (56 - 58)	35 (35 - 36)	21% (21% - 21%)	13% (13% - 13%)
Illinois	Lower Omicron Severity	8,672 (8,609 - 8,736)	2,984 (2,939 - 3,036)	68 (68 - 69)	24 (23 - 24)	25% (25% - 25%)	9% (9% - 9%)

Indiana	Higher Omicron Severity	7,138 (7,069 - 7,193)	2,775 (2,713 - 2,837)	105 (104 - 106)	41 (40 - 42)	30% (30% - 30%)	12% (11% - 12%)
Indiana	Lower Omicron Severity	7,874 (7,830 - 7,914)	2,029 (1,998 - 2,064)	116 (115 - 116)	30 (29 - 30)	33% (33% - 33%)	9% (8% - 9%)
Iowa	Higher Omicron Severity	2,486 (2,461 - 2,510)	875 (852 - 896)	78 (77 - 79)	27 (27 - 28)	26% (26% - 26%)	9% (9% - 9%)
Iowa	Lower Omicron Severity	2,717 (2,698 - 2,735)	645 (631 - 660)	85 (84 - 86)	20 (20 - 21)	28% (28% - 29%)	7% (7% - 7%)
Kansas	Higher Omicron Severity	2,749 (2,717 - 2,778)	1,102 (1,073 - 1,131)	94 (93 - 95)	38 (37 - 39)	31% (30% - 31%)	12% (12% - 13%)
Kansas	Lower Omicron Severity	3,044 (3,024 - 3,064)	805 (789 - 824)	104 (103 - 104)	27 (27 - 28)	34% (34% - 34%)	9% (9% - 9%)

Kentucky	Higher Omicron Severity	6,070 (5,997 - 6,141)	2,700 (2,635 - 2,763)	135 (133 - 136)	60 (58 - 61)	36% (36% - 37%)	16% (16% - 17%)
Kentucky	Lower Omicron Severity	6,402 (6,341 - 6,456)	2,366 (2,329 - 2,413)	142 (141 - 143)	52 (52 - 54)	38% (38% - 39%)	14% (14% - 15%)
Louisiana	Higher Omicron Severity	4,073 (4,034 - 4,116)	1,796 (1,768 - 1,820)	88 (87 - 89)	39 (38 - 39)	26% (25% - 26%)	11% (11% - 11%)
Louisiana	Lower Omicron Severity	4,351 (4,299 - 4,404)	1,504 (1,468 - 1,549)	94 (93 - 95)	33 (32 - 33)	27% (27% - 28%)	9% (9% - 10%)
Maine	Higher Omicron Severity	1,025 (1,012 - 1,036)	419 (406 - 430)	75 (74 - 76)	31 (30 - 31)	41% (41% - 42%)	17% (16% - 17%)
Maine	Lower Omicron Severity	1,099 (1,089 - 1,109)	344 (334 - 353)	80 (79 - 81)	25 (24 - 26)	44% (44% - 45%)	14% (13% - 14%)

Maryland	Higher Omicron Severity	2,824 (2,781 - 2,871)	2,164 (2,116 - 2,207)	46 (45 - 47)	35 (34 - 36)	18% (18% - 19%)	14% (14% - 14%)
Maryland	Lower Omicron Severity	3,440 (3,377 - 3,494)	1,522 (1,470 - 1,580)	56 (55 - 57)	25 (24 - 26)	22% (22% - 23%)	10% (9% - 10%)
Massachusetts	Higher Omicron Severity	2,625 (2,595 - 2,651)	1,931 (1,906 - 1,959)	38 (37 - 38)	28 (27 - 28)	14% (14% - 15%)	11% (10% - 11%)
Massachusetts	Lower Omicron Severity	3,132 (3,109 - 3,154)	1,423 (1,401 - 1,443)	45 (45 - 45)	20 (20 - 21)	17% (17% - 17%)	8% (8% - 8%)
Michigan	Higher Omicron Severity	9,608 (9,539 - 9,670)	3,546 (3,489 - 3,598)	96 (95 - 96)	35 (35 - 36)	30% (30% - 30%)	11% (11% - 11%)
Michigan	Lower Omicron Severity	10,609 (10,554 - 10,651)	2,540 (2,501 - 2,578)	106 (105 - 106)	25 (25 - 26)	33% (33% - 33%)	8% (8% - 8%)

Minnesota	Higher Omicron Severity	3,616 (3,591 - 3,640)	1,350 (1,325 - 1,373)	63 (63 - 64)	24 (23 - 24)	28% (28% - 29%)	11% (10% - 11%)
Minnesota	Lower Omicron Severity	3,942 (3,923 - 3,960)	1,023 (1,004 - 1,043)	69 (69 - 69)	18 (18 - 18)	31% (31% - 31%)	8% (8% - 8%)
Mississippi	Higher Omicron Severity	3,473 (3,425 - 3,518)	1,853 (1,815 - 1,908)	118 (116 - 119)	63 (62 - 65)	26% (26% - 27%)	14% (14% - 14%)
Mississippi	Lower Omicron Severity	3,731 (3,662 - 3,800)	1,582 (1,532 - 1,652)	126 (124 - 129)	54 (52 - 56)	28% (28% - 29%)	12% (12% - 13%)
Missouri	Higher Omicron Severity	6,551 (6,490 - 6,607)	2,614 (2,565 - 2,669)	106 (105 - 107)	42 (42 - 43)	32% (32% - 32%)	13% (12% - 13%)
Missouri	Lower Omicron Severity	7,303 (7,255 - 7,345)	1,856 (1,826 - 1,883)	118 (118 - 119)	30 (30 - 31)	36% (35% - 36%)	9% (9% - 9%)

Montana	Higher Omicron Severity	1,446 (1,430 - 1,464)	351 (333 - 366)	131 (129 - 133)	32 (30 - 33)	41% (41% - 42%)	10% (9% - 10%)
Montana	Lower Omicron Severity	1,501 (1,488 - 1,518)	295 (281 - 306)	136 (135 - 137)	27 (25 - 28)	43% (42% - 43%)	8% (8% - 9%)
Nebraska	Higher Omicron Severity	1,386 (1,370 - 1,402)	562 (545 - 578)	71 (70 - 71)	29 (28 - 29)	28% (28% - 28%)	11% (11% - 12%)
Nebraska	Lower Omicron Severity	1,537 (1,526 - 1,549)	411 (399 - 423)	78 (78 - 79)	21 (20 - 22)	31% (31% - 31%)	8% (8% - 9%)
Nevada	Higher Omicron Severity	3,555 (3,530 - 3,585)	1,546 (1,521 - 1,568)	113 (112 - 114)	49 (48 - 50)	32% (32% - 33%)	14% (14% - 14%)
Nevada	Lower Omicron Severity	3,831 (3,807 - 3,856)	1,270 (1,248 - 1,294)	122 (121 - 123)	40 (40 - 41)	35% (35% - 35%)	12% (11% - 12%)

New Hampshire	Higher Omicron Severity	748 (734 - 764)	302 (285 - 316)	54 (53 - 55)	22 (21 - 23)	30% (30% - 31%)	12% (12% - 13%)
New Hampshire	Lower Omicron Severity	828 (818 - 839)	222 (210 - 232)	60 (59 - 60)	16 (15 - 17)	34% (33% - 34%)	9% (9% - 9%)
New Jersey	Higher Omicron Severity	3,605 (3,560 - 3,654)	3,501 (3,450 - 3,546)	39 (38 - 39)	38 (37 - 38)	11% (11% - 11%)	11% (11% - 11%)
New Jersey	Lower Omicron Severity	4,533 (4,453 - 4,599)	2,528 (2,463 - 2,597)	49 (48 - 50)	27 (27 - 28)	14% (14% - 14%)	8% (8% - 8%)
New Mexico	Higher Omicron Severity	2,583 (2,560 - 2,604)	868 (848 - 890)	122 (121 - 123)	41 (40 - 42)	33% (33% - 34%)	11% (11% - 11%)
New Mexico	Lower Omicron Severity	2,731 (2,712 - 2,751)	718 (703 - 736)	129 (128 - 130)	34 (33 - 35)	35% (35% - 35%)	9% (9% - 9%)

New York State (non-NYC)	Higher Omicron Severity	6,096 (6,032 - 6,155)	4,404 (4,348 - 4,469)	54 (53 - 54)	39 (38 - 39)	17% (17% - 17%)	12% (12% - 12%)
New York State (non-NYC)	Lower Omicron Severity	7,232 (7,163 - 7,290)	3,254 (3,199 - 3,319)	64 (63 - 64)	29 (28 - 29)	20% (20% - 20%)	9% (9% - 9%)
New York City	Higher Omicron Severity	2,552 (2,508 - 2,599)	2,985 (2,945 - 3,031)	30 (30 - 31)	35 (35 - 36)	7% (7% - 7%)	9% (8% - 9%)
New York City	Lower Omicron Severity	3,434 (3,376 - 3,483)	2,089 (2,047 - 2,145)	41 (40 - 41)	25 (24 - 25)	10% (10% - 10%)	6% (6% - 6%)
North Carolina	Higher Omicron Severity	8,522 (8,480 - 8,566)	3,743 (3,705 - 3,782)	81 (80 - 81)	35 (35 - 36)	30% (30% - 30%)	13% (13% - 13%)
North Carolina	Lower Omicron Severity	9,104 (9,066 - 9,144)	3,152 (3,120 - 3,185)	86 (86 - 87)	30 (30 - 30)	32% (32% - 32%)	11% (11% - 11%)

North Dakota	Higher Omicron Severity	712 (697 - 728)	233 (217 - 247)	92 (90 - 94)	30 (28 - 32)	26% (25% - 26%)	8% (8% - 9%)
North Dakota	Lower Omicron Severity	751 (723 - 762)	194 (183 - 222)	97 (93 - 98)	25 (24 - 29)	27% (26% - 28%)	7% (7% - 8%)
Ohio	Higher Omicron Severity	13,800 (13,695 - 13,899)	6,258 (6,156 - 6,363)	117 (116 - 118)	53 (52 - 54)	32% (32% - 32%)	14% (14% - 15%)
Ohio	Lower Omicron Severity	15,796 (15,678 - 15,911)	4,206 (4,116 - 4,299)	134 (133 - 135)	36 (35 - 36)	36% (36% - 37%)	10% (9% - 10%)
Oklahoma	Higher Omicron Severity	4,863 (4,782 - 4,944)	2,376 (2,296 - 2,443)	122 (120 - 124)	60 (58 - 61)	30% (30% - 31%)	15% (14% - 15%)
Oklahoma	Lower Omicron Severity	5,238 (5,178 - 5,288)	1,995 (1,962 - 2,030)	131 (130 - 133)	50 (49 - 51)	33% (32% - 33%)	12% (12% - 13%)

Oregon	Higher Omicron Severity	3,074 (3,046 - 3,096)	1,213 (1,193 - 1,236)	72 (72 - 73)	29 (28 - 29)	43% (42% - 43%)	17% (17% - 17%)
Oregon	Lower Omicron Severity	3,277 (3,251 - 3,302)	1,008 (989 - 1,028)	77 (77 - 78)	24 (23 - 24)	46% (45% - 46%)	14% (14% - 14%)
Pennsylvania	Higher Omicron Severity	11,937 (11,840 - 12,019)	5,480 (5,398 - 5,572)	92 (91 - 93)	42 (42 - 43)	26% (26% - 26%)	12% (12% - 12%)
Pennsylvania	Lower Omicron Severity	13,390 (13,307 - 13,469)	4,005 (3,946 - 4,084)	103 (103 - 104)	31 (30 - 32)	29% (29% - 29%)	9% (9% - 9%)
Puerto Rico	Higher Omicron Severity	788 (770 - 803)	811 (796 - 829)	24 (24 - 25)	25 (24 - 25)	19% (18% - 19%)	19% (19% - 20%)
Puerto Rico	Lower Omicron Severity	906 (879 - 927)	679 (653 - 706)	28 (27 - 28)	21 (20 - 22)	22% (21% - 22%)	16% (16% - 17%)

Rhode Island	Higher Omicron Severity	487 (475 - 500)	311 (297 - 323)	44 (43 - 46)	28 (27 - 29)	14% (13% - 14%)	9% (8% - 9%)
Rhode Island	Lower Omicron Severity	579 (568 - 589)	219 (209 - 229)	53 (52 - 54)	20 (19 - 21)	16% (16% - 16%)	6% (6% - 6%)
South Carolina	Higher Omicron Severity	5,378 (5,335 - 5,417)	2,426 (2,392 - 2,460)	104 (103 - 104)	47 (46 - 47)	30% (30% - 30%)	14% (13% - 14%)
South Carolina	Lower Omicron Severity	5,711 (5,665 - 5,749)	2,078 (2,045 - 2,115)	110 (109 - 111)	40 (39 - 41)	32% (32% - 32%)	12% (11% - 12%)
South Dakota	Higher Omicron Severity	674 (664 - 689)	265 (251 - 276)	75 (74 - 77)	30 (28 - 31)	22% (22% - 22%)	9% (8% - 9%)
South Dakota	Lower Omicron Severity	717 (707 - 727)	221 (212 - 230)	80 (79 - 81)	25 (24 - 26)	23% (23% - 24%)	7% (7% - 7%)

Tennessee	Higher Omicron Severity	8,581 (8,524 - 8,638)	4,033 (3,979 - 4,088)	123 (122 - 124)	58 (57 - 59)	33% (33% - 33%)	15% (15% - 16%)
Tennessee	Lower Omicron Severity	9,221 (9,149 - 9,274)	3,380 (3,335 - 3,434)	132 (131 - 133)	48 (48 - 49)	35% (35% - 35%)	13% (13% - 13%)
Texas	Higher Omicron Severity	26,451 (26,383 - 26,529)	11,854 (11,798 - 11,916)	90 (89 - 90)	40 (40 - 40)	28% (28% - 28%)	12% (12% - 13%)
Texas	Lower Omicron Severity	28,495 (28,383 - 28,592)	9,752 (9,671 - 9,858)	97 (96 - 97)	33 (33 - 33)	30% (30% - 30%)	10% (10% - 10%)
Utah	Higher Omicron Severity	1,936 (1,917 - 1,954)	609 (595 - 624)	58 (57 - 59)	18 (18 - 19)	38% (37% - 38%)	12% (12% - 12%)
Utah	Lower Omicron Severity	2,052 (2,035 - 2,068)	485 (474 - 498)	61 (61 - 62)	15 (14 - 15)	40% (40% - 40%)	9% (9% - 10%)

Vermont	Higher Omicron Severity	164 (158 - 170)	110 (104 - 116)	25 (24 - 26)	17 (16 - 18)	26% (25% - 27%)	18% (17% - 18%)
Vermont	Lower Omicron Severity	189 (184 - 195)	85 (79 - 90)	29 (28 - 30)	13 (12 - 14)	30% (29% - 31%)	14% (13% - 14%)
Virginia	Higher Omicron Severity	5,063 (5,018 - 5,109)	3,041 (2,995 - 3,084)	59 (58 - 59)	35 (35 - 36)	26% (26% - 26%)	16% (15% - 16%)
Virginia	Lower Omicron Severity	5,742 (5,675 - 5,797)	2,346 (2,297 - 2,414)	66 (66 - 67)	27 (27 - 28)	30% (29% - 30%)	12% (12% - 12%)
Washington	Higher Omicron Severity	4,262 (4,235 - 4,289)	1,973 (1,950 - 2,003)	55 (55 - 55)	26 (25 - 26)	35% (35% - 35%)	16% (16% - 17%)
Washington	Lower Omicron Severity	4,685 (4,649 - 4,714)	1,551 (1,520 - 1,582)	61 (60 - 61)	20 (20 - 20)	39% (38% - 39%)	13% (13% - 13%)

West Virginia	Higher Omicron Severity	2,685 (2,655 - 2,716)	1,091 (1,063 - 1,122)	151 (149 - 152)	61 (60 - 63)	39% (39% - 40%)	16% (16% - 16%)
West Virginia	Lower Omicron Severity	2,865 (2,843 - 2,886)	912 (893 - 932)	161 (159 - 162)	51 (50 - 52)	42% (42% - 42%)	13% (13% - 14%)
Wisconsin	Higher Omicron Severity	4,505 (4,473 - 4,533)	1,775 (1,749 - 1,803)	76 (76 - 77)	30 (30 - 31)	31% (30% - 31%)	12% (12% - 12%)
Wisconsin	Lower Omicron Severity	5,058 (5,034 - 5,081)	1,219 (1,199 - 1,243)	86 (85 - 86)	21 (20 - 21)	34% (34% - 35%)	8% (8% - 8%)
Wyoming	Higher Omicron Severity	749 (736 - 758)	143 (135 - 155)	129 (127 - 131)	25 (23 - 27)	46% (45% - 47%)	9% (8% - 10%)
Wyoming	Lower Omicron Severity	767 (756 - 774)	126 (119 - 136)	132 (131 - 134)	22 (21 - 23)	47% (47% - 48%)	8% (7% - 8%)

