The Perfect Storm of 2019: An immunological and phylodynamic analysis of Cambodia's unprecedented dengue outbreak

Cara E. Brook[1], Yimei Li[1], Christina Yek[2], Graham R. Northrup[3], Sreyngim Lay[4,5], Sophana Chea[4,5], Vida Ahyong[6], Daniel M. Parker[7], Somnang Man[4,5], Andrea R. Pacheco[4], Oum Mengheng[4,5], Fabiano Oliveira[2], Liz Fahsbender[8], Rithea Leang[5], Rekol Huy[5], Chea Huch[5], Chanthap Lon[4,5], Cristina M. Tato[6], Joseph L. DeRisi[6], Michael Boots[9], Jennifer A. Bohl[2], Jessica E. Manning[2,4*]

[1] Department of Ecology and Evolution, University of Chicago, Chicago, Illinois, USA

[2] Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, Maryland, USA

[3] Center for Computational Biology, University of California, Berkeley, California, USA

[4] International Center of Excellence in Research, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Phnom Penh, Cambodia

[5] National Center for Parasitology, Entomology, and Malaria Control, Phnom Penh, Cambodia

[6] Chan Zuckerberg Biohub, San Francisco, California, USA

[7] Department of Population Health and Disease Prevention, University of California, Irvine, California, USA

[8] Chan Zuckerberg Initiative, San Francisco, California, USA

[9] Integrative Biology, University of California, Berkeley, California, USA

*Corresponding author:

Jessica E. Manning

Email: Jessica.manning@nih.gov

Mailing address: 1 Christopher Howes Place, Phnom Penh, CAMBODIA

Abstract

The year 2019 witnessed the highest number of dengue cases ever reported globally. We

analyzed epidemiological, serological, and phylogenomic data to investigate the drivers of the

2019 epidemic in Cambodia. Using epidemiological models fit to a 19-year national dataset, we

identified an overall trend of declining annual force of infection (FOI) for dengue virus (DENV)

in Cambodia, interspersed with FOI spikes corresponding to epidemic year caseloads that

exceeded demographic predictions. We constructed time-resolved phylogenetic trees with 105

DENV genomes sequenced from the 2019 Cambodian epidemic, paired with historical Southeast

Asian data, to document the first-recorded introduction of DENV-2 Cosmopolitan genotype into

Cambodia. This introduction yielded highly localized transmission and decreased genomic

diversity when compared to endemic DENV-1, supporting the hypothesis of epidemic invasion.

Introduction of this genetically distinct lineage into a population with limited prior immunity—

paired with a spike in FOI—was a key driver of the 2019 Cambodian epidemic.

These studies were registered at clinicaltrials.gov under NCT04034264 and NCT03534245.

Introduction

Dengue virus (DENV) transmission has increased dramatically over the past two decades,

culminating in the year 2019 with the highest number of global cases ever reported (>5.2

million) to the World Health Organization (WHO)[1]. Since nearly three-quarters of DENV

infections are estimated to be clinically inapparent (thereby unreported), these counts represent a

vast underestimate of the true scale of dengue burden on public health systems[2]. DENV is a

flavivirus primarily transmitted by the *Aedes aegypti* mosquito, a ubiquitous arthropod vector in

tropical and subtropical regions[2]. DENV is comprised of four antigenically distinct serotypes—

each of which is further subdivided among four to seven distinct genotypes; infection with one

serotype is thought to result in lifelong homotypic immunity to that same serotype (homotypic

immunity) but only temporary (up to two-year) protection against different serotypes

(heterotypic immunity)[3]. Heterotypic secondary infections are often more clinically severe due to

an interaction with pre-existing flavivirus-specific antibodies known as antibody-dependent

enhancement (ADE)[4,5]. In regions with multiple circulating endemic flaviviruses, this interplay

makes dengue outbreaks difficult to predict at local, subnational, or national levels. Outbreak

forecasting is further challenged by the multifaceted disease ecology characteristic of any

arbovirus—for which infections are also impacted by changes in human land use, behavior, or

movement, as well as the distribution and abundance of arthropod vectors. Nonetheless, national

health systems in DENV-endemic regions rely on forecasting models to inform resource

allocation on a year-to-year basis—ranging from vector control to hospital capacity for

supportive care, given no available treatments for dengue. The important consequences of

forecasting predications emphasize the need for improved understanding of outbreak drivers,

particularly in resource-scarce settings.

Southeast Asia (SEA) represents ~70% of global reported dengue cases[1]. Surprisingly,

while dengue is largely classified as an urban disease, only 46% of the SEA population is

considered urban. For most SEA countries, roughly 35-55% of the total population is

concentrated in urban areas co-localizing with the highest dengue burdens; Singapore reports the

largest urbanized population percentage in SEA (100%), while Cambodia—with 16 million

people—reports the lowest (24%)[6]. Like many SEA countries, Cambodia nonetheless exhibits

significant peri-urban sprawl surrounding major metropolitan centers like Phnom Penh[7,8]. As part

of the global dengue phenomenon in 2019, Cambodia also suffered its worst dengue epidemic on

record, with approximately 40,000 total cases reported across all 26 provinces—a likely drastic

underestimation of the true disease burden[9]. Similar to other resource-scarce countries in SEA,

Cambodia hosts a national disease surveillance system that is limited to clinicosyndromic

diagnoses, and reporting is highly variable outside of the capital city. Clinical diagnoses often

correspond to more severe heterotypic secondary DENV infections[10], and in Cambodia, these are

reported collectively without the ability to systematically distinguish between serotypes.

The past two decades have witnessed explosive dengue outbreaks alongside extensive,

rapid, heterogenous development across much of SEA, with significant peri-urban sprawl

expanding beyond central mega-city limits and high-density population centers springing up

around industry hubs (often factories) in otherwise rural areas[7,8]. Longitudinal phylodynamic

studies of georeferenced DENV sequences from Thailand have demonstrated the importance of

microscale transmission—particularly at the household level—in generating DENV diversity in

urban Bangkok, predicting that DENV transmission will intensify as peri-urban settings in SEA

become better connected[11]. Previously, dengue epidemics have been linked to turnover in the

dominant, regional serotype, or—more recently—to replacement of a dominant viral genotype

with a serologically homotypic but nonetheless divergent viral lineage[12,13]. Indeed, recent work

links the magnitude of periodic dengue epidemics to antigenic evolution; large epidemics tend to

result from the takeover of those DENV lineages most antigenically distinct from previously

circulating strains of the same serotype or most closely-related to sequences of a different

serotype[13]. Investigations into the dynamical drivers of dengue transmission in rural and peri-

urban settings have been limited to date. As climatic changes drive increases in the population at-

risk for dengue infection in SEA and elsewhere[14], better-designed active surveillance programs

and improved targeting and evaluation of vector control interventions are greatly needed[15].

Here, we explore the dynamics of dengue transmission in peri-urban Cambodia,

investigating potential mechanisms underlying the 2019 epidemic, the highest year of total

reported dengue cases in the 19-year duration of the Cambodian national surveillance program.

We hypothesized that the 2019 epidemic spike in dengue case incidence could be due to either:

(a) a heightened force of infection (FOI) resulting from improved conditions for and elevated

abundance of the DENV mosquito vector, *Ae. aegypti* or (b) a constant or declining FOI coupled

with a higher proportion of susceptible individuals lacking prior immunity to a recently

introduced and antigenically novel strain of the virus or (c) some combination of the two. To

investigate these hypotheses, we queried a 19-year dataset (2002-2020) of serotype-agnostic

dengue case counts from the national surveillance program to estimate the annual FOI across this

period, then compared results at the national scale with our own febrile surveillance data in peri-

urban Kampong Speu province in 2019 and 2020. Whole genome sequencing of DENV from

serum samples during the 2019 outbreak identified the first record of the DENV-2 Cosmopolitan

lineage ever documented in Cambodia. Phylodynamic analysis of the spread of the DENV-2

Cosmopolitan lineage in this region suggests that this new genotype likely invaded a population

with limited prior immunity, driving the largest documented dengue outbreak in Cambodia

concurrent with the 2019 global epidemic.

Results

Epidemic year (2007, 2012, 2019) dengue case counts exceeded demographic predictions in

Cambodia.

Serotype-agnostic clinicosyndromic surveillance data for Cambodian dengue from 2002-
2020 demonstrated cyclic annual spikes in infection which peaked during the wet season,
typically May to October, and demonstrated three major epidemic spikes in 2007, 2012, and
2019 (Fig. 1a, 1b). To query the mechanistic drivers underpinning each annual epidemic, we first
fit a simple time series Susceptible-Infected-Recovered (TSIR) model[16–18] to the national data
(Fig. 1c, 1d)(Supplementary Text 1). The TSIR model can successfully capture the transmission
dynamics of perfectly immunizing childhood infections, such as measles, for which the
population of naïve, susceptible hosts can be reconstructed from year to year based on the annual
host population birth rate[17,18]. Though the TSIR model has been applied to dengue dynamics
previously[19–22], the dynamics of DENV infection—in particular, the role of reinfection with
secondary serotypes—are sufficiently complex such that the model is only expected to be
predictive over very short time horizons. We first sought to identify the extent to which the TSIR
model failed to predict dengue dynamics in epidemic years, highlighting a need for either a
higher FOI or a larger susceptible population to explain the data.

Fig. 1 TSIR modeling insights into epidemic dynamics of DENV in Cambodia. a National-level clinicosyndromic reported DENV cases (*dashed line*) with fitted TSIR output to the three inter-epidemic periods (2002-2006, 2008-2011, 2013-2018; *blue lines*) and TSIR projections for epidemic years (2007, 2012, 2019) under trained parameter estimates (*green lines*) and incorporating a factorial increase in the susceptible population (*purple line*) or, conversely, the FOI (*red line*). Parameter estimates are available for viewing in Table S1. **b** Biweekly transmission rate estimates across the calendar year, as recovered from TSIR fits to the three inter-epidemic periods. Solid line corresponds to mean estimated biweekly transmission, thin segments to 95% confidence intervals. **c** Annual susceptible population reconstructions for Cambodia, colored by World Bank reported birth rates[23], highlighting susceptible population increases needed to recover epidemic year case counts, assuming no factorial increase in FOI in epidemic years. **d** Reported cases vs. predicted cases from TSIR-fitted biweeks and epidemic year biweeks under trained parameter assumptions (*green*) and assumptions of either increased S (*purple*) or increased FOI (*red*).

We fit the TSIR model[16–18] separately to three discrete inter-epidemic intervals within our 19-year national time series, assuming circulation of a single major serotype in the years preceding each outbreak (respectively, 2002-2006, 2008-2011, and 2013-2018), then assessed the model's efficacy in predicting infections in the outbreak year (2007, 2012, 2019). The TSIR model predicted the timing of most annual outbreaks, though it struggled to fit the idiosyncratic dynamics witnessed in the 2013-2018 inter-epidemic period, and as expected, grossly underestimated the magnitude of the epidemic years (Fig. 1a). Transmission rate estimates for the three inter-epidemic training periods all recovered comparable seasonal dynamics, with transmission peaking annually in May, preceding the June-July annual spike in cases (Fig. 1b). We estimated the average biweekly TSIR transmission rate to be largely similar across the three inter-epidemic periods in the dataset and marginally lower for the most recent (2013-2018) inter-epidemic period than for the 2008-2011 period that preceded it, contradicting hypotheses of steadily increasing FOI driving more explosive epidemics with time.

The TSIR model effectively captured epidemic year incidence when the demographic reconstruction of the susceptible population was enhanced by a multiplicative factor of increase in the susceptible population of 1.25, 1.42, and 1.50, respectively, for the epidemic years of 2007, 2012, and 2019—or when the susceptible population was held constant as predicted from birth rates but the time-varying annual FOI was elevated by a corresponding factorial increase of 1.27, 1.45, and 1.54 (Fig. 1a). We hypothesized that a higher FOI might result from an increase in mosquito vector density and/or an increase in mosquito biting rates. While a larger susceptible population could in principle result from a boom in births[24] or a flux in immigration[25], there is no demographic support for either of these assumptions in Cambodia across the time period of study; indeed, on average, births have declined across the past two decades in Cambodia[23]—though concurrently decreasing death rates resulting from improved sanitation, better healthcare, and overall rapid socioeconomic development yielded a steadily increasing total population size

(Fig. S1)[23]. Additionally, invasion by a genetically distinct DENV lineage to which a larger

cohort of the resident population lacks prior immunity can manifest as an expansion in the

susceptible population[22]. While these dynamics might result from the introduction of a

previously unseen serotype in a given region, recent work on DENV antigenic evolution suggests

that, if sufficiently genetically distinct from prior circulating strains, a newly introduced viral

genotype could yield secondary infections of hosts previously exposed to distantly related

homotypic lineages, as well[13]. Thus, we considered the hypothesis that new serotype invasion or

genotype substitution into a cohort of "susceptible" hosts lacking prior immunity to the novel

lineage could explain the epidemic years. Prior reports from serotype-specific molecular analyses

of the 2007 and 2012 dengue epidemics in Cambodia suggest that these outbreaks can be at least

partially attributed to invasion and expansion of DENV-3 and DENV-1 lineages, respectively, in

the corresponding epidemic years[26–29].

Increasingly explosive DENV epidemics despite declining Force of Infection through time.

TSIR modeling of longitudinal DENV dynamics crystallized our two major hypotheses

for the possible factors driving recent epidemics in our system: those of elevated FOI or an

increased susceptible population due to novel antigen invasion. Clinicosyndromic dengue

surveillance data from the Cambodian national time series reported age at time of infection as

metadata, offering an opportunity to quantify changes in the mean annual FOI. We observed that,

on average, since 2010, the annual proportion of dengue case incidence in Cambodia has been

concentrated in a progressively older patient population with each passing year (Fig. 2a),

consistent with reports from Thailand, Singapore, and other SEA countries undergoing rapid

socioeconomic development and urbanization[30]. Indeed, the mean age of DENV infection in

Cambodia has increased by >.5 years per year since 2018 (Fig. 2b, Table S2).

Fig. 2 Trends in age-structured incidence and insights from FOI models. a Annual cumulative proportion of incidence by age
of patient from national-level reported dengue cases in Cambodia, binned by year and colored in progressively darker shades of
gray while highlighting epidemic years in 2007 (*orange*), 2012 (*red*), and 2019 (*brown*). **b** Mean age of DENV infection by year
across our 2002-2020 dataset with 95% confidence intervals by standard error. Data are depicted as open circles and GAM fits to
the monthly annual data are shown as red line (mean) with 95% CI in red shading. GAM results and slope derivatives by year can
be viewed in Table S2. **c** National annual per capita force of infection (FOI,) for Cambodia DENV from two-serotype, age-
structured incidence model fitted to 2002-2020 time series surveillance data. The model estimates back in time 22 years prior to
the onset of the data series, corresponding to the cohort of the oldest-age individual in the first (2002) year of the dataset. 95%
CIs by standard error are shown in gray shading, based on profile likelihood of each annual FOI while holding all other
parameters constant. FOI estimates are depicted from 1987, the first year for which reasonable confidence was established (full
time series of FOI estimates back to 1981 can be viewed in Table S3). Vertical dashed line highlights the onset of the data series.
Inset panel depicts across the data-driven period. Epidemic years are highlighted by color as in panel a. **d** Model projections
(*triangles, dashed lines*) fitted to age-cumulative incidence data (*circles, solid lines*) for each epidemic year. The full time series
of model fits can be viewed in Fig. S2.

Building beyond the simple TSIR, we next followed previously developed methods[30,31] to
construct an FOI model recapitulating the age-structured cumulative incidence for each year in
our dataset and for the 22 years preceding our time series, which corresponded to the onset of
exposure for the oldest individuals in our dataset (Supplementary Text 2). Our FOI model
assumes that reported dengue cases were derived from secondary infections (which typically
exhibit increased clinical severity prompting healthcare-seeking behavior) and that individuals
eventually become exposed to at least two serotypes in a lifetime. We used this model to

estimate the annual FOI separately for each year in the dataset under assumptions of two

simultaneously circulating DENV serotypes in the system, consistent with prior reporting of

dengue incidence in Cambodia. Historically, all four DENV serotypes have been identified in

Cambodia, but annual circulation is typically dominated by co-circulation of two serotypes and a

near-negligent proportion of cases derived from a third serotype in any given year[26].

On average, the estimated annual FOI for Cambodia DENV decreased steadily across the

19-year time series—and compared to the 22 years preceding it—but short-term, one-to-three-

year spikes were observed corresponding to the 2007, 2012, and 2019 epidemic spikes (Fig. 2c).

These results supported the hypothesis of elevated FOI—perhaps linked to changes in vector

abundance or competency—as one possible driver of the 2019 epidemic in Cambodia. Broadly,

our model effectively recaptured the data across the entire time series (Fig. S2), including in

epidemic years (Fig. 2c); the steadily decreasing annual FOI resulted in a higher proportion of

cases captured in older age classes and a slower accumulation of proportional case load with age

over time (Fig. 2d).

Regional active surveillance in the 2019 Cambodia dengue epidemic.

To better understand the drivers of the 2019 dengue outbreak in Cambodia, we enrolled

febrile patients and performed DENV RT-PCR and rapid diagnostic tests for confirmation of

DENV viremia and DENV-specific IgM/IgG antibodies, respectively. This work was carried out

in part with our ongoing active febrile surveillance study in Kampong Speu province, adjacent to

the Cambodian capital city of Phnom Penh, between July 2018 and December 2020. In total, we

screened 760 samples collected from 697 unique individuals who reported to our clinic with

fever during the study period, identifying 106 PCR-positive DENV infections. All patients

reported within a self-identified window of five days since fever onset; as such, we assumed that

any positive infections with DENV-specific IgG antibodies represented a secondary heterotypic

infection; in total, we identified 46 IgG-positive DENV infections at presentation and 57 IgG-

negative infections, which likely represented primary infections in previously DENV-naïve

individuals (no IgG test was administered at point of care for the remaining 3 positive dengue

cases). We next applied the same methods adopted for FOI estimation from age-structured

cumulative incidence data at the national level to our 2019-2020 regional data, this time fitting

our model to both the age-structured cumulative proportion of secondary infections (our prior

assumption for all nationally reported cases), in addition to the age-structured proportion of

primary cases. Resulting model fits reliably recapitulated the 2019 regional data for both

secondary and primary infections (Fig. 3a). We then compared our resulting estimates for the

2019 FOI with our prior estimates derived from the national data (Fig. 3b). Estimated FOIs for

2019 were remarkably similar between regional and national data, again showing evidence of

elevated FOI in the 2019 epidemic year. Though our regional data did not predate 2019, we were

nonetheless able to estimate FOI back in time (following the same methods applied to the

national level[30,31]) from the age profile of infections in 2019 and 2020. Amazingly, despite an

absence of data in these years, epidemic year peaks corresponding to the 2007 and 2012

epidemic years were perceptible in the regional dataset as well, though they tended to peak in the

year following the epidemic—rather than coincident with the year itself (Fig. 3b). These trends

may reflect delayed reporting to the national system, regional differences in the extent of each

epidemic across all of Cambodia, or uncertainty generated by a limited regional dataset.

Fig. 3 Fits of FOI model to 2019-2020 age-structured incidence data for DENV from our own active febrile surveillance
study in Kampong Speu province, Cambodia. a Top: cumulative proportion of multitypic infections by age of patient, binned
by year, with data shown as solid lines and circles and model projections as triangles and dashed lines; 95% confidence intervals
by standard error from profile likelihood of the 2019 force of infection ( ) estimate are shown in translucent shading around
mean projections. Bottom: proportion of primary infections by age of patient, binned by year, with data and model projections
colored as with multitypic infections. **b** Estimated Cambodia dengue per capita   from 2002-2020 based on fitting of age-
cumulative incidence model to 2019-2020 age-structured incidence data from our Kampong Speu active surveillance study (*solid*
line). 95% confidence intervals are shown by translucent shading; *dashed line* indicates national   fit for comparison. Epidemic
years are highlighted in colored points after Fig. 2. Red vertical line indicates the onset of data series for Kampong Speu cohort
model fitting (2019 and 2020 only). All FOI estimates from model fitting to 2019-2020 active febrile surveillance study can be
viewed in Table S4.

Following the elevated FOI which both national and regional data identified in epidemic
years, we next tested for the possible role of increased vector abundance or activity as a driver of
the 2019 dengue epidemic. Because the traditional entomological indicators for *Aedes* spp. (e.g.
Bretau Index, larval surveys) are not reliable predictors of dengue risk[32], we instead measured
host-vector contact via quantification of antibodies to *Ae. aegypti* salivary extract in all 760
samples at time of presentation to hospital[33]. We found no evidence of association of *Ae. aegypti*
saliva antibodies at point of care with DENV infection (Table S5), though, recently, increased
levels of *Ae. aegypti* saliva antibodies (prior to infection) were associated with increased dengue
risk in a DENV-naïve population[33]. Here, in a population with a mixed history of past exposure
to DENV, this effect may be attenuated by the presentation of more severe secondary DENV
infections. At time of infection, clear vector-associated risk factors—including habits of
mosquito net use, larvicide use, and mosquito coil use—demonstrated no association with DENV
infection status either. Though vector-driven selection has a significant role in overall
transmission potential[34], epidemic years may be more intensely driven by changes in serotype or
genotype that underpin viral evasion of existing host immunity[13]. In our global predictor model,
patient age and upper/middle vs. lower socioeconomic status were negatively correlated with
DENV positivity, highlighting the preponderance of infections in children and those in poorer
income brackets (Table S6). Household living vs. apartment living was also positively correlated
with DENV positivity, potentially indicative of the predominance of peri-urban infections
associated with more favorable *Ae. aegypti* habitats. On the whole, statistical risk factor analysis
demonstrated only limited support for the hypothesis of heightened FOI from improved vector
conditions as a driver of the 2019 dengue outbreak, despite age-structured model estimations.
These findings thus indirectly supported our alternative hypothesis that the epidemic could be
partly attributable to the introduction of a new, antigenically distinct DENV lineage in a
susceptible population. This alternate hypothesis inspired our subsequent genomics work.
302
Identification of a DENV-2 Cosmopolitan genotype introduction in Cambodia.
To interrogate the hypothesis of an antigenically novel DENV lineage introduction in
Cambodia as a potential driver of the 2019 epidemic, we sequenced and published 105 whole
DENV genomes (51 DENV-1, 51 DENV-2, and 3 DENV-4), representing one-quarter of full
genome DENV-1 sequences and nearly half of DENV-2 sequences currently available for

Cambodia—thus emphasizing the considerable challenges faced in undertaking genomic

epidemiology in resource-scarce settings[35]. Maximum likelihood phylogenetic analysis of the

resulting genomes demonstrated that most DENV-2 sequences recovered in 2019 and 2020

belonged to the DENV-2 Cosmopolitan III lineage, the first record of this genotype reported in

Cambodia; all DENV-1 sequences clustered in the Genotype 1 lineage, consistent with

previously reported genotype records in Cambodia (Fig. S3). Sequences collected from 2019

were largely split between DENV-1 and DENV-2 serotypes (60.3% vs. 39.4% respectively,

N=78), while sequences collected in 2020 were dominated by DENV-2 (20/24, 83.3%). In 2019

and 2020, respectively, 23/31 (74.2%) and 18/20 (90%) of DENV-2 sequences belonged to the

Cosmopolitan III lineage, with the remaining sequences clustering in the Asian-1 DENV-2

lineage previously reported in Cambodia.

Recent analyses of the Thailand dengue system link transitions in serotype dominance

and clade replacement of genotype sublineages within a single serotype to epidemic magnitude[13].

This work suggests that large epidemics typically result from the invasion, expansion, and

evolution of viral lineages that are very distinct from previously resident lineages of the same

serotype but (in the majority of cases) more closely related to co-circulating viruses of endemic

heterotypic serotypes, reflecting the viral fitness advantage afforded by ADE[36]. In Thailand, the

most severe epidemics were linked to clade replacement of a resident viral genotype by a more

evolutionarily fit DENV lineage within the same serotype, resulting in a 'selective sweep', which

subsequently reduced overall viral diversity and, consequently, diminished between-serotype

antigenic differences[13]. In the Cambodian epidemic spike of 2019, the Cosmopolitan genotype

introduction, followed by serotypic and genotypic homogeneity in 2020, are consistent with the

dynamics of clade replacement for an endemic DENV serotype. Clade replacement dynamics

have been previously witnessed in conjunction with epidemic outbreaks in Cambodia[12,37] and

elsewhere[22,34,38–40].

We next constructed serotype-specific Bayesian timetrees[41] from DENV sequences to

assess the divergence time of 2019-2020 lineages from sequences previously reported from

Cambodia and from neighboring SEA countries across our 2002-2020 study period (Fig. 4). The

majority of DENV-1 lineages (detailed in Fig. S4) diverged relatively recently from 2015 and

2016 Cambodian sequences recovered for this serotype (Fig. 4b), with a time to Most Recent

Common Ancestor (tMRCA) of approximately 7.6 years (MRCA at November 2012, 95% HPD:

August 2013 – April 2012). BLAST analysis indicated that this cluster of DENV-1 Cambodia

sequences demonstrated the highest identity to sequences recovered from China in 2019, while

another, rarer subset of DENV-1 sequences more closely resembled those recovered from

Thailand in 2019 and 2020[42]. By contrast, the majority of DENV-2 sequences in the

Cosmopolitan III lineage demonstrated high divergence from previously reported sequences

(only in the Asian-1 lineage) for Cambodia, with a tMRCA of approximately 83.8 years (MRCA

at 1936, 95% HPD: 1932-1941). DENV-2 Cosmopolitan sequences recovered from Cambodia in

2019 and 2020 showed the highest similarity to sequences previously derived from recent

DENV-2 Cosmopolitan outbreaks in Singapore, Malaysia, Sri Lanka, and Thailand[22,39,42].

Locally, within our regional data subset for SEA, sequences tightly clustered in both geographic

space and time demonstrated high phylogenetic relatedness (Fig. S4), consistent with previous

studies emphasizing the importance of microscale transmission dynamics for DENV[11].

351
352
Fig. 4 Bayesian timetrees highlight geospatial structuring in evolutionary relationships for Cambodian dengue. a Map of
Southeast Asia with countries colored corresponding to sequences derived from each country, as shown in tip points on
phylogenetic timetrees constructed using BEAST 2 for **b** DENV-1 and **c** DENV-2. X-axis highlights divergence times between
corresponding sequences. Historical sequences are represented as triangle tips and sequences contributed by active febrile
surveillance in this study as circles. Cambodia and corresponding sequences are shaded purple. Clade bars indicate the genotype
of corresponding sequences within each serotype: genotype-1 for DENV-1 and Asian-1 and Cosmopolitan for DENV-2. A
detailed inset of geographic localities for 2019-2020 Cambodia sequences can be viewed in Fig.S4.
360
361

Hyper-localized transmission of DENV-2 Cosmopolitan sequences in Cambodia in the 2019 outbreak.

Previous genomic analysis of geolocated DENV sequences recovered over a 16-year period in Thailand (1994 – 2010) demonstrated the importance of microscale, household-level transmission in driving DENV dynamics, particularly in the highly urbanized setting of Bangkok[11]. In Bangkok, population size was shown to be tightly, positively correlated with the number of distinct transmission chains—a measure of phylogenetic diversity—driving DENV cases for the region (Fig. 5a). In more rural settings in Thailand, this relationship was less pronounced, but the authors predicted that patterns of increasing urbanization would drive DENV diversity to more closely approximate dynamics witnessed in Bangkok. We compared patterns in DENV genomic diversity captured in geolocated sequences from 2019 and 2020 in peri-urban Cambodia with those previously investigated in Thailand (Fig. 5), building transmission chains from sequence pairs which shared a MRCA within the past six months. The number of transmission chains recovered for endemic DENV-1 in our study region closely matched that predicted by population size in urban Bangkok. For DENV-2, which showed less overall diversity than DENV-1—in keeping with a hypothesis of recent invasion—the lower number of transmission chains per population size better approximated that recovered from rural regions in Thailand (Fig. 5a). We further demonstrated a tighter coupling between phylogenetic relatedness and physical, geographic distance for DENV-2 vs. DENV-1 sequences, consistent with epidemic invasion behavior and the tight clustering of a novel antigenic introduction in space and time (Fig. 5b). Finally, prior analysis of Bangkok DENV sequences demonstrated a tighter coupling of phylogenetic relatedness and spatial distance in less mobile populations, particularly young children and women[11]. These patterns held constant for DENV-1 sequences in our peri-urban setting in Cambodia but were reversed for DENV-2 sequences, consistent with a hypothesis of a novel sequence introduction that generated secondary infections and heightened local transmission in older individuals (Fig. 5c-f).

Fig. 5 Geographic proximity structures evolutionary relatedness for Cambodian DENV. a Number of effective transmission
chains for circulating DENV estimated across populations of varying densities. Black (urban) and gray (rural) circles with
corresponding 95% confidence intervals depict estimates for Thailand from Salje et al. 2017[11], while triangles depict estimates
from our Kampong Speu active febrile surveillance study for DENV-1 (*green*) and DENV-2 (*blue*). **b** Proportion of sequence
pairs for Kampong Speu DENV-1 sequences (*green*), all DENV-2 (*dark blue*), and DENV-2 Cosmopolitan sequences (*light blue*)
which are derived from the same transmission chain (*i.e.* share a MRCA <6 months since the earlier sequence in the pair) across
progressively longer Euclidean distances separating the localities from which the sequences were recovered. Proportion of
sequence pairs within the same transmission chain for children <5 (*purple*) vs. 5+ (*green*) years of age for **c** DENV-1 and **d**
DENV-2 Cosmopolitan sequence pairs across progressively longer distances between cases. Proportion of sequence pairs within
the same transmission chain for female (*pink*) vs. male (*blue*) hosts of **e** DENV-1 and **f** DENV-2 Cosmopolitan sequence pairs
across progressively longer distances between cases.

Discussion

To date, few studies have been published on the worldwide 2019 dengue surge; further

studies of this global phenomenon are needed to forecast and mitigate future dengue epidemics.

Here, we report the novel introduction of the DENV-2 Cosmopolitan genotype, coupled with

elevated FOI, as one potential explanation for Cambodia's largest recorded dengue epidemic to

date. By contrast, recent analyses of the 2019 dengue epidemic from Brazil suggest that DENV-1

and DENV-2 circulated cryptically in-country for five years before resurging in 2019[43].

Consistent with our findings in Cambodia, the study in Brazil identified no change in mosquito

virus transmission potential to explain the 2019 resurgence, but rather attributed the DENV

epidemic to transient immune protection, new public health interventions, and behavioral

modifications arising in the wake of the Zika virus outbreak[43]. Reports from other 2019 dengue

epidemics in Bhutan and Bangladesh point to increased monsoon activity coincident with mass

movement of people for school and religious holidays as key drivers of the surge in South

Asia[44,45]—while others still highlight a role for reintroduction of previously extinct serotypes,

combined with high rainfall and insecticide resistance in these same regions[46]. Global climactic

factors may play a role in these coincident epidemics but have not yet been systematically

studied in the context of the 2019 global dengue surge.

Our identification of the invasion of the DENV-2 Cosmopolitan genotype in Cambodia,

coincident with the 2019 epidemic, echoes studies from the late 1990s in the Americas, which

demonstrated that the introduction and subsequent takeover of the DENV-2 Asian-1 genotype

were associated with higher frequencies of dengue hemorrhagic fever than had been previously

witnessed under the endemic American genotype. Indeed, the invading DENV-2 Asian-1

genotype exhibited a higher viral replication rate within the *Ae. aegypti* mosquito than did

previous American genotypes. This replication increase, combined with a shorter extrinsic

incubation period, resulted in a 65-fold increase in vectorial capacity that drove dengue

outbreaks in the Americas throughout the late 1990s[47]. Previous modeling studies have suggested

that novel genotypes may circulate at low prevalence in a host population for many years prior to

detection, expansion, and displacement of resident genotypes[48]. Our study emphasizes the

extraordinary dearth of publicly available DENV sequence data for Southeast Asia; indeed,

DENV is sequenced so infrequently in Cambodia that it is impossible to know whether 2019

truly marked the first year of DENV-2 Cosmopolitan introduction to the country, or simply the

year of intensified expansion and consequential epidemic dynamics. More broadly, our work

illustrates the importance of the combined forces of pre-existing immunity and viral genetics in

driving increasingly severe dengue epidemics. As the global burden of dengue continues to

expand, ongoing serological and genomic surveillance is needed to improve epidemic forecasting
in Southeast Asia and around the world.

Methods

Ethics

This study was approved by the National Ethics Committee on Human Research and the

National Institutes of Health (NIH) Institutional Research Board. Written informed consent was

obtained from the individual participant or the parent or guardian of the child participants

enrolled in this study. This study was registered at clinicaltrials.gov as NCT04034264 and

NCT03534245.

Enrollment and sample collection

Nursing staff at the Kampong Speu (KPS) Referral Hospital identified, consented,

enrolled, and collected demographic data from study participants. Participants, aged 6 months to

65 years of age, presented to the outpatient department with a documented fever of 38°C or

greater in the previous 24 hours. DENV infection was first screened using SD Bioline

DengueDuo rapid tests for NS1 antigen, pan-dengue IgM and IgG. Sera was collected and

processed as described elsewhere for RNA extraction and confirmatory qRT-PCR testing for

DENV-1 – 4[33]. To assess direct evidence of vector-host contact in patient sera, quantification of

total IgG to *Ae. aegypti* whole salivary gland homogenate was performed via ELISA in 96-well

Immulon® plates in duplicate via antibody binding to whole *Ae. aegypti* SGH 2ul/ug

concentration diluted in carbonate-bicarbonate buffer, reported as arbitrary ELISA units as

described in Manning et al. 2021[33].

Cambodia national data

We obtained a 2002-2020 time series of age-structured dengue cases reported at the

national level from clinicosyndromic surveillance efforts administered by the National Dengue

Control Program of the Cambodian Ministry of Health. Each of the 26 provinces in Cambodia

reports cases on a monthly basis to the national authorities who compile these reports for public

health use and reporting to the World Health Organization since inception of the national

surveillance system in 2002. Daily cases and the corresponding age and gender of each patient

were reported. We binned cases to every-two-week (biweekly) intervals for application to TSIR

modeling and summarized by annual age class within each year for application to FOI models.

TSIR modeling

We first fit a time series Susceptible-Infected-Recovered model[16–18] to the three inter-

epidemic periods (2002-2006, 2008-2011, and 2013-2018) of the national time series, in order to

highlight the extent to which the case load in each epidemic year deviated from that projected

from birth rates alone. The TSIR model leverages an input time series of case counts, births, and

total population to estimate the susceptible population and disease transmission rate at intervals

corresponding to the generation time of the pathogen in question (here, roughly two weeks[49]).

Under TSIR assumptions, the transmission rate is held constant from year to year, though

allowed to vary intra-annually to reflect seasonal dynamics corresponding to the pathogen

transmission rate. Using the R package tsiR[16], we fit the TSIR model to an input time series of

dengue case counts from the national surveillance data and an evenly distributed input vector of

birth and total population size per biweek derived from publicly available demographic data for

Cambodia published by the World Bank[23]. For each inter-epidemic period, we reconstructed the

Susceptible population from the regression of cumulative cases on cumulative births, allowing

the transmission rate (β) to vary across 26 biweekly intervals in a single year and the

homogeneity parameter (α) to vary with each model fit. Using the model trained on each inter-

epidemic period, we then projected cases in the subsequent epidemic years (2007, 2012, 2019)

while reconstructing the Susceptible population from births alone.

This first exercise demonstrated the ineffectiveness with which these assumptions were

able to recover case counts reported from epidemic years; therefore, we next sought to quantify

the increase in the assumed Susceptible count needed to capture reported cases. Previously,

Wagner et al. 2020[22] demonstrated analytically that, under short time horizons, the Susceptible

population (S') in a two-strain DENV system, in which a new secondary serotype invades a

population previously dominated by a single endemic serotype, can be approximated by the

equation:

$$S' \approx S + \frac{R_1 \rho_2 N}{I_1} \approx S^* \left[1 + \frac{\rho_2 \gamma \beta}{\mu(\gamma + \mu)} \right]$$

in which S corresponds to individuals naïve to all prior infection, while R_1 and I_1 correspond to

individuals recovered from or infected with the endemic strain only, and N and ρ_2 respectively

represent the total population size and the rate of exogenous importation of a new serotype. S^*

then corresponds to the steady-state value of the Susceptible population under assumptions of a

single endemic serotype only, while μ, γ, and β encompass the rates of birth, death, and

transmission of exogenous infections of the new serotype. Building on these assumptions, we

first calculated the factorial increase in the reconstructed Susceptible population needed to

recover case counts for the three epidemic years. We then estimated the factorial increase in FOI

that would be needed to recover the same result if the Susceptible population was held constant,

thereby highlighting the two major hypotheses under consideration.

Quantifying mean age of infection

We quantified annual variation in the mean age of DENV infection across our 2002-2020

time series using a generalized additive model (GAM) in the 'mgcv' package in R[50]. We fit a

GAM with one thinplate smoothing spline for year and one cyclic cubic smoothing spline for

month to the response variable of numerical age across our time series. This approach allowed us

to distinguish intra- and interannual seasonal patterns in our data and project model predictions

by year excluding the seasonal effects of month. To quantify the rate of change in the age of

DENV infection in Cambodia across our time series, we then calculated the derivative of our

model projection of age at annual intervals in the R package 'gratia'.

Force of Infection estimation

We next used the age-stratified, national-level surveillance data to estimate the annual

FOI for DENV in Cambodia across the 19-year time series from 2002-2020. Methods for

estimating FOI from age-stratified serological data for single-strain pathogens are well

established[51–55], and prior work has adapted these methods to account for the role of pre-existing

heterotypic immunity in DENV infection[31] and modified them for application to age-structured

incidence data, in lieu of serology[30]. We applied the model developed by Cummings et al. 2009[30]

to age-structured incidence recovered from the Cambodian national time series, assuming

reported cases to represent secondary infections and all individuals in the dataset to eventually

experience exposure to multiple DENV serotypes across their lifetimes. We allowed for a unique

FOI across each year in the time series but assumed a constant FOI across all age cohorts within

a single year; as in Cummings et al. 2009[30] and Ferguson et al. 1999[31], this method additionally

supported the estimation of FOIs that predated our data time series, albeit with diminished

confidence in inference as compared to our data-associated years. Specifically, we estimated one

FOI per year for all 22 years predating the onset of the 2002 time series and corresponding to the

year of birth cohort for the oldest individual in the first year of our dataset. Subsequently, we

applied this same model to age-structured incidence data recovered through our own active

surveillance study in Kampong Speu province in 2019 and 2020. For the latter estimation, we

computed FOI estimates 37 years back from 2019 (again, corresponding to the oldest infection in

our 2019 data), this time simultaneously fitting model projections of both age-stratified primary

and secondary cases to those identified in our datasets from IgG antibody tests at point of care.

Risk factor analysis of DENV positivity

We next sought to identify predictive variables associated with DENV positivity from our

2019-2020 active febrile surveillance study in Kampong Speu province. We first tested for a

relationship between *Ae. aegypti* saliva antibodies and DENV positivity (based on rapid test at

point of care with follow-up PCR confirmation) in 621 fevers from 580 unique patients (a subset

of individuals reported with fevers multiple times). We used a generalized linear mixed model in

the binomial family to test the association between the response variable of DENV-positive

infection (by PCR) and the predictor variable of saliva antibody titer, controlling for individual

ID. We subsequently queried the entire dataset of 760 fevers (106 of which were DENV-

positive) from 697 unique patients for any association between DENV-positive infection and a

suite of potential risk factors. We again used a generalized linear mixed model in the binomial

family, controlling for individual ID, to test the association between the response variable of

DENV-positive infection and the predictor variables of: age (numerical), sex (categorical:

male/female), housing type (categorical: house/apartment/other), socioeconomic status

(categorical: upper-middle/lower), bed net use (categorical: rarely/regularly), insecticide use

(categorical: yes/no), and mosquito coil use (categorical: yes/no).

Viral sequencing

Metagenomic Next-Generation Sequencing (mNGS) was applied to serum samples

collected from all patients reporting with symptoms in our febrile cohort study, whether or not

PCR or rapid testing determined these individuals to be DENV-positive. Briefly, pathogen

mNGS libraries were prepared from isolated pathogen RNA and converted to cDNA Illumina

libraries using the NEBNext Ultra II DNA Library Prep Kit (E7645) according to manufacturer's

instructions. Library size and concentration were determined using the 4150 Tapestation system,

Agilent, and Qubit 4 Flurometer, Invitrogen (for quantitation only). External RNA Controls 103

Consortium collection, ERCC, ThermoFisher, were used as indicators of potential library

preparation errors and for input RNA mass calculation. Samples were sequenced on a NovaSeq

(Illumina) instrument and an iSeq100 (Illumina) instrument using 150 nucleotide paired-end

sequencing. A water ("no template") control was included in each library preparation.

Raw fastq files were uploaded to the CZID portal, a cloud-based, open-source

bioinformatics platform, to identify microbes from metagenomic data (https://czid.org)[56].

Potential pathogens were distinguished from commensal flora and contaminating microbial

sequences from the environment by establishing a Z-score metric based on a background

distribution derived from 16 non-templated "water-only" control libraries. Data were normalized

to reads mapped per million input reads for each microbe at both species and genus levels. Taxa

with Z-score less than 1, base pair alignment less than 50 base pairs, NT log(1/e) less than 10 and

reads per million (rpM) less than 10 were removed from analysis. Microbial sequences from the

samples are located in the National Center for Biotechnology Information (NCBI) Sequence

Read Archive.

Construction of consensus genomes

We attempted to construct full-genome DENV sequences from any samples which were

confirmed to be DENV-positive by RT-qPCR and which generated at least one reliable contig

mapping to any serotype of DENV in the CZID pipeline. To generate these full genome

consensus sequences, we ran the ARTIC network's Nextflow consensus genome pipeline[57],

mapping each sequence to the closest GenBank accession number hit in the original mNGS run

of CZID, using a cutoff of 5 reads per nucleotide site to make a consensus call (sites with <5

reads were called as "N"). Sequences were additionally run through the CZID integrated

consensus genome pipeline, again mapping to the closest hit identified in GenBank from the

original mNGS assembly. Resulting consensus sequences from both assembly pipelines were
then aligned with reference sequences and visually examined in Geneious Prime. Raw reads
from mNGS were then mapped to each full genome contig in turn and examined manually to
determine the correct call for each base pair.

Using these methods, we generated full or near-full genome sequences for 51 DENV-1,
51 DENV-2, and 3 DENV-4 samples, representing 105 of the 106 DENV-positive patients
identified in our dataset. All contributed genomes were >10000 bps in length and had a
maximum of 90 Ns (corresponding to <1% of the DENV genome). Resulting sequences were
uploaded to NCBI as individual FASTA files under the following accession numbers: (DENV-1)
OK159935-OK159976, OL411495-OL411499, and OL412140, OL412678, and OL412703;
(DENV-2) OL414717-OL414765, OL412740, OL420733 and OL435143; (DENV-4):
MZ976858-MZ976860.

Phylogenetic analysis
We supplemented our own Cambodia sequences with all other Cambodian sequences for
DENV-1 and DENV-2 available in GenBank at the time of analysis, selecting all full or partial
genome nucleotide sequences >10000 bp in length up to a collection date of December 31, 2020
(DENV-1: tax id 11053, 160 sequences, including 51 contributed by this study; DENV-2: tax id
11060, 100 sequences, including 51 contributed by this study). We further supplemented these
Cambodia sequences with genomes collected from other major Southeast Asian countries (nine),
which were Laos, Myanmar, Malaysia, Thailand, Vietnam, Brunei, Indonesia, the Philippines,
and Singapore. All countries were represented in both the DENV-1 and DENV-2 datasets. To
avoid overrepresenting certain countries outside of Cambodia, we limited sequence selection to a
maximum of three randomly selected genomes collected per year from each available year per
country, beginning in 2002, the year in which we began our national time series.

After selection, sequences were aligned separately by serotype in the program MAFFT[58],
and the best fit nucleotide substation model for each set of data was evaluated in the program
ModelTest-NG[59]. For both DENV-1 and DENV-2, a GTR+I+G4 nucleotide was determined to
offer the best fit to the data. Using this best fit nucleotide substitution model, we next built a
Bayesian phylogenetic tree for each set of DENV genomes in the program BEAST 2[60]. We
incorporated the date of sample collection for each sequence (or the midpoint of the year of

collection if the date was not reported), and, after Salje et al. 2017[11], we specified a strict

molecular clock at a rate of 7.9×10^{-4} s/s/y[61] and a Coalescent Bayesian skyline prior in our

models. We ran Markov Chain Monte Carlo chains in BEAST 2 for 150 million iterations,

logging results every 10,000 iterations. After chains completed, we removed the initial 10% of

iterations as burn-in and evaluated parameter convergence in Tracer v1.6. We summarized

resulting phylogenetic trees in TreeAnnotator and visualized summary trees in the R package

ggtree[62].

Finally, we computed a Maximum Likelihood phylogenetic tree to illustrate the

phylogenetic placement of our new Cambodia sequences in relation to all previously described

genotypes of DENV-1 and DENV-2. For this analysis, we included all Cambodian sequences of

DENV-1 and DENV-2 available in NCBI, as well as a broadly representative subset of

sequences within all known genotypes of DENV-1 (genotypes I, II, and III) and DENV-2

(genotypes Asian-I, Asian-II, Asian-American, American, Cosmopolitan I, Cosmopolitan II, and

Cosmopolitan III). As with Bayesian timetrees, sequences were first aligned in MAFFT[58], and

the best fit nucleotide substitution model was computed in ModelTest-NG[59]. As before, we

found that a GTR+I+G4 best represented both sequence subsets; using this model, corresponding

phylogenetic trees were then constructed in the program RAxML[63] . Following best practices

outlined in the RAxML-NG manual, 20 ML inferences were made on each original alignment

and bootstrap replicate trees were inferred using Felsenstein's method[64], with the MRE-based

bootstopping test applied after every 50 replicates[65]. Resulting phylogenies were then visualized

in ggtree[62].

Estimating transmission chains from sequence data

Finally, we followed methods outlined in Salje et al. 2017[11] to calculate the proportion of

sequences within each DENV serotype, genotype, or qualitative categorization that could be

attributed to the same transmission chain on our Bayesian timetrees, defined as having a most

recent common ancestor within the past six months in the same season. We then compared the

resulting measures of evolutionary divergence against the Euclidean distance separating the

precise GPS coordinates of the collection points of each sequence pair. The reciprocal of the

proportion of sequences sharing a transmission chain corresponds to the effective number of

transmission chains circulating in a given population. Thus, to compare our estimates of

transmission chain density against those previously reported for Thailand[11], we computed the

total effective number of transmission chains observed in our 2019-2020 sequence dataset,

separately for DENV-1 and DENV-2, at the World Bank reported population density for

Kampong Speu province[23].

Data Availability

657  All genome sequence data from this study have been submitted to the NCBI Sequence

Read Archive under Bioproject ID PRJNA681566. Consensus DENV sequences are also

available in GenBank, under the following accession numbers: DENV-1: OK159935-OK159976,

OL411495-OL411499, and OL412140, OL412678, and OL412703; DENV-2: OL414717-

OL414765, OL412740, OL420733 and OL435143; DENV-4: MZ976858-MZ976860). All

bioinformatics code for the initial mNGS assembly is available at

https://github.com/chanzuckerberg/idseq-workflows and all wet lab bench protocols are updated

at

https://docs.google.com/document/d/1RtNQc1D4or_ys7OxCCBjh4SDIdy7JaI4IE7if8EkHgE/edi

t. Further information on how to use ½ reaction volumes and FastSelect® are available upon

request. Code for generation of consensus genomes is available from the Artic Network at

https://artic.readthedocs.io/en/latest/, and detailed instructions of all specific analyses undertaken

to produce the figures and data presented here are available on our open access github repository

at: https://github.com/brooklabteam/cambodia-dengue-national. All other reasonable data

requests can be made to authors directly.

<u>References</u>

1. World Health Organization. Dengue and severe dengue. *WHO Fact Sheets* (2022).

2. Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, 504–507 (2013).

3. Sabin, A. B. Research on dengue during World War II. *American Journal of Tropical*
Medicine and Hygiene **1**, 30–50 (1952).

4. Katzelnick, L. C. *et al.* Antibody-dependent enhancement of severe dengue disease in
humans. *Science* **932**, 929–932 (2017).

5. Katzelnick, L. C. *et al.* Zika virus infection enhances future risk of severe dengue disease.
Science (1979) **369**, 1123–1128 (2020).

6. statista. Cambodia: Urbanization from 2010 to 2020.
https://www.statista.com/statistics/455789/urbanization-in-cambodia/. (2022).

7.   Peng, T. N. Southeast Asia's demographic situation, regional variations, and national
685   challenges. in (eds. Singh, D. & Cook, M.) 55–82 (ISEAS Publishing, 2017).
686   doi:doi:10.1355/9789814762878-007.
687   8.   Jones, G. W. *The Population of Southeast Asia. Asia Research Institute Working Paper*
688   *Series* www.nus.ari.edu.sg/pub/wps.htm. (2013).
689   9.   Vong, S. *et al.* Under-recognition and reporting of dengue in Cambodia: A capture-
690   recapture analysis of the National Dengue Surveillance System. *Epidemiology and*
691   *Infection* **140**, 491–499 (2012).
692   10.   Burke, D. S., Nisalak, A. & Johnson, D. E. A prospective study of dengue infections in
693   Bangkok. *The American Journal of Tropical Medicine and Hygiene* **38**, 172–180 (1988).
694   11.   Salje, H. *et al.* Dengue diversity across spatial and temporal scales: Local structure and the
695   effect of host population size. *Science* **355**, 1302–1306 (2017).
696   12.   O'Connor, O. *et al.* Potential role of vector-mediated natural selection in dengue virus
697   genotype/lineage replacements in two epidemiologically contrasted settings. *Emerging*
698   *Microbes and Infections* **10**, 1346–1357 (2021).
699   13.   Katzelnick, L. C. *et al.* Antigenic evolution of dengue viruses over 20 years. *Science*
700   *(1979)* (2021).
701   14.   Messina, J. P. *et al.* The current and future global distribution and population at risk of
702   dengue. *Nature Microbiology* **4**, 1508–1515 (2019).
703   15.   Cattarino, L., Rodriguez-Barraquer, I., Imai, N., Cummings, D. A. T. & Ferguson, N. M.
704   *Mapping global variation in dengue transmission intensity. Sci. Transl. Med* vol. 12
705   https://www.science.org (2020).
706   16.   Becker, A. D. & Grenfell, B. T. TSIR: An R package for time-series susceptible-infected-
707   recovered models of epidemics. *PLoS ONE* **12**, 1–10 (2017).
708   17.   Grenfell, B. T., Bjørnstad, O. N., Finkenstädt, B. F., Monographs, S. E. & May, N.
709   Dynamics of measles epidemics: Scaling noise, determinism, and predictability with the
710   TSIR model. *Ecological Monographs* **72**, 185–202 (2002).
711   18.   Bjornstad, O. N., Finkenstadt, B. F. & Grenfell, B. T. Dynamics of measles epidemics:
712   Estimating scaling of transmission rates using a times series SIR model. *Ecological*
713   *Monographs* **72**, 169–184 (2002).
714   19.   Oidtman, R. J. *et al.* Inter-annual variation in seasonal dengue epidemics driven by
715   multiple interacting factors in Guangzhou, China. *Nature Communications* **10**, (2019).
716   20.   Rehman, N. A. *et al.* Quantifying the localized relationship between vector containment
717   activities and dengue incidence in a real-world setting: A spatial and time series modelling
718   analysis based on geo-located data from Pakistan. *PLoS Neglected Tropical Diseases* **14**,
719   1–22 (2020).
720   21.   Kraemer, M. U. G. *et al.* Inferences about spatiotemporal variation in dengue virus
721   transmission are sensitive to assumptions about human mobility: a case study using
722   geolocated tweets from Lahore, Pakistan. *EPJ Data Science* **7**, (2018).
723   22.   Wagner, C. E. *et al.* Climatological, virological and sociological drivers of current and
724   projected dengue fever outbreak dynamics in Sri Lanka. *Journal of the Royal Society*
725   *Interface* **17**, (2020).
726   23.   World Bank. Cambodia. *World Development Indicators*
727   https://data.worldbank.org/country/KH (2021).

728   24.   Finkenstadt, B. F., Grenfell, B. T. & Grenfell, T. Time series modelling of childhood
729   diseases: a dynamical systems approach. *Journal of the Royal Statistical Society, Series C*
730   *(Applied Statistics)* **49**, 187–205 (2000).
731   25.   Ferrari, M. J. *et al.* The dynamics of measles in sub-Saharan Africa. *Nature* **451**, 679–84
732   (2008).
733   26.   Vong, S. *et al.* Dengue incidence in urban and rural Cambodia: Results from population-
734   based active fever surveillance, 2006-2008. *PLoS Neglected Tropical Diseases* **4**, (2010).
735   27.   Lover, A. A. *et al.* Spatial epidemiology and climatic predictors of paediatric dengue
736   infections captured via sentinel site surveillance, Phnom Penh Cambodia 2011-2012. *BMC*
737   *Public Health* **14**, (2014).
738   28.   Ly, S. *et al.* Asymptomatic dengue virus infections, cambodia, 2012–2013. *Emerging*
739   *Infectious Diseases* **25**, 1354–1362 (2019).
740   29.   Arima, Y., Chiew, M. & Matsui, T. Epidemiological update on the dengue situation in the
741   Western Pacific Region, 2012. *Western Pac Surveill Response J* **6**, 82–89 (2015).
742   30.   Cummings, D. A. T. *et al.* The impact of the demographic transition on dengue in
743   Thailand: Insights from a statistical analysis and mathematical modeling. *PLoS Medicine*
744   **6**, (2009).
745   31.   Ferguson, N. M., Donnelly, C. A. & Anderson, R. M. Transmission dynamics and
746   epidemiology of dengue: insights from age-stratied sero-prevalence surveys. *Proceedings*
747   *of the Royal Society B* **354**, 757–768 (1999).
748   32.   Garjito, T. A. *et al.* Stegomyia indices and risk of dengue transmission: A lack of
749   correlation. *Frontiers in Public Health* **8**, (2020).
750   33.   Manning, J. E. *et al.* Development of inapparent dengue associated with increased
751   antibody levels to *Aedes aegypti* salivary proteins: A longitudinal dengue cohort in
752   Cambodia. *The Journal of Infectious Diseases* (2021) doi:10.1093/infdis/jiab541.
753   34.   Lambrechts, L. *et al.* Dengue-1 virus clade replacement in Thailand associated with
754   enhanced mosquito transmission. *Journal of Virology* **86**, 1853–1861 (2012).
755   35.   Bohl, J. A. *et al.* Discovering disease-causing pathogens in resource-scarce Southeast Asia
756   using a global metagenomic pathogen monitoring system. *Proceedings of the National*
757   *Academy of Sciences* **119**, e2115285119 (2022).
758   36.   Cummings, D. A. T., Schwartz, I. B., Billings, L., Shaw, L. B. & Burke, D. S. Dynamic
759   effects of antibody-dependent enhancement on the fitness of viruses. *Proceedings of the*
760   *National Academy of Sciences* **102**, 15259–15264 (2005).
761   37.   Duong, V. *et al.* Genetic diversity and lineage dynamic of dengue virus serotype 1
762   (DENV-1) in Cambodia. *Infection, Genetics and Evolution* **15**, 59–68 (2013).
763   38.   de Jesus, J. G. *et al.* Genomic detection of a virus lineage replacement event of dengue
764   virus serotype 2 in Brazil, 2019. *Memorias do Instituto Oswaldo Cruz* **115**, (2020).
765   39.   Wijewickrama, A. *et al.* Emergence of a Dengue virus serotype 2 causing the largest ever
766   dengue epidemic in Sri Lanka. *bioRxiv* (2018) doi:10.1101/329318.
767   40.   Suzuki, K. *et al.* Genotype replacement of dengue virus type 3 and clade replacement of
768   dengue virus type 2 genotype Cosmopolitan in Dhaka, Bangladesh in 2017. *Infection,*
769   *Genetics and Evolution* **75**, (2019).
770   41.   Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics with
771   BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* **29**, 1969–1973 (2012).

772   42.   Poltep, K. *et al.* Genetic diversity of dengue virus in clinical specimens from Bangkok,
773   Thailand, during 2018–2020: Co-circulation of all four serotypes with multiple genotypes
774   and/or clades. *Tropical Medicine and Infectious Disease* **6**, (2021).
775   43.   Brito, A. F. *et al.* Lying in wait: the resurgence of dengue virus after the Zika epidemic in
776   Brazil. *Nature Communications* **12**, 1–13 (2021).
777   44.   Tsheten, T. *et al.* Epidemiological analysis of the 2019 Dengue epidemic in Bhutan.
778   *International Journal of Environmental Research and Public Health* **18**, 1–13 (2021).
779   45.   Hossain, M. S. *et al.* Dengue in a crowded megacity: Lessons learnt from 2019 outbreak
780   in Dhaka, Bangladesh. *PLoS Neglected Tropical Diseases* **14**, 1–5 (2020).
781   46.   Ahsan, A., Haider, N., Kock, R. & Benfield, C. Possible drivers of the 2019 Dengue
782   outbreak in Bangladesh: the need for a robust community-level surveillance system.
783   *Journal of Medical Entomology* **58**, 37–39 (2021).
784   47.   Rico-Hesse, R. *et al.* Origins of Dengue type 2 viruses associated with increased
785   pathogenicity in the Americas. *Virology* **230**, 244–251 (1997).
786   48.   Lourenço, J. & Recker, M. Viral and epidemiological determinants of the invasion
787   dynamics of Novel Dengue Genotypes. *PLoS Neglected Tropical Diseases* **4**, (2010).
788   49.   Salje, H. *et al.* Reconstructing unseen transmission events to infer dengue dynamics from
789   viral sequences. *Nature Communications* 1–10 (2021) doi:10.1038/s41467-021-21888-9.
790   50.   Wood, S. N. mgcv: GAMs and Generalized Ridge Regression for R. *R News* **1/2**, 20–24
791   (2001).
792   51.   Grenfell, B. T. & Anderson, R. M. The estimation of age-related rates of infection from
793   case notifications and serological data. *The Journal of Hygiene* **95**, 419–36 (1985).
794   52.   Pomeroy, L. W. *et al.* Serotype-specific transmission and waning immunity of endemic
795   foot-and-mouth disease virus in Cameroon. *PLoS ONE* **10**, 1–16 (2015).
796   53.   Long, G. H. *et al.* Identifying the age cohort responsible for transmission in a natural
797   outbreak of Bordetella bronchiseptica. *PLoS Pathogens* **6**, (2010).
798   54.   Heisey, D. M., Joly, D. O. & Messier, F. The fitting of general force-of-infection models
799   to wildlife disease prevalence data. *Ecology* **87**, 2356–2365 (2006).
800   55.   Muench, Hugo. *Catalytic models in epidemiology.* (Harvard University Press, 1959).
801   56.   Kalantar, K. L. *et al.* IDseq-An open source cloud-based pipeline and analysis service for
802   metagenomic pathogen detection and monitoring. *Gigascience* **9**, 1–14 (2021).
803   57.   ARTIC-network. The ARTIC field bioinformatics pipeline. https://github.com/artic-
804   network/fieldbioinformatics (2021).
805   58.   Katoh, K., Rozewicki, J. & Yamada, K. D. MAFFT online service: Multiple sequence
806   alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* **20**,
807   1160–1166 (2018).
808   59.   Darriba, Di. *et al.* ModelTest-NG: A new and scalable tool for the selection of DNA and
809   protein evolutionary models. *Molecular Biology and Evolution* **37**, 291–294 (2020).
810   60.   Bouckaert, R. *et al.* BEAST 2: A software platform for Bayesian evolutionary analysis.
811   *PLoS Computational Biology* **10**, (2014).
812   61.   Sall, A. A. *et al.* Yellow fever virus exhibits slower evolutionary dynamics than dengue
813   virus. *Journal of Virology* **84**, 765–772 (2010).
814   62.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. Ggtree: an R Package for
815   visualization and annotation of phylogenetic trees with their covariates and other
816   associated data. *Methods in Ecology and Evolution* **8**, 28–36 (2017).

63. Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: A fast,
scalable and user-friendly tool for maximum likelihood phylogenetic inference.
Bioinformatics **35**, 4453–4455 (2019).
64. Felsenstein, J. Confidence limits on phylogenies: An approach using the bootstrap.
Evolution (N Y) **39**, 783–791 (1985).
65. Pattengale, N. D., Alipour, M., Bininda-Emonds, O. R. P., Moret, B. M. E. & Stamatakis,
A. How many bootstrap replicates are necessary? *Journal of Computational Biology* **17**,
337–354 (2010).
825

Acknowledgements

This research is supported by the Division of Intramural Research at the National Institute of Allergy and Infectious Diseases at the National Institutes of Health and the Bill and Melinda Gates Foundation [grant numbers OPP1211806, OPP1211841]. We thank patients and families of Kampong Speu District Referral Hospital who participated in this study. We thank members of the National Dengue Control Program not listed in the author byline and the Provincial Health Department of Kampong Speu province in Cambodia. We thank all the other employees at the Chan Zuckerberg Biohub and Chan Zuckerberg Initiative not listed in the author byline. We thank Brian Moyer and the NIAID Office of Cyberinfrastructure and Computational Biology (OCICB) for their assistance in improving the cyberinfrastructure of our Cambodian field sites.

Author contributions

CEB and JEM conceived the study. JEM, JAB, HR, RL, CH, and SC carried out the febrile surveillance study and collected samples for mNGS in 2019. VA, CMT, JLD, JAB, SC, SL, CY, OM, SM, and JEM carried out RT-qPCR to identify dengue-positive samples, as well as library preparation and mNGS for raw sequencing. Other ancillary demographic data collection and analyses were performed by RL, KS, CY, FO, and ARP. DP, CL, and JEM set up geospatial system for patients. CEB and JAB generated consensus dengue genomes for upload to NCBI. CEB carried out TSIR and FOI modeling, as well as all phylogenetic analyses, with help from YL, GN, and MB. CEB and JEM wrote the first draft of the manuscript. All authors approved and contributed to subsequent drafts of the manuscript and agree with the results presented.

Competing interests

The authors declare no competing interests.

852 **Materials & Correspondence**

853 Correspondence and request for materials should be addressed to JEM.

854