Climate, demography, immunology, and virology combine to drive two decades of

dengue virus dynamics in Cambodia

Cara E. Brook[1*], Carly Rozins[2], Jennifer A. Bohl[3], Vida Ahyong[4], Sophana Chea[5], Liz

Fahsbender[6], Rekol Huy[7], Sreyngim Lay[5], Rithea Leang[7], Yimei Li[1], Chanthap Lon[5], Somnang

Man[5,7], Mengheng Oum[5], Graham R. Northrup[8], Fabiano Oliveira[3], Andrea R. Pacheco[5], Daniel

M. Parker[9,10], Katherine Young[11], Michael Boots[12], Cristina M. Tato[4], Joseph L. DeRisi[4], Christina

Yek[3], Jessica E. Manning[3,5]

<u>**Affiliations**</u>:

[1] Department of Ecology and Evolution, University of Chicago, Chicago, Illinois, USA

[2] Department of Science and Technology Studies, York University, Toronto, Canada

[3] Laboratory of Malaria and Vector Research, National Institute of Allergy and Infectious

Diseases, National Institutes of Health, Rockville, Maryland, USA

[4] Chan Zuckerberg Biohub, San Francisco, California, USA

[5] International Center of Excellence in Research, National Institute of Allergy and Infectious

Diseases, National Institutes of Health, Phnom Penh, Cambodia

[6] Chan Zuckerberg Initiative, San Francisco, California, USA

[7] National Center for Parasitology, Entomology, and Malaria Control, Phnom Penh, Cambodia

[8] Center for Computational Biology, University of California, Berkeley, California, USA

[9] Department of Population Health and Disease Prevention, University of California, Irvine,

California, USA

[10] Department of Epidemiology and Biostatistics, University of California, Irvine, California, USA

[11] Department of Biological Sciences, University of Texas, El Paso, Texas, USA

[12] Department of Integrative Biology, University of California, Berkeley, California, USA

*Corresponding author: Cara E. Brook

Email: cbrook@uchicago.edu

Author Contributions: CEB and JEM conceived the study. JEM, JAB, HR, RL, and SC

conducted the febrile surveillance study and collected samples for mNGS in 2019-2022. VA,

CMT, JLD, JAB, SC, SL, CY, OM, SM, and JEM carried out RT-qPCR to identify dengue-positive

samples, as well as library preparation and mNGS. Other ancillary demographic data collection

and analyses were performed by RL, KS, CY, FO, and ARP. DP, CL, and JEM set up geospatial

system for patients. CEB and JAB generated consensus dengue genomes for upload to NCBI.

CEB carried out all TSIR, FOI, phylogenetic, and simulation analysis, with help from KY, YL, GN,

CR, and MB. CR and CEB extended the catalytic model to include waning multitypic immunity.

CEB and JEM wrote the first draft of the manuscript. All authors approved and contributed to

subsequent drafts of the manuscript and agree with the results presented.

Competing Interest Statement: No competing interests.

Classification: Biological Sciences / Population Biology

Keywords: dengue, genomic epidemiology, force of infection, wavelet decomposition

This PDF file includes:

46   Main Text

47   Figures 1 to 5

ABSTRACT

The incidence of dengue virus disease has increased globally across the past half-century, with highest number of cases ever reported in 2019. We analyzed climatological, epidemiological, and phylogenomic data to investigate drivers of two decades of dengue in Cambodia, an understudied endemic setting. Using epidemiological models fit to a 19-year dataset, we first demonstrate that climate-driven transmission alone is insufficient to explain three epidemics across the time series. We then use wavelet decomposition to highlight enhanced annual and multiannual synchronicity in dengue cycles between provinces in epidemic years, suggesting a role for climate in homogenizing dynamics across space and time. Assuming reported cases correspond to symptomatic secondary infections, we next use an age-structured catalytic model to estimate a declining force of infection for dengue through time, which elevates the mean age of reported cases in Cambodia. Reported cases in >70 year-old individuals in the most recent 2019 epidemic are best explained when also allowing for waning multitypic immunity and repeat symptomatic infections in older patients. We support this work with phylogenetic analysis of 192 dengue virus (DENV) genomes that we sequenced between 2019-2022, which document emergence of the DENV-2 Cosmopolitan genotype into Cambodia. This lineage demonstrates phylogenetic homogeneity across wide geographic areas, consistent with invasion behavior and in contrast to high phylogenetic diversity exhibited by endemic DENV-1. Finally, we simulate an age-structured, mechanistic model of dengue dynamics to demonstrate how expansion of an antigenically distinct lineage that evades preexisting multitypic immunity effectively reproduces the older-age infections witnessed in our data.

SIGNIFICANCE STATEMENT

The year 2019 witnessed the highest number of dengue cases ever reported, including in Cambodia, a Southeast Asian country with endemic transmission. We analyzed 19 years of national dengue surveillance data for Cambodia to demonstrate how increasing temperature and precipitation enhance spatiotemporal synchronicity in epidemic years. We document how two decades of demographic transition has increased the age of reported dengue infection and correspondingly depressed the force of infection through time. We show that introduction of a divergent DENV-2 genotype in Cambodia in 2019 likely drove repeated symptomatic infections in older-age individuals that contributed to a high burden epidemic. As climates warm, we anticipate more synchronized dynamics globally and a shifting burden of symptomatic disease into older cohorts.

MAIN TEXT

INTRODUCTION

Dengue virus (DENV) transmission has increased dramatically over the past two decades, culminating in 2019 with the highest number of global cases ever reported (>5.2 million) to the World Health Organization (WHO) (1). Since nearly three-quarters of DENV infections are estimated to be clinically inapparent (thereby unreported), these counts represent a vast underestimate of the true scale of dengue burden (2). DENV is a flavivirus primarily transmitted by the *Aedes aegypti* mosquito, an ubiquitous arthropod vector in tropical and subtropical regions (2). DENV is comprised of four antigenically distinct serotypes—each of which is further subdivided among four to seven distinct genotypes; infection with one serotype is thought to result in lifelong immunity to that same serotype (homotypic immunity) but only temporary (up to two-year) protection against different serotypes (heterotypic immunity) (3). Heterotypic secondary infections are often more clinically severe due to an interaction with pre-existing flavivirus-specific antibodies known as antibody-dependent enhancement (ADE) (4, 5). Southeast Asia (SEA) reports ~70% of dengue cases globally (1). Historically, dengue has been classified as an urban disease, though recent studies highlighting the importance of microscale transmission in generating DENV diversity in urban Bangkok predict that DENV transmission will intensify across SEA as peri-urban settings become better connected (6–8). Cambodia, at a population size of 16 million, is currently the least urbanized country in SEA (9) but exhibits significant peri-urban sprawl surrounding major metropolitan centers like the capital city, Phnom Penh (6, 7, 9). Urbanization and climatic changes are increasing the population at-risk for dengue in Cambodia and elsewhere (10–12).

Dengue was first detected in Cambodia in 1963 (13), though political instability and civil war in the late 1970s delayed formalization of a passive surveillance policy until 1980 (14). Once adopted, this policy required reporting of clinically diagnosed cases from public health centers and hospitals to the national level. In 2001, the National Dengue Control Program (NDCP) was

inaugurated in Cambodia, and in 2002, the NDCP formally implemented the WHO clinical case

definitions of dengue and its complications as criteria for the surveillance program (14, 15). The

NDCP surveillance system is largely limited to clinicosyndromic diagnoses, meaning that

documented cases often correspond to more severe heterotypic secondary DENV infections (16),

which are reported collectively without the ability to systematically distinguish serotypes (15).

Nonetheless, the NDCP also instituted some active surveillance efforts in Cambodia in 2001,

which included limited molecular testing to identify distinct serotypes at sentinel sites in four

provinces, which were expanded to 15 provinces by 2021 (14, 15).

Over the past two decades, all four dengue serotypes have been detected in Cambodia

by molecular surveillance (15), though cases have been largely dominated by one or two

serotypes in a given year—with DENV-1 and DENV-3 most common in the early 2000s and

DENV-1 and DENV-2 predominant in the last decade (15). Cambodia witnessed three major

dengue epidemics across this period–in 2007, 2012, and 2019 (15). The first epidemic, in 2007,

occurred coincidentally with a genotype replacement event in DENV-1 (17, 18), though cases

were dominated by DENV-3, marking the last year of this serotype's dominance in the region (14,

18). The second epidemic, in 2012, has been largely attributed to DENV-1 (15, 19–21), and the

third, in 2019, appears to have been driven by co-circulation of both DENV-1 and DENV-2 (15).

Consistent with the global trends (1), Cambodia suffered its worst dengue epidemic on record in

2019, with approximately 40,000 total cases reported across all 25 provinces—a likely drastic

underestimation of the true disease burden (22). The past twenty years of surveillance in

Cambodia have witnessed, on average, steadily increasing dengue incidence, coupled with a

steady increase in the mean age of reported dengue infection (15). The increased age of reported

infection has been anecdotally attributed to an aging population following demographic transition,

as similar phenomena have been documented previously in neighboring Thailand (23, 24) and in

Nicaragua (25).

Explosive periodic outbreaks are a hallmark of dengue virus disease, though the multiple

drivers of these phenomena have long been debated (24, 26–29). Seasonal climate cycles are a

strong predictor of annual cycling for many arboviruses, including dengue (10, 11), and climate

has been implicated as a possible driver of multiannual dengue periodicity, as well. In Thailand,

multiannual dengue cycles demonstrate coherence with El Niño phenomena (29), and epidemic

years exhibit more synchronized dynamics across latitudes (29), as well as higher correlation with

local temperature than do inter-epidemic periods (28). The interaction of demography and

heterotypic immunity is also thought to play a role in driving multiannual dengue cycles (24, 26,

30), which, in Thailand, show elongated periodicity as a result of declining birth rates and slower

build-up of the susceptible population over the past half-century (24). Virology also contributes to

dengue epidemics, which have, historically, been linked to turnover in the dominant regional

serotype (31–33) or to replacement of a dominant viral genotype within a single serotype (18, 34).

Genotype-specific antigenic diversity is increasingly recognized within traditional serotype

classifications for dengue (35), and recent work links the magnitude of periodic dengue epidemics

to this antigenic evolution (34, 35). Large epidemics tend to result from the takeover of DENV

lineages most antigenically distinct from previously circulating strains of the same serotype or

most antigenically similar to lineages of a different serotype (34).

Here, we explore the dynamics of dengue virus transmission across the past two

decades in peri-urban Cambodia, investigating the potential mechanisms that underly periodic

epidemics, particularly the epidemic of 2019. We queried a 19-year dataset (2002-2020) of

serotype-agnostic dengue case counts from the NDCP, aggregated at the province level, to, first,

interrogate the role that climate played in driving epidemics in 2007, 2012, and 2019, and second,

more generally, explore the impact of climate on annual and multiannual cycling across the time

series. We, third, fit catalytic models to the age-structured incidence of reported dengue disease

to estimate the annual force of infection (FOI), at the province level, across this time series (24,

36, 37), then considered the national data in the context of our own active febrile surveillance

study carried out in peri-urban Kampong Speu and surrounding provinces from 2019-2022 (38).

Whole genome sequencing of DENV from serum samples collected in our active surveillance

program, combined with additional sequences from NDCP surveillance, identified the first record

of the DENV-2 Cosmopolitan lineage ever documented in Cambodia (39). Phylodynamic analysis

of the spread of the DENV-2 Cosmopolitan lineage in this region suggests that this genotype

caused pathogenic infections in older age individuals, contributing to the largest documented

dengue outbreak on record for Cambodia in 2019. Finally, we constructed an age-structured

discrete-time, dynamical model to simulate the interplay of climate, demography, immunology,

and virology which combine to structure two decades of dengue dynamics in Cambodia.

RESULTS

Though warming temperatures correlate with increasing dengue incidence in Cambodia,

epidemic years are not consistent climate anomalies.

Because of the widely acknowledged role of climate as a driver of arboviral disease

globally (10, 11), coupled with recent work from Thailand that highlights coherence between El

Niño climate anomalies and dengue epidemics (29), we first explored the influence of changing

temperature and precipitation on dengue caseload in Cambodia. To this end, we aggregated

high-resolution temperature and precipitation data at the Cambodian province level, across two-

week intervals from 2002-2019 and sought to address the extent to which epidemic years

represented climatic anomalies over this time period; we defined 'anomalies' as any years in

which temperature or precipitation records demonstrated significant deviations from average

trends through time, as calculated using generalized additive models (GAMs) (40). Across the

time series, temperature peaked in the first half of each year, between April and July, preceding

the peak in dengue caseload (Fig. S1). Precipitation peaked in the latter half of each year,

between August and October (Fig. S2), preceding the onset of the following year's dengue

season. GAMs (40) demonstrated that, after controlling for intra-annual variation, temperature

has increased significantly across the past two decades in all provinces; no significant interannual

changes were detected for precipitation (Fig. S3-S4; Table S1). Additional GAMs and climate

data normalized into z-scores indicated that the epidemic years of 2012 and 2019 were

anomalously warm, while 2007 was anomalously cool, consistent with the observed interannual

increase in temperature (Fig. S5). Years 2015-2016, which spanned a major El Nino event in

SEA (but did not correspond to a dengue epidemic in Cambodia) (41), were also significantly

warmer than average. By contrast, precipitation showed no anomalies in the year preceding each

of the three major dengue epidemics (Fig. S6). We concluded that, while both temperature and

dengue incidence increased across our time series, epidemic years could not be consistently

distinguished by any consistently aberrant climatic profile.

Climate-informed transmission rates failed to recover epidemic dynamics in a TSIR model.

To formally interrogate the role of climate as a driver of dengue epidemics in Cambodia,

we developed a simple, climate-informed time series Susceptible-Infected-Recovered (TSIR)

model (42–44), which we fit to dengue case counts, aggregated over two week intervals at the

province level, for the three inter-epidemic periods (2002-2006, 2008-2011, and 2013-2018).

Typically used to describe the transmission dynamics of perfectly immunizing childhood

infections, such as measles (43, 44), TSIR has been applied to dengue dynamics previously and

offers an effective means by which to isolate the impact of climate on transmission, despite

oversimplifying the multiple serotype dynamics of dengue virus disease (45–49).

We used biweekly transmission rates recovered from TSIR fits to the inter-epidemic

periods, as well as climate-informed transmission rates for epidemic years projected from

province-specific lagged temperature and precipitation data, to predict epidemic-year cases (Fig.

1; *Methods; SI Appendix*). We found that TSIR successfully recaptured the timing of annual

dengue epidemics across the 22 provinces considered, with the recovered transmission rate

peaking between May and August, slightly preceding reported cases. The magnitude and timing

of transmission varied by province and between the three inter-epidemic periods, showing no

consistent pattern of directional change in magnitude with time (Table S2). TSIR-estimated

transmission was significantly positively associated with higher temperature and precipitation

(lagged such that climate variables preceded transmission by a median 3.5 months for

temperature and 1 month for precipitation) in the corresponding province across all inter-epidemic

periods (Table S3-S4). More rapid transmission gains were observed for corresponding increases

in temperature vs. precipitation (Fig. S7-S9; Table S4). Nonetheless, climate-informed

transmission rates for epidemic years were not substantially different from rates fitted to inter-

epidemic periods, reflecting the absence of major climate anomalies across the time series (Fig.

S10).

Figure 1. Climate-informed TSIR insights into epidemic dynamics of DENV in Cambodia. Inset panels
show province level clinicosyndromic reported DENV cases (*dashed black lines*) with fitted TSIR output to
the three inter-epidemic periods (2002-2006, 2008-2011, 2013-2018; *blue lines*) and TSIR projections for
epidemic years (2007, 2012, 2019) under trained biweekly transmission rate (β) estimates (*pink lines*),
incorporating a factorial increase in the susceptible population (*gold lines*), using a climate-projected β
estimated from lagged temperature and precipitation data by province (*green lines*), or using both climate-
projected β and a factorial increase in the susceptible population (*red lines*) (*SI Appendix,* Table S2, S5).
The center map shows province level administrative boundaries for Cambodia, shaded by the mean
biweekly temperature in 2020.

As expected, TSIR underpredicted the magnitude of epidemic year caseloads for all

provinces; climate-informed transmission improved TSIR projections but still failed to recover

epidemic peaks (Fig. 1). Experimentally, we amplified the susceptible population input at the

beginning of each epidemic year in our model, which facilitated recovery of DENV caseloads

(*Methods*) (49). In the absence of climate input into the transmission rate, the proportional

increase in susceptible population at the beginning of each epidemic year needed to recover the

epidemic peak was a median 1.8x for 2007, 2.10x for 2012, and 2.17x for 2019 (Table S5). The

proportional increase in the susceptible population needed to recover the epidemic peak was not

significantly lower when epidemic year cases were projected using climate-informed transmission

rates for 2007 and 2012 (paired student's t-test: [2007] t=0.33, p>0.1; [2012] t=-0.52;p>0.1) and

only marginally lower for 2019 (t=1.31, p=0.1), suggesting (at most) a minimal role for

temperature and precipitation in driving epidemic dynamics (Table S5). For many provinces, the

combination of climate-informed transmission rate with susceptible augmentation still fell short of

effectively capturing epidemic caseloads, highlighting the need for alternative explanations for

these high transmission years.

Wavelet analysis showed enhanced synchrony in dengue dynamics across provinces and

climate time series in epidemic years.

Though TSIR modeling failed to recover a clear role for climate in driving major dengue

epidemics in Cambodia, the analysis did highlight a critical influence of climate on the intraannual

timing of cases and the interannual periodicity of dengue dynamics. Prior work in Thailand has

demonstrated enhanced synchronicity between dengue incidence and climate, and between

geographically distant localities, during major epidemics (28, 29), as well as a link between

multiannual climate cycles and higher burden epidemics (29). We hypothesized that these

multiannual climate cycles not considered in TSIR might also be important in the Cambodia

system. To this end, we undertook wavelet decomposition of the biweekly dengue time series and

the corresponding climate data, both at the province level, to extract and disaggregate annual

and multiannual trends. We found elevated amplitude and statistically significant peaks in

average wavelet power of both annual (Fig. 2A, S11A) and multiannual (Fig. 2B, S11B) cycles in

dengue incidence during the three epidemic years (50). Elevated amplitudes for multiannual

cycles were also observed during the 2015-2016 El Niño event in more northerly provinces (Fig.

2B), including those abutting Thailand, where associations between dengue and El Niño have

been previously established (29). We additionally observed significantly elevated synchronicity

between provinces in the timing and amplitude of yearly dengue incidence across epidemic

years, as measured both by pairwise Pearson's correlation coefficient (Fig. 2C) and cross-

wavelet power (Fig. S11). Synchronicity in the annual incidence data was negatively associated

with geographic distance between provinces, though patterns were less clear than have been

described elsewhere (29) (Fig. S12). High synchronicity between annual case data for paired

provinces was significantly positively associated with high temperature, precipitation, and

population size of a focal province (Fig. S12; Table S6).

300
301

302
Figure 2. Wavelet reconstructions show heightened synchrony in epidemic years. Reconstructed **A**
annual and **B** multiannual dengue cycles by province, by year from NDCP incidence per 100,000 population.
C Mean pairwise Pearson's correlation coefficient (ρ) for annual dengue incidence between focal province
and all other provinces through time. **D** Mean ρ comparing province-to-province reconstructed multiannual
dengue cycles across a 5-year sliding window, with overlapping window frames plotted (partially translucent)
atop one another. In all panels, epidemic years are highlighted by vertical red or black bars. Top panels
indicate the distribution of corresponding values (median = solid line; max-to-min range = gray shading)
observed across all provinces within each timestep. X-axis labels are marked on January 1 of the
corresponding year.

For multiannual cycles, we observed a trend of steadily increasing synchronicity (Fig. 2D)

and cross-wavelet power (Fig. S11D) between provinces through time, though no peaks occurred

in epidemic years. Cross-wavelet power between raw dengue incidence and the corresponding,

province-level mean temperature and total precipitation (grouped biweekly) also peaked in

epidemic years (Fig. S13 AB). At the multiannual scale, cross-wavelet power between province-

level reconstructed dengue cycles and temperature and precipitation largely increased across the

time series (Fig. S13CD), though values were highest slightly preceding the 2019 epidemic and

overlapping the 2015-2016 El Niño event (41). The same pattern was observed when comparing

monthly reconstructed multiannual dengue cycles and the Oceanic Niño Incidence (ONI) (Fig.

S13E). For temperature, precipitation, and ONI, cross-wavelet power with multiannual dengue

cycles peaked earlier in the time series in more southern provinces and moved gradually

northward (Fig. S13C-E). All told, wavelet analyses suggested a role for climate in synchronizing

annual dengue epidemics across provinces, but only minimal support for a hypothesis of

multiannual climate cycles driving periodic epidemics across our time series. In contrast to reports

from two decades prior in Thailand (24), the periodicity of multiannual dengue cycles across

provinces did not increase in duration throughout our time series, despite decreasing birth rates

(Fig. S14). This perhaps reflects the relative slowdown in both birth and death rate declines in

Cambodia over the past two decades, when compared with the dramatic declines witnessed

twenty years prior (Fig. S14).

Mean age of reported dengue infection increased across the study period, corresponding

to a declining force of infection.

The muted effects of climate on our epidemic time series led us to next investigate

demographic and immunological drivers of dengue dynamics in Cambodia. Previous work has

documented a trend of increasing mean age of reported dengue infection across the past two

decades in Cambodia at the national level (15); we confirmed this to be consistent at the province

level (Fig. 3A,B; Fig. S15; Table S7). Nationally, the mean age of reported dengue infection

increased from 6.79 years (95% CI: 5.85-7.72) in 2002 to 10.34 years (95% CI: 9.41-11.27) in

2020 (p<0.001). Increases in age of reported cases were even more dramatic in more remote

provinces: in the distant northeastern province of Mondul Kiri, for example, the mean reported

age of infection increased from 3.72 years (95% CI: 1.97-5.47) in 2002 to 21.18 years (95% CI:

20.84-21.53) in 2020 (Fig. 3A; Fig. S15; Table S7).

Figure 3. Demographic transition underpins declining force of infection and increasing age of reported dengue incidence in Cambodia. A Mean age of reported dengue infection, by province in the last year of the NDCP time series (2020). **B** Age distribution of reported dengue cases by year, with violin plots highlighting changes in the interquartile range of cases. The interannual trend in the mean age of dengue infection is plotted as a solid red line, with 95% confidence intervals by standard error shown as a narrow, translucent band behind it (Table S7). Epidemic years (2007, 2012, 2019) are highlighted by a light blue, dashed line in the background. **C** National (black) and province level (colored) estimates for the annual, per serotype, per capita force of infection from 1981 to 2020. 95% confidence intervals from the hessian matrix are shown as translucent shading. FOI estimates are compared against national birth and death rates for Cambodia across the time series, with epidemic years highlighted by vertical dashed lines. **D** Age modifiers to the FOI fit as shared across all provinces for 2002-2010 and 2011-2020 subsets of the data, with 95% confidence intervals from the hessian matrix are shown as translucent shading. **E** Cumulative increase in the proportion of cases reported by age at the national level, colored by year. Data are shown as dotted lines and model output as solid lines. Model includes national FOI estimates from **C**, age modification terms from **D**, and time-varying waning multitypic immunity as shown in the inset.

To interrogate the mechanistic drivers of this pattern, we estimated the annual force of

infection (FOI, or λ), the rate at which susceptibles become infected, by fitting a catalytic model

with multiple serotype exposures to the province level data for every year in the dataset (2002-

2020) and the 22 years preceding the onset of the time series, dating back to the birth year of the

oldest individual in the first year of the data (Fig. 3C) (24, 36, 37). As in previous models for

dengue (24, 36), we assumed that reported cases represented secondary, symptomatic

infections most likely to report to public hospitals and clinics. Resulting patterns in FOI by

province largely mirrored those recovered nationally (Fig. S16), demonstrating a consistently high

FOI in the early 1990s, corresponding to the years in which individuals born in Cambodia's 1980s

birth pulse (which followed severe population reductions from civil war in the 1970s) would likely

experience secondary infections. The FOI subsequently demonstrated a gradual decline across

the time series, interspersed with local peaks in epidemic years and in 2010 and 2015.

Age-structured FOI modifications and waning multitypic immunity in 2019-2020 improved

the catalytic model's ability to recapture observed data.

We next fit 19 multiplicative age modification parameters, shared across all provinces, to

allow for modulation of annual FOI across different age categories; eight age class were

considered for the first half of the time series and eleven for the second, corresponding to the

expanded age range of reported cases (Fig. 3D). Incorporation of age modifiers improved model

fits to the data across all provinces (Table S8). Consistent with recent work in Thailand (23), we

identified a high hazard of infection in adolescents (13-15 year-olds) across the time series. In the

second decade, we additionally noted an elevated age-specific hazard of infection in 30-39 year-

olds and >50 year-olds, corresponding to case reports in older cohorts.

Because the age range of these older cases greatly exceeded average age increases

predicted by declining FOI, we reconsidered our original assumption by which reported cases

corresponded to clinically apparent secondary infections only. Visualization of the age distribution

of reported cases by year at both the national (Fig. 3B) and province levels (Fig. S15) indicated

that, in addition to increases in the mean reported age of dengue infection, the past two decades

of dengue incidence in Cambodia have also witnessed expansion in the age *range* of reported

cases—such that approximately 1% of reported cases (712/68597) in 2019 occurred in

individuals over the age of 45, with 61 infections reported in individuals >70 years in age. We also

observed an isolated spike in the age distribution of reported cases in some provinces in the

epidemic years of 2007 and 2012 (Fig. S15). To recapture these symptomatic infections in older

age-individuals, we modified the multitypic catalytic modeling framework presented in previous

work (24, 25, 36) to permit individuals to wane from a state of multitypic immunity back to that of

monotypic immunity subject to renewed hazard of symptomatic re-infection (*SI Appendix*). Using

this new modeling framework, we estimated a shared parameter across all provinces, which

corresponded to the rate of waning multitypic immunity (σ). When we allowed σ to vary by year,

we estimated a significant signature of waning multitypic immunity in the 2019 and 2020 data only

(Fig. 3E, *inset*). Incorporating time-varying σ improved model fit when added both to the FOI-only

model and the FOI model modified by age class (Table S8). Our best fit FOI model including both

age modifications and time-varying multitypic waning immunity effectively recaptured the

observed cumulative proportion of cases per age class per year at national (Fig. 3E) and province

(Fig. S17) levels. In general, cases accumulated more slowly across age classes with advancing

years, tracking declining FOI.

DENV genotyping from 2019-2022 suggests a likely DENV-2 clade replacement event in

Cambodia.

The results of our FOI analysis led us to question whether regional turnover in the

dominant DENV virus lineage could underpin the 2019 signature of immunological nativity

identified in our data. Prior work emphasizes the key role that antigenic novelty, influenced by

both serotype and genotype, plays in determining fitness advantages for co-circulating DENV

variants (34, 35). To address this question in our system, we leveraged serum samples amassed

during an active febrile surveillance study that we carried out in Kampong Speu province,

adjacent to the Cambodian capital city of Phnom Penh, between July 2018 and December 2022

(38, 39). In this study, serum collected from patients reporting to the provincial referral hospital

within 24 hours of fever was subject to RNA extraction and metagenomic Next Generation

Sequencing (mNGS) to identify the etiology of infection (38, 39). We successfully identified

serotype and corresponding genotype from mNGS data for 239 DENV-positive Kampong Speu

cases. We supplemented these with samples collected from 11 cases identified in our more

limited active surveillance in five neighboring provinces in 2021-2022 and 20 cases identified in

routine NDCP surveillance across six provinces, also in 2021-2022 (Fig. 4A; Table S9). From this

data subset of positive DENV samples, we sequenced and submitted 192 whole DENV genomes

to NCBI (63 DENV-1, 120 DENV-2, and 9 DENV-4; Table S9), representing one-third of full

genome DENV-1 sequences and two-thirds of full genome DENV-2 sequences currently available

for Cambodia.

Figure 4. Bayesian time trees highlight geospatial structuring in evolutionary relationships for Cambodian dengue. A Map of Cambodia with locations of DENV serum samples genotyped from 2019-2022 in part with this study. Cases are grouped together within 10km radii. The centroid of each case cluster is plotted, with circle size corresponding to sample number and shading corresponding to serotype and genotype (DENV-1=green; DENV-2 Asian 1= light blue; DENV-2 Cosmopolitan = navy; DENV-4=orange). The age distribution of cases by serotype and genotype is shown in the upper left; DENV-2 Cosmopolitan and DENV-4 infections occurred in significantly older age individuals than reference DENV-1 infections

(linear regression; p<0.001; Table S10). **B.** Number of genotyped DENV cases by year from febrile
surveillance in this study, colored by serotype and genotype as showing in panel A. **C** Map of Southeast
Asia with countries colored corresponding to sequences, as shown in tip points on phylogenetic timetrees
constructed using BEAST 2 for DENV-1 and **D** DENV-2. X-axis highlights divergence times between
corresponding sequences. Reference sequences from GenBank are represented as circle tips and
sequences contributed by active febrile surveillance in this study as triangles. Cambodia and corresponding
sequences are shaded purple. Clade bars indicate the genotype of corresponding sequences within each
serotype: genotype-1 for DENV-1 and Asian-1 and Cosmopolitan III for DENV-2. See Fig. S19 for a detailed
inset of geographic localities for 2019-2022 Cambodia sequences. **E** Proportion of geolocated sequence
pairs from panel A for DENV-1 (*green*) and DENV-2 (*blue*) genomes derived from the same transmission
chain across progressively longer Euclidean distances. **F** Number of effective transmission chains for
circulating DENV estimated across populations of varying densities. Black (urban) and gray (rural) circles
with corresponding 95% confidence intervals depict estimates for Thailand from Salje et al. 2017 (8), while
triangles depict estimates from our Kampong Speu active febrile surveillance study for DENV-1 (*green*) and
DENV-2 (*blue*).
451

Maximum likelihood phylogenetic analysis of the resulting genomes demonstrated that

most DENV-2 sequences recovered from our 2019-2022 study belonged to the DENV-2

Cosmopolitan III lineage, the first record of this genotype in Cambodia, though its emergence has

been reported recently in several neighboring SEA countries (Fig. S18) (51–54). All DENV-1

sequences clustered in the Genotype 1 lineage, consistent with previous genotype records in

Cambodia (Fig. S18). Genotyped cases from 2019 were largely split between DENV-1 (N=56/93,

60.2%) and DENV-2 (N=35/93, 37.6%) serotypes, while those from 2020-2022 were dominated

by DENV-2 (2020: N=25/32, 78.1%; 2021: 25/28, 89.3%; 2022: 107/117, 91.4%) (Fig. 4B). A

small number of DENV-4 cases were also sequenced across the time series (Fig. 4B). The

Cosmopolitan III lineage of DENV-2 became increasingly prevalent across the time series, as

compared with the Asian 1 genotype previously recorded in Cambodia (Fig. 4B). The

Cosmopolitan lineage increased in prevalence from 74.3% (26/35) of DENV-2 cases in our study

in 2019 to 99.1% (106/107) of cases in 2022 (Fig. 4B). We additionally observed that the mean

age of infection was significantly higher for DENV-2 Cosmopolitan cases genotyped in our study

(while the mean age of DENV-2 Asian 1 infection showed no difference), as compared with

reference DENV-1 case (Fig. 4A; Table S10, p<0.001). Mean age of infection was also

significantly elevated for DENV-4, likely because it circulated at much lower prevalence.

We constructed serotype-specific Bayesian timetrees (55, 56) from full genome DENV

sequences to assess the divergence time of 2019-2022 lineages from sequences previously

reported from Cambodia and neighboring SEA countries between 2002-2022 (Fig. 4C-D; Table

S9). The majority of DENV-1 lineages recovered from our study in 2019-2022 clustered closely

with other DENV-1 sequences reported to NCBI from Cambodia across the past decade (Fig.

4C). By contrast, DENV-2 sequences in the Cosmopolitan III lineage demonstrated high

divergence from the most recently reported Asian-1 sequences for Cambodia, with a time to most

recent common ancestor (tMRCA) of approximately 77.1 years (Fig. 4D; MRC: 1945, 95% HPD:

1942-1949). Locally, within our regional data subset, sequences collected from cases coincident

in geographic space and time were highly phylogenetically related (Fig. S19), supporting previous

studies emphasizing the importance of microscale transmission for DENV (8). Nonetheless, we

observed reduced phylogenetic diversity overall and a higher degree of phylogenetic relatedness

across wider geographic distances for DENV-2 vs. DENV-1 sequences, consistent with a

hypothesis of recent genotype invasion (Fig. 4E). Indeed, some 60% of paired DENV-2

sequences separated by up to 400m of geographic distance were estimated to share a MRCA

within six months of the onset of the earlier case (corresponding to a shared 'transmission chain'),

as compared with only 22% of DENV-1 sequences. At a distance of up to 1 km separation, some

40% of DENV-2 pairs still belonged to the same transmission chain, in contrast to only 13% of

DENV-1 sequences. Specifically for the Kampong Speu data, we additionally estimated the

number of distinct DENV-2 vs. DENV-1 effective transmission chains in circulation in the region

(Fig. 4F). When compared with prior work from neighboring Thailand, the number of effective

transmission chains per population size for DENV-1 in Kampong Speu closely matched that

predicted by population size in the endemic transmission setting of urban Bangkok (8). By

contrast, the lower number of transmission chains per population size estimated for DENV-2 was

closer to that previously observed in rural regions in Thailand with less continuous endemic

transmission (Fig. 4F).

All told, our phylogenetic analyses describe the emergence of a previously unrecorded

genotype of highly divergent DENV-2 in at least ten different Cambodian provinces. This

genotype greatly increased in prevalence following its first record in Cambodia in 2019 and

caused infections in older-age individuals. The limited phylogenetic diversity observed from

DENV-2 sequences suggest that the Cosmopolitan lineage arrived relatively recently to the

region. Its high genetic divergence from previously reported genotypes in Cambodia may be

associated with antigenic novelty that played a role in the 2019 epidemic.

Mechanistic simulations of clade replacement with antigenic escape in an age-structured

model recapture patterns witnessed in the data.

505   Combined, our FOI analysis and genotyping suggest that the DENV-2 Cosmopolitan

lineage caused a genotype replacement event in Cambodia, which likely played a part in the

2019 epidemic in the region. The higher burden of cases in older age individuals more likely to

have some form of pre-existing dengue immunity additionally suggested that the Cosmopolitan

genotype may be sufficiently antigenically divergent to overcome this immunity, resulting in

symptomatic infection. To formally evaluate this hypothesis, we constructed a mechanistic, age-

structured discrete time deterministic epidemic model in biweekly timesteps (*SI Appendix*) (57–

59), with which we simulated diverse mechanisms hypothesized to underpin observed patterns in

our data. We sought to explain both the general pattern of increasing average age of reported

infection through time, as well as the high caseload and wide distribution in age range of cases

associated with the 2019 epidemic (and, to a lesser extent, prior epidemic years). Previous work

in Thailand shows similar patterns of drastically older age infections in more recent years, which

the authors explain via a mechanism of increased detectability of tertiary or quaternary infections

through time (23). In practice, this mechanism is similar to our extension of Ferguson et al.

(1999)'s catalytic model for dengue, which incorporates waning multitypic immunity, as both

approaches allow for case detection in individuals experiencing more than two dengue infections

across their lifetime. We used our simulation model to evaluate whether these two mechanisms

were functionally distinguishable and, if so, which best replicated Cambodian data.

523   For simplicity, we restricted ourselves to a three-serotype system and initiated

simulations incorporating seasonal variation in transmission as estimated by TSIR (Fig. 1) and

annual variation in FOI corresponding to national estimates (Fig. 3). As with our FOI analysis, we

modeled secondary infections as reported cases for (H0) normal demographic simulations. We

then compared these results against simulations of (H1) hypotheses that allowed for continuously

increasing detectability of tertiary cases through time or (H2) hypotheses of genotype

replacement paired with waning multitypic immunity and detectable reinfection within a serotype

(Fig. 5). We also compared (H3) hypotheses of genotype invasion with no lineage-specific

changes in immunology or (H4) hypotheses of genotype invasion with continuous increasing

tertiary case detection, again not specific to the invading lineage (Fig. S21). We focused our

analyses on drivers of the 2019 epidemic but also conducted a subset of simulations to highlight

insights gleaned from the 2007 epidemic in Cambodia (Fig. 5, *bottom*).

536
Figure 5. Simulations of genotype clade replacement recapitulate observed expansion in the age
structure of reported dengue cases from Cambodian data. A Total reported cases (solid line = mean
FOI; translucent shading = 95% confidence interval for FOI), **B** age distribution of reported cases by year
(black = secondary; blue = repeat secondary or tertiary), and **C** cumulative proportion of reported cases by
age, from deterministic model simulations indicated to the left of panel A.

All simulations successfully recovered the pattern of increasing mean age of reported

infection, as witnessed in the data. This increase was steeper under (H1) assumptions of

increasing detectability for tertiary infections (Fig. 5). Because we parameterized FOI from the

national time series for Cambodia, for which the 2019 FOI was locally elevated (Fig. 3), all

simulations also effectively reproduced a higher burden of cases in 2019. Critically, we found that

sudden, sharp expansions in the age distribution of reported cases, as witnessed broadly across

provinces in epidemic years for Cambodia (Fig. S15), were only possible when modeling some

form of antigenic novelty for the invading lineage (H2; Fig. 5). Simulations of this novel antigenic

invasion in 2007 (Fig. 5 H2, *bottom*) also demonstrated how the expanded age distribution should

subsequently narrow in the years following clade replacement with immune evasion, again

consistent with Cambodian data (Fig. S15). By contrast, (H1) hypotheses of constant increasing

tertiary case detection through time will only continue to expand this distribution over longer time

horizons. Importantly, modeling of genotype turnover had no demonstrable impact on the age

distribution of cases in the absence of *specific* enhanced detectability for the invading clade, even

when allowing for generally heightened tertiary case detection through time (Fig. S21). Thus, we

conclude that only introduction of antigenically divergent genotypes (or, of course, serotypes) can

be expected to generate this dramatic short-term expansion in the age distribution of cases.

Notably, our results were agnostic to classification of repeat infections with an invading lineage as

tertiary detections (e.g. detection of DENV-2 infection following prior exposure to DENV-1 and

DENV-4) vs. repeat detections within a serotype (e.g. detection of DENV-2 following prior

exposure to a previous lineage of DENV-2). The epidemic consequences of this invasion were

comparable, so long as enhanced detectability (e.g. antigenic escape) was specific to invading

lineage.

DISCUSSION

Drawing from two decades of national surveillance data and a more recent genomic

cohort study, we queried the mechanisms that underly dengue virus transmission and drive

periodic dengue epidemics in Cambodia. All told, our study highlights the complex interplay of

climate, demography, immunology, and virology that dictates the dynamics of dengue virus

disease in endemic settings.

572   Our investigations of climate effects on dengue transmission in Cambodia mirror those

previously reported in Thailand (28, 29), Sri Lanka (49), and China (45), emphasizing the role of

temperature, and—to a lesser extent—precipitation, in synchronizing annual dynamics in

epidemic years. Consistent with prior work that highlights a role for El Niño in driving multiannual

dengue cycles in other systems (29), we observed synchrony between reconstructed multiannual

dengue cycles in Cambodia and the ONI; however, this synchrony appeared to peak during the

robust 2015-2016 El Niño, which did not correspond to one of the three major dengue epidemic

years in our dataset—though some signature of elevated amplitude for multiannual dengue

cycles in northern Cambodia (adjacent to the Thai border) suggest a local epidemic may have

occurred during this time. The muted impact of climate on dengue dynamics in Cambodia could

reflect overall higher caseloads in association with elevated temperatures in more recent years

(28), making any further transmission gains comparatively less dramatic. In addition, climate

impacts on DENV transmission may be generally less discernable in Cambodia, which has

reduced latitudinal variation (and corresponding climate differences) than other SEA countries,

such as Thailand. Nonetheless, our analysis suggests that warmer temperatures play a role in

driving epidemic dynamics for Cambodian dengue and likely contributed to high caseloads in

2019. As temperatures increase globally, dengue transmission is likely to accelerate, and

variation in both timing of annual dengue transmission and multiannual peaks in caseload may

become more homogenous.

591   In addition to climate, our data and corresponding analysis emphasizes the importance of

human demography and prior immunity in structuring dengue transmission globally. Consistent

with studies conducted elsewhere (23–25), we observed a pattern of increasing mean age of

reported dengue infection that we effectively recaptured by modeling a declining FOI through

time—which we attributed to declining birth and death rates consistent with Cambodia's

demographic transition. Certainly, other mechanisms of demographic change (e.g. immigration

and emigration) could also impact infection dynamics. Cambodia experienced a net negative

migration rate across the time series investigated (60); as the majority of Cambodian migrants are

young adults seeking employment in neighboring Thailand (61), it is possible that their emigration

could lower the dengue FOI even further. Though the Cambodian migrant population comprises

<0.5% of the total national population (60), migration rates can be 10-fold higher in border

provinces and could have correspondingly elevated impacts in these regions (61).

The declining FOI through time is somewhat counterintuitive considering recent explosive

dengue epidemics in Cambodia and across the globe. Nonetheless, we provide evidence for a

high frequency of reported cases in older age individuals that underlies the high case burden in

epidemic years. Our study is unique from previous investigations in recognition, not only of

escalation of the mean age of reported dengue infection, but also of the expanding age range of

reported disease. Indeed, symptomatic individuals >70 years in age during the 2019 epidemic are

difficult to reconcile under assumptions by which reported cases correspond to clinically apparent

secondary infections only. In our modeling framework, we take a novel approach to account for

these infections by allowing for waning multitypic immunity and repeated symptomatic infections

in older age individuals.

Recent analyses out of the Thailand system highlight a similar uptick in case reports

among older individuals (23), which the authors attribute to increased detectability in tertiary and

quaternary infections. The authors offer a series of hypothetical explanations for increasing case

detectability through time: that tertiary and quaternary dengue infections might be more

pathogenic (and therefore more detectable) due to an abundance of comorbidities in older

individuals, that immunopathology may be exacerbated in older patients who experienced longer

durations between repeat infections; or that waning multitypic immunity could allow for repeat

infections in the oldest age cohorts (our hypothesis). Our analyses in the Cambodia system are

not mutually exclusive with any of these explanations; however, we hypothesize that, were

comorbidities or immunopathogenesis in older individuals driving patterns in the observed data,

we would expect to see an hour-glass shape in symptomatic cases, with reduced reporting in

middle-aged individuals who have progressed beyond secondary exposures but are at lower risk

for both comorbidities and immunopathology. Instead, we see a gradual tapering in the age-

frequency of infection, which expands in range with time (Fig. S15), consistent with the

hypothesis of waning multitypic immunity. This latter hypothesis, when considered in association

with genotype replacement, can also explain isolated expansion and subsequent contraction in

the age range of reported cases witnessed in epidemic years, while the first two hypotheses

cannot. In addition, hypotheses of heightened pathology in more frequent older age infections are

difficult to reconcile in Cambodia with nationally-reported declines in the dengue mortality burden

despite increasing caseload through time (15).

Our deterministic model simulations demonstrate how short-term spikes in the age

distribution of reported cases, independent of the mean annual trend in the time series, can be

achieved via introduction of an antigenically distinct lineage (either a divergent genotype within an

established serotype, or a novel serotype) into an endemic, 3-serotype system. Though limited in

scope, PCR-based sentinel surveillance data provide no support for serotype turnover associated

with the 2019 dengue epidemic in Cambodia (15). By contrast, genome sequencing data from our

own febrile cohort study offer support for a genotype replacement event. Recent analyses of the

Thailand system link transitions in serotype dominance and clade replacement of genotype

sublineages within a single serotype to epidemic magnitude (34, 35). In Thailand, the most

severe epidemics were linked to clade replacement of a resident viral genotype by an

evolutionarily fitter DENV lineage within the same serotype, resulting in a 'selective sweep', which

subsequently reduced overall viral diversity and, consequently, diminished between-serotype

antigenic differences (34). In the 2019 epidemic in Cambodia, emergence of the Cosmopolitan

genotype, followed by serotypic and genotypic homogeneity in 2020-2022, are consistent with the

dynamics of clade replacement within DENV-2. Indeed, prior clade replacements have been

associated with epidemics in Cambodia (17, 18) and elsewhere (49, 53, 62–64).

As one limitation of our study, the bulk of our genotyping efforts were concentrated in

peri-urban Kampong Speu province, allowing for the possibility that the clade replacement

phenomenon documented here may be regionally localized, with consequentially limited impacts

on national caseloads. Nonetheless, the predominance of the DENV-2 Asian 1 lineage in the mid-

2000s has been well-documented across most provinces in Cambodia (65, 66), and our extended

sequencing efforts in 2021-2022 suggest that the Cosmopolitan takeover covered a similarly

widespread geographic extent. Additionally, reemergence of DENV-2 Cosmopolitan lineage has

also been documented in neighboring Vietnam and associated with a 2019 epidemic (52),

suggesting a more generalized regional phenomenon.

Our study faces additional limitations in that we were unable to obtain virus isolates and

undertake subsequent antigenic cartography (67), which would be needed to resolve whether the

emerging Cosmopolitan genotype in Cambodia is, indeed, antigenically distinct from previously

circulating Asian 1 lineages of DENV-2 and, potentially, more antigenically similar to endemic

heterotypic viruses. Indeed, because much of our inference is derived from serotype-agnostic

national surveillance data, we are unable to rigorously evaluate the role of a possible genotype

replacement event in particularly driving a high burden of disease in the oldest-age (>70 years)

individuals in NDCP dataset. Nonetheless, the expansion of this lineage following the 2019

epidemic offers support for the enhanced fitness of this genotype. In addition, our own febrile

surveillance study demonstrates a significantly older mean age of infection for DENV-2

Cosmopolitan cases than for DENV-1 cases, which further supports our hypothesis of invasion

paired with immune escape. Critically, our simulation analyses demonstrate how the expanded

age distribution of cases witnessed across epidemic years in Cambodia can only be recovered

under assumptions of lineage-specific immune invasion.

Several published studies offer regional explanations for the global dengue phenomenon

of 2019. Recent analyses from Brazil, for example, argue that a low FOI in 2017 and 2018

resulting from new public health interventions and behavioral modifications implemented in the

wake of the Zika virus epidemic drove a resurgence in cases in 2019 and an expansion of

specific lineages of DENV-1 and DENV-2 that had been circulating cryptically for much of the

past decade (68). Independent work in the same region indicates that the DENV-2 lineage

responsible for the Brazilian outbreak additionally caused a clade replacement event (63).

Dengue surged worldwide in 2019, though the factors driving this surge appear to be somewhat

heterogenous across ecosystems. Nonetheless, our study lends support for the role of climate in

synchronizing epidemic dynamics across landscapes; it is possible that optimal climate conditions

may facilitate expansion of low prevalence lineages with intrinsic fitness advantages (e.g. higher

replication rate, shorter incubation period (69)) over resident genotypes, thus driving epidemics.

Previous modeling studies have suggested that invading dengue lineages may circulate at low

prevalence in a host population for many years prior to detection, expansion, and displacement of

resident clades in the same serotype (70).

Here, we propose the emergence of the DENV-2 Cosmopolitan genotype, coupled with a

subtle climate-driven increase in FOI, and overlaid on the background of an aging population with

waning multitypic immunity, as one possible explanation for Cambodia's largest recorded dengue

epidemic to date. Our study emphasizes the extraordinary dearth of publicly available DENV

sequence data for SEA; indeed, DENV is sequenced so infrequently in Cambodia that it is

impossible to know whether 2019 truly marked the first year of DENV-2 Cosmopolitan

introduction to the country, or simply the year of intensified expansion and consequential

epidemic dynamics. More broadly, our work illustrates the importance of the combined forces of

climate, demography, immunology, and virology in driving increasingly severe dengue epidemics.

As the global burden of dengue continues to expand, ongoing serological and genomic

surveillance is needed to improve epidemic forecasting in Southeast Asia and around the world.

METHODS

Data

Detailed descriptions of NDCP case data, climate data, and Kampong Speu febrile surveillance data used in these analyses are reported in the *SI Appendix.*

Climate analyses

We used GAMs to quantify intra- and interannual trends and identify any years that might be considered climatic anomalies, defined as a statistically significant partial effect of a single year on the long-term time series for temperature and precipitation (Fig. S1-4; Table S1; *SI Appendix*). In addition, we reduced temperature and precipitation time series by province into z-scores for visualization (Fig. S5-6).

Climate-informed TSIR modeling

Using the R package tSIR (42), we fit time series Susceptible-Infected-Recovered models (42–44) to the province level dengue data for the three inter-epidemic periods (2002-2006, 2008-2011, and 2013-2018), compiled at biweekly intervals corresponding to the generation time of the pathogen (*SI Appendix*). Using TSIR transmission parameters, we projected cases in subsequent epidemic years (2007, 2012, 2019) by province. We then followed Wagner et al. 2020 (49) to estimate the factorial increase in reconstructed susceptible population needed to recover case counts for the three epidemic years.

We next used cross correlation analysis to determine the optimal lag between biweekly mean temperature and total precipitation versus transmission at the province level; we estimated a median 3.5 month lag for temperature and 1 month lag for precipitation, consistent with prior analyses (Table S3; *SI Appendix*) (49). We then constructed a suite of regression models (including GAMs) for each interepidemic period, incorporating a response variable of the log of biweekly transmission with the corresponding predictors of optimally lagged biweekly mean

temperature and total precipitation per province (45, 49) (Fig. S8-10; Table S4). We used lagged

temperature and precipitation from each epidemic year in each province in a fitted GAM to project

climate-informed transmission rates for 2007, 2012, and 2019 (*SI Appendix*). We then simulated

TSIR using climate-informed transmission rates to recover epidemic year predictions of province-

level caseload. Because climate-derived transmission rates still failed to recover epidemic peaks,

we allowed another susceptible amplification term to determine the relative increase in

susceptible population needed to recover epidemic dynamics even after accounting for climate

(Table S5).

Wavelet analyses

We next used wavelet decomposition in the R package 'WaveletComp' (50) to explore

annual and multiannual periodicity in dengue epidemics and climate variables. We converted

biweekly case totals by province from the NDCP data into incidence rates per 100,000

population, then used a Morlet wavelet with nondimensional frequency ($\omega = 6$) to extract,

detrend, and reconstruct annual cycles in dengue incidence with a maximum period of two years

and multiannual cycles with periods ranging from two to 20 years (Fig. 2AB) (28, 29). We

additionally calculated the average wavelet power (squared amplitude, here averaged across all

significant wavelets within a specified timestep) for significant cycles within each biweekly

timestep for annual and multiannual cycles per province (Fig. S11AB).

We investigated synchronicity in dengue incidence across space by computing the

Pearson's correlation coefficient (ρ) and the mean cross-wavelet power spectrum between

province pairs, using the annual raw incidence and reconstructed annual and multiannual cycles

(the latter averaged over a sliding 5-year interval) (Fig. 2CD, S11CD; *SI Appendix*). We

constructed a GAM to identify statistical correlates of high synchronicity pairs (Fig. S12; Table S6;

SI Appendix). To investigate synchrony between dengue and climate, we computed mean cross-

wavelet power between biweekly dengue incidence and mean temperature (Fig. S13A) and total

precipitation (Fig. S13B) per province, then repeated between reconstructed multiannual dengue

cycles over a 5-year interval and the same two climate variables, again per province (Fig.

S13CD). Finally, we reconstructed *monthly* multiannual dengue cycles per province to compute

average cross-wavelet power with the Oceanic Niño Index (ONI), a time series quantifying the

intensity of the El Niño Southern Oscillation (Fig. S13E). Lastly, we extracted the mean period

from reconstructed multiannual dengue cycles by province and nationally, to compare with

demographic data (birth rates, death rates, population size (60)) (Fig. S14).

Increasing age of reported infection and FOI estimation

To quantify interannual trends in the mean age of reported DENV infection, we fit a GAM

with a response variable of age to a fixed predictor of the interaction of year and province, with a

random effect of province, allowing both slope and y-intercept to vary by locality. We visualized

trends across the age distribution of reported cases by province (Fig. S15) and summarized

nationally (Fig. 3AB; Table S7).

To estimate annual FOI for DENV in Cambodia from 2002-2020, we applied the model

developed by Ferguson et al. 1999 (36) and Cummings et al. 2009 (24) to age-structured

incidence recovered from the NDCP data, at the province level, assuming reported cases to

represent secondary infections and individuals in the dataset to experience exposure to multiple

DENV serotypes across their lifetimes. We allowed for a unique FOI across each year in the time

series but first assumed constant FOI across all age cohorts within a year. We estimated one FOI

per year per province (excepting Tboung Khmum), in addition to a summary national FOI, for all

years in the dataset and extending back to the year of birth for the oldest individual in the first

year of the corresponding data subset (40 years for national data) (Fig. S16; Table S8; *SI*

Appendix). We estimated mean FOI per serotype assuming four circulating dengue serotypes

and plotted FOI estimates in comparison to national reported birth and death rates (Fig. 3C) (60).

After fixing FOI by province, we followed prior work (24) to estimate FOI modifiers by age,

shared across all provinces. We fit nineteen age-specific modifiers to the fixed FOI values,

corresponding to eight age classes in first decade of the dataset and eleven in the second

decade, where the age distribution was larger (Table S8). We then modified our model to allow

waning from multitypic back to monotypic immunity (σ), such that older individuals could

experience renewed symptomatic infections (*SI Appendix*). As with age-modification terms, we

estimated waning multitypic immunity shared across all provinces but allowed σ to vary annually.

We compared fits of FOI-only, FOI with age modification, FOI with waning multitypic immunity,

and FOI with both age modification and waning multitypic immunity models to the data (Table

S8), then simulated the resulting accumulation of cases with age from the best fit model at

national (Fig. 3E) and province levels (Fig. S17).

Viral sequencing

mNGS was applied to RNA extracted from 239 serum samples collected from patients

reporting with symptoms in our 2019-2022 febrile cohort study in Kampong Speu province, with

an additional subset of 11 samples sourced from five nearby provinces (39). A further 20 samples

were received for RNA extraction and mNGS from six provinces (including 8 from Kampong

Speu) in part with 2021-2022 NDCP surveillance (Table S9; Fig. 4AB). See *SI Appendix* for

detailed methods for library preparation, sequencing, and consensus genome generation*.* We

successfully generated 192 full or near-full genome sequences from the original 270 samples,

which we submitted to NCBI (63 DENV-1, 120 DENV-2, and 9 DENV-4). All contributed genomes

were >10,000 bps in length and had a maximum of 90 Ns (corresponding to <1% of the DENV

genome). Samples for which full genome recovery was not possible were identified to genotype

using BLAST (71) and visualized in Fig. 4AB (Table S9).

Phylodynamic and phylogenetic analysis

See *SI Appendix* for details of sequence selection for Bayesian phylodynamic timetrees

and maximum likelihood phylogenies*.* After selection, sequences were aligned by serotype in the

program MAFFT (72), and the best fit nucleotide substitution model for each serotype was

evaluated in ModelTest-NG (73). Using the best-fit model (GTR+I+G4 in all cases), we built a

Bayesian phylogenetic tree for each serotype in BEAST 2 (56), incorporating the date of sample

collection for each sequence (or the midpoint of the collection year if date was not reported), and

specifying a strict molecular clock rate of 7.9×10^{-4} s/s/y (8) and a Coalescent Bayesian skyline

prior. We ran Markov Chain Monte Carlo chains for 150 million iterations, logging results every

10,000 iterations. After chains completed, we removed the initial 10% of iterations as burn-in and

evaluated parameter convergence (ESS$\geq$ 200) in Tracer v1.6. We summarized resulting

phylogenetic trees in TreeAnnotator and visualized summary trees in ggtree (74) (Fig. 4CD; Fig.

S18, S19). Serotype-specific maximum likelihood phylogenetic trees were constructed in RAxML

(75). Twenty ML inferences were made on each original alignment and bootstrap replicate trees

were inferred using Felsenstein's method (76), with the MRE-based bootstopping test applied

after every 50 replicates (77). Resulting phylogenies were visualized in ggtree (74) (Fig. S18).

We followed Salje et al. 2017 (8) to calculate the proportion of sequences within each

DENV serotype that could be attributed to the same transmission chain on our Bayesian

timetrees, defined as having an MRCA within the past six months in the same season (see Fig.

S20 for alternative MRCA cutoffs). We compared the proportion of DENV-1 vs. DENV-2

sequences sharing a transmission chain to the Euclidean distance separating the GPS

coordinates for each sequence pair (Fig. 4E). We then computed the total effective number of

transmission chains (the reciprocal of the proportion of sequences sharing a chain) in our 2019-

2022 Kampong Speu sequence dataset for DENV-1 and DENV-2 at the World Bank reported

population density for Kampong Speu province (60) (Fig. 4F).

Mechanistic modeling of age-structured dengue dynamics

Finally, we constructed a mechanistic, age-structured discrete time deterministic model in

biweekly timesteps (57–59) to simulate three-serotype dengue dynamics in a population

demographically structured to mimic that of Cambodia over the past half-century (*SI Appendix*).

We compared simulations under (H0) baseline demographic assumptions with simulations of (H1)

increasing tertiary case detections through time and (H2) genotype invasion and clade-

replacement with waning immunity, timed both in 2019 and in 2007. We additionally simulated

novel genotype invasions under (H3) assumptions of no antigenic escape, or (H4) lineage-

agnostic increasing tertiary case detection (Fig. S21). Finally, we fit the Ferguson-Cummings

four-serotype catalytic model to the simulated data for each hypothesis to recapture the input FOI

(Fig. S22). Holding FOI constant, we estimated a time-varying signature of waning monotypic

immunity for all simulated time series (Fig. S22).

ACKNOWLEDGEMENTS

This research is supported by the Division of Intramural Research at the National Institute

of Allergy and Infectious Diseases at the National Institutes of Health and the Bill and Melinda

Gates Foundation [OPP1211806, OPP1211841]. We thank patients and families of Kampong

Speu District Referral Hospital who participated in this study, members of the National Dengue

Control Program not listed in the author byline, and the Provincial Health Department of Kampong

Speu province in Cambodia. We thank Brian Moyer and the NIAID Office of Cyberinfrastructure

and Computational Biology (OCICB) for assistance in improving the cyberinfrastructure of our

Cambodian field sites. We thank employees at the Chan Zuckerberg Biohub and Chan

Zuckerberg Initiative not listed in the author byline. This work was completed with resources

provided by the University of Chicago Research Computing Center.

REFERENCES

1.   World Health Organization, Dengue and severe dengue. *WHO Fact Sheets* (2022).

2.   S. Bhatt, *et al.*, The global distribution and burden of dengue. *Nature* **496**, 504–507 (2013).

3.   A. B. Sabin, Research on dengue during World War II. *American Journal of Tropical*
Medicine and Hygiene **1**, 30–50 (1952).

4.   L. C. Katzelnick, *et al.*, Antibody-dependent enhancement of severe dengue disease in
humans. *Science (New York, N.Y.)* **932**, 929–932 (2017).

5.   L. C. Katzelnick, *et al.*, Zika virus infection enhances future risk of severe dengue disease.
Science **369**, 1123–1128 (2020).

6.   T. N. Peng, "Southeast Asia's demographic situation, regional variations, and national
challenges" in D. Singh, M. Cook, Eds. (ISEAS Publishing, 2017), pp. 55–82.

7.   G. W. Jones, "The Population of Southeast Asia" (2013).

8.   H. Salje, *et al.*, Dengue diversity across spatial and temporal scales: Local structure and the
effect of host population size. *Science (New York, N.Y.)* **355**, 1302–1306 (2017).

9.   statista, Cambodia: Urbanization from 2010 to 2020.
https://www.statista.com/statistics/455789/urbanization-in-cambodia/. (2022).

10.   J. H. Huber, M. L. Childs, J. M. Caldwell, E. A. Mordecai, Seasonal temperature variation
influences climate suitability for dengue, chikungunya, and Zika transmission. *PLoS Negl*
Trop Dis **12**, e0006451 (2018).

11.   E. A. Gould, S. Higgs, Impact of climate change and other factors on emerging arbovirus
diseases. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **103**, 109–121
(2009).

12.   J. P. Messina, *et al.*, The current and future global distribution and population at risk of
dengue. *Nature Microbiology* **4**, 1508–1515 (2019).

13.   Hahn, H., Chastel, C., Dengue in Cambodia in 1963. Nineteen laboratory-proved cases.
American Journal of Tropical Medecine and Hygiene **19**, 106–109 (1970).

14.   R. Huy, *et al.*, National dengue surveillance in Cambodia 1980–2008: epidemiological and
virological trends and the impact of vector control. *Bull. World Health Organ.* **88**, 650–657
(2010).

15.   C. Yek, *et al.*, National dengue surveillance, Cambodia 2002–2020. *Bull World Health Org*
101, 605–616 (2023).

16.   D. S. Burke, A. Nisalak, D. E. Johnson, A prospective study of dengue infections in Bangkok.
The American Journal of Tropical Medicine and Hygiene **38**, 172–180 (1988).

17.   V. Duong, *et al.*, Genetic diversity and lineage dynamic of dengue virus serotype 1 (DENV-
1) in Cambodia. *Infection, Genetics and Evolution* **15**, 59–68 (2013).

18.   O. O'Connor, *et al.*, Potential role of vector-mediated natural selection in dengue virus
genotype/lineage replacements in two epidemiologically contrasted settings. *Emerging*
Microbes and Infections **10**, 1346–1357 (2021).

19.   A. A. Lover, *et al.*, Spatial epidemiology and climatic predictors of paediatric dengue
infections captured via sentinel site surveillance, Phnom Penh Cambodia 2011-2012. *BMC*
Public Health **14** (2014).

20.   Y. Arima, M. Chiew, T. Matsui, Epidemiological update on the dengue situation in the
Western Pacific Region, 2012. *Western Pacific surveillance and response journal : WPSAR*
6, 82–89 (2015).

21.   S. Ly, *et al.*, Asymptomatic dengue virus infections, Cambodia, 2012–2013. *Emerging*
Infectious Diseases **25**, 1354–1362 (2019).

22.   S. Vong, *et al.*, Under-recognition and reporting of dengue in Cambodia: A capture-
recapture analysis of the National Dengue Surveillance System. *Epidemiology and Infection*
140, 491–499 (2012).

23.   A. T. Huang, *et al.*, Assessing the role of multiple mechanisms increasing the age of dengue
cases in Thailand. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2115790119 (2022).

24.   D. A. T. Cummings, *et al.*, The impact of the demographic transition on dengue in Thailand:
Insights from a statistical analysis and mathematical modeling. *PLoS Medicine* **6** (2009).

25.   L. C. Katzelnick, *et al.*, Dynamics and determinants of the force of infection of dengue virus
from 1994 to 2015 in Managua, Nicaragua. *Proceedings of the National Academy of*
Sciences of the United States of America **115**, 10762–10767 (2018).

26.   N. Ferguson, R. Anderson, S. Gupta, The effect of antibody-dependent enhancement on
the transmission dynamics and persistence of multiple-strain pathogens. *Proc. Natl. Acad.*
Sci. U.S.A. **96**, 790–794 (1999).

27.   S. I. Hay, *et al.*, Etiology of interepidemic periods of mosquito-borne disease. *Proc. Natl.*
Acad. Sci. U.S.A. **97**, 9335–9339 (2000).

28.   B. García-Carreras, *et al.*, Periodic synchronisation of dengue epidemics in Thailand over
the last 5 decades driven by temperature and immunity. *PLoS Biol* **20**, e3001160 (2022).

29.   W. G. van Panhuis, *et al.*, Region-wide synchrony and traveling waves of dengue across
eight countries in Southeast Asia. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 13069–13074 (2015).

30.   D. A. T. Cummings, I. B. Schwartz, L. Billings, L. B. Shaw, D. S. Burke, Dynamic effects of
antibody-dependent enhancement on the fitness of viruses. *Proceedings of the National*
Academy of Sciences **102**, 15259–15264 (2005).

31.   K. L. McElroy, *et al.*, Endurance, refuge, and reemergence of dengue virus type 2, Puerto
Rico, 1986–2007. *Emerg. Infect. Dis.* **17**, 64–71 (2011).

32.   B. Adams, *et al.*, Cross-protective immunity can account for the alternating epidemic
pattern of dengue virus serotypes circulating in Bangkok. *Proceedings of the National*
Academy of Sciences of the United States of America **103**, 14234–9 (2006).

33.   C. V. F. Carrington, J. E. Foster, O. G. Pybus, S. N. Bennett, E. C. Holmes, Invasion and
maintenance of Dengue virus type 2 and type 4 in the Americas. *Journal of Virology* **79**,
14680–14687 (2005).

34.   L. C. Katzelnick, *et al.*, Antigenic evolution of dengue viruses over 20 years. *Science* (2021).

35.   S. M. Bell, L. Katzelnick, T. Bedford, Dengue genetic divergence generates within-serotype
antigenic variation, but serotypes dominate evolutionary dynamics. *eLife* **8**, e42496 (2019).

36.   N. M. Ferguson, C. A. Donnelly, R. M. Anderson, Transmission dynamics and epidemiology
of dengue: insights from age-stratied sero-prevalence surveys. *Proceedings of the Royal*
Society B **354**, 757–768 (1999).

37.   Hugo. Muench, *Catalytic models in epidemiology.* (Harvard University Press, 1959).

38.   J. E. Manning, *et al.*, Development of inapparent dengue associated with increased
antibody levels to *Aedes aegypti* salivary proteins: A longitudinal dengue cohort in
Cambodia. *The Journal of Infectious Diseases* (2021).
https://doi.org/10.1093/infdis/jiab541.

39.   J. A. Bohl, *et al.*, Discovering disease-causing pathogens in resource-scarce Southeast Asia
using a global metagenomic pathogen monitoring system. *Proceedings of the National*
Academy of Sciences **119**, e2115285119 (2022).

40.   S. N. Wood, mgcv: GAMs and Generalized Ridge Regression for R. *R News* **1/2**, 20–24
(2001).

41.   NOAA: National Weather Service, *Climate Predication Scenter: Cold & Warm Episodes by*
Season (2023). Available at:
https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php.

42.   A. D. Becker, B. T. Grenfell, TSIR: An R package for time-series susceptible-infected-
recovered models of epidemics. *PLoS ONE* **12**, 1–10 (2017).

43.   O. N. Bjornstad, B. F. Finkenstadt, B. T. Grenfell, Dynamics of measles epidemics:
Estimating scaling of transmission rates using a times series SIR model. *Ecological*
Monographs **72**, 169–184 (2002).

44. B. T. Grenfell, O. N. Bjornstad, B. F. Finkenstadt, Dynamics of measles epidemics: Scaling
noise, determinism, and predictability with the TSIR Model. *Ecological Monographs* **72**,
185–202 (2002).

45. R. J. Oidtman, *et al.*, Inter-annual variation in seasonal dengue epidemics driven by
multiple interacting factors in Guangzhou, China. *Nature Communications* **10** (2019).

46. N. A. Rehman, *et al.*, Quantifying the localized relationship between vector containment
activities and dengue incidence in a real-world setting: A spatial and time series modelling
analysis based on geo-located data from Pakistan. *PLoS Neglected Tropical Diseases* **14**, 1–
22 (2020).

47. M. U. G. Kraemer, *et al.*, Big city, small world: density, contact rates, and transmission of
dengue across Pakistan. *J. R. Soc. Interface.* **12**, 20150468 (2015).

48. M. U. G. Kraemer, *et al.*, Inferences about spatiotemporal variation in dengue virus
transmission are sensitive to assumptions about human mobility: a case study using
geolocated tweets from Lahore, Pakistan. *EPJ Data Science* **7** (2018).

49. C. E. Wagner, *et al.*, Climatological, virological and sociological drivers of current and
projected dengue fever outbreak dynamics in Sri Lanka. *Journal of the Royal Society
Interface* **17** (2020).

50. Roesch, Angi, Schmidbauer, Harald, WaveletComp: Computational Wavelet Analysis. R
package version 1.1. (2018). Deposited 2018.

51. J. Phadungsombat, *et al.*, Emergence of genotype cosmopolitan of dengue virus type 2 and
genotype III of dengue virus type 3 in Thailand. *PLoS ONE* **13**, 1–26 (2018).

52. V. T. Tran, *et al.*, Reemergence of Cosmopolitan genotype dengue virus serotype 2,
southern Vietnam. *Emerg. Infect. Dis.* **29** (2023).

53. A. Wijewickrama, *et al.*, Emergence of a Dengue virus serotype 2 causing the largest ever
dengue epidemic in Sri Lanka. *bioRxiv* (2018). https://doi.org/10.1101/329318.

54. H. A. Tissera, *et al.*, Severe dengue epidemic, Sri Lanka, 2017. *Emerg. Infect. Dis.* **26**, 682–
691 (2020).

55. A. J. Drummond, M. A. Suchard, D. Xie, A. Rambaut, Bayesian phylogenetics with BEAUti
and the BEAST 1.7. *Molecular Biology and Evolution* **29**, 1969–1973 (2012).

56. R. Bouckaert, *et al.*, BEAST 2: A software platform for Bayesian evolutionary analysis. *PLoS
Computational Biology* **10** (2014).

57. P. Klepac, H. Caswell, The stage-structured epidemic: Linking disease and demography with
a multi-state matrix approach model. *Theoretical Ecology* **4**, 301–319 (2011).

58.   C. J. E. Metcalf, *et al.*, Structured models of infectious disease: Inference with discrete
data. *Theoretical Population Biology* **82**, 275–282 (2012).

59.   P. Klepac, *et al.*, Stage-structured transmission of phocine distemper virus in the Dutch
2002 outbreak. *Proceedings. Biological sciences / The Royal Society* **276**, 2469–2476
(2009).

60.   World Bank, Cambodia. *World Development Indicators* (2021). Available at:
https://data.worldbank.org/country/KH [Accessed 12 April 2022].

61.   I. A. Sampson, G. M. Miles, E. Piano, "Undocumented, unregistered and invisible": an
exploratory study of the reasons for and effects of migrating to Thailand of Cambodian
young people. *IJSSP* **41**, 862–874 (2021).

62.   L. Lambrechts, *et al.*, Dengue-1 virus clade replacement in Thailand associated with
enhanced mosquito transmission. *Journal of Virology* **86**, 1853–1861 (2012).

63.   J. G. de Jesus, *et al.*, Genomic detection of a virus lineage replacement event of dengue
virus serotype 2 in Brazil, 2019. *Memorias do Instituto Oswaldo Cruz* **115** (2020).

64.   K. Suzuki, *et al.*, Genotype replacement of dengue virus type 3 and clade replacement of
dengue virus type 2 genotype Cosmopolitan in Dhaka, Bangladesh in 2017. *Infection,*
Genetics and Evolution **75** (2019).

65.   V. Duong, *et al.*, Complex dynamic of dengue virus serotypes 2 and 3 in Cambodia
following series of climate disasters. *Infection, Genetics and Evolution* **15**, 77–86 (2013).

66.   V. T. T. Hang, *et al.*, Emergence of the Asian 1 genotype of dengue Virus serotype 2 in Viet
Nam: *In Vivo* fitness advantage and lineage replacement in South-East Asia. *PLoS*
Neglected Tropical Diseases **4** (2010).

67.   L. C. Katzelnick, *et al.*, Dengue viruses cluster antigenically but not as discrete serotypes.
Science **349**, 1338–1343 (2015).

68.   A. F. Brito, *et al.*, Lying in wait: the resurgence of dengue virus after the Zika epidemic in
Brazil. *Nature Communications* **12**, 1–13 (2021).

69.   R. Rico-Hesse, *et al.*, Origins of Dengue type 2 viruses associated with increased
pathogenicity in the Americas. *Virology* **230**, 244–251 (1997).

70.   J. Lourenço, M. Recker, Viral and epidemiological determinants of the invasion dynamics of
Novel Dengue Genotypes. *PLoS Neglected Tropical Diseases* **4** (2010).

71.   S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic local alignment search
tool. *Journal of Molecular Biology* **215**, 403–410 (1990).

72. K. Katoh, J. Rozewicki, K. D. Yamada, MAFFT online service: Multiple sequence alignment,
interactive sequence choice and visualization. *Briefings in Bioinformatics* **20**, 1160–1166
(2018).

73. Di. Darriba, *et al.*, ModelTest-NG: A new and scalable tool for the selection of DNA and
protein evolutionary models. *Molecular Biology and Evolution* **37**, 291–294 (2020).

74. G. Yu, D. K. Smith, H. Zhu, Y. Guan, T. T. Y. Lam, Ggtree: an R Package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods*
in Ecology and Evolution **8**, 28–36 (2017).

75. A. M. Kozlov, D. Darriba, T. Flouri, B. Morel, A. Stamatakis, RAxML-NG: A fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**,
4453–4455 (2019).

76. J. Felsenstein, Confidence limits on phylogenies: An approach using the bootstrap.
Evolution **39**, 783–791 (1985).

77. N. D. Pattengale, M. Alipour, O. R. P. Bininda-Emonds, B. M. E. Moret, A. Stamatakis, How
many bootstrap replicates are necessary? *Journal of Computational Biology* **17**, 337–354
(2010).

1046

1047

1048

A biweekly annual reconstructed cycle distribution

B biweekly multi-annual reconstructed cycle distribution

C annual distribution pairwise province Pearson's correlation

D multi-annual distribution pairwise province Pearson's correlation

Reconstructed annual cycles
−50 −1 0 1 3 10 30 70

Reconstructed multiannual cycles
−50 −5 0 3 10 30 50 70 100

Average pairwise province Pearson's correlation coefficient (top: annual; bottom: multi−annual)
0.00 0.25 0.50 0.75 1.00

	A total reported cases from deterministic simulation	B age distribution of reported cases by year	C cumulative proportion of cases by age
H0: Normal Demographic Simulation			
H1: Increasing Tertiary Case Detection			
H2: Genotype Replacement + Waning Immunity (2019)			
H2: Genotype Replacement + Waning Immunity (2007)			