Pruning and thresholding approach for methylation risk scores in multi-ancestry populations

Junyu Chen[1], Evan Gatev[2], Todd Everson[3,1], Karen N. Conneely[4], Nastassja Koen[5,6,7], Michael P. Epstein[4], Michael S. Kobor[8,9,10], Heather J. Zar[11,12], Dan J. Stein[5,6,7], Anke Huels[1,3]

1. Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA
2. Institute of Molecular Biology "Acad. Roumen Tsanev", Sofia, Bulgaria
3. Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, USA
4. Department of Human Genetics, School of Medicine, Emory University, Atlanta, GA, USA
5. Neuroscience Institute, University of Cape Town, Cape Town, South Africa
6. Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa
7. South African Medical Research Council (SAMRC) Unit on Risk and Resilience in Mental Disorders, University of Cape Town, Cape Town, South Africa
8. Department of Medical Genetics, University of British Columbia, Vancouver, Canada
9. BC Children's Hospital Research Institute, Vancouver, Canada
10. Centre for Molecular Medicine and Therapeutics, Vancouver, Canada
11. Department of Pediatrics and Child Health, Red Cross War Memorial Children's Hospital, University of Cape Town, Cape Town, South Africa
12. South African Medical Research Council (SAMRC) Unit on Child and Adolescent Health, University of Cape Town, Cape Town, South Africa

Corresponding author:
Anke Hüls, PhD
Department of Epidemiology and Gangarosa Department of Environmental Health
Rollins School of Public Health, Emory University
1518 Clifton Road, Atlanta, GA 30322, USA
E-mail: anke.huels@emory.edu

Funding:

The Drakenstein Child Health Study was funded by the Bill & Melinda Gates Foundation (OPP 1017641), Discovery Foundation, South African Medical Research Council, National Research Foundation South Africa, CIDRI Clinical Fellowship and Wellcome Trust (204755/2/16/z). Additional support for the DNA methylation work was by the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health (NICHD) under Award Number R21HD085849, and the Fogarty International Center (FIC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. AH is supported the HERCULES Center (NIEHS P30ES019776). DJS and HJZ are supported by the South African Medical Research Council (SAMRC). The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of manuscript.

Disclosure statement:

All authors declare they have no actual or potential competing financial interest.

Data Availability Statement:

The data for simulation studies were derived from the following dataset (GSE55763, GSE84727, GSE80417, GSE111629 and GSE72680) from NCBI Gene Expression Omnibus (https://www.ncbi.nlm.nih.gov/geo/). The data for real data application are available from the corresponding author, [AH], upon reasonable request.

Abstract

Recent efforts have focused on developing methylation risk scores (MRS), a weighted sum of the individual's DNAm values of pre-selected CpG sites. Most of the current MRS approaches that utilize Epigenome-wide association studies (EWAS) summary statistics only include genome-wide significant CpG sites and do not consider co-methylation. New methods that relax the p-value threshold to include more CpG sites and account for the inter-correlation of DNAm might improve the predictive performance of MRS. We paired informed co-methylation pruning with P-value thresholding to generate pruning and thresholding (P+T) MRS and evaluated its performance among multi-ancestry populations. Through simulation studies and real data analyses, we demonstrated that pruning provides an improvement over simple thresholding methods for prediction of phenotypes. We demonstrated that European-derived summary statistics can be used to develop P+T MRS among other population such as African population. However, the prediction accuracy of P+T MRS may differ across multi-ancestry population due to environmental/cultural/social differences.

Introduction

DNA methylation (DNAm), one of the most studied epigenetic mechanisms,

regulates the mode of expression of DNA segments independent of alterations of

their sequence by adding a methyl group at cytosine residues, hence contributing to

variation in cellular phenotypes[1]. With current advances in and reduction of cost of

array-based profiling technologies, increasing numbers of large-scale epigenome-

wide association studies (EWAS) have been conducted to study DNAm in

association with complex human diseases as well as environmental and social

factors[2,3]. EWAS has thus far been successful in identifying dozens of cytosine

guanine dinucleotides (CpGs) associated with various diseases and exposures,

which could potentially be used for disease diagnosis and prediction, development of

drug targets, and monitoring of drug response[3-8]. However, differential DNAm in

individual CpGs often shows a weak prediction capacity and can only explain a small

fraction of phenotype variance. Polyepigenetic approaches that aggregate

information on differential DNAm from multiple CpGs might produce a more accurate

biomarker for clinical usage[9,10].

A well-known polygenic approach for genotype data is polygenic risk scores (PRS),

which are weighted sum of risk alleles of a pre-selected number of genetic variants[11].

Recently, many efforts have focused on transferring PRS approaches to DNA

methylation data to construct methylation risk scores (MRS), which are defined as

weighted sums of the individuals' DNAm values of a pre-selected number of CpGs[10].

However, there are many methodological challenges in constructing DNA

methylation risk scores[10,12,13]. One of the problems is that DNAm is influenced by

ancestry, which captures genetic ancestry (differences in the genome related to

ancestry) as well as social determinants of health such as racism and discrimination,

socioeconomic status, and environmental effects[14]. Thus, ideally, when external

weights are used for the calculation of MRS, these weights should be assessed in a

population with the same ancestry as the study samples. However, current

epigenetic literature remains limited by the lack of diversity, with most focusing on

European populations[15], therefore making it difficult to identify appropriate weights

for MRS for other populations. While it is well known that PRSs are not applicable

across different ancestries[16,17], little is known about the performance of MRS across

multi-ancestry populations.

Currently, there are two popular approaches to construct MRS. The first one is to

use penalized regression models such as Elastic Net and LASSO

regularization[10,18,19], which usually requires individual-level DNAm data. When only

summary-level statistics are available, individual CpGs that reached genome-wide

significance in an external epigenome-wide association studies (EWAS) are selected

and the beta-coefficients estimated from EWAS are used as weights to generate

MRS[10]. However, research in PRS has shown that the optimal p-value threshold

strongly depends on the data[20], and including a larger proportion of variants could

potentially capture more of the phenotype variation[21]. Moreover, most MRS from the

second approach do not consider DNA co-methylation, defined as proximal CpGs

with correlated DNAm across individuals[22], which could potentially bias the

generation of MRS. Shah et al 2015 proposed to remove redundant CpGs by

keeping the most significant CpGs in co-methylation[23,24], however, a window to

define DNA co-methylation needs to be pre-defined and the effect of accounting for

DNA co-methylation was not evaluated.

One of the most widely used PRS approaches to deal with single nucleotide

polymorphisms (SNPs) in high linkage disequilibrium (LD) and to identify p-value

thresholds with the best prediction accuracy is the pruning and threshold (P+T)

method[25]. In the P+T approach, the correlation square (R^2) for SNPs within a close

genetic distance is calculated and less significant SNPs that are correlated with an

R^2 greater than a particular value (LD pruning)[26] are removed. Next, several p-value

thresholds are tested to maximize the prediction accuracy of the derived PRS (p-

value thresholding)[26,27]. Theoretically, the P + T approach could be applied to

generate MRS, however, there is no standard procedure on how to conduct pruning

for DNAm data and the performance of such MRS across multi-ancestry populations

remains unknown.

Here, we propose to use the Co-Methylation with genomic CpG Background

(CoMeBack), a tool that uses a sliding window to estimate DNA co-methylation, to

account for correlations of DNAm at proximal CpG sites[22], and pair it with p-value

thresholding to construct P+T CoMeBack MRS. CoMeBack uses unmeasured

intermittent CpGs from the human reference genome to link array probes in hope of

reducing false positives while improving the identification of biologically relevant co-

methylation[22]. We conducted simulation studies based on data from an adult

population consisting of three groups of different ancestries (Indian, White and Black,

n = 1,199) to evaluate the prediction performance of P+T CoMeBack MRS and how

it changes across multi-ancestry population. Next, we applied the P+T CoMeBack

approach to DNAm data from the Drakenstein Child Health Study (n=270)[28], a multi-

ancestry birth cohort from South Africa, to evaluate the performance of MRS for

maternal smoking status. Our simulation study and real data application

demonstrated that the P+T approach improves the predictive accuracy of MRS over

methods that do not account for co-methylation and has similar performance as

LASSO regression, which requires access to the raw DNAm data. We also showed

that MRS built upon the data from a population of one genetic ancestry could

achieve high prediction performance among populations of other genetic ancestries,

but the performance might differ in the presence of environmental/cultural/social

differences associated with ancestry.

Materials and methods

P+T CoMeBack approach for MRS

P+T CoMeBack method refers to the calculation of MRS using informed co-methylation pruning (P) with CoMeBack and P-value thresholding (T). First, summary statistics from an EWAS (typically include the participant ID, effect size, standard error and P-value of each CpG site) need to be estimated in an independent dataset (training dataset) to avoid overfitting, and then applied to generate MRS in a testing dataset (the samples used to evaluate the performance of MRS).

In our P+T CoMeBack method, co-methylation pruning is completed by applying CoMeBack to DNAm data of the testing dataset or a reference panel[22]. Specifically, CoMeBack chains two adjacent array probes if the following requirements are met: 1. two probes are less than 2kb apart; 2. the reference human genome annotation shows a set of unmeasured genomic CpGs between them; 3. the density of unmeasured genomic CpGs between them is at least one CpG every 400bp. Chaining of adjacent array probes continues until an array probe does not meet the requirements, which will form a unit where multiple CpGs are chained together. Correlations between DNAm levels will then be calculated for all array probes inside each unit. If all pairs of adjacent probes in a unit have a correlation square (R^2) greater than 0.3, such unit will be declared as a co-methylated region (CMR).

Pruning is conducted by only keeping one CpG site per CMR in the dataset, the one

with the lowest (most significant) P-value in the EWAS summary statistics.

P + T CoMeBack will be compared to the standard pruning approach, in which less

significant SNPs that are correlated with an $R^2 > 0.3$ and located within 2000bp of

each other are being removed.

Next, P-value thresholding step (T) is performed for the pruned set of CpG sites.

Specifically, the P-value thresholding step (T) is performed by applying different P-

value thresholds (e.g., P-value thresholds $\in$ [0.05, 0.005, 5 x 10^{-4}, 5 x 10^{-5}, …]) and

only including those CpG sites in the final MRS calculation that reached a P-value

below those thresholds in the EWAS summary statistics.

Finally, for each P-value threshold, MRS are calculated as a weighted sum of DNAm

β values (β value = methylated allele intensity / (unmethylated allele intensity +

methylated allele intensity + 100), ranging from 0 representing unmethylation to 1 for

complete methylation) of the selected CpGs, where the weights are the

corresponding effect sizes for each CpG from the EWAS summary statistics. The

squared correlation (R^2) between the phenotype of interest and MRS obtained using

each P-value threshold is calculated to represent the prediction accuracy. The P-

value threshold that produces MRS with the highest prediction accuracy in the

testing data set is selected as the optimal P-value and the corresponding MRS is

used for downstream analysis. The pipeline for generating P+T MRS is written in an

R script, which is available at GitHub (https://github.com/jche453/Pruning-

[Thresholding-MRS.git](https://github.com/jche453/Pruning-Thresholding-MRS.git)).

In our simulation studies and real data application, we compare the P+T CoMeBack

MRS approach to the standard P+T and T approach, which refers to an approach in

which the MRS is calculated by only thresholding, not accounting for correlations

between included CpGs (no pruning).

Simulation studies

To validate the performance of the proposed P+T MRS approach, we conducted

simulation studies based on whole blood Illumina Infinium Human Methylation 450K

BeadChip data from an ethnically heterogeneous discovery cohort composed of

several publicly available datasets (GSE55763, GSE84727, GSE80417, GSE111629

and GSE72680)[22]. Intra-dataset normalization and batch effects correction were

performed using ComeBat function in R-package sva[29], followed by merging of

datasets and correction for inter-dataset batch effects using the same function. After

the removing XY chromosome binding, non-CpG, cross-hybridizing probes and

probes that are in close distance with common SNPs, there were 386,362 CpGs left

for MRS analysis. We randomly selected 1,199 adults (898 Indians, 136 Blacks and

165 Whites) to conduct the simulation studies.

CoMeBack was applied to the DNA methylation β values of the 386,362 CpGs to

obtain CMR. In each simulation, 10 of the 386,362 CpGs were randomly selected to

be causal, k% (k = 30, 50, 70 or 100) of which are in a CMR with other CpGs. At

most, one CpG would be causal in each CMR.

The causal CpGs were randomly assigned a "true" effect size from a uniform

distribution as $w_i \sim U(-0.5, 0.5)$. We then simulated a phenotype for the j-th subject

as follow:

158
$$Y_j = \sum_{i=1}^{10} w_i m_{ij} + \varepsilon_j, \varepsilon_j \sim N(0, \delta^2),$$

where m_{ij} is the DNAm β value of causal CpG site i of the j-th subject, and ε_j is an

error term that follows a normal distribution. Different δ^2 were set to ensure that the

targeted variance of phenotype explained by DNAm alone equals 10%, 30% 50% or

80%.

We also simulated a second phenotype Y_j^* for the j-th subject, which was directly

affected by ancestry using European ancestry as reference:

166
$$Y_j^* = \sum_{i=1}^{10} w_i m_{ij} + a * (if\ Indian) + b * (if\ Black) + \varepsilon_j^*, \varepsilon_j^* \sim N(0, \delta^{*2})$$

Effect of ancestry in our simulations is simulated as the effect of genetic ancestry

assuming there were no complex social determinants involved in the causal

pathway. Different δ^{*2} were used so that the variance of phenotype that was

explained by DNAm and ancestry together equals 20%, 50% or 80. For our

simulations, effect a was set to 0.1 and b to 0.2. In each simulation, both simulated

phenotypes Y_j and Y_j^* share the same epigenetic liability ($\sum_{i=1}^{10} w_i\, m_{ij}$).

In each simulation, for fair comparison, 762 Indians were randomly chosen as the

training dataset so that there were at least 136 people left for each race group in the

testing dataset. Associations between CpGs and each of the two simulated

phenotypes were assessed by robust linear regression model using limma R

package[30] in the training dataset. We calculated top 10 principal components (PCs)

from DNAm of 386,362 CpGs[31] and used EpiDISH to estimate cell type proportions

of each CpGs[32]. We observed that in our simulation dataset, top 10 PCs are highly

correlated with cell type proportions (Supplement figure 1A), and using either

summary statistics adjustment for top 10 PCs or summary statistics adjusted for cell

type proportions would lead to almost identical prediction performance of MRS

(Supplement figure 1B). Thus, to account for population stratification and cell type

difference, we adjusted for the top 10 PCs in our main analyses. The summary

statistics (effect size and P-values) obtained from association tests in the training

data were saved and later used to construct MRS in the testing dataset. We

repeated 1000 simulations per scenario to evaluate the prediction accuracy (R^2),

power and type 1 error rate of the P+T MRS. Linear regression analysis was used to

access the association between MRS and simulated phenotypes, and power is

defined as the proportion of simulations where MRS were significantly associated

with the simulated phenotype with at α level of 0.05. To estimate type 1 error rate,

we first obtained a null association between MRS values and simulated phenotype

values by permutation of MRS values. Linear regression analysis was used to

access the association between permutated MRS and simulated phenotypes, and

type 1 error rate is defined as the proportion of simulations where permutated MRS

were significantly associated with the simulated phenotype.

We evaluated the performance of the MRS not only in scenarios of A) same ancestry

in training and test data, but also B) across different ancestry groups (training data:

Indian, test data: European or African) and C) in multi-ancestry populations (training

data: Indian, test data: Indian, European and African). For scenario C), we evaluated

two analysis strategies: 1. Joint-analysis: perform MRS analyses in the whole testing

dataset where subjects from all racial groups were merged; 2. Standardization: scale

MRS to have a standard normal distribution within each racial group before merging

all subjects for analyses.

Application study of smoking MRS

To evaluate the performance of the P+T CoMeBack approach in a real data setting,

we applied the P+T CoMeBack approach to calculate a MRS for maternal smoking

status during pregnancy using cord blood DNAm data from newborns in the South

African Drakenstein Child Health Study (DCHS), a multi-ancestry longitudinal study

investigating determinants of early child development[33]. There were 145 Black

African infants and 115 Mixed ancestry infants in the DCHS. A detailed description of

the enrollment process, inclusion criteria, variables measurement and ethical

approval of the study have been previously published[33,34].

Cotinine levels were measured in urine provided by mothers within four weeks of

enrollment and classified as <499 ng/ml (non-smoker), or ≥500 ng/ml (active

smoker)[28]. Cord blood was collected at time of delivery and used to measure DNA

methylation by either MethylationEPIC BeadChips (EPIC, n=145) or the Illumina

Infinium HumanMethylation450 BeadChips (450K, n=103)[33,34], followed by quality

control and normalization to calculate β values (details have been published

elsewhere) [35].

Summary statistics for the calculation of MRS were obtained from a study that meta-

analyzed the associations between newborn blood DNA methylation and sustained

maternal smoking during pregnancy among 5,648 mother-child pairs as part of the

Pregnancy and Childhood Epigenetics (PACE) Consortium (Table 1)[36]. The

participants of all cohorts used in the meta-analysis except one were of European

ancestry.

In addition, we compared P+T CoMeBack MRS to three previously published MRS

for maternal smoking during pregnancy (Reese MRS, Richmond 19 MRS, Richmond

568 MRS; Table 1). Reese MRS model was trained among 1,068 newborns of

European ancestry in the Norwegian Mother and Child Cohort Study, while

Richmond 568 MRS and Richmond 19 MRS was trained in multi-ancestry newborns

(N=6,685) and children around 6.8 years old (N=3,187) in PACE Consortium

respectively. The training population for Reese MRS and Richmond 19 MRS

overlapped with the training population for summary statistics used in P+T

CoMeBack MRS in our study. Reese et al. used a LASSO regression to select CpGs

for Reese MRS, which is a weighted sum of DNAm β values of 28 CpGs with

weights estimated from the LASSO regression[37]. Richmond 19 MRS is a weighted

sum of DNAm β values of 19 CpGs that were significantly associated with prenatal

smoking in an EWAS conducted in peripheral blood from children of averaged 6.8

years age (Richmond 19 MRS)[38,39]. In the same study, Richmond 568 MRS was

proposed based on 568 CpGs that were significantly associated with prenatal

smoking in cord blood[38,39]. We obtained the weights of reported CpGs from the

mentioned studies and applied them to DNAm data in DCHS to generate Reese

MRS, Richmond 19 MRS and Richmond 568 MRS.

Linear regressions were used to assess the associations between maternal smoking

status and each MRS, controlling for ancestry (in pooled samples), cell type

proportions and top 5 PCs calculated from genotypes. In order to obtain comparable

beta-coefficients and standard errors across different MRS, each MRS was divided

by their interquartile range (IQR) before linear regression analysis.

Results

Simulation results

We compared the prediction performance of P+T CoMeBack MRS to the T method

among 136 Indians in the test data across different simulation scenarios (Figure 1).

Figure 1A shows that P+T CoMeBack MRS that account for co-methylation between

CpGs have stable prediction performance when proportion of causal CpGs located in

a CMR (k%) varies. P+T without CoMeBack had similar prediction performance while

the T method had a slightly lower prediction performance. While the P+T CoMeBack

MRS showed subtle improvement over T method when V^2_{DNAm} is 80% (Figure 1A),

the difference between P+T CoMeBack MRS and the T method decreases as the

V^2_{DNAm} decreases. This is likely because as variance explained by DNAm decreases,

there is less power for association testing, and it becomes increasingly difficult to

distinguish real signals from statistical noise while generating the summary statistics.

Next, we assessed the performance of P+T CoMeBack, P+T and T method across

different ancestries and among multi-ancestry populations. All three methods

achieved a high power (> 95%) and a low type 1 error rate (~ 5%) within each

ancestry in most scenarios for both phenotypes except when the phenotype variance

explained by DNA methylation is 10% or 30% (Supplement table 1-4).

Whether the simulated phenotypes were independent of ancestry or not, MRS

among Whites and Blacks achieved a prediction R^2 as high as among Indians, which

should have the best prediction of the simulated phenotypes since weights were

obtained from Indian training samples (Figure 2). Findings were similar when the

phenotype variance explained by DNA methylation was reduced from 80% to 10%,

30% or 50% (Supplement Figure 2). When the phenotypes are not associated with

ancestry (Figure 2A), the three MRS analyses strategies (stratification, joint analysis

and standardization) lead to nearly identical results. However, when the phenotypes

are ancestry-dependent, both joint analysis and standardization of MRS showed very

poor prediction of the phenotypes (Figure 2B).

MRS of maternal smoking status

Figure 3 shows the prediction performance of MRS for maternal smoking status

among DCHS newborns. As the p-value threshold decreases, the prediction

accuracy of the resulting MRS increases before reaching a plateau, demonstrating

the importance of P-value thresholding in MRS to control for noise. Among mixed

ancestry newborns, P+T CoMeBack MRS of smoking status excluded 22 CpGs in

pruning and achieved a prediction R^2 of 29.5% using P-value threshold of 5×10^{-22},

while the standard P+T without CoMeBack had a lower prediction R^2 of 26.2% using

P-value threshold of 5×10^{-10} and the best T method MRS had the lowest prediction

accuracy (24.5%) using P-value threshold 5×10^{-9}, confirming the benefits of pruning

in MRS calculation (Figure 3A). All three MRS had lower prediction performance for

maternal smoking among Black African infants (10.9%, and 8.0% respectively)

(Figure 3B), which is likely due to the low prevalence of smokers among mothers of

Black African infants in DCHS (13%) compared to mothers of mixed ancestry infants

(49%) (Supplement Figure 3). Additionally, the distributions of all MRS in Black

African infants and mixed ancestry infants were similar within each category of

maternal smoking status (Supplement Figure 4), confirming that the difference of

prediction R^2 between Mixed and Black infants is less likely due to ancestry-related

factors other than prevalence of maternal smoking. Joint-analysis of P+T CoMeBack

MRS showed a prediction accuracy of 20.4%, which is between the prediction

accuracy of P+T CoMeBack MRS among Black African infants and mixed ancestry

infants (Figure 3C). Standardization approach did not improve the performance of

MRS (Figure 3D).

We next compared the prediction accuracy and distribution of P+T CoMeBack MRS

to other established MRS for maternal smoking during pregnancy and newborn

DNAm (Figure 4). Overall, P+T CoMeBack and Reese MRS had stable and similar

classification performance in all analyses compared to other MRS. P+T CoMeBack

MRS and Reese MRS showed a similar prediction R^2 among both Black and Mixed

ancestry infants, which are better than other smoking MRS (Figure 4A). P+T

CoMeBack MRS had the largest AUC (0.820) in the ROC curve among mixed infants

(Figure 4B) but a smaller AUC than Reese MRS in Black infants and joint-analysis

(Figure 4C-D). Further, all 6 MRS showed significant association with smoking status

in Mixed infants, Black infants and joint-analysis (Table 2), showing the promise of

using MRS to capture the overall DNAm signals in association testing.

Discussion

Based on the well-established P+T CoMeBack framework in PRS, we developed

P+T CoMeBack MRS, which aggregates EWAS signals and could potentially be

used as a biomarker in association studies where single CpGs do not achieve

significance[4,40,41]. The proposed P+T CoMeBack MRS approach uses CoMeBack for

co-methylation pruning and evaluates multiple P-value thresholds to maximize

prediction performance. Such MRS could potentially serve as a powerful dimension

reduction approach for mediation and multi-omics integration analyses[4,40-43] as well

as biomarkers of individual disease risk in a clinical setting[44-46].

Overall, our simulation studies demonstrated good performance of P+T CoMeBack

MRS for predicting phenotypes of interest with good statistical power and well-

controlled type 1 error. We demonstrated that the prediction accuracy of MRS

reflects the variance of phenotype that is explained by DNAm. By accounting for

inter-correlation between CpGs, P+T CoMeBack MRS and P+T without CoMeBack

showed a slightly better performance than the standard T method. In the real data

application, we observed the best prediction of maternal smoking status when using

P+T CoMeBack, which confirms the usefulness of accounting for co-methylation and

demonstrates the ability of CoMeBack to control for false discover of CMR and

usefulness in constructing MRS[22]. However, we note that P+T CoMeBack MRS

could still have poor prediction performance if the external EWAS is underpowered

or subject to bias.

In the prediction of maternal smoking status, P+T CoMeBack MRS showed

comparable performance to Reese MRS, which was derived using the LASSO

method[37]. When predictors are highly correlated, LASSO typically selects one of the

correlated predictors and shrinks the effect size of the rest to zero, which might

produce similar results to our pruning procedure in developing MRS. One of the

advantages of P+T CoMeBack MRS is that it is based on EWAS summary statistics

which are often publicly available, hence making it a valuable approach, as it is often

difficult to obtain individual DNAm data from an external cohort. Additionally, P+T

CoMeBack MRS can make use of meta-analysis-type summary statistics, which

aggregates results from multiple studies to improve association estimates. In

contrast, to construct MRS like Reese MRS, individual DNAm data are usually

required to perform a LASSO regression, and these are often not accessible.

Recently, novel penalized regressions have been proposed to generate PRS with

only GWAS summary statistics and publicly available reference data[47], but their

applications to EWAS summary statistics for MRS have not been investigated. To

develop MRS for different exposures and outcomes, we urge EWAS studies to make

their genome-wide summary statistics publicly available.

In our simulation studies, weights obtained from Indian training samples were

applied to generate P+T CoMeBack MRS, thus MRS among Indian testing samples

were assumed to have the best prediction of the simulated phenotypes. However,

MRS among Whites and Blacks also achieved a prediction accuracy as high as

among Indians for both simulated phenotypes suggesting that genetic ancestry does

not contribute to difference in prediction abilities of MRS across multi-ancestry

population. This is likely because we assumed all ancestries share the same causal

CpGs and effect sizes. However, in the real world, this assumption could possibly be

violated for many phenotypes. Unlike ancestry in our simulation studies, ancestry in

the real world is complex. The meaning of ancestry could be different in different

regions/nations, and "effect of ancestry" involves the joint effects of ancestry-

associated social determinants of health and environmental effects , and cultural

context[48]. Ancestry, along with environment and social differences associated with it,

could affect both MRS and phenotypes in numerous causal pathways and potentially

modify the effect of MRS on the phenotypes. Thus, even if all ancestries indeed

share the same causal CpGs and effect sizes, it might still not be sufficient to

disentangle the relationship between ancestry, DNAm and phenotype of interest.

This may greatly impact the transferability of MRS across different ancestries, which

could be the reason why we observed an inconsistency of performance of P+T

CoMeBack MRS in terms of their distributions and predictions across multi-ancestry

population in the real data analyses. In practice, we recommend that researchers

conduct MRS analyses stratified by ancestry first and evaluate the effect of ancestry

on MRS analyses before pooling participants together for a joint analysis.

In our real data application, summary statistics for smoking were obtained from a

cohort with mainly people of European ancestry[36]. MRS of smoking among mixed

ancestry infants achieved a prediction accuracy of nearly 30%. However, the

prediction accuracy of P+T CoMeBack MRS among Black African infants was only

10.9%. We suspect that the difference was largely due to the prevalence of active

smoking among mothers of Black African infants being lower than those of mixed

ancestry infants (13% vs 49%), which is similar to how the prevalence of outcome

affects the predictive ability of PRS[49].

To the best of our knowledge, this is the first study to propose using CoMeBack for

pruning MRS among multi-ancestry populations. However, there are several

potential limitations that warrant mention. First, the sample size of both simulation

studies and real data analyses was relatively small, thus our results might not fully

capture the strengths and limitations of P+T CoMeBack MRS. Second, lack of

different ancestry-specific summary statistics made it impossible to compare the use

of external weights from population of different ancestries (e.g. European ancestry vs

other ancestries). Third, the prevalence of active maternal smoking is different in

different ancestries and has influenced the performance of P+T CoMeBack MRS. As

a result, real data analysis of smoking MRS could not provide firm evidence about

the transferability of MRS between Black African and mixed ancestry infants. Fourth,

we mainly focused on the prediction performance of P+T CoMeBack MRS. Further

studies are needed to assess the performance of P+T CoMeBack MRS in mediation

analysis.

In conclusion, P+T in general and P+T using CoMeBack in particular, provides an

improvement for prediction of phenotype of interest, over T method that does not

account for co-methylation between CpGs. In contrast to PRS, using existing

summary statistics that were derived from European populations can be used to

calculate MRS in other ancestries, thus reducing the ancestry/ethnicity disparity in

medical research. However, caution is needed in the analyses and interpretation of

MRS results across multi-ancestry populations. More investigations of MRS are

urged to further improve their prediction accuracy and translational values, also in

combination with other clinical and non-clinical variables, especially among multi-

ancestry population. With the current increase of large consortia-led EWAS for

different exposures and health outcomes (e.g., the PACE consortium), we believe

the predictive performance of MRS will continue to increase, and the P+T CoMeBack

method has the potential to be widely used for risk prediction and association testing.

Acknowledgments:

The authors thank the study and clinical staff at Paarl Hospital, Mbekweni and TC

Newman clinics, as well as the CEO of Paarl Hospital, and the Western Cape Health

Department for their support of the study. The authors thank the families and

children who participated in this study.

Table 1. Overview of included EWAS, their phenotypes, training sample and methods.

MRS	Training dataset publication	Training Population	Phenotype	MRS publication	P-value threshold/ Method	No. of CpG sites (joint-analysis)
P+T CoMeBack MRS	Sikdar et al. 2019[36]	Multi-ethnic newborns (mainly White, N=5,648)	Most cohorts ascertained sustained smoking during pregnancy by questionnaires; two cohorts incorporated cotinine-based smoking measure	-	5×10^{-22} (Mixed)	21 (43 passed P-value threshold and 22 excluded by pruning)
					5×10^{-24} (Black)	20 (42 passed P-value threshold and 22 excluded by pruning)
					5×10^{-22} (Pooled)	21 (43 passed P-value threshold and 22 excluded by pruning)
P+T MRS	Sikdar et al. 2019[36]	Multi-ethnic newborns (mainly White, N=5,648)	Same as above	-	5×10^{-10} (Mixed)	198 (233 passed P-value threshold and 35 excluded by pruning)
					5×10^{-36} (Black)	4 (26 passed P-value threshold and 22 excluded by pruning)
					5×10^{-24} (Pooled)	8 (42 passed P-value threshold and 34 excluded by pruning)

					5×10^{-9} (Mixed)	344
T MRS	Sikdar et al. 2019[36]	Multi-ethnic newborns (mainly White, N=5,648)	Same as above	-	5×10^{-24} (Black)	42
					5×10^{-16} (Pooled)	72
Reese MRS	Reese et al. 2017[37]	White newborns (N=1,068)*	Sustained smoking during pregnancy obtained from combined information of cotinine-based and self-report based classification	Reese et al. 2017[37]	Logistic LASSO regression	28
Richmond 568 MRS	Joubert et al. 2016[39]	Multi-ethnic newborns (N=6,685)*	Maternal smoking during pregnancy via questionnaires	Richmond et al. 2018[38]	Robust linear regression; Bonferroni corrected P-value < 0.05	568
Richmond 19 MRS	Joubert et al. 2016[39]	Multi-ethnic older children (average age = 6.8 years) (N=3,187)	Maternal smoking during pregnancy via questionnaires	Richmond et al. 2018	Robust linear regression; Bonferroni corrected P-value < 0.05	19

* These training populations overlapped with training population for summary statistics used for P+T MRS.

Table 2. Association between maternal smoking status and MRS in DCHS.

MRS	Mixed			Black			Pooled (Joint-analysis)		
	Beta-coefficient*	Standard Error	P-value	Beta-coefficient	Standard Error	P-value	Beta-coefficient	Standard Error	P-value
P+T CoMeBack MRS	0.88	0.12	5.65×10^{-11}	0.76	0.18	5.74×10^{-5}	0.82	0.10	3.01×10^{-14}
P+T MRS	0.71	0.11	1.05×10^{-8}	0.52	0.14	3.29×10^{-4}	0.69	0.10	1.32×10^{-10}
T MRS	0.80	0.13	7.35×10^{-9}	0.70	0.19	2.82×10^{-4}	0.64	0.09	2.55×10^{-11}
Reese MRS	0.97	0.13	3.77×10^{-11}	0.73	0.16	2.06×10^{-5}	0.77	0.09	2.50×10^{-15}
Richmond 568 MRS	0.85	0.14	9.84×10^{-9}	0.54	0.16	7.82×10^{-4}	0.65	0.10	1.57×10^{-10}
Richmond 19 MRS	0.76	0.14	$5.37E \times 10^{-7}$	0.70	0.19	3.78×10^{-4}	0.70	0.11	2.05×10^{-9}

*Beta-coefficients indicate change in interquartile range (IQR) of MRS between smokers and non-smokers.

Reference

1. Berger SL, Kouzarides T, Shiekhattar R, Shilatifard A. An operational definition of epigenetics. *Genes Dev*. Apr 1 2009;23(7):781-3. doi:10.1101/gad.1787609

2. Rakyan VK, Down TA, Balding DJ, Beck S. Epigenome-Wide Association Studies for common human diseases. *Nature reviews Genetics*. 07/12 2011;12(8):529-541. doi:10.1038/nrg3000

3. Wei S, Tao J, Xu J, et al. Ten Years of EWAS. *Adv Sci (Weinh)*. Aug 11 2021:e2100727. doi:10.1002/advs.202100727

4. Wahl S, Drong A, Lehne B, et al. Epigenome-wide association study of body mass index, and the adverse outcomes of adiposity. *Nature*. Jan 5 2017;541(7635):81-86. doi:10.1038/nature20784

5. McCartney DL, Hillary RF, Stevenson AJ, et al. Epigenetic prediction of complex traits and death. *Genome biology*. 2018;19(1):136-136. doi:10.1186/s13059-018-1514-1

6. Heiss JA, Brenner H. Epigenome-wide discovery and evaluation of leukocyte DNA methylation markers for the detection of colorectal cancer in a screening setting. *Clinical epigenetics*. 2017;9:24-24. doi:10.1186/s13148-017-0322-x

7. Gelato KA, Shaikhibrahim Z, Ocker M, Haendler B. Targeting epigenetic regulators for cancer therapy: modulation of bromodomain proteins, methyltransferases, demethylases, and microRNAs. *Expert Opin Ther Targets*. Jul 2016;20(7):783-99. doi:10.1517/14728222.2016.1134490

8. Krushkal J, Silvers T, Reinhold WC, et al. Epigenome-wide DNA methylation analysis of small cell lung cancer cell lines suggests potential chemotherapy targets. *Clinical epigenetics*. 2020;12(1):93-93. doi:10.1186/s13148-020-00876-8

9. Guan Z, Yu H, Cuk K, Zhang Y, Brenner H. Whole-Blood DNA Methylation Markers in Early Detection of Breast Cancer: A Systematic Literature Review. *Cancer Epidemiol Biomarkers Prev*. Mar 2019;28(3):496-505. doi:10.1158/1055-9965.Epi-18-0378

10. Hüls A, Czamara D. Methodological challenges in constructing DNA methylation risk scores. *Epigenetics*. Jan-Feb 2020;15(1-2):1-11. doi:10.1080/15592294.2019.1644879

11. Wray NR, Lee SH, Mehta D, Vinkhuyzen AA, Dudbridge F, Middeldorp CM. Research review: Polygenic methods and their application to psychiatric traits. *J Child Psychol Psychiatry*. Oct 2014;55(10):1068-87. doi:10.1111/jcpp.12295

12. Martin EM, Fry RC. Environmental Influences on the Epigenome: Exposure-Associated DNA Methylation in Human Populations. *Annual Review of Public Health*. 2018/04/01 2018;39(1):309-333. doi:10.1146/annurev-publhealth-040617-014629

13. Notterman DA, Mitchell C. Epigenetics and Understanding the Impact of Social Determinants of Health. *Pediatr Clin North Am*. Oct 2015;62(5):1227-40. doi:10.1016/j.pcl.2015.05.012

14. Borrell LN, Elhawary JR, Fuentes-Afflick E, et al. Race and Genetic Ancestry in Medicine — A Time for Reckoning with Racism. *New England Journal of Medicine*. 2021/02/04 2021;384(5):474-480. doi:10.1056/NEJMms2029562

15. Cronjé HT, Elliott HR, Nienaber-Rousseau C, Pieters M. Replication and expansion of epigenome-wide association literature in a black South African population. *Clinical Epigenetics*. 2020/01/07 2020;12(1):6. doi:10.1186/s13148-019-0805-z

16. Khera AV, Chaffin M, Zekavat SM, et al. Whole-Genome Sequencing to Characterize Monogenic and Polygenic Contributions in Patients Hospitalized With Early-Onset Myocardial Infarction. *Circulation*. Mar 26 2019;139(13):1593-1602. doi:10.1161/circulationaha.118.035658

17. Martin AR, Kanai M, Kamatani Y, Okada Y, Neale BM, Daly MJ. Clinical use of current polygenic risk scores may exacerbate health disparities. *Nature Genetics*. 2019/04/01 2019;51(4):584-591. doi:10.1038/s41588-019-0379-x

18. Horvath S. DNA methylation age of human tissues and cell types. *Genome biology*. 10/21

06/10/received

10/21/accepted 2013;14(10):R115-R115. doi:10.1186/gb-2013-14-10-r115

19. Thompson M, Hill BL, Rakocz N, et al. Methylation risk scores are associated with a collection of phenotypes within electronic health record systems. *npj Genomic Medicine*. 2022/08/25 2022;7(1):50. doi:10.1038/s41525-022-00320-1

20. Goldstein BA, Yang L, Salfati E, Assimes TL. Contemporary Considerations for Constructing a Genetic Risk Score: An Empirical Approach. *Genet Epidemiol*. Sep 2015;39(6):439-45. doi:10.1002/gepi.21912

21. Euesden J, Lewis CM, O'Reilly PF. PRSice: Polygenic Risk Score software. *Bioinformatics (Oxford, England)*. 2015;31(9):1466-1468. doi:10.1093/bioinformatics/btu848

22. Gatev E, Gladish N, Mostafavi S, Kobor MS. CoMeBack: DNA methylation array data analysis for co-methylated regions. *Bioinformatics*. May 1 2020;36(9):2675-2683. doi:10.1093/bioinformatics/btaa049

23. Shah S, Bonder MJ, Marioni RE, et al. Improving Phenotypic Prediction by Combining Genetic and Epigenetic Associations. *Am J Hum Genet*. Jul 2 2015;97(1):75-85. doi:10.1016/j.ajhg.2015.05.014

24. Odintsova VV, Rebattu V, Hagenbeek FA, et al. Predicting Complex Traits and Exposures From Polygenic Scores and Blood and Buccal DNA Methylation Profiles. *Front Psychiatry*. 2021;12:688464. doi:10.3389/fpsyt.2021.688464

25. Wu J, Pfeiffer RM, Gail MH. Strategies for developing prediction models from genome-wide association studies. *Genet Epidemiol*. Dec 2013;37(8):768-77. doi:10.1002/gepi.21762

26. Privé F, Vilhjálmsson BJ, Aschard H, Blum MGB. Making the Most of Clumping and Thresholding for Polygenic Scores. *The American Journal of Human Genetics*. 2019/12/05/ 2019;105(6):1213-1221. doi:https://doi.org/10.1016/j.ajhg.2019.11.001

27. Choi SW, O'Reilly PF. PRSice-2: Polygenic Risk Score software for biobank-scale data. *GigaScience*. 2019;8(7)doi:10.1093/gigascience/giz082

28. Vanker A, Barnett W, Workman L, et al. Early-life exposure to indoor air pollution or tobacco smoke and lower respiratory tract illness and wheezing in African infants: a longitudinal birth cohort study. *Lancet Planet Health*. Nov 2017;1(8):e328-e336. doi:10.1016/s2542-5196(17)30134-1

29. Leek JT, Johnson WE, Parker HS, Jaffe AE, Storey JD. The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics*. Mar 15 2012;28(6):882-3. doi:10.1093/bioinformatics/bts034

30.	Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*. Apr 20 2015;43(7):e47. doi:10.1093/nar/gkv007

31.	Barfield RT, Almli LM, Kilaru V, et al. Accounting for population stratification in DNA methylation studies. *Genet Epidemiol*. Apr 2014;38(3):231-41. doi:10.1002/gepi.21789

32.	Teschendorff AE, Breeze CE, Zheng SC, Beck S. A comparison of reference-based algorithms for correcting cell-type heterogeneity in Epigenome-Wide Association Studies. *BMC Bioinformatics*. Feb 13 2017;18(1):105. doi:10.1186/s12859-017-1511-5

33.	Zar HJ, Barnett W, Myer L, Stein DJ, Nicol MP. Investigating the early-life determinants of illness in Africa: the Drakenstein Child Health Study. *Thorax*. Jun 2015;70(6):592-4. doi:10.1136/thoraxjnl-2014-206242

34.	Stein DJ, Koen N, Donald KA, et al. Investigating the psychosocial determinants of child health in Africa: The Drakenstein Child Health Study. *J Neurosci Methods*. Aug 30 2015;252:27-35. doi:10.1016/j.jneumeth.2015.03.016

35.	Hüls A, Wedderburn CJ, Groenewold NA, et al. Newborn differential DNA methylation and subcortical brain volumes as early signs of severe neurodevelopmental delay in a South African Birth Cohort Study. *World J Biol Psychiatry*. Jan 12 2022:1-12. doi:10.1080/15622975.2021.2016955

36.	Sikdar S, Joehanes R, Joubert BR, et al. Comparison of smoking-related DNA methylation between newborns from prenatal exposure and adults from personal smoking. *Epigenomics*. Oct 2019;11(13):1487-1500. doi:10.2217/epi-2019-0066

37.	Reese SE, Zhao S, Wu MC, et al. DNA Methylation Score as a Biomarker in Newborns for Sustained Maternal Smoking during Pregnancy. *Environ Health Perspect*. Apr 2017;125(4):760-766. doi:10.1289/ehp333

38.	Richmond RC, Suderman M, Langdon R, Relton CL, Davey Smith G. DNA methylation as a marker for prenatal smoke exposure in adults. *International Journal of Epidemiology*. 2018;47(4):1120-1130. doi:10.1093/ije/dyy091

39.	Joubert Bonnie R, Felix Janine F, Yousefi P, et al. DNA Methylation in Newborns and Maternal Smoking in Pregnancy: Genome-wide Consortium Meta-analysis. *The American Journal of Human Genetics*. 2016/04/07/ 2016;98(4):680-696. doi:https://doi.org/10.1016/j.ajhg.2016.02.019

40.	Yu H, Raut JR, Schöttker B, Holleczek B, Zhang Y, Brenner H. Individual and joint contributions of genetic and methylation risk scores for enhancing lung cancer risk stratification: data from a population-based cohort in Germany. *Clin Epigenetics*. Jun 18 2020;12(1):89. doi:10.1186/s13148-020-00872-y

41.	Westerman K, Fernández-Sanlés A, Patil P, et al. Epigenomic Assessment of Cardiovascular Disease Risk and Interactions With Traditional Risk Metrics. *J Am Heart Assoc*. Apr 21 2020;9(8):e015299. doi:10.1161/jaha.119.015299

42.	Guan Z, Raut JR, Weigl K, et al. Individual and joint performance of DNA methylation profiles, genetic risk score and environmental risk scores for predicting breast cancer risk. *Mol Oncol*. 2020;14(1):42-53. doi:10.1002/1878-0261.12594

43.	Grant CD, Jafari N, Hou L, et al. A longitudinal study of DNA methylation as a potential mediator of age-related diabetes risk. *Geroscience*. Dec 2017;39(5-6):475-489. doi:10.1007/s11357-017-0001-z

44.   García-Calzón S, Perfilyev A, Martinell M, et al. Epigenetic markers associated with metformin response and intolerance in drug-naïve patients with type 2 diabetes. *Sci Transl Med*. Sep 16 2020;12(561)doi:10.1126/scitranslmed.aaz1803

45.   Deng Y, Wan H, Tian J, et al. CpG-methylation-based risk score predicts progression in colorectal cancer. *Epigenomics*. Apr 2020;12(7):605-615. doi:10.2217/epi-2019-0300

46.   Kilanowski A, Chen J, Everson T, et al. Methylation risk scores for childhood aeroallergen sensitization: Results from the {LISA} birth cohor. *Authorea*. Nov 2021;doi:10.22541/au.163620398.85835627/v1

47.   Pattee J, Pan W. Penalized regression and model selection methods for polygenic scores on summary statistics. *PLoS Comput Biol*. Oct 2020;16(10):e1008271. doi:10.1371/journal.pcbi.1008271

48.   VanderWeele TJ, Robinson WR. On the causal interpretation of race in regressions adjusting for confounding and mediating variables. *Epidemiology*. Jul 2014;25(4):473-84. doi:10.1097/ede.0000000000000105

49.   Gibson G. On the utilization of polygenic risk scores for therapeutic targeting. *PLoS Genet*. Apr 2019;15(4):e1008060. doi:10.1371/journal.pgen.1008060

Figure 1. Simulation study. Prediction R² of P+T CoMeBack, P+T and T method in dependence of (A) the proportion of causal CpG sites in CMRs and (B) proportion of phenotype variance explained by DNA methylation, among Indian participants. For each simulation, the discovery cohort was repeatedly and randomly split into a training set comprising 762 Indians and a testing set comprising 136 people of the same ancestry. Phenotypes were simulated without an influence of ancestry. Results are shown for (**A**) different proportions of causal CpGs located in CMR (30%, 50%, 70%, 100%) and (**B**) different proportions of phenotype variance explained by DNA methylation (10%, 30%, 50%, 80%). Each box represents the distribution of prediction accuracy across 1000 simulations, where the central mark is the median and the edges of the box are the 25th and 75th percentiles.

Figure 2. Simulation study. Prediction R^2 of P+T CoMeBack and T approach across different racial groups and among multi-ancestry populations. For each simulation, the discovery cohort was repeatedly and randomly split into a training set comprising 762 Indians and a testing set comprising 136 people of each ancestry group. The proportion of causal CpGs located in CMR is 70% and the proportion of phenotype variance explained by DNA methylation (and ancestry) is 80%. Results are shown for the prediction of simulated phenotypes (**2A**) without an influence of ancestry and (**2B**) influenced by ancestry. Joint-analysis refers to MRS analyses of all participants pooled from all ancestry groups and standardization refers to standardizing MRS within each ancestry group and then merging all participants before analyses. Each box represents the distribution of prediction accuracy across 1000 simulations, where the central mark is the median and the edges of the box are the 25th and 75th percentiles.

Figure 3. Real data application. MRS for the prediction of maternal smoking during pregnancy using cord blood DNA methylation data from newborns in the South African Drakenstein Child Health Study (DCHS). Prediction R^2 of maternal smoking status is shown stratified for **A.** Mixed infants. **B.** Black infants. **C.** joint-analysis (all subjects pooled from all ancestries) **D.** Standardization (standardizing MRS within each ancestry and merging all subjects before analyses

Figure 4. Real data application. Comparison of P+T CoMeBack method to P+T, T and 3 other published MRS for predicting maternal smoking status in the South African Drakenstein Child Health Study (DCHS). A Prediction R^2 of all 6 MRS methods for Mixed infants, Black infants and pooled samples (joint-analysis). A receiver operating characteristic (ROC) curve comparing prediction performance of all 6 MRS among (**B**) Mixed infants, (**C**) Black infants and (**D**) pooled samples (joint-analysis).