Strategies for using antigen rapid diagnostic tests to reduce transmission of SARS-CoV-2 in low- and middle-income countries: a mathematical modelling study applied to Zambia

Alvin X. Han[1,*], Sarah Girdwood[2], Shaukat Khan[3], Jilian A. Sacks[4], Amy Toporowski[4], Naushin Huq[4], Emma Hannay[4], Colin A. Russell[1,5,†], Brooke E. Nichols[1,4,5,†]

[1]Department of Medical Microbiology & Infection Prevention, Amsterdam University Medical Center, University of Amsterdam, Amsterdam, The Netherlands

[2]Health Economics and Epidemiology Research Office, Department of Internal Medicine, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

[3]Clinton Health Access Initiative, Boston, MA, USA

[4]Foundation for Innovative New Diagnostics (FIND), Geneva, Switzerland

[5]Department of Global Health, School of Public Health, Boston University, Boston, MA, USA

†Contributed equally

*Correspondence to Alvin X. Han (x.han@amsterdamumc.nl)

Keywords: COVID-19, SARS-CoV-2, Diagnostic testing, Low- and middle-income countries

Summary (229/250 words)

Background

Increasing the availability of antigen rapid diagnostic tests (Ag-RDTs) in low- and middle-

income countries (LMICs) is key to alleviating global SARS-CoV-2 testing inequity (median

testing rate in December 2021-March 2022 when the Omicron variant was spreading in

multiple countries; high-income countries=600 tests/100,000 people/day; LMICs=14 tests/

100,000 people/day). However, target testing levels and effectiveness of asymptomatic

community screening to impact SARS-CoV-2 transmission in LMICs are unclear.

Methods

We used PATAT, an LMIC-focused agent-based model to simulate COVID-19 epidemics,

varying the amount of Ag-RDTs available for symptomatic testing at healthcare facilities and

asymptomatic community testing in different social settings. We assumed that testing was a

function of access to healthcare facilities and availability of Ag-RDTs. We explicitly

modelled symptomatic testing demand from non-SARS-CoV-2 infected individuals and

measured impact based on the number of infections averted due to test-and-isolate.

Results

Testing symptomatic individuals yields greater benefits than any asymptomatic community

testing strategy until most symptomatic individuals who sought testing have been tested.

Meeting symptomatic testing demand likely requires at least 200-400 tests/100,000

people/day on average as symptomatic testing demand is highly influenced by non-SARS-

CoV-2 infected individuals. After symptomatic testing demand is satisfied, excess tests to

proactively screen for asymptomatic infections among household members yields the largest

additional infections averted.

Conclusions

Testing strategies aimed at reducing transmission should prioritize symptomatic testing and

incentivizing test-positive individuals to adhere to isolation to maximize effectiveness.

Main Text

Introduction

Since the emergence of SARS-CoV-2 in 2019, the COVID-19 pandemic has resulted in over 500 million confirmed cases and 6 million deaths worldwide as of May 2022.[1] While vaccination is the key medical intervention to mitigate the pandemic, SARS-CoV-2 testing remains an important public health tool for case identification and transmission reduction. Testing is especially important in many low- and middle-income countries (LMICs) that continue to struggle with gaining equitable access to global vaccine supplies.[2] Testing is also the backbone of surveillance systems to monitor the emergence of novel variants of concern (VOCs)[3] that may escape immunity acquired from previous infections and vaccination.[4]

At the same time, the global imbalance in SARS-CoV-2 testing rates is substantial.[5] Between December 2021 and March 2022 when the Omicron (BA.1) VOC was spreading in multiple countries, the median testing rate in LMICs was 14 (IQR = 7-41) tests per 100,000 persons per day (/100k/day), whereas HICs tested >43 times more, with a median rate of 603 (IQR = 317-1181) tests/100k/day.[6] Limited testing has likely led to substantial underestimation of SARS-CoV-2 prevalence and COVID-19 attributable mortality in LMICs.[7] The diagnostics pillar of the Global Access to COVID-19 Tools (ACT) Accelerator, co-convened by FIND and the Global Fund in partnership with the World Health Organization to enhance access to COVID-19 tests and sequencing, has set a minimum testing rate of 100 tests/100k/day.[8] This minimum testing rate is thought to be a "critical threshold to facilitate effective public health interventions".[8] Additionally, asymptomatic testing in a community setting (i.e. community testing) was identified by this initiative as a crucial step for LMICs to close in on the global equity gap.[8]

Real-time reverse transcription polymerase chain reaction (PCR) tests remain the gold

standard for COVID-19 diagnostic testing, as they are the most sensitive testing method.[9]

However, PCR-based testing can be plagued by long turnaround times and necessitate

relatively costly laboratory infrastructures, robust sample transport networks and well-trained

personnel that are lacking in many low-resourced settings.[10] Furthermore, RNA can still be

detected even after infectiousness has declined, rendering PCR tests imperfect for

determining the infectious potential of an infected person.[11] While the sensitivity of

antigen rapid diagnostic tests (Ag-RDTs) is lower than PCR(>80%)[12], Ag-RDTs are

cheaper, capable of producing results in under 30 minutes and can be performed easily at

point-of-care.[13] As such, when used in a timely fashion, Ag-RDTs can identify potentially

infectious people more quickly. Ag-RDTs offer a practical alternative diagnostic tool to

enable massive scale up of testing in all countries. In resource-limited settings, Ag-RDTs

could potentially reduce the testing equity gap between HICs and LMICs.[5]

To date, there is no robust evidence-base on how scaling-up Ag-RDTs to 100 tests/100k/day

would impact community transmissions. There is also a lack of information on the

effectiveness of community testing programs when used to complement symptomatic testing

under the constraints of limited test availability. Furthermore, while there have been studies

investigating the impact of comprehensive test-and-trace programs on transmission

reductions,[14,15], it is less clear to what extent a test-and-isolation strategy (i.e. the only

intervention is the required isolation of individuals with a positive diagnosis) would impact

total infections. It is thus important to estimate the impact of test-and-isolation only since

most low-resource settings did not implement resource-intensive contact tracing

programs.[16]

In this study, we developed and used the Propelling Action for Testing and Treating

(PATAT) simulation model, an agent-based modelling framework to investigate the impact

of using Ag-RDTs for healthcare facility-based symptomatic testing. We considered testing

programs both with and without additional asymptomatic testing programs in the community,

using a population with demographic profiles, contact mixing patterns, and levels of public

health resources akin to those in many LMICs. We used PATAT to interrogate how different

Ag-RDT distribution availability and testing strategies, including the implementation of

community testing in households, schools, formal workplaces and regular mass gatherings

such as religious gatherings, could impact onward disease transmission. In turn, we aimed to

identify key priorities and gaps that should be addressed when implementing mass testing

programs using Ag-RDTs in low-resource settings.

Methods

*The **P**ropelling **A**ction for **T**esting **A**nd **T**reating (PATAT) simulation model*

PATAT first creates an age-structured population of individuals within contact networks of

multi-generational households, schools, workplaces, religious gatherings (i.e. regular mass

gatherings) and random community with the given demographic data here based on

archetypal LMIC estimates (Figure 1). The simulation starts with a user-defined proportion of

individuals infected with SARS-CoV-2. Given that viral loads of an infected individual at the

time of testing affect Ag-RDT sensitivity,[12] PATAT randomly draws a within-host viral

load trajectory over the course of each individual's infection from known distribution of

trajectories[17,18] using previously developed methods.[19] Given conflicting evidence,[20]

similar viral load trajectories were drawn for both asymptomatic and symptomatic infected

individuals.

Figure 1: **Schematic of <u>P</u>ropelling <u>A</u>ction for <u>T</u>esting <u>A</u>nd <u>T</u>reating (PATAT) simulation model**.

The simulation computes transmission events across different contact networks each day and

updates the disease progression of infected individuals based on the Susceptible-Exposed-

Infected-Recovered-Death (SEIRD) epidemic model, stratifying them based on symptom

presentation (asymptomatic, mild or severe). Symptomatic individuals may seek symptomatic

testing at clinics after symptom onset. Given that most LMICs are currently testing at rates

far below 100 tests/100k/day to the extent that only a small proportion of COVID-19 positive

deaths were identified in life (e.g. <10% in Zambia),[21] we assumed that all clinic-provided

testing demand by mild symptomatic individuals are satisfied by Ag-RDTs while PCR tests

are restricted for testing severe patients only. Positively-tested individuals may go into

isolation and their household members may also be quarantined.

Simulation variables

We assumed a populations size of 1,000,000 individuals, creating contact networks and

healthcare facilities based on demographic data collected from Zambia.[22] We initialized

each simulation with 1% of the population being infected by SARS-CoV-2 and ran the model

over a 90-day period. We permutated a range of R_e values (i.e. 0.9, 1.1, 1.2, 1.5, 2.0, 2.5 and

3.0) against varying Ag-RDT stock availability (i.e. 100, 200-1,000 (in 200 increments),

1,000-5,000 (in 1,000 increments) tests per 100,000 persons per day). Various test

distribution strategies were simulated: (1) 85% of weekly allocated tests were used for

routine asymptomatic community testing at a social setting and the remaining used for

symptomatic testing at healthcare facilities; (2) all weekly allocated test stocks are distributed

to healthcare facilities for symptomatic testing only with no asymptomatic community testing

and (3) weekly allocated tests are used for symptomatic testing at healthcare facilities first

before any remaining tests at the end of the week are used for asymptomatic community

testing in the next week. As a baseline, we simulated a set of runs using the same range of R_e

with no testing at all.

To determine impact of testing on reducing infections, we assumed that the only public health

intervention measure is by test-and-isolate of positive-tested individuals. We also performed

a separate set of simulations that require a same-day quarantine of asymptomatic household

members of positively-tested individuals. This distinction is important because quarantine

should change contact patterns of more individuals per positive test, thereby increasing test

utility. We did not consider the quarantine of close contacts outside of household members as

contact tracing programs are often resource intensive and discontinued in most countries.

Distribution of routine asymptomatic community test

Due to their fixed nature and potential accessibility, routine asymptomatic community testing

may be implemented in households, schools, formal workplaces or religious gatherings.

Community tests stocks may be distributed in each setting in two ways: (1) even distribution

to as many entities as possible once per week (e.g. if we have 10 tests available for 10

households per week, then one member of each household would be tested); (2) concentrated

distribution to test all individuals in selected entities twice a week who will continue to get

tested throughout the epidemic (e.g. if we have 10 tests available for 10 households per week

but only one housing of 5 members, then all 10 tests will be distributed to this selected

household of 5 for testing on Monday and Thursday of every week).

Healthcare provided symptomatic testing demand

Symptomatic testing demand estimation is particularly challenging for SARS-CoV-2 because

COVID-19 symptoms overlaps with other respiratory infections, thus increasing testing

demand. We assumed that symptomatic individuals would seek testing at clinics based on a

probability distribution that inversely correlate with the distance between their homes and the

nearest clinic[23] (Table S1). We assumed that the time delay between testing and symptom

onset follows lognormal distribution of with mean of one day and standard deviation of 0.5

day. Additionally, we simulated daily demand of clinic tests from individuals that were not

infected by SARS-CoV-2 but sought symptomatic testing as they presented with COVID-19-

like symptoms. This non-COVID-19 related demand was estimated by assuming a 10% test

positivity rate at the start as well as end of an epidemic curve and 20% test positivity rate at

the peak, linearly interpolating the demand for periods between these time points (Figure 2).

These assumptions are based on observed test positivity rates in multiple countries

experiencing infection waves during the second half of 2021.[24] If there are limited clinic

test stocks for the day, the available tests are randomly distributed among symptomatic

SARS-CoV-2-infected patients and those seeking tests for non-COVID-19 related reasons.

We assumed that any individual who failed to receive a test due to test shortage would not

seek clinic-provided testing again for the rest of their infection. If these individuals had

previously decided to self-isolate upon presenting symptoms, they may continue to do so (see

Supplementary Data). Otherwise, we assumed that they would continue to mix with the

community. We also assumed that SARS-CoV-2 infected individuals who were tested but

received a false negative result continue mixing with the community. In turn, any false

positive tested individual would then go into isolation.

All key parameters are tabulated in Table S1 and full details of PATAT are described in the

Supplementary Data. The PATAT model source code is available at

https://github.com/AMC-LAEB/PATAT-sim.

Figure 2: **Projected symptomatic testing demand based on assumed case positivity rate**. This projected
demand includes both SARS-CoV-2 infected persons who were tested and reported as well as those who seek
symptomatic testing for other reasons (e.g. individuals presenting COVID-19-like symptoms but were not
infected with SARS-CoV-2).
210

Results

Healthcare provided symptomatic testing demand should be fulfilled first

We first investigated if routine asymptomatic community testing programs could substantially reduce SARS-CoV-2 transmissions in the population. We compared scenarios where either all Ag-RDT stocks were used only for symptomatic testing (with no community-based testing of asymptomatic populations), or that most tests were used for community testing and only 15% of weekly available stocks were allocated for symptomatic testing. The proportion of infections averted under each test distribution strategy was computed as a measure of impact. Regardless of community testing distribution strategy or R_e, we found that setting aside large proportions of Ag-RDTs for community testing led to lower proportion of infections averted than if all tests were solely used for symptomatic testing (Figure 3). A far greater number of tests is needed under the community testing scenarios relative to the only-symptomatic one to result in an equal or larger proportion of infections averted. This conclusion remains the same when household members of all positively tested individuals were quarantined (Figure S1).

Figure 3: **Impact of either using all available Ag-RDT for symptomatic testing or a majority of them (85%) for community testing in various settings (even distribution only; <u>without</u> quarantine of household members)**. The proportion of secondary infections averted after 90 days relative to the no testing baseline for different number of tests available per 100,000 persons per day and assumed R_e value is plotted for each test distribution strategy. The vertical red line denotes the number of tests required to saturate symptomatic testing demand (Figure 4).

Number of tests needed to saturate healthcare provided symptomatic testing demand

Given the importance of saturating symptomatic testing demand, we then estimated the number of tests needed to saturate symptomatic testing demand under different R_e (Figure 3 and 4A). Besides symptomatic SARS-CoV-2 infected individuals who sought testing, our simulations factored in that 80-90% of symptomatic test stocks were used by individuals who were not infected by SARS-CoV-2 by assuming that test positivity rate ranges at 10%-20% over the course of the infection wave. Under these assumptions, even when $R_e \leq 1.2$ (Figure 4B), at least 200-400 tests/100k/day was needed to ensure all test-seeking individuals were tested. If $R_e \geq 1.5$, at least 10 times more tests, in the range of 2,000-5,000 tests/100k/day, was needed to satisfy all symptomatic testing demand. These conclusions were similar when household members of positive-tested individuals were quarantined (Figure S2).

Figure 4: **Symptomatic testing demand during an epidemic (<u>without</u> quarantine of household members)**.
(**A**) Number of symptomatic tests performed per 100,000 persons per day over time for different R_e. Each differently colored shaded curve denotes a different number of tests available per 100,000 persons per day. We assumed that all healthcare facilities in the community will have new stocks of one week's worth of Ag-RDTs every Monday. The symptomatic testing demand include both symptomatic SARS-CoV-2 infected individuals who seek testing at healthcare facilities and those who seek symptomatic testing for other reasons based on assumed case positivity rates (see Methods). The area between the curve plotting the testing rate needed to saturate symptomatic testing demand (N_{sat}) and the curve for testing rate $< N_{sat}$ is the amount of symptomatic testing shortage accumulated over time between those two testing rates. (**B**) 7-day moving average of time-varying effective reproduction number (R_e) over simulated epidemic period (90 days) assuming that testing demand is fully satisfied.

Marginal impact of symptomatic testing prior to saturating demand

We linearly regressed the number of infections averted against test availability to compute

the number of additional infections averted with increasing Ag RDT availability, before

saturating symptomatic testing demand (Figures 5A-B). Assuming only symptomatic cases

that test positive isolate, the largest marginal benefit of increasing Ag-RDT availability for

symptomatic testing prior to demand saturation is achieved when R_e = 1.1-1.2, with close to

20,000 additional infections averted for every increase of 100 more Ag-RDTs available for

268　symptomatic testing (Figure 5B; Table 1). When operating at tests availability that meet all

269　symptomatic testing demand, the greatest impact of test-and-isolate is also achieved when R_e

270　= 1.1-1.2 with ~40% of total infections averted (Figure 5A).

Figure 5: **Marginal impact of symptomatic testing prior to saturating demand (<u>without</u> quarantine of household members).** (**A**) Contour plots depicting infections averted relative to the no testing baseline for simulations with different R_e values and varying number of Ag-RDTs availability. Number of infections averted relative to no testing baseline after 90 days (left panel); Proportion of secondary infections averted relative to no testing baseline after 90 days (right panel). (**B**) Number of additional infections averted for every 100 more Ag-RDTs available prior to saturating symptomatic testing demand for different R_e values. Dashed red line shows marginal benefit with quarantine of household members while solid black line depicts that without quarantine. (**C**) Mean daily percentage reduction in transmissions while time-varying R_e of simulated epidemic is still > 1 for different initial R_e values and varying number of Ag-RDTs available for symptomatic testing only. (**D**) Reduction in number of days when time-varying R_e of simulated epidemic is > 1 for different initial R_e values and varying number of Ag-RDTs available for symptomatic testing only.

w/ quarantine of household members	R_e	No. of additional infections averted per 100 more tests
No	0.9	1,772
	1.1	19,807
	1.2	19,372
	1.5	3,655
	2.0	1,149
	2.5	401
	3.0	216
Yes	0.9	2,205
	1.1	23,444
	1.2	23,250
	1.5	5,702
	2.0	1,999
	2.5	853
	3.0	441

Table 1: **Number of additional infections averted for every 100 more Ag-RDTs available prior to**

saturating symptomatic testing demand for different R_0 values.

Both the marginal benefit and the maximum infection reduction at demand saturation,

however, diminish exponentially with increasing values of R_e (Figure 5A-B and Table 1).

Nonetheless, there are other impacts that could be gained from performing more symptomatic

testing at values of $R_e > 1.2$. For instance, for R_e values between 1.5 and 2.0 without

quarantining household members, it is possible to reduce daily transmissions by up to 11%

with increasing levels of test availability during the growth phase of the epidemic ($R_e > 1$;

Figure 5C). Additionally, when $R_e \sim 1.5$ and test availability is in the range of 2,000

tests/100K/day or more, it is possible to shorten the duration of the epidemic's growth phase

(and in turn, the epidemic itself) by about one week (Figure 5D).

The marginal benefit of symptomatic testing can be further augmented if asymptomatic

household members of positively-tested individuals quarantine as well (Figure S3). However,

depending on R_e and level of test availability, the percentage of infections averted only

improved modestly by 2-10%. As we assumed that individuals would isolate and quarantine

in their own homes, infectious individuals in isolation may infect healthy household members

in quarantine with them.

A symptomatic-testing-first strategy to community testing

Figure 6: **Symptomatic-testing-first strategy to community testing (<u>without</u> quarantine of household
members)**. When community testing is performed under this strategy, the leftover tests from the previous
week's stock allocated for symptomatic testing are used for community testing in various setting in the current
week. Two different types of community test distributions approaches (even or concentrated; see Methods) were
simulated. The proportion of secondary infections averted after 90 days relative to the no testing baseline for
different number of tests available per 100,000 persons per day and assumed R_e value is plotted for each test
distribution strategy. The vertical red line denotes the number of tests required to saturate symptomatic testing
demand.

Given the importance of symptomatic testing, we then simulated an alternate community

testing strategy that prioritizes saturating symptomatic testing demand first every week. If

there were leftover tests from clinics in the previous week, they were used for community

testing in the following week. We also investigated two ways in which community tests were

either evenly and randomly distributed in the social setting to as many individuals as possible

or concentrate the available tests to a fixed number of persons throughout the epidemic

period.

Even under this symptomatic-testing-first approach, other than households, community

testing in almost all social settings only yields greater reduction in infections when test

availability is higher than what is needed to saturate symptomatic testing needs (Figure 6).

Overall, household community testing yielded the greatest reduction in transmissions for all

simulated R_e values, followed by schools if $R_e \leq 1.5$. Community testing in religious

gatherings and formal workplaces only results in modest improvements over symptomatic

testing. An even distribution of community tests tends to produce larger reduction in

infections. The difference between even and concentrated community test distributions also

increases with larger test availability. These results were similarly observed when household

members of positively-tested individuals were quarantined (Figure S4).

Figure 7: Routine community testing in households outperforms other settings. (**A**) Average breakdown of infections based on the social setting where transmissions occurred for the simulations presented in this work. (**B**) Results from simulations using different testing strategies where $R_e = 1.5$, no quarantine of household members of positively-tested individuals assumed, and Ag-RDT availability of 5,000 tests per 100,000 persons per day. Community testing (even distribution) was performed with a symptomatic-testing-first approach. The average total number of diagnosed cases (left), time-varying reproduction number (R_t; middle) and number of infections averted (right) over the epidemic period are plotted. (**C**, **D**) Transmissions across distinct social settings. The top row of stacked plots shows the proportion of infections stratified by the source settings where infectors were infected for each sink setting where their infectees were infected. The stacked bars are colored by the source settings as per the bottom row of bar plots. The bottom row of bar plots shows the contribution of transmission exports into other settings (i.e. transmission events where the infectee were infected in a setting that is different from their infectors) from different source settings where the infectors were infected. (**C**) No testing baseline results from the example case as in (B). (**D**) Results from either implementing a symptomatic-testing-first community testing in households (left column) or religious gathering (right column). The dashed bar outlines are the no testing baseline results as in (C).

In the symptomatic-testing-first approach, household community testing can achieve greater

reduction in transmissions before saturating symptomatic testing demand but only at high

levels of test availability (>1,000 tests/100k/day; Figure 6 and S4). There are several reasons

why household community testing outperformed other settings. First, large multigenerational

homes (mean household size = 5 people) were simulated to mirror what is often found in

many LMICs. Second, population in LMICs tend to skew young (i.e. 48% of the population

are expected to be ≤ 15 years in age).[22] Furthermore, overall employment rates are low

(i.e. assumed 39% and 23% among men and women respectively)[22] and a large majority of

employed individuals likely work in informal employment settings (i.e. assumed 64% and

76% among employed men and women respectively)[22] where test distribution is assumed

to be difficult or infeasible. Third, dedicated isolation and quarantine facilities are likely rare

in low-resource settings. Thus, positively-tested individuals and their close contacts could

only isolate/quarantine themselves in their own homes. In turn, almost 60% of all infections

observed in a typical simulation arose from transmissions in households. Random community

transmissions aside, schools are then the second most common setting where transmissions

occurred (~14%) and workplaces, be if formal or informal, the least common (<3%) (Figure

7A).

Interestingly, even though we assumed that 70% of all households regularly attended large

religious gatherings weekly, they contributed to a limited proportion of total infections (~5%)

(Figure 7A). Yet, if we compare the results between household and religious gathering testing

at levels of test availability large enough to satisfy symptomatic testing demand (e.g. $N =$

5,000), the total number of diagnosed cases over time is actually similar for both community

testing strategies (Figure 7B). In fact, testing in religious gatherings yielded relatively larger

number of cumulative diagnoses by the end of the epidemic but household testing suppressed

R_e more during the growth phase of the epidemic, resulting in greater number of infections

averted over time. This is because household testing not only reduces the already higher

number of infections taking place in households, it also decreased transmissions between

different distinct social settings (e.g. if an infector infected an infectee in the household

setting but the infector was infected in school) (Figure 7C-D).

Discussion

Community asymptomatic testing only achieves high levels of infection reduction after

symptomatic testing demand has been saturated. However, the current minimum target of 100

tests/100k/day is unlikely to saturate symptomatic testing demand even with scenarios where

$R_e \leq 1.2$. Saturating symptomatic testing demand in realistic epidemic wave scenarios where

$R_e > 1.2$ likely requires >1,000 tests/100k/day. This is because testing demand is largely

shaped by non-SARS-CoV-2 infected individuals due to the overlap of SARS-CoV-2

infection symptoms with other respiratory tract infections. In other words, even before

implementing any form of community testing, it is crucial to increase investments in testing

capacities that meet symptomatic testing demand first. For instance, if Zambia had ~10

million Ag-RDTs available through the first three months (i.e. ~600 tests/100k/day over three

months for 18 million people) of its first Omicron BA.1 wave (R_e~2.5) that were only used

for symptomatic testing, ~37,000 infections could likely be averted. However, if testing rate

was at 100 tests/100k/day, the number of infections averted drops nearly 10-fold to ~3700

cases averted on average despite only a 6-fold reduction in testing.

If $R_e < 1.5$, or can be reduced to that point through other public health interventions,

increasing testing capacity from 100 tests/100k/day to 200-400 tests/100k/day provides the

greatest proportional reduction in secondary transmissions. Furthermore, testing has the

potential to be most effective at reducing transmission when $R_e < 1.5$. We would also obtain

the greatest reduction in transmissions through increased testing volumes if R_e~1.0 (Table 1).

As SARS-CoV-2 outbreaks can have R_e appreciably above 1.5, it is important to combine

testing with other public health measures such as vaccination, physical distancing and

masking so as to maximize impact of testing programs. It is also important to note that the

utility of testing in averting infections is predicated on people changing and maintaining their

behaviour to reduce contacts following a positive test.[25] Encouraging and incentivizing

these changes of behaviour are essential for the effectiveness of any test-and-isolate program,

particularly individuals of lower socioeconomic status[26] and communities in low-resource

settings.[27]

As a corroboration of our results, we compared the weekly average testing rate[6] to the

average R_e values estimated from COVID-19 case counts

(https://github.com/epiforecasts/covid-rt-estimates)[28] of 134 countries between December

2021 and March 2022 when the Omicron BA.1 VOC spread rapidly across multiple countries

(Figure 8). Although the demographic profiles differ between high-income countries (HICs)

and LMICs, we found that some HICs were expectedly testing at rates that were sufficient or

even higher than what was likely needed to saturate the symptomatic testing demand we had

estimated for LMICs at similar epidemic intensity (i.e. R_e values). However, as Omicron

(BA.1) cases surged, some HICs such as the United States, Germany and Australia were still

reportedly facing test shortages.[29–31] Based on our results, these countries were indeed

falling short of meeting symptomatic testing demand (Figure 8). Finally, if we assume that

most HICs are testing at rates that sufficiently meet symptomatic testing demand, we found

that most of them were testing >100 tests/100k/day.

Figure 8: **Global reported COVID-19 testing rate between December 2021 and March 2022 when the Omicron BA.1 variant of concern spread rapidly across multiple countries**. Each data point denotes the average weekly reported COVID-19 testing rate of a country against the average time-varying reproduction number (R_t) computed in the same week and is colored by the income level of the country while sized by time (i.e. month/year). The shaded area denotes the level of test availability we had estimated to saturate symptomatic testing demand given different equivalent initial R_e values (Figure 4). The red vertical line at 100 tests per 100,000 persons per day is the minimum testing rate target set by the ACT-Accelerator diagnostics pillar. Testing rate data were sourced from the SARS-CoV-2 Test Tracker by FIND (https://www.finddx.org/covid-19/test-tracker/) while R_t was computed from reported COVID-19 case counts using EpiNow2 (https://github.com/epiforecasts/covid-rt-estimates).[28]

If there are excess tests available after meeting symptomatic testing demand, it is important to critically consider where and how routine community testing of asymptomatic populations is implemented to maximize impact. Given that a larger proportion of infections is expected to occur within households, household community testing after meeting symptomatic testing demand in the previous week would yield greater total infections averted. While testing at regular mass gatherings such as religious gatherings every week, for instance, could lead to comparable number of diagnosed infections, doing so only effectively tallies the number of infections that had happened in the week prior and limited infections at these gatherings.

Disseminating tests across households, on the other hand, is more effective in not just

lowering transmissions occurring in households but likely lessens the number of

transmissions between different contact networks as well.

There are limitations with our study. First, we assumed that all healthcare facilities will have

access to all Ag-RDT stocks available each week. However, there could be disparities in

stock allocation between different clinics such as prioritizing stock allocation for hospitals.

Such disproportionate distributions could lead to uneven fulfilment of symptomatic testing

demand and consequently affect levels of infections. Furthermore, we assumed that there was

a sufficient number of qualified health workforce and implementation support available to

implement the various testing strategies. The strained healthcare system, especially in more

remote regions of the country, poses a major limiting factor in implementing a widespread

testing program. While our key finding that >100 tests/100k/day is needed to saturate

symptomatic testing demand before rolling out community testing programs is unlikely to

change, randomized community trials in LMICs using Ag-RDTs for community testing can

provide better impact estimates under realistic scenarios.

Second, we only modelled scenarios where test-and-isolation was the only public health

intervention. Symptomatic testing demand would expectedly be lower if other non-

pharmaceutical interventions (NPIs) were introduced, and thus potentially improve the utility

of community testing at lower test availability. However, the impact of NPIs is confounded

by temporal effects[32] and thus difficult to parameterize their mean effects on infection

control and in turn, testing demand. Since NPIs effectively decrease the number of secondary

transmissions and in turn, R_e, we expect that the testing demand for a population subjected to

NPIs and testing would mirror the demand we had estimated for a population subject to

testing only but at lower R_e values. Analogously, we also did not model how vaccination-

and infection-acquired immunity affect testing demand explicitly. However, by the same

reasoning that increased population immunity lowers R_e, the testing demand for a partially

immune population should be similar to that of a naïve population at lower R_e values as well.

Third, we parameterized incubation and virus shedding periods using those empirically

measured from infections by wild-type (Wuhan-like) SARS-CoV-2[17,33] for this work.

However, generation intervals have shortened considerably for recent VOCs such as

Delta[34] and Omicron BA.1[35] and could impact the utility of testing in identifying an

infection before it becomes infectious. We thus repeated the symptomatic-testing only

simulations using incubation and virus shedding periods estimated for Omicron BA.1. There

is effectively no difference in the amount of infections averted between the wild-type and

Omicron BA.1 variant across all testing rates at $R_e \geq 1.5$ (i.e. the expected initial effective

reproduction number of the Omicron variant; Figure S5 and Supplementary Data).

To conclude, Ag-RDTs are a valuable diagnostic tool for COVID-19 testing capacities in

LMICs. The target on minimal testing rate of 100 tests/100k/day should be seen as a true

minimum if testing is going to be used for reducing transmission but substantially higher

testing rates are needed to fulfil likely symptomatic testing demand or effective

implementation of community testing.

Data availability

All data relevant to the study are included in the Article, the Supplementary Data and the

GitHub repository (https://github.com/AMC-LAEB/PATAT-sim). The PATAT model source

code can also be found in the GitHub repository

(https://github.com/AMC-LAEB/PATAT-sim).

Funding

503   This work was supported by the European Research Council [NaviFlu 818353 to A.X.H. and

504   C.A.R.], the National Institutes of Health [5R01AI132362-04 to C.A.R.] and the Dutch

505   Research Council (Nederlandse Organisatie voor Wetenschappelijk Onderzoek) [Vici

506   09150182010027 to C.A.R.].

Acknowledgements

509   The authors are pleased to acknowledge that all computational work reported in this paper

510   was performed on the Shared Computing Cluster which is administered by Boston

511   University's Research Computing Services (www.bu.edu/tech/support/research/).

Authors' contributions

A.X.H. contributed to the conceptualization, data curation, formal analysis, investigation,

methodology, software, validation and visualization of the study. B.E.N. and C.A.R.

contributed to the conceptualization, data curation, funding acquisition, investigation,

methodology, project administration, resources, validation and supervision of the study.

J.A.S., A.T., N.H. and E.H. contributed to the conceptualization, investigation, validation and

visualization of the study. S.G. and S.K. contributed to the investigation, validation and

visualization of the study. A.X.H. and C.A.R. wrote the original draft of the manuscript. All

authors are involved in the review and editing of the manuscript. All authors had full access

to all data of the study and the final responsibility for the decision to submit for publication.

Potential conflicts of interest

References

1. Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time. Lancet Infect Dis **2020**; 20:533–534. Available at: http://www.thelancet.com/article/S1473309920301201/fulltext. Accessed 20 February 2022.

2. Ye Y, Zhang Q, Wei X, Cao Z, Yuan H-Y, Zeng DD. Equitable access to COVID-19 vaccines makes a life-saving difference to all countries. Nature Human Behaviour 2022 **2022**; :1–10. Available at: https://www.nature.com/articles/s41562-022-01289-8. Accessed 20 February 2022.

3. Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature 2022 **2022**; :1–10. Available at: https://www.nature.com/articles/s41586-022-04411-y. Accessed 20 February 2022.

4. Straten K van der, Guerra D, Gils MJ van, et al. Mapping the antigenic diversification of SARS-CoV-2. medRxiv **2022**; :2022.01.03.21268582. Available at: https://www.medrxiv.org/content/10.1101/2022.01.03.21268582v1. Accessed 20 February 2022.

5. Batista C, Hotez P, Amor Y ben, et al. The silent and dangerous inequity around access to COVID-19 testing: A call to action. EClinicalMedicine **2022**; 43. Available at: http://www.thelancet.com/article/S2589537021005113/fulltext. Accessed 20 February 2022.

6.   FIND. Test tracker - FIND. 2022. Available at: https://www.finddx.org/covid-19/test-

tracker/. Accessed 28 April 2022.

7.   Tessema SK, Nkengasong JN. Understanding COVID-19 in Africa. Nature Reviews

Immunology 2021 21:8 **2021**; 21:469–470. Available at:

https://www.nature.com/articles/s41577-021-00579-y. Accessed 20 February 2022.

8.   ACT-Accelerator Strategic Plan & Budget: October 2021 to September 2022.

Available at: https://www.who.int/publications/m/item/act-accelerator-strategic-plan-

budget-october-2021-to-september-2022. Accessed 19 February 2022.

9.   Vandenberg O, Martiny D, Rochas O, van Belkum A, Kozlakidis Z. Considerations for

diagnostic COVID-19 tests. Nature Reviews Microbiology 2020 19:3 **2020**; 19:171–

183. Available at: https://www.nature.com/articles/s41579-020-00461-z. Accessed 20

February 2022.

10.   Mulu A, Bekele A, Abdissa A, et al. The challenges of COVID-19 testing in Africa:

the Ethiopian experience. PAMJ 2021; 38:6 **2021**; 38:1–4. Available at:

https://www.panafrican-med-journal.com/content/article/38/6/full. Accessed 20

February 2022.

11.   Jefferson T, Spencer EA, Brassey J, Heneghan C. Viral Cultures for Coronavirus

Disease 2019 Infectivity Assessment: A Systematic Review. Clinical Infectious

Diseases **2021**; 73:e3884–e3899. Available at:

https://academic.oup.com/cid/article/73/11/e3884/6018217. Accessed 11 September

2022.

12.   Brümmer LE, Katzenschlager S, Gaeddert M, et al. Accuracy of novel antigen rapid

diagnostics for SARS-CoV-2: A living systematic review and meta-analysis. PLoS

Med **2021**; 18:e1003735-. Available at: https://doi.org/10.1371/journal.pmed.1003735.

13.   Onsongo SN, Otieno K, van Duijn S, et al. Performance of a rapid antigen test for

573   SARS-CoV-2 in Kenya. Diagn Microbiol Infect Dis **2022**; 102:115591.

574   14.   Park Y, Huh IS, Lee J, et al. Application of Testing-Tracing-Treatment Strategy in

575   Response to the COVID-19 Outbreak in Seoul, Korea. J Korean Med Sci **2020**; 35.

576   Available at: https://doi.org/10.3346/jkms.2020.35.e396. Accessed 9 September 2022.

577   15.   Kendall M, Milsom L, Abeler-Dörner L, et al. Epidemiological changes on the Isle of

578   Wight after the launch of the NHS Test and Trace programme: a preliminary analysis.

579   Lancet Digit Health **2020**; 2:e658–e666. Available at:

580   http://www.thelancet.com/article/S2589750020302417/fulltext. Accessed 9 September

581   2022.

582   16.   Vecino-Ortiz AI, Congote JV, Bedoya SZ, Cucunuba ZM. Impact of contact tracing on

583   COVID-19 mortality: An impact evaluation using surveillance data from Colombia.

584   PLoS One **2021**; 16:e0246987. Available at:

585   https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0246987. Accessed 9

586   September 2022.

587   17.   Linton NM, Kobayashi T, Yang Y, et al. Incubation Period and Other Epidemiological

588   Characteristics of 2019 Novel Coronavirus Infections with Right Truncation: A

589   Statistical Analysis of Publicly Available Case Data. Journal of Clinical Medicine

590   2020, Vol 9, Page 538 **2020**; 9:538. Available at: https://www.mdpi.com/2077-

591   0383/9/2/538/htm. Accessed 26 January 2022.

592   18.   Kissler SM, Fauver JR, Mack C, et al. Viral dynamics of acute SARS-CoV-2 infection

593   and applications to diagnostic and public health strategies. PLoS Biol **2021**;

594   19:e3001333-. Available at: https://doi.org/10.1371/journal.pbio.3001333.

595   19.   Quilty BJ, Clifford S, Hellewell J, et al. Quarantine and testing strategies in contact

596   tracing for SARS-CoV-2: a modelling study. Lancet Public Health **2021**; 6:e175–e183.

597   Available at: https://doi.org/10.1016/S2468-2667(20)30308-X.

598   20.   Boyton RJ, Altmann DM. The immunology of asymptomatic SARS-CoV-2 infection:

599   what are the key questions? Nature Reviews Immunology 2021 21:12 **2021**; 21:762–

600   768. Available at: https://www.nature.com/articles/s41577-021-00631-x. Accessed 10

601   April 2022.

602   21.   Gill CJ, Mwananyanda L, MacLeod W, et al. Sustained high prevalence of COVID-19

603   deaths from a systematic post-mortem study in Lusaka, Zambia: one year later.

604   medRxiv **2022**; :2022.03.08.22272087. Available at:

605   https://www.medrxiv.org/content/10.1101/2022.03.08.22272087v2. Accessed 28

606   March 2022.

607   22.   Zambia Statistics Agency. 2019 Labour Force Survey Report. 2019. Available at:

608   https://www.zamstats.gov.zm/index.php/publications/category/7-labour. Accessed 26

609   January 2022.

610   23.   Dovel K, Balakasi K, Gupta S, et al. Frequency of visits to health facilities and HIV

611   services offered to men, Malawi. Bull World Health Organ **2021**; 99:618–626.

612   Available at: https://pubmed.ncbi.nlm.nih.gov/34475599/. Accessed 5 April 2022.

613   24.   Hasell J, Mathieu E, Beltekian D, et al. A cross-country database of COVID-19

614   testing. Scientific Data 2020 7:1 **2020**; 7:1–7. Available at:

615   https://www.nature.com/articles/s41597-020-00688-8. Accessed 17 February 2022.

616   25.   Pavelka M, Van-Zandvoort K, Abbott S, et al. The impact of population-wide rapid

617   antigen testing on SARS-CoV-2 prevalence in Slovakia. Science (1979) **2021**;

618   372:635–641. Available at: https://www.science.org/doi/abs/10.1126/science.abf9648.

619   Accessed 20 February 2022.

620   26.   Sweeney S, Capeding TPJ, Eggo R, et al. Exploring equity in health and poverty

621   impacts of control measures for SARS-CoV-2 in six countries. BMJ Glob Health

622   **2021**; 6:e005521. Available at: https://gh.bmj.com/content/6/5/e005521. Accessed 21

623   February 2022.

624   27.   Torres-Rueda S, Sweeney S, Bozzani F, et al. Stark choices: exploring health sector

625   costs of policy responses to COVID-19 in low-income and middle-income countries.

626   BMJ Glob Health **2021**; 6:e005759. Available at:

627   https://gh.bmj.com/content/6/12/e005759. Accessed 21 February 2022.

628   28.   Abbott S, Hickson J, Badr HS, et al. epiforecasts/EpiNow2: Beta release. **2021**;

629   Available at: https://zenodo.org/record/5036949. Accessed 7 April 2022.

630   29.   Reuters. U.S. testing struggles to keep up with Omicron. 2021. Available at:

631   https://www.reuters.com/world/us/us-testing-struggles-keep-up-with-omicron-2021-

632   12-22/. Accessed 7 April 2022.

633   30.   DW News. Germany faces PCR test bottlenecks. 2022. Available at:

634   https://www.dw.com/en/germany-faces-pcr-test-bottlenecks/av-60431333. Accessed 7

635   April 2022.

636   31.   BBC News. Covid: Australians desperate for tests amid Omicron surge. 2022.

637   Available at: https://www.bbc.com/news/world-australia-59864428. Accessed 7 April

638   2022.

639   32.   Sharma M, Mindermann S, Rogers-Smith C, et al. Understanding the effectiveness of

640   government interventions against the resurgence of COVID-19 in Europe. Nature

641   Communications 2021 12:1 **2021**; 12:1–13. Available at:

642   https://www.nature.com/articles/s41467-021-26013-4. Accessed 21 February 2022.

33.   Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized

patients with COVID-2019. Nature 2020 581:7809 **2020**; 581:465–469. Available at:

https://www.nature.com/articles/s41586-020-2196-x. Accessed 26 January 2022.

34.   Kang M, Xin H, Yuan J, et al. Transmission dynamics and epidemiological

characteristics of Delta variant infections in China. medRxiv **2021**;

:2021.08.12.21261991. Available at:

https://www.medrxiv.org/content/10.1101/2021.08.12.21261991v1. Accessed 25

February 2022.

35.   Hay JA, Kissler SM, Fauver JR, et al. Viral dynamics and duration of PCR positivity

of the SARS-CoV-2 Omicron variant. medRxiv **2022**; :2022.01.13.22269257.

Available at: https://www.medrxiv.org/content/10.1101/2022.01.13.22269257v1.

Accessed 17 March 2022.