Dietary Exposure to Oxidized Lipids in US Infants

Lisaura Maldonado-Pereira[1,3], Carlo Barnaba[2], Ilce Gabriela Medina-Meza[1,3] *

[1]Department of Chemical Engineering and Materials Science, Michigan State University, East Lansing, MI, USA. *maldon60@msu.edu*
[2]Institute of Quantitative Health Science and Engineering, Michigan State University, East Lansing, MI, USA. *barnabac@msu.edu*
[3]Department of Biosystems and Agricultural Engineering, East Lansing, MI, USA. *ilce@msu.edu*

Corresponding Author
Medina-Meza, Ilce Gabriela, Ph.D.
Department of Biosystems and Agricultural Engineering
469 Wilson Rd.| Room 302C
East Lansing, MI
ilce@msu.edu
Phone: 517-884-1971

Abstract

Baby Foods (BFs) and Infant formulas (IFs) are the main sources of nutrition for an infant throughout the 1st year of life. A variety of enriched products are commercially available for parents seeking to fulfill their baby's nutritional needs. Consequently, different bioactive lipids are present in BFs and IFs including dietary oxidative substances (DOxS) whose known toxicity has been associated with mutagenicity, cancer, and other chronic diseases. In this work, we performed an exposure assessment of 25 bioactive lipids on a total of 63 baby food samples (50 – IFs and 13 – BFs) commercially available in the US. To determine the dietary exposure to DOxS, we used EPA's SHEDS-HT probabilistic model. Even though β-Sitosterol was the most exposed bioactive lipid with 75,4 µg/day, cholesterol was the most absorbed compound during the entire first year (19,320 µg/day). Additionally, we found 7α-hydroxycholesterol (7α-OH) as a potential DOxS biomarker of the BFs manufacturing process. This is the first time that an exposure assessment including DOxS ingested by infants after BFs and IFs consumption is performed, enabling much-needed information regarding these hazardous compounds and their potential effects on infants' health.

Keywords: Oxidized lipids, Dietary exposure, Toxicity, Baby foods, Infant formula, Oxysterols

1. Introduction

Diet, during the first year of life of a newborn, plays a crucial role to ensure an infant's proper growth and brain development (Babawale et al., 2018; CDC, 2021b; Harris & Pomeranz, 2020; Timby, Domellöf, Hernell, Lönnerdal, & Domellöf, 2014). Despite the U. S. Dietary Guidelines for Americans recommendation for breastfeeding during the first 6 months, reports show that only 25% of babies are exclusively breastfed in the US (CDC, 2021a), and globally, only 40% have been estimated (Collective, 2017; Timby et al., 2021). Different circumstances influence the mother's decision to rely mainly on infant formulations (IFs) instead of breast milk. After the baby is 6 months old, baby foods (BFs), also known as complementary foods, are commonly introduced to the infant, and it is not until the infant's 1st or 2nd year that they will be eating almost the same foods as any other adult (CDC, 2021b).

Cholesterol content in human milk is typically comprised of 0.34 to 1.3 wt% based on total lipids (Jensen, 1999). On the contrary, standard IF doesn't contain cholesterol but it is high in plant-based oils such as palm and soybean oil (Zubin Maslov, Hill, Lüscher, & Narula, 2021) which provide other lipid classes such as phytosterols and tocopherols also needed for a healthy growth and development processes. During weaning (4-6 months), BFs are gradually introduced. BFs consist of unsweetened, unsalted, no added starch or artificial flavors/colors puree of fruits, vegetables, grains, and blends of meats or poultry. All these ingredients are good sources of protein, iron, zinc, and other much-needed nutrients (Gerber, 2021a, 2021b).

The main purpose of IFs and BFs manufacturing relies on providing the best, most nutritious, and safest food for the most vulnerable population, infants. Both IF and BFs undergo industrial processing to guarantee their microbiological safety. Processing, in particular thermal processing, act on macro and micronutrients of raw material, potentially favoring changes in the

nutritional properties of these foods. For instance, several lipid oxidation molecules, such as

dietary oxysterols (DOxS), are formed because of food's over-processing during their

confection, storage, and even transportation processes.

The oxidation of cholesterol results in the accumulation of DOxS. The formation of DOxS is

triggered by high temperature, hydrostatic pressure used for nonthermal pasteurization, light

exposure, and aging of food (Cais-Sokolińska, Walkowiak-Tomczak, & Rudzińska, 2022; Chudy

& Teichert, 2021; Lisaura Maldonado-Pereira, 2021; Risso et al., 2022; Sabolová et al., 2017).

Several studies have summarized the biological activities of these compounds which have been

extensively associated with different chronic diseases including pro-inflammatory and pro-

oxidant activities, as well as promote carcinogenesis (Garcia-Llatas, Mercatante, López-García,

& Rodríguez-Estrada, 2021; L. Maldonado-Pereira, Schweiss, Barnaba, & Medina-Meza, 2018;

Poli et al., 2022; Willinger, 2019; Zmysłowski & Szterk, 2019). Our group recently studied the

metabolomic signatures of liquid and powder infant formulations and proposed that 7-keto could

be considered a biomarker of powder IFs (A. Kilvington, Barnaba, Rajasekaran, Laurens

Leimanis, & Medina-Meza, 2021). Additionally, Risso and coworkers pointed out that the shelf-

life is the primary determinant of DOxS after analyzing whole milk powders (Risso et al., 2022).

Their study serves as another example of how DOxS could be used as markers to further quantify

and characterize the nutritional quality and freshness, not only of ingredients but also of

composite products such as commercial chocolate milk and whole powder milk. These findings

highlight the potential relationship of DOxS loads with specific unit operations used in food

processing. This knowledge may be a critical factor in identifying processing steps that may

increase their formation in foods and could prevent infants' exposure to these compounds and

their long-term effect on their health. Previously, we assessed the dietary exposure of DOxS on

infant formula-fed infants in the first 6 months of life (Alice Kilvington, Barnaba, Rajasekaran,

Laurens, & Gabriela Medina-Meza, 2021). To date, the estimated exposure level of DOxS in

infants (0-1 years old) has not been addressed.

The present study provides a dietary assessment of sterols and their oxidized derivatives

using a probabilistic exposure assessment model during the first year of infant development for

several sterols species (previously analyzed in our laboratory) present in various IFs and BFs

commercially available in the US market (A. Kilvington et al., 2021; A. Kilvington,

Maldonado☐Pereira, Torres☐Palacios, & Medina☐Meza, 2019). The dietary exposure

assessment will provide an estimation of the exposure extent of these compounds in infants

which is the first step in conducting a risk assessment of these sterols and their oxidative species

that could become a potential health risk in the infant's community (Wei et al., 2022).

2. Methods

2.1. Stochastic Human Exposure and Dose Simulation (SHEDS-HT)

SHEDS-HT was coded in the R language (v. 2.15.3). Several modifications were performed

to the code's input files to enable the dietary exposure analysis of bioactive lipids including

sterols (both animal and plant origin), squalene, and tocopherols. A complete description of the

SHEDS-HT model can be found in Isaacs' work (Isaacs et al., 2014). The following sections will

explain the modifications employed in this analysis.

2.1.1. Chemical Pathway and Scenario

A list of the compounds analyzed in this study is shown in **Table 1**. The chemical and

physical properties of the compounds required by SHEDS-HT were obtained from The

Metabolomics Innovation Centre and the ChemSpider webpages (Centre, 2021; ChemSpider,

2021). Concentration values of these chemicals in Infant Formulas (IFs) were obtained from

previous work performed in the Food and Health Engineering Laboratory (FHEL) at Michigan

State University (A. Kilvington et al., 2019).

Table 1: Chemical properties used in SHEDS-HT to evaluate exposure of the compounds of interest.
133

Compound	Water Solubility (mg/L)	Absorption factor (f_{abs})
7keto	0.0027[1]	0.12[3]
Triol	0.000156[1]	0.15[3]
5,6alpha-Epoxy	0.005104[2]	0.28[3]
5,6beta-Epoxy	0.005104[2]	0.32[3]
7beta-OH	0.48[1]	0.42[3]
7alpha-OH	0.48[1]	0.30[3]
20alpha-OH	0.57[1]	0.40*
22-OH	0.65[1]	0.40[4]
24-OH	0.40[1]	0.40*
25-OH	0.24[1]	0.40*
Campesterol	0.05803[5]	0.1431[5]
Beta-sitosterol	0.02739[5]	0.05074[5]
Stigmasterol	0.0000112[5]	0.040[5]
Brassicasterol	0.06420[5]	0.3121[5]
Alpha-Tocopherol	0.003317[5]	0.03394[5]
Beta-Tocopherol	0.02826[5]	0.03423[5]
Delta-Tocopherol	0.2477[5]	0.03477[5]
Gamma-Tocopherol	0.02852[5]	0.03424[5]
Cholesterol	0.077038[5]	0.314705[5]
Squalene	0.004897[5]	0.986721[5]
Desmosterol	0.049427[5]	0.378243[5]

134
[1]The Metabolomics Innovation Centre (https://www.metabolomicscentre.ca/)
[2]ChemSpider – Search and share chemistry (www.chemspider.com)
[3]Value obtained from the paper: Osada, Sasaki, & Sugano, 1994
[4]Value obtained from the paper: Tranheim Kase et al., 2012
[5]Properties provided by SHEDS-HT
*Absorption value of 22-OH was used for these compounds since no estimated value could be found for these
specific DOxS (20alpha-OH, 24-OH, and 25-OH)

DOxS concentrations from Baby Foods (BFs) were obtained experimentally as is explained

in sections 2.2-2.4. Since this study was focused on ingestion of DOxS because of BFs and IFs

consumption, SHEDS-HT only needs to consider the dietary pathway and the direct ingestion

scenario of the samples tested. In this study, the dietary exposures are calculated by determining

the total daily mass of chemical intake for each simulated person via different foods.

Concentration distributions for each food are provided as input. For each individual, daily

concentrations (µg of sterol per g of food) are sampled (one for each food item). Dietary

exposures are calculated as the sum (over food groups) of the food item's concentration and

mass of food consumed (as determined by the assigned food diary for the person). Exposure

equation used by SHEDS-HT:

153
$$Exposure\left(\frac{\mu g}{day}\right) = Consumption\left(\frac{g\ of\ food}{day}\right) * Residue\ of\ the\ Compound\ in\ Food\left(\frac{\mu g}{g\ food}\right)$$

154   Intake dose estimates are calculated using distributions of route-specific fractional

155   absorptions as it is shown below:

156
157
$$Dose\ Intake\left(\frac{mg}{kg-day}\right) = \frac{abs\ dermal\left(\frac{mg}{day}\right) + exp\ ingest\left(\frac{mg}{day}\right) + dose\ inhale\left(\frac{mg}{day}\right)}{1000 * BW}$$

158   Where *abs dermal* if the absorption of the compound through the skin, *exp ingest* is the

159   exposure of the compound through ingestion of the food item, and *dose inhale* is the dose of the

160   compound inhaled by the infant.

161   Absorption values are obtained using the estimated absorption factor (f_{abs}) of each bioactive

162   lipid in the following equation:

163
$$Absorption\left(\frac{ug}{day}\right) = Exposure\left(\frac{ug}{day}\right) * f_{abs}$$

A full description of the exposure equations used in SHEDS-HT and their corresponding

input parameter distributions are provided in Isaacs' work (Isaacs et al., 2014).

2.1.2. Population Module

Input-based data from the U.S. Census was used to generate a simulated population

representative of the U.S. population in terms of age and gender. A population of 10,000

individuals was selected for this study. Afterward, Monte Carlo methods were used to assign

relevant exposure factors and cohort-matched activity and food intake diaries to each person.

Food diaries were based on the National Health and Nutrition Examination Survey-What We

Eat in America (NHANES-WWEIA) 1999–2018 two-day food intake diaries (USDA, 2018).

Food diaries calculated the mass of each food group consumed by the individual.

2.2. Bioactive Lipids Extraction in Baby Foods

Bioactive lipids were extracted from 13 baby foods according to a slightly modified Folch's

(Folch, Lees, & Sloane Stanley, 1957) cold extraction method as is described in Maldonado-

Pereira et al. (2022). Thirty grams of sample were minced and placed in a 500 ml glass bottle

with a screwcap with 200 mL of chloroform:methanol solution (1:1, v/v). The sample was mixed

for 15 min at 300 rpm. Homogenization was performed using an Ultra-Tourrax for 3 minutes.

The bottle was kept in an oven at 60°C for 20 min before adding 100 mL chloroform. Then, the

bottle was vortexed for 2 min following by filtration. The filtrate was mixed thoroughly with 100

mL of 1 M KCl solution. Samples were left overnight at 4 °C. Then, the lower phase containing

lipids was collected and dried at 60°C with a vacuum evaporator at 25 in Hg. Total fat content

was determined gravimetrically (L. Maldonado-Pereira, Barnaba, De los Campos, & Medina-

Meza, 2021).

2.3. Determination of Total Cholesterol, Tocopherols, and Phytosterols Content in Baby

Foods by GC-MS

A simultaneous quantification method for cholesterol, tocopherols, and phytosterols

developed by our group was used (A. Kilvington et al., 2021). Cold saponification at room

temperature was performed following Bonoli's protocol with some modifications (Matteo Bonoli

et al., 2008). Two hundred mg of lipid were mixed with 50 µg of 19-hydroxycholesterol and 140

μg of 5α-cholestane as internal standards for the determination of dietary COPs and total

cholesterol, respectively. Subsequently, the sample was mixed with 10 mL of 1 N KOH solution

in methanol and left under gentle agitation, and covered from the light for at least 15 h (Sander,

Addis, Park, & Smith, 1989). One-tenth of the unsaponifiable matter was subjected to silylation.

The sample was mixed with 100 μL of pyridine and 100 μL of the Sweeley's reactive mixture

(pyridine/hexamethyldisilazane/trimethylchlorosilane, 10:2:1, v/v/v) at 75°C for 45 min, dried

under nitrogen stream and dissolved in 1 mL of n-hexane. One microliter of the silylated solution

was injected into a gas chromatograph using a fitted with a Zebron™ZB-5HT (Phenomenex,

Torrance, CA) capillary column (30 m × 0.25 mm × 0.25 μm). Oven temperature conditions

were set up as follows: 260 to 300°C at 2.5°C/min, then from 300 to 320°C at 8°C/min, and

finally hold at 320°C for 1 min. The injector and detector were both set at 320°C. Helium was

used as the carrier gas at a flow rate of 54.0 mL/min, the split ratio at 1:50, and the pressure

constant at 134 kPa.

2.4. DOxS profiling in Baby Foods

The other nine-tenths of the sample was used for the quantification of dietary COPs.

Purification by NH_2-SPE was performed according to the literature (Rose-Sallin, 1995). The

enriched fraction was recovered and evaporated to dryness under a nitrogen stream.

Subsequently, the purified fraction was silylated with Sweeley's reactive mixture (at 75 °C for

45 min), dried under a nitrogen stream and dissolved in 1 mL of n-hexane.

Dietary DOxS were identify using the selected ion molecule (SIM) method with a GCMS-

QP2010 SE single quadrupole Shimadzu Corporation (Kyoto, Japan). One μL of the silylated

DOxS was injected into a single quadrupole (GCMS-QP2010 SE) with the following conditions:

from 250 to 280 °C at 2°C/min, hold at 280 °C for 7 min, and from 280 to 315 °C at 1.5 °C/min.

Helium was used as a carrier gas (flow rate of 0.37 mL/min); the split ratio was 1:15 and the

pressure was 49.2 kPa. The interface temperature for the GC-MS was 320 °C, with the electron

multiplier voltage set at 0 kV and the injector at 320 °C. A fused silica column (30 m x 0.25 mm

i.d. x 0.25 um thickness) coated with 5% phenyl polysiloxane (Phenomenex Zebron ZB-5) was

used.

2.5. *Statistical Analysis*

Descriptive statistics were obtained from the SHEDS-HT model. Mean, standard deviation,

and the following quantiles: 5%, 25%, 50%, 75%, 90%, and 95%, were reported. Since the data

did not follow a normal distribution after a Shapiro-Wilk test a non-parametric Matt-Whitney *U*-

test was performed at a $p < 0.05$ significance level. Data normalization was performed for

exposure, absorption, and dose intake results for comparison purposes. A sparse PLS-DA (sPLS-

DA) was used to evaluate relationships between a single 'omics data set. Additionally, a

biomarker analysis, including the ROC curves, were performed using MetaboAnalyst in its web

server version (Chong et al., 2018) All the statistical analysis were computed using RStudio

(version 1.4.1717 © 2009-2021 RStudio, PBC)

3. Results and discussion

3.1 Bioactive Lipids Profiling in BF

A complete list of the BF samples, their codes, and results can be found in the Supplemental

Material – **Table 1S and Table 2S**. Cholesterol was the most abundant compound in BF (**Figure**

1S), reaching a concentration value of 1,247.2 mg/100g fat for BF1 (**Table 1S**). A higher

cholesterol value compared to the one reported in the nutritional facts label was observed for

samples: BF1, BF6, and BF11. The items presenting higher fat content had mainly beef, turkey,

and dairy products as the main ingredient in their preparation process. Plain meat BFs after the

first 6 months of age is recommended by the CDC to avoid iron deficiency (CDC, 1998; Fox,

Pac, Devaney, & Jankowski, 2004; Miller, 2013). Even though a higher fat content was observed

for those specific BFs, fat consumption is normally promoted during the infants' first years of

life because of its essential role during the infants' growth phase. Restricting fat before the age of

two is not recommended and could be harmful during the infants' growing stage (Allen &

Myers, 2006; Hardy & Kleinman, 1994; Johnson, 1998; Öhlund, Hörnell, Lind, & Hernell,

2008).

Phytosterols comprised 39% of the total bioactive lipid's distribution, followed by DOxS

with only 2%. Tocopherols, desmosterol, and squalene were not detected in our BF samples.

β-Sitosterol was the dominant phytosterol which agrees with previous studies (Guadalupe

García-Llatas et al., 2008; G. García-Llatas & Rodríguez-Estrada, 2011; A. Kilvington et al.,

2021). The presence of high amounts of β-Sitosterol reflect the use of vegetable oils in the

preparation of most BFs.

3.2 Dietary Exposure from IFs and BFs during Infant's 1st Year of Life

The equation used by SHEDS-HT to calculate the exposure value of each compound is

shown above. Exposure results are reported in units of µg/day (**Table 3S**) and are based on the

quantities of IFs and BFs ingested by infants between 0 and 1 year old reported by the

NHANES-WWEIA 1999-2018. Overall, 59% of the most exposed bioactive lipids were

phytosterols (**Figure 1**) - specifically, β-Sitosterol with 75,410 ug/day (**Figure 2**) - which is the

most abundant group and lipid reported in our previous study (A. Kilvington et al., 2021).

Cholesterol and campesterol were the 2nd and 3rd most exposed bioactive lipids. Among DOxS,

5,6β-epoxycholesterol, 7β-OH, and 7α-OH obtained the 8th, 9th, and 10th highest exposure levels

with 1,244.4, 191.8, and 165.5 µg/day, respectively. These DOxS have been related to several

adverse effects on human health in previous studies such as cardiovascular diseases -

atherosclerosis, hypertension, ischemic stroke-, neurological diseases, viral infections, and

cancer (A. Kilvington et al., 2021; Lisaura Maldonado-Pereira, 2021; L. Maldonado-Pereira et

al., 2018; Zmysłowski & Szterk, 2019), which arise concerns related to their consumption by

infants.

266
267   **Figure 1:** Bioactive Lipids Exposure in IFs and BFs Consumption during 1st Year of Life
268

Figure 2: Percentages of the Bioactive Lipids Exposure Values in IFs and BFs after their Consumption during the 1st Year of Life according to Supplemental Material - Table 3S

3.1 Dietary Dose Intake and Absorption of Bioactive Lipids by Infants (0-1 years old)

The dietary dose intake of phytosterols (**Figure 3B**) was higher than any other bioactive lipid group which agrees with previous studies (Claumarchirant, Matencio, Sanchez-Siles, Alegría, & Lagarda, 2015; Hamdan, Claumarchirant, Garcia-Llatas, Alegría, & Lagarda, 2017).

β-Sitosterol showed the largest dose intake among all bioactive lipids reaching a value of 9.64 mg/kg day in females (Hamdan, Sanchez-Siles, Garcia-Llatas, & Lagarda, 2018; Hamdan, Sanchez-Siles, Matencio, et al., 2018).

Interestingly, dietary dose intakes (mg/kg-day) were higher in male infants compared to females for most of the DOxS (5,6β-epoxycholesterol, 6-ketocholesterol, 7-ketocholesterol, 7α-OH, and 7β-OH) and α-tocopherol (**Figure 3A and 3C**). On the other hand, dietary dose intake of β-sitosterol, cholesterol, δ-tocopherol, and 25-OH were slightly higher for female infants.

Even though this difference in dose intake values between sexes was not significant, it highlights

the effect of body weight's average variability between females and males (body weight usually

varies by only 0.79 kg) (CDC, 2001). In our previous study focused only on IFs consumption,

the difference in dietary dose intake values between males and females was also not statistically

significant (A. Kilvington et al., 2021).

The estimated absorption factors (f_{abs}) provided a predicted absorption value (mg/kg-day) for

each specific bioactive lipid analyzed in this model (**Figure 4A and 4B**). Cholesterol showed the

highest absorption value with 1.90 mg/kg-day. Subsequently, β-sitosterol, campesterol, and

desmosterol showed values of 0.38, 0.34, and 0.33 mg/kg-day, respectively.

Figure 3: Dose Intake (mg/kg-day) of A) DOxS, B) Phytosterols, C) Tocopherols, and D) Cholesterol, Desmosterol, and Squalene

Even though β-sitosterol was the compound with the highest exposure value, cholesterol was

absorbed by 80% more. The bioaccessibility of β-sitosterol has been reported to be lower than

other sterols, which explains the reason for being less absorbed than other sterols, despite its

higher exposure level, and dietary dose intake (Claumarchirant et al., 2015).

All DOxS together showed an absorption value of 0.062 mg/kg-day, being 5,6β-

epoxycholesterol the DOxS with the highest absorption value (0.040 mg/kg-day). There's a clear

distinction between DOxS absorption comparing 0-6 and 6-12 months of life of an infant. The

presence of 7-keto in IFs was previously seen in metabolomics studies such as the one performed

by our group (A. Kilvington et al., 2021), where 7-keto was proposed as a processing biomarker

for powder IFs. Here, it is observed that it is not the most exposed, ingested, and absorbed sterol

(**Figure 4A**) once solids foods are introduced into the infant's diet. Therefore, it could be stated

that it's highly absorbed by infants, probably during just the first 6 months of life based on their

dietary intake through *only* IFs consumption. With the introduction of solid foods past the first 6

months of life, 5,6β-epoxycholesterol resulted to be more absorbed. Absorption of these two

DOxS raises concerns since their toxicity has been reported in previous studies in concentrations

as low as 30.0 and 39.7 ng/mg of biological tissue, specifically in atherosclerosis patients

(Micheletta et al., 2004). Maldonado-Pereira and coworkers provided a comprehensive review

that reported the concentration of DOxS that exerted biological activities (L. Maldonado-Pereira

et al., 2018). It is crucial to study in-depth the relationship between infants' exposure and

absorption of these DOxS with the future development of different chronic diseases.

3.2 Comparison of Bioactive Lipids Exposure, Dose Intake, and Absorption

Normalized values of exposure, dose intake, and absorption for all compounds are shown in

318
319

Figure 4: A) Comparison of absorption and Dose Intake. B) Normalization of Exposure, Dose Intake, and Absorption

320
321

Figure 4B. Cholesterol's normalized absorption value was almost doubled compared to either its

normalized exposure or dose intake values. It is known that massive amounts of cholesterol are

needed for an infant's growth (especially during the first 0-24 months of life) since every

membrane requires cholesterol with especially high amounts in neuronal cells (Lawrence &

Lawrence, 2022; Woollett & Heubi, 2000; Zubin Maslov et al., 2021). Therefore, such a high

absorption rate of cholesterol was already expected.

Overall, 5,6β-epoxycholesterol was the most absorbed DOxS. It is noteworthy to mention

that DOxS were absorbed in a significantly less proportion than cholesterol (**Figure 5**) as a result

of their lower solubility in the mixed micelles and/or lower esterification inside the enterocytes

(Garcia-Llatas et al., 2021). Additionally, the dominant presence of phytosterols in BF and IF

could be exerting an antioxidant effect that promotes such high cholesterol absorption rate while

it hinders further absorption of COPs. This high presence of phytosterols could result in a

positive attribute of IF and BF by reducing the absorption of the DOxs and promoting cholesterol

absorption which is much needed by infants at this early stage of life.

Figure 5: Normalization of Exposure, Dose Intake, and Absorption Only for COPs during 1[st] Year of Life

For β-sitosterol, a higher dose intake was obtained after comparing the exposure level. As it

was mentioned above, β-sitosterol is an abundant phytosterol in vegetable oils which is the main

ingredient of IF and certain BFs. Additionally, it could be the exerting an antioxidant effect

during the bioactive lipids absorption process after BFs and IFs consumption.

Lastly, since the absorption values for 25-OH, triol, and 7-keto were estimated values

obtained from 2 different scientific sources (Centre, 2021; ChemSpider, 2021), an Absorption,

Digestion, Metabolism, and Excretion (ADME) study is needed to obtained realistic absorption

rates that will facilitate more accurate exposure assessments.

3.3 Dietary Dose Intake Comparison between SHEDS-HT and Theoretical Equation

After comparing the results from this exposure assessment with our previous study

(Kilvington et al., 2021), a better idea of the dietary intake and potential absorption values was

obtained. Kilvington and her co-worker estimated dose intake values using the theoretical

equation:

$$DI = \frac{[C] \times Intake \times Abs}{BW}$$

This equation was modified to analyze the ingestion and dose intake during the first 6 months

of life of several bioactive lipids. Our previous work identified 7-keto as a potential biomarker

for IFs. Lately, Risso and coworkers also suggested 7-keto as a biomarker in their study using

powder milk samples (Risso et al., 2022).

Our results -which are derived from the equation previously mentioned used by SHEDS-HT-

revealed that compared to the first 6 months of an infant where their diet is mainly comprised of

IFs, 5,6β-epoxycholesterol, 7β-OH, and 7α-OH were the most absorbed DOxS from BFs

consumption. 5,6β-epoxycholesterol, 7β-OH, and 7α-OH have been previously found in several

types of meat and poultry samples (Chiu, 2018; Gehring et al., 2020; Grau, Codony, Grimpa,

Baucells, & Guardiola, 2001; Khan et al., 2015; Musto, Faraone, Cellini, & Musto, 2014; Osada,

Hoshina, Nakamura, & Sugano, 2000; Pie, 1991; Rant et al., 2019). Compared to IFs, BFs'

protein content commonly comes from different meat and poultry ingredients used during their

confection. Interestingly, based on unpublished results from our lab, 5,6β-epoxycholesterol has

been found present in human placenta samples. This finding could somehow relate the higher

absorption rate of this DOxS with prenatal factors such as hormones present during the infant's

fetal development (Prins et al., 2022; Rodríguez-Rodríguez et al., 2018; Wu, Tian, & Lin, 2015)

Biomarker analysis for our BF and IF data revealed that a 7α-OH could be a DOxS

processing biomarker for BFs (**Figure 6**). From these results and considering our previous study,

we suggest that 7α-OH is a biomarker for BFs while 7-keto serves as a biomarker for powder IF.

It is worth noting that 7-OH isomers are direct precursors of 7-keto. Differences in the type of

thermal processing of each type of food (IF and BF) are the main reason that leads to the

formation of certain DOxS, which suggests them as potential biomarkers of either BF or powder

IF. This hypothesis is strongly supported by previous studies that have reported differences in

DOxS dominance on the same food matrices cooked under different conditions (Francesc

Guardiola, 2002; F. Guardiola, Bou, Boatella, & Codony, 2004; Lisaura Maldonado-Pereira,

2021; L. Maldonado-Pereira et al., 2018).

An sPLS-DA was performed to evaluate the accuracy of our models. Data were classified by

BF, Liquid IFs (L-IFs), and Powder IF (P-IFs) to highlight any potential differences in IF

processing techniques. **Figures 7A** show that there is indeed a noticeable difference between the

confection process of BF, L, and P that affects the values of the DOxS present in the samples.

Figure 6: 7α-OH as a Potential Biomarkers of BFs

Additionally, **Figure 7B** shows the prediction zone for other samples based on our data. In

our model, only 3 out of 63 samples were incorrectly classified, reaching a 95.2% accuracy. Both

of these figures provide intuition on how novel samples would be classified according to the

generated model. Lastly, **Figure 8** shows a ROC for all 3 models (BF, L-IFs, and P-IFs) with

highly positive sensitivity rates for each one (BF: 0.1, P-IFs: 0.9905, and L-IFs: 0.981, allowing

a good prediction rate for future data.

Figure 7: A) IF and BF data with confidence ellipses and B) Prediction Zones

Besides the use of 7α-OH and 7-keto as biomarkers of thermal processing techniques

employed during the confection of BFs and IFs, it is important to keep in mind their biological

toxicity that could become a potential hazard among infants. Comparing both DOxS, 7α-OH and

7-keto have been found in patients suffering different chronical diseases such as hypertension,

multiple sclerosis, cancer, atherosclerosis, among others (L. Maldonado-Pereira et al., 2018). The

lowest concentrations exerting a toxicological effect previously reported for 7α-OH and 7-keto

were 0.004 and 0.006 µg/mL of blood plasma, respectively, in atherosclerosis patients. Since

there is so little available information regarding the relationship between these compounds and

the development of these diseases as a result of the consumption of IF s and BFs by infants, this

vulnerable population could be potentially at a health risk. More accurate information regarding

DOxS absorption values, their metabolism, and excretion rates is needed to start reducing this

knowledge gap. Therefore, we suggest the development of an Absorption, Digestion,

Metabolism, and Excretion (ADME) study followed by the completion of a risk assessment to

identify the real hazard and protect our infants.

Figure 8: ROC for BF, Liquid Formula, and Infant Formula Models

4. Conclusions

Bioactive lipid distribution of BF is dominated by cholesterol (59%) from animal based

foods, followed by phytosterols (~39%) from plant based foods. β-Sitosterol was the most

abundant phytosterol as a result of its dominant presence in vegetable oils, which is the BF's

main ingredient.

Dietary exposure assesment showed that phytosterols had the highest exposure

percentages (59%) compared to DOxS, tocopherols, cholesterol, squalene, and desmosterol. β-

Sitosterol was the most exposed compound (0.38 mg/kg-day) but the 2nd most absorbed

compound because of its lower bioaccessibility which has been previously reported

(Claumarchirant et al., 2015). Cholesterol was the most absorbed lipid with a 1.90 mg/kg-day.

5,6β-epoxycholesterol was the DOxS with the highest absorption value (0.040 mg/kg-day)

which places it overall in the 7^{th} position among all bioactive lipids absorbed. Even though

absorbance of all DOxS together accounted for only 1.86% (0.062 mg/kg-day), there's evidence

of their biological effect which raises concerns related to their consumption by infants. There is a

critical need to deeper investigate the relationship of infants diet – especially during the 1 year of

life – with future development of chronic diseases.

One of our model's limitations includes the lack of information regarding the absorption

fractions (f_{abs}) of 25-OH, triol, and 7-keto. For the purpose of our study, these values were

estimated from the available data found from 2 different scientific sources (Centre, 2021;

ChemSpider, 2021). This is why the FHEL suggests and is currently conducting an Absorption,

Digestion, Metabolism, and Excretion (ADME) study – especifically of DOxS - to obtain more

accurate absorption values to enable a more precise exposure assessment. Despite this weakness,

this is the first time that an exposure assessment including DOxS ingested by infants after BFs

and IFs consumption is performed, enabling information regarding these hazardous compounds

and their potential future effects on infants' health.

Based on the biomarker analysis, 7α-OH could be a potential biomarker of BFs which could

not be predicted by just analyzing the exposure and absorption data obtained from SHEDS-HT.

The difference in DOxS distribution between BF, IFs-L, and IFs-P as a result of processing

techniques variability, was confirmed after an sPLS-DA was performed. All 3 groups showed

segregation among samples of the same classification group. Therefore, based on our results

which also compared processing techniques between BF, L-IFs, and P-IFs and their significance,

we support Kilvington's proposal of 7-ketocholesterol as a biomarker for powder IF based on its

specific spray drying and canning techniques employed during its confection (A. Kilvington et

al., 2019; Risso et al., 2022).

Acknowledgments:

This study was funded by the Center for Research Ingredients Safety (CRIS) with the

GR100229 grant, and the USDA National Institute of Food and Agriculture, Hatch project

MICL02526 to I.G.M.M. The authors are grateful to Kristin Isaacs, Center for Computational

Toxicology and Exposure, U.S. Environmental Protection Agency for her help with the SHEDS-

HT software package. The authors are grateful to Nama Naseem, Nicole Urrea, Ashley Xu,

Rosemary Laurito, Grant Gmitter, and Lisa Zou for their assistance during lipid extraction.

Conflict of interest

The authors have declared no conflict of interest.

References

Allen, R. E., & Myers, A. L. (2006). Nutrition in toddlers. *Am Fam Physician, 74*(9), 1527-1532.

Babawale, E. A., Jones, P. J., Mercer, K. E., Lin, H., Yeruva, L., Bar Yoseph, F., & Rutherfurd, S. M. (2018). Modulating Sterol Concentrations in Infant Formula Influences Cholesterol Absorption and Synthesis in the Neonatal Piglet. *Nutrients, 10*(12). doi:10.3390/nu10121848

Cais-Sokolińska, D., Walkowiak-Tomczak, D., & Rudzińska, M. (2022). Photosensitized oxidation of cholesterol and altered oxysterol levels in sour cream: Effects of addition of cucumber pickles. *J Dairy Sci, 105*(6), 4760-4771. doi:10.3168/jds.2022-21856

CDC. (1998). *Recommendations to prevent and control iron deficiency in the United States.* MMWR

CDC. (2001). *Data Table of Infant Weight-for-age Charts.* Retrieved from https://www.cdc.gov/growthcharts/html_charts/wtageinf.htm

CDC. (2021a). *Maternity Practices in Infant Nutrition and Care (mPINC) Survey.* Retrieved from https://www.cdc.gov/breastfeeding/data/mpinc/index.htm

CDC. (2021b). *Vitamins & Minerals.* Retrieved from https://www.cdc.gov/nutrition/infantandtoddlernutrition/vitamins-minerals/index.html

Centre, T. M. I. (2021). The Metabolomics Innovation Centre. Retrieved from https://www.metabolomicscentre.ca/

ChemSpider. (2021). ChemSpider. Retrieved from http://www.chemspider.com/

Chiu, C.-W. a. K. T.-H. a. C. B.-H. (2018). Improved Analytical Method for Determination of Cholesterol-Oxidation Products in Meat and Animal Fat by QuEChERS Coupled with Gas Chromatography–Mass Spectrometry. *Journal of Agricultural and Food Chemistry, 66*(13), 3561-3571. doi:10.1021/acs.jafc.8b00250

Chong, J., Soufan, O., Li, C., Caraus, I., Li, S., Bourque, G., . . . Xia, J. (2018). MetaboAnalyst 4.0: towards more transparent and integrative metabolomics analysis. *Nucleic Acids Res, 46*(W1), W486-W494. doi:10.1093/nar/gky310

Chudy, S., & Teichert, J. (2021). Oxysterols in stored powders as potential health hazards. *Scientific Reports, 11*(1), 21192. doi:10.1038/s41598-021-00636-5

Claumarchirant, L., Matencio, E., Sanchez-Siles, L. M., Alegría, A., & Lagarda, M. J. (2015). Sterol Composition in Infant Formulas and Estimated Intake. *J Agric Food Chem, 63*(32), 7245-7251. doi:10.1021/acs.jafc.5b02647

Collective, G. B. (2017). *Global Breastfeeding Scorecard 2017, tracking progress for breastfeeding policies and programmes.* Paper presented at the UNICEF and World Health Organization, New York and Geneva.

Folch, J., Lees, M., & Sloane Stanley, G. (1957). A simple method for the isolation and purification of total lipides from animal tissues. *J biol Chem, 226*(1), 497-509.

Fox, M. K., Pac, S., Devaney, B., & Jankowski, L. (2004). Feeding infants and toddlers study: What foods are infants and toddlers eating? *J Am Diet Assoc, 104*(1 Suppl 1), s22-30. doi:10.1016/j.jada.2003.10.026

Garcia-Llatas, G., Mercatante, D., López-García, G., & Rodríguez-Estrada, M. T. (2021). Oxysterols – how much do we know about food occurrence, dietary intake and absorption? *Current Opinion in Food Science.* doi:https://doi.org/10.1016/j.cofs.2021.08.001

García-Llatas, G., Cercaci, L., Rodriguez-Estrada, M. T., Lagarda, M. J., Farré, R., & Lercker, G.
(2008). Sterol Oxidation in Ready-to-Eat Infant Foods During Storage. *Journal of*
Agricultural and Food Chemistry, 56(2), 469-475. doi:10.1021/jf072475g
García-Llatas, G., & Rodríguez-Estrada, M. T. (2011). Current and new insights on phytosterol
oxides in plant sterol-enriched food. *Chem Phys Lipids, 164*(6), 607-624.
doi:10.1016/j.chemphyslip.2011.06.005
Gehring, J., Touvier, M., Baudry, J., Julia, C., Buscail, C., Srour, B., . . . Allès, B. (2020).
Consumption of Ultra-Processed Foods by Pesco-Vegetarians, Vegetarians, and Vegans:
Associations with Duration and Age at Diet Initiation. *The Journal of Nutrition, 151*(1),
120-131. doi:10.1093/jn/nxaa196
Gerber. (2021a). Baby Food. In.
Gerber. (2021b). Daily nutrition goals & food groups for Crawlers. Retrieved from
https://www.gerber.com/learning-center/crawler-nutrition-basics
Grau, A., Codony, R., Grimpa, S., Baucells, M. D., & Guardiola, F. (2001). Cholesterol oxidation in
frozen dark chicken meat: influence of dietary fat source, and alpha-tocopherol and
ascorbic acid supplementation. *Meat Science, 57*(2), 197-208. doi:10.1016/s0309-
1740(00)00094-2
Guardiola, F. (2002). *Cholesterol and phytosterol oxidation products: analysis, occurrence, and*
biological effects: The American Oil Chemists Society.
Guardiola, F., Bou, R., Boatella, J., & Codony, R. (2004). Analysis of sterol oxidation products in
foods. *Journal of Aoac International, 87*(2), 441-466.
Hamdan, I. J. A., Claumarchirant, L., Garcia-Llatas, G., Alegría, A., & Lagarda, M. J. (2017). Sterols
in infant formulas: validation of a gas chromatographic method. *Int J Food Sci Nutr,*
68(6), 695-703. doi:10.1080/09637486.2017.1287883
Hamdan, I. J. A., Sanchez-Siles, L. M., Garcia-Llatas, G., & Lagarda, M. J. (2018). Sterols in Infant
Formulas: A Bioaccessibility Study. *J Agric Food Chem, 66*(6), 1377-1385.
doi:10.1021/acs.jafc.7b04635
Hamdan, I. J. A., Sanchez-Siles, L. M., Matencio, E., Bermúdez, J. D., Garcia-Llatas, G., & Lagarda,
M. J. (2018). Sterols in human milk during lactation: bioaccessibility and estimated
intakes. *Food Funct, 9*(12), 6566-6576. doi:10.1039/c8fo01795f
Hardy, S. C., & Kleinman, R. E. (1994). Fat and cholesterol in the diet of infants and young
children: implications for growth, development, and long-term health. *J Pediatr, 125*(5
Pt 2), S69-77. doi:10.1016/s0022-3476(06)80739-0
Harris, J. L., & Pomeranz, J. L. (2020). Infant formula and toddler milk marketing: opportunities
to address harmful practices and improve young children's diets. *Nutr Rev, 78*(10), 866-
883. doi:10.1093/nutrit/nuz095
Isaacs, K. K., Glen, W. G., Egeghy, P., Goldsmith, M. R., Smith, L., Vallero, D., . . . Ozkaynak, H.
(2014). SHEDS-HT: An Integrated Probabilistic Exposure Model for Prioritizing Exposures
to Chemicals with Near-Field and Dietary Sources. *Environmental Science & Technology,*
48(21), 12750-12759. doi:10.1021/es502513w
Jensen, R. G. (1999). Lipids in human milk. *Lipids, 34*(12), 1243-1271.
doi:https://doi.org/10.1007/s11745-999-0477-2
Johnson, R. (1998). Young children and dietary fat intake: how much is too little? *J Am Diet*
Assoc, 98(4), 402. doi:10.1016/S0002-8223(98)00091-1

Khan, M. I., Min, J. S., Lee, S. O., Yim, D. G., Seol, K. H., Lee, M., & Jo, C. (2015). Cooking, storage,
and reheating effect on the formation of cholesterol oxidation products in processed
meat products. *Lipids Health Dis, 14*, 89. doi:10.1186/s12944-015-0091-5
Kilvington, A., Barnaba, C., Rajasekaran, S., Laurens Leimanis, M. L., & Medina-Meza, I. G.
(2021). Lipid profiling and dietary assessment of infant formulas reveal high intakes of
major cholesterol oxidative product (7-ketocholesterol). *Food Chem, 354*, 129529.
doi:10.1016/j.foodchem.2021.129529
Kilvington, A., Barnaba, C., Rajasekaran, S., Laurens, M. L. L., & Gabriela Medina-Meza, I. (2021).
Lipid profiling and dietary assessment of infant formulas reveal high intakes of major
cholesterol oxidative product (7-ketocholesterol). *Food Chemistry*, 129529.
doi:https://doi.org/10.1016/j.foodchem.2021.129529
Kilvington, A., Maldonado-Pereira, L., Torres-Palacios, C., & Medina-Meza, I. (2019).
Phytosterols and their oxidative products in infant formula. *Journal of Food Process
Engineering*, e13151.
Lawrence, R. M., & Lawrence, R. A. (2022). 10 - Normal Growth, Growth Faltering, and Obesity
in Breastfed Infants. In R. A. Lawrence & R. M. Lawrence (Eds.), *Breastfeeding (Ninth
Edition)* (pp. 298-325). Philadelphia: Elsevier.
Maldonado-Pereira, L. (2021). *Comprehensive Study of the Oxidative Stability of the Most
Consumed Ultra-Processed Foods in the USA.* (Ph.D.). Michigan State University, Ann
Arbor. Retrieved from
http://ezproxy.msu.edu/login?url=https://www.proquest.com/dissertations-
theses/comprehensive-study-oxidative-stability-most/docview/2536440379/se-
2?accountid=12598 Retrieved from
http://resolver.ebscohost.com/openurl?ctx_ver=Z39.88-
2004&ctx_enc=info:ofi/enc:UTF-
8&rfr_id=info:sid/ProQuest+Dissertations+%26+Theses+Global&rft_val_fmt=info:ofi/fm
t:kev:mtx:dissertation&rft.genre=dissertations+%26+theses&rft.jtitle=&rft.atitle=&rft.au
=Maldonado-Pereira%2C+Lisaura&rft.aulast=Maldonado-
Pereira&rft.aufirst=Lisaura&rft.date=2021-01-
01&rft.volume=&rft.issue=&rft.spage=&rft.isbn=9798738634932&rft.btitle=&rft.title=C
omprehensive+Study+of+the+Oxidative+Stability+of+the+Most+Consumed+Ultra-
Processed+Foods+in+the+USA&rft.issn=&rft_id=info:doi/ Dissertations & Theses @ CIC
Institutions; ProQuest Dissertations & Theses Global database. (28497722)
Maldonado-Pereira, L., Barnaba, C., De los Campos, G., & Medina-Meza, I. (2021). Evaluation of
the Nutritional Quality of Ultra-Processed Foods (Ready to Eat + Fast Food): Fatty Acid
Composition. *medRxiv*, 2021.2004.2016.21255610. doi:10.1101/2021.04.16.21255610
Maldonado-Pereira, L., Schweiss, M., Barnaba, C., & Medina-Meza, I. G. (2018). The role of
cholesterol oxidation products in food toxicity. *Food and Chemical Toxicology, 118*, 908-
939. doi:10.1016/j.fct.2018.05.059
Micheletta, F., Natoli, S., Misuraca, M., Sbarigia, E., Diczfalusy, U., & Iuliano, L. (2004). Vitamin E
supplementation in patients with carotid atherosclerosis: reversal of altered oxidative
stress status in plasma but not in plaque. *Arterioscler Thromb Vasc Biol, 24*(1), 136-140.
doi:10.1161/01.ATV.0000104028.07929.72

Miller, J. L. (2013). Iron deficiency anemia: a common and curable disease. *Cold Spring Harbor*
perspectives in medicine, 3(7), a011866. doi:10.1101/cshperspect.a011866
Musto, M., Faraone, D., Cellini, F., & Musto, E. (2014). Changes of DNA quality and meat
physicochemical properties in bovine supraspinatus muscle during microwave heating.
Journal of the Science of Food and Agriculture, 94(4), 785-791. doi:10.1002/jsfa.6441
Osada, K., Hoshina, S., Nakamura, S., & Sugano, M. (2000). Cholesterol Oxidation in Meat
Products and Its Regulation by Supplementation of Sodium Nitrite and Apple Polyphenol
before Processing. *Journal of Agricultural and Food Chemistry, 48*(9), 3823-3829.
doi:10.1021/jf991187k
Pie, J. E. a. S. K. a. S. C. (1991). Cholesterol oxidation in meat products during cooking and
frozen storage. *Journal of Agricultural and Food Chemistry, 39*(2), 250-254.
doi:10.1021/jf00002a005
Poli, G., Leoni, V., Biasi, F., Canzoneri, F., Risso, D., & Menta, R. (2022). Oxysterols: From redox
bench to industry. *Redox Biol, 49*, 102220. doi:10.1016/j.redox.2021.102220
Prins, J. R., Schoots, M. H., Wessels, J. I., Campmans-Kuijpers, M. J. E., Navis, G. J., van Goor, H.,
. . . Gordijn, S. J. (2022). The influence of the dietary exposome on oxidative stress in
pregnancy complications. *Molecular Aspects of Medicine, 87*, 101098.
doi:https://doi.org/10.1016/j.mam.2022.101098
Rant, W., Radzik-Rant, A., Świątek, M., Niżnikowski, R., Szymańska, Ż., Bednarczyk, M., . . .
Ślęzak, M. (2019). The effect of aging and muscle type on the quality characteristics and
lipid oxidation of lamb meat. *Arch Anim Breed, 62*(2), 383-391. doi:10.5194/aab-62-383-
2019
Risso, D., Leoni, V., Canzoneri, F., Arveda, M., Zivoli, R., Peraino, A., . . . Menta, R. (2022).
Presence of cholesterol oxides in milk chocolates and their correlation with milk powder
freshness. *PLoS One, 17*(3), e0264288. doi:10.1371/journal.pone.0264288
Rodríguez-Rodríguez, P., Ramiro-Cortijo, D., Reyes-Hernández, C. G., López de Pablo, A. L.,
González, M. C., & Arribas, S. M. (2018). Implication of Oxidative Stress in Fetal
Programming of Cardiovascular Disease. *Frontiers in physiology, 9*, 602-602.
doi:10.3389/fphys.2018.00602
Rose-Sallin, C. (1995). Quantification of Cholesterol Oxidation Products in Milk Powders Using
[2H7]Cholesterol To Monitor Cholesterol Autoxidation Artifacts. *Journal of agricultural*
and food chemistry, 43(4), 935-941.
Sabolová, M., Pohořelá, B., Fišnar, J., Kouřimská, L., Chrpová, D., & Pánek, J. (2017). Formation
of oxysterols during thermal processing and frozen storage of cooked minced meat. *J Sci*
Food Agric, 97(15), 5092-5099. doi:10.1002/jsfa.8386
Sander, B. D., Addis, P. B., Park, S. W., & Smith, D. E. (1989). Quantification of Cholesterol
Oxidation-Products in a Variety of Foods. *Journal of food protection, 52*(2), 109-114.
Timby, N., Adamsson, M., Domellöf, E., Grip, T., Hernell, O., Lönnerdal, B., & Domellöf, M.
(2021). Neurodevelopment and growth until 6.5 years of infants who consumed a low-
energy, low-protein formula supplemented with bovine milk fat globule membranes: a
randomized controlled trial. *Am J Clin Nutr, 113*(3), 586-592. doi:10.1093/ajcn/nqaa354
Timby, N., Domellöf, E., Hernell, O., Lönnerdal, B., & Domellöf, M. (2014). Neurodevelopment,
nutrition, and growth until 12 mo of age in infants fed a low-energy, low-protein

627   formula supplemented with bovine milk fat globule membranes: a randomized
628   controlled trial. *Am J Clin Nutr, 99*(4), 860-868. doi:10.3945/ajcn.113.064295
629 USDA. (2018). *What We Eat in America, NHANES 1999–2018*. National Center for Health
630   Statistics: Hyattsville, MD.: Centers for Disease Control and Prevention Retrieved from
631   https://www.cdc.gov/nchs/nhanes/wweia.htm
632 Wei, S., Wei, Y., Gong, Y., Chen, Y., Cui, J., Li, L., . . . Yi, L. (2022). Metabolomics as a valid
633   analytical technique in environmental exposure research: application and progress.
634   *Metabolomics, 18*(6), 35. doi:10.1007/s11306-022-01895-7
635 Willinger, T. (2019). Oxysterols in intestinal immunity and inflammation. *J Intern Med, 285*(4),
636   367-380. doi:10.1111/joim.12855
637 Woollett, L. A., & Heubi, J. E. (2000). Fetal and Neo-natal Cholesterol Metabolism. In *Endotext*.
638   South Dartmouth, MA: MDText.com, Inc.
639 Wu, F., Tian, F.-J., & Lin, Y. (2015). Oxidative Stress in Placenta: Health and Diseases. *BioMed*
640   *Research International, 2015*, 293271. doi:10.1155/2015/293271
641 Zmysłowski, A., & Szterk, A. (2019). Oxysterols as a biomarker in diseases. *Clin Chim Acta, 491*,
642   103-113. doi:10.1016/j.cca.2019.01.022
643 Zubin Maslov, P., Hill, J. A., Lüscher, T. F., & Narula, J. (2021). High-sugar feeding and increasing
644   cholesterol levels in infants. *Eur Heart J, 42*(12), 1132-1135.
645   doi:10.1093/eurheartj/ehaa868
646 Öhlund, I., Hörnell, A., Lind, T., & Hernell, O. (2008). Dietary fat in infancy should be more
647   focused on quality than on quantity. *European Journal of Clinical Nutrition, 62*(9), 1058-
648   1064. doi:10.1038/sj.ejcn.1602824
649

Table 1S: Food Codes for Baby Foods Used in this Study

Sample ID	Baby Food	Brand
BF1 - RTE	Beef and broth/gravy	Beech-Nut
BF2 - RTE	Chicken and broth/gravy	Gerber
BF3 - RTE	Vegetables and beef	Gerber
BF4 - RTE	Vegetables and chicken	Gerber
BF5 - RTE	Chicken noodle dinner	Gerber
BF6 - RTE	Macaroni, tomato, and cheese	Gerber
BF7 - RTE	Turkey, and rice	Gerber
BF8 - RTE	Turkey and broth/gravy	Beech-Nut
BF9 - RTE	Fruit yogurt	Gerber
BF10 - RTE	Chicken with rice	Gerber
BF11 - RTE	Vegetables and turkey	Gerber
BF12 - RTE	Macaroni and cheese with vegetables	Gerber
BF13 - RTE	Pasta pick-ups (cheese ravioli)	Gerber

Table 2S: DOxS Profiling in BF Samples

	BF1	BF2	BF3	BF4	BF5	BF6	BF7	BF8	BF9	BF10	BF11	BF12	BF13	Total Average ±STD
Serving Size (g)	71.00	113.40	113.40	113.40	113.40	128.00	113.40	71.00	113.40	113.40	113.40	113.40	85.00	-
Cholesterol (mg/100 g fat) ± STD	1,247.23 ± 227.45	673.84 ± 98.50	58.17 ± 8.50	319.43 ± 83.74	348.65 ± 82.26	123.24 ± 29.61	263.77 ± 76.69	683.55 ± 310.63	162.73 ± 95.67	251.02 ± 57.95	313.34 ± 27.54	172.87 ± 17.84	327.40 ± 87.35	380.40 ± 307.87
Cholesterol Reference (mg/100 g fat)	1,000.00	833.33	250.00	500.00	400.00	7.81	400.00	875.00	333.33	333.33	285.71	333.33	500.00	-
Phytosterols (mg/100 g fat) ±STD														
Brassicasterol	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	19.72 ± 3.21	29.60 ± 3.30	ND	24.66 ± 4.94
Campesterol	ND	ND	10.48 ± 5.14	182.76 ± 53.20	136.48 ± 47.29	21.98 ± 6.84	190.88 ± 21.15	ND	ND	48.06 ± 9.69	ND	99.33 ± 24.73	44.86 ± 3.55	91.85 ±66.85
Stigmasterol	ND	ND	6.52 ± 1.87	75.16 ± 41.86	33.77 ± 7.19	ND	ND	ND	ND	25.41 ± 14.33	48.31 ± 2.23	ND	ND	37.83 ± 23.04
β-Sitosterol	ND	ND	47.10 ± 13.21	498.98 ± 129.71	391.77 ± 93.78	100.31 ± 19.76	421.59 ± 137.27	ND	ND	160.14 ± 26.56	435.62 ± 41.06	236.18 ± 18.19	ND	286.46 ± 160.93
Fucosterol	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.046 ± 0.014	0.046 ± 0.00
Total Phytosterols	ND	ND	64.1	756.9	562.02	122.29	612.47	0	0	233.61	503.65	365.11	44.91	251.16 ± 264.06

*P-value < 0.001

Table 2S: DOxS Profiling in BF Samples (cont'd)

	BF1	BF2	BF3	BF4	BF5	BF6	BF7	BF8	BF9	BF10	BF11	BF12	BF13	Total Average ±STD
	COPs (mg/100 g fat) ±STD													
7α-OH	8.97 ± 5.42	2.92 ± 0.73	0.10 ± 0.031	1.80 ± 0.22	2.10 ± 0.50	3.73 ± 1.98	2.12 ± 0.16	5.85 ± 1.82	0.68 ± 0.12	2.08 ± 0.63	0.66 ± 0.092	0.41 ± 0.10	4.22 ± 3.77	2.74 ± 2.40
7β-OH	24.61 ± 15.21	2.27 ± 0.64	0.065 ± 0.037	1.61 ± 0.24	1.94 ± 0.37	0.66 ± 0.12	2.38 ± 0.15	4.91 ± 1.12	0.59 ± 0.17	1.62 ± 0.46	0.58 ± 0.020	0.40 ± 0.078	1.27 ± 0.89	3.30 ± 6.27
5,6α-epoxy	0.69 ± 0.36	0.044 ± 0.012	ND	ND	0.052 ± 0.021	ND	ND	0.030 ± 0.019	ND	0.027 ± 0.016	ND	ND	0.062 ± 0.023	0.15 ± 0.24
5,6β-epoxy	1.39 ± 0.81	0.033 ± 0.0062	ND	ND	0.045 ± 0.027	ND	ND	0.073 ± 0.039	ND	0.042 ± 0.032	1.05 ± 0.12	ND	0.10 ± 0.025	0.39 ± 0.53
7-keto	16.69 ± 5.45	0.84 ± 0.34	ND	0.57 ± 0.065	0.59 ± 0.18	0.67 ± 0.42	0.48 ± 0.15	1.68 ± 0.61	0.24 ± 0.075	0.65 ± 0.087	ND	ND	1.31 ± 0.83	2.37 ± 4.79
Triol	2.63 ± 0.46	0.22 ± 0.031	ND	0.27 ± 0.086	0.31 ± 0.091	0.20 ± 0.034	0.39 ± 0.056	0.27 ± 0.10	0.21 ± 0.050	0.35 ± 0.041	0.12 ± 0.036	0.18 ± 0.013	0.15 ± 0.03	0.44 ± 0.66
6-keto	ND	0.11 ± 0.064	ND	ND	ND	ND	ND	0.12 ± 0.018	ND	ND	ND	0.10 ± 0.044	ND	0.11 ± 0.01
20α-OH	ND	ND	0.65 ± 0.17	ND	ND	ND	ND	ND	ND	ND	5.29 ± 0.40	ND	ND	2.97 ± 2.32
22-OH	ND	ND	0.07 ± 0.03	ND	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.07 ± 0.00
24-OH	ND	ND	ND	ND	ND	0.11 ± 0.06	0.13 ± 0.02	0.45 ± 0.20	ND	ND	ND	ND	0.25 ± 0.19	0.24 ± 0.14

25-OH	3.60 ± 1.22	0.24 ± 0.055	ND	0.16 ± 0.056	0.27 ± 0.16	ND	ND	0.67 ± 0.16	ND	0.27 ± 0.047	ND	ND	0.47 ± 0.32	0.81 ± 1.15
Total COPs	58.59 ± 27.02	6.68 ± 1.80	0.88 ± 0.11	4.61 ± 0.64	5.31 ± 1.22	5.37 ± 2.51	5.51 ± 0.15	14.06 ± 3.95	1.71 ± 0.37	5.04 ± 1.25	7.69 ± 0.42	1.08 ± 0.23	7.85 ± 6.04	9.57 ± 14.53

Table 3S: Values of Exposure, Dose Intake, and Absorption Provided by SHEDS-HT

	Females			Males			Total		
	Exposure (ug/day)	*Dose Intake (ug/day)*	*Absorption (mg/kg-day)*	*Exposure (ug/day)*	*Dose Intake (ug/day)*	*Absorption (mg/kg-day)*	*Exposure (ug/day)*	*Dose Intake (ug/day)*	*Absorption (mg/kg-day)*
Cholesterol	77,575.04	7.75	2.44	46,101.42	4.44	1.40	61,391.3	6.04	1.90
Desmosterol	9,689.138	0.99	0.38	8,286.66	0.79	0.30	8,967.98	0.90	0.34
Squalene	1,339.386	0.13	0.13	1,576.29	0.15	0.15	1,461.20	0.14	0.14
Phytosterols									
Brassicasterol	0.087003	9.10E-06	2.84E-06	42.59	0.0041	0.0013	21.94	0.0021	0.00066
β-Sitosterol	94,659.75	9.64	0.49	57,224.17	5.46	0.28	75410.37	7.49	0.38
Campesterol	27,609.45	2.85	0.41	20,512.97	1.97	0.28	23,960.44	2.40	0.34
Fucosterol	10,530.81	1.10	0.044	7,890.36	0.76	0.030	9,173.092	0.93	0.037
Stigmasterol	18,091.73	1.88	0.075	13,095.2	1.27	0.051	15,522.51	1.56	0.063
Tocopherols									
α-Tocopherol	333.66	0.034	0.0011	844.76	0.082	0.0028	596.47	0.059	0.0020
β-Tocopherol	604.36	0.064	0.0022	866.75	0.085	0.0029	739.28	0.075	0.0026
δ-Tocopherol	12,780.61	1.34	0.047	8,782.58	0.86	0.030	10,724.83	1.09	0.038
γ-Tocopherol	2,75.94	0.22	0.0075	1,656.91	0.16	0.0055	1,860.48	0.19	0.0065
DOxS									
20-OH	0.38	4.03E-05	1.61E-05	1.30	0.00013	5.03E-05	0.86	8.42E-05	3.37E-05
22-OH	0.30	3.12E-05	1.25E-05	0.21	2.04E-05	8.16E-06	0.25	2.56E-05	1.03E-05
24-OH	0.32	3.29E-05	1.32E-05	0.31	2.98E-05	1.19E-05	0.31	3.13E-05	1.25E-05
25-OH	132.46	0.014	0.0056	98.85	0.0096	0.0038	115.17	0.012	0.0047
4-cholesten-3,6-dione	0	0	0	0.0055	5.28E-07	2.15E-07	0.0028	2.71E-07	1.1E-07
6-Keto	8.64	0.00091	0.00037	103.38	0.010	0.0042	57.35	0.0057	0.0023
7α-OH	120.85	0.013	0.0038	207.76	0.020	0.0060	165.54	0.017	0.0050
7β-OH	127.20	0.013	0.0056	252.86	0.024	0.010	191.81	0.019	0.0080
7-Keto	36.18	0.0038	0.00046	80.57	0.0078	0.00094	59.01	0.0059	0.00071
5,6α-Epoxy	4.67	0.00049	0.00014	10.25	0.00099	0.00028	7.54	0.00075	0.00021
5,6β-Epoxy	732.54	0.076	0.024	1,727.98	0.17	0.054	1,244.39	0.12	0.040
Triol	45.87	0.0048	0.00072	78.35	0.0076	0.0011	62.57	0.0063	0.00094

Figure 1S: Distribution of Bioactive Lipids Concentrations Among BF