Multi-omic association study identifies DNA methylation-mediated genotype and smoking exposure effects on lung function in children living in urban settings

Matthew Dapas[1], Emma E. Thompson[1], William Wentworth-Sheilds[1], Selene Clay[1], Cynthia M. Visness[2], Agustin Calatroni[2], Joanne E Sordillo[3], Diane R. Gold[4,5], Robert A. Wood[6], Melanie Makhija[7], Gurjit K. Khurana Hershey[8,9], Michael G. Sherenian[8,9], Rebecca S. Gruchalla[10], Michelle A. Gill[11], Andrew H. Liu[12], Haejin Kim[13], Meyer Kattan[14], Leonard B. Bacharier[15], Deepa Rastogi[16], Matthew C. Altman[17], William W. Busse[18], Patrice M. Becker[19], Dan Nicolae[20], George T. O'Connor[21], James E. Gern[18], Daniel J. Jackson[18], Carole Ober[1]

[1] Department of Human Genetics, University of Chicago, Chicago IL
[2] Rho Inc., Durham, NC
[3] Department of Population Medicine, Harvard Medical School, Boston, MA
[4] Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA
[5] Channing Division of Network Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA
[6] Department of Pediatrics, Johns Hopkins University Medical Center, Baltimore, MD
[7] Division of Allergy and Immunology, Ann & Robert H. Lurie Children's Hospital, Chicago, IL
[8] Department of Pediatrics, University of Cincinnati College of Medicine, Cincinnati, OH
[9] Division of Asthma Research, Cincinnati Children's Hospital Medical Center, Cincinnati, OH
[10] Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, TX
[11] Department of Pediatrics, Washington University School of Medicine, St. Louis, MO
[12] Department of Allergy and Immunology, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, CO
[13] Department of Medicine, Henry Ford Health System, Detroit, MI
[14] Columbia University College of Physicians and Surgeons, New York, NY
[15] Monroe Carell Jr. Children's Hospital at Vanderbilt University Medical Center, Nashville, TN
[16] Children's National Health System, Washington, DC
[17] Department of Allergy and Infectious Diseases, University of Washington, Seattle, WA
[18] Department of Pediatrics and Medicine, University of Wisconsin School of Medicine and Public Health, Madison, WI
[19] National Institute of Allergy and Infectious Diseases, Bethesda, MD
[20] Department of Statistics, University of Chicago, Chicago, IL
[21] Pulmonary Center, Boston University School of Medicine, Boston, MA

ABSTRACT

Impaired lung function in early life is associated with the subsequent development of chronic respiratory disease. Most genetic associations with lung function have been identified in adults of European descent and therefore may not represent those most relevant to pediatric populations and populations of different ancestries. In this study, we performed genome-wide association analyses of lung function in a multiethnic cohort of children (n=1035) living in low-income urban neighborhoods. We identified one novel locus at the *TDRD9* gene in chromosome 14q32.33 associated with percent predicted forced expiratory volume in one second (FEV_1) (p=2.4x10^{-9}; β_z= -0.31, 95% CI= -0.41- -0.21). Mendelian randomization and mediation analyses revealed that this genetic effect on FEV_1 was partially mediated by DNA methylation levels at this locus in airway epithelial cells, which were also associated with environmental tobacco smoke exposure (p=0.015). Promoter-enhancer interactions in airway epithelial cells revealed chromatin interaction loops between FEV_1-associated variants in *TDRD9* and the promoter region of the *PPP1R13B* gene, a stimulator of p53-mediated apoptosis. Expression of *PPP1R13B* in airway epithelial cells was significantly associated the FEV_1 risk alleles (p=1.26x10^{-5}; β=0.12, 95% CI=0.06- 017). These combined results highlight a potential novel mechanism for reduced lung function in urban youth resulting from both genetics and smoking exposure.

AUTHOR SUMMARY

Lung function is determined by both genetic and environmental factors. Impairment of lung function can result from harmful environmental exposures in early life, which disproportionally affect children living in low-income, urban communities. However, most genetic association studies of lung function have been performed in adults and without regard for socioeconomic status. Therefore, genetic risk factors discovered to date may not reflect those most relevant to high-risk populations. In this study, we sought to identify genetic variants correlated with lung function in a multiethnic cohort of children living in low-income, urban neighborhoods and analyze how tobacco smoke exposure may influence any genetic effects. We discovered a common genetic variant associated with lower lung function in this population, and we found that the association was mediated by nearby epigenetic changes in DNA methylation, which were in turn correlated with smoking exposure. We then identified a nearby gene, *PPP1R13B*, which is known to aid in the deactivation of damaged cells, whose expression in airway cells aligned with these genetic and epigenetic effects. This study reveals a potential mechanism through which genetic risk and environmental exposures can affect airway development, perhaps leading to interventions that can help reduce the burden of asthma in socioeconomically disadvantaged children.

INTRODUCTION

Reduced lung function is a hallmark of asthma and chronic obstructive pulmonary disease (COPD). Lung function measures, such as forced expiratory volume in one second (FEV_1) and forced vital capacity (FVC), are strong predictors of future all-cause mortality [1-6]. Airway obstruction often begins in early life [7-10], with lower lung function in infancy being a risk factor for the development of asthma in childhood [11] and COPD in late adulthood [12].

Genetic factors contribute to differences in lung function among individuals, with heritability estimates ranging from 0.50 for FEV_1 to 0.66 for FEV_1/FVC ratio [13]. The many genome-wide association studies (GWAS) of lung function measures [14-25] have implicated pathways related to lung development [20,26-28], inflammation [26], and tissue repair [29], among others [29]. Lung function is also affected by environmental exposures, such as smoking [30-32] and air pollution [33], which can disrupt airway development in early life, increasing the risk of childhood asthma and perhaps other chronic obstructive diseases [8-12,34,35]. For example, exposure to second hand smoke in utero and through childhood is associated with increased risk of childhood asthma [36], lower lung function in adolescence [37], and larger declines in lung function later in life [38,39]. Such adverse exposures are known to alter the epigenetic landscape in exposed individuals [40,41], potentially mediating downstream biological effects [42-44] and modifying genetic associations with lung function [45,46].

Environmental risk factors disproportionally affect socioeconomically disadvantaged children, particularly those living in urban environments [47,48]. In fact, socioeconomic effects contribute to disparities in lung health [49], including the higher burden of chronic respiratory disease among Black and Hispanic children compared to non-Hispanic white children [49-52]. Most genetic association studies of lung function, however, have been limited to adults of European descent. Therefore, genetic risk factors discovered to date may not reflect those most relevant to high-risk populations, which can further exacerbate health disparities [53,54]. Identifying genetic variants and epigenetic variation associated with lung function in high-risk, multiethnic, pediatric populations may provide more direct insights into the early development of impaired lung function.

In this study, we analyzed measures of lung function from the Asthma Phenotypes in the Inner

City (APIC) [55,56] and Urban Environment and Childhood Asthma (URECA) cohorts [57], which

consist of children living in low-income neighborhoods in 10 U.S. cities. We performed whole-genome

sequencing (WGS) on 1,035 participants from APIC and URECA (ages 5-17 years; 67% non-Hispanic

Black, 25% Hispanic; 66% with doctor-diagnosed asthma) and performed a GWAS with FEV_1 and the

FEV_1/FVC ratio. We then performed expression quantitative trait locus (eQTL) and methylation

quantitative trait locus (meQTL) mapping in airway epithelial cells and peripheral blood mononuclear

cells (PBMCs) from a subset of the URECA children. We further tested for genotype and DNA

methylation interactions with smoking exposure. We aimed to identify methylation-mediated genetic and

smoking exposure associations with lung function, linking environmental effects, epigenetic

modifications, and specific genetic risk alleles to reduced pulmonary health in urban youth.

RESULTS

Genetic variants at the *TDRD9* locus are associated with lung function

We completed WGS and variant calling on 1,035 participants from the APIC and URECA studies

(APIC=508, URECA=527; **Table 1**). The mean sequencing depth was 31.6x per sample (**S1A Fig**). On

average, 95.3%, 90.3% and 62.6% of each genome was mapped with at least 10x, 20x and 30x

sequencing read depth, respectively (**S1B Fig**). Approximately 3.8 million high-confidence autosomal

variants were called per sample. Variant call concordance between replicate sample pairs (n=3) was

>99.9% for single nucleotide polymorphisms (SNPs) and was 98.9% for insertions and deletions (**S1**

Table).

Table 1. Demographic characteristics of sequenced APIC and URECA participants.

Characteristic	All	APIC	URECA
Number	1035	508	527
Age, years, mean (SD)	10.3 (2.5)	10.9 (3.1)	9.6 (1.1)
Female sex	477 (46%)	216 (43%)	261 (50%)
Race/Ethnicity			
Black (non-Hispanic)	696 (67%)	319 (63%)	377 (72%)
White (non-Hispanic)	14 (1%)	7 (1%)	7 (1%)
Hispanic	258 (25%)	153 (30%)	105 (20%)
Other/mixed	64 (6%)	26 (5%)	38 (7%)
Unknown	3 (<1%)	2 (<1%)	1 (<1%)
Site			
Baltimore	234 (23%)	85 (17%)	149 (28%)
Boston	189 (18%)	65 (13%)	124 (23%)
Chicago	62 (6%)	62 (12%)	-
Cincinnati	45 (4%)	45 (9%)	-
Dallas	38 (4%)	38 (9%)	-
Denver	59 (6%)	59 (12%)	-
Detroit	50 (5%)	50 (10%)	-
New York	164 (16%)	64 (13%)	100 (19%)
St. Louis	155 (15%)	-	155 (29%)
Washington, D.C.	39 (4%)	39 (8%)	-
Household income < $15k	550 (54%)	273 (54%)	277 (54%)
Caretaker completed HS	756 (73%)	364 (72%)	392 (74%)
Caretaker smokes*	315 (33%)	123 (27%)	192 (39%)
Asthma	681 (75%)	508 (100%)	173 (43%)
BMI, Z-score, mean (SD)	0.9 (1.2)	1.0 (1.2)	0.8 (1.1)
FEV_1, % predicted, mean (SD)	94.9 (16.3)	91.9 (17.6)	98.5 (14.5)
FEV/FVC, mean (SD)	0.80 (0.09)	0.78 (0.10)	0.83 (0.07)

Results are presented as counts and percentages or as means with standard deviations. Missing data were not included in percentage calculations. Ages for URECA correspond to the year the genome-wide association study lung function data were collected. *Caretaker smoking status in URECA was collected at age 10. APIC: Asthma Phenotypes in the Inner City; BMI: body mass index; FEV_1: forced expiratory volume in one second; FEV_1/FVC: ratio of FEV_1 to forced vital capacity; HS: high school; URECA: Urban Environment and Childhood Asthma.

The sequenced cohort included 696 (67%) participants who self-identified as non-Hispanic Black and 258 (25%) who self-identified as Hispanic (**Table 1**). Principal component and admixture analyses using genotypes were used to characterize the ancestry of the participants (**Fig 1**). This revealed that the

124   genetic ancestry of our sample was 66% African, 26% European, 7% Native American, and 1% East

125   Asian. The cohort was 54% male and included 681 (66%) children diagnosed with asthma (**Table 1**).

126
Fig 1. Ancestry composition of sequenced APIC & URECA participants. A) The top two principal
components (PCs) of ancestry are plotted for sequenced APIC & URECA participants, colored by self-
identified race/ethnicity, along with the four ancestral reference populations used for determining
ancestry. NS= not specified. B) The proportion of genetic variance explained by each of the top 10 PCs.
C) The relative values of the top 10 PCs are plotted for each sample, colored by reference population. D)
The estimated proportion of admixture from each ancestral population is shown for each sequenced APIC
& URECA participant. Each vertical line corresponds to one sample. 1KG, 1000 Genomes project; HGDP,
Human Genome Diversity Project; YRI, Yoruba in Ibadan, Nigeria; CEU, Utah residents with Northern and
Western European ancestry; CHB, Han Chinese in Beijing, China; JPT, Japanese in Tokyo, Japan; NAT,
Native Americans from HGDP; EAS, East Asian ancestry; AFR, African ancestry; EUR, European
ancestry.
138

Using the WGS variant calls for 14.1 million variants with minor allele frequency (MAF) $\geq$0.01,

we performed a GWAS of two lung function traits: FEV_1 (% predicted) and FEV_1/FVC (Z-scores),

measured between ages 5-17 (**Table 1**, **S2 Fig**), adjusting for age, sex, asthma diagnosis, the first 10

principal components (PCs) of ancestry, and sample relatedness using a linear mixed model [58]. The

FEV_1 GWAS included 896 participants from APIC (n=504) and URECA (n=392), and the FEV_1/FVC

GWAS included 886 participants from APIC (n=497) and URECA (n=389). The genomic control factor,

λ_{GC}, for both GWAS results was 1.02 (**S3 Fig**), indicating adequate control for population stratification.

We identified one locus on chromosome 14q32.33 that was associated with FEV_1 at genome-wide

significance ($p<2.5 \times 10^{-8}$); no other variants were associated with FEV_1 and no variants were associated

with FEV_1/FVC at genome-wide levels of significance (**Fig 2**). The FEV_1 locus on chromosome 14

consisted of a 200 kb region of associated variants in high linkage disequilibrium (LD) across the *TDRD9*

(Tudor Domain Containing 9) gene (**Fig 3**, **S2 Table**). The minor allele at the lead SNP (rs10220464;

MAF=0.30) was significantly associated with lower FEV_1 ($p=2.4 \times 10^{-9}$; β_z= -0.31, 95% confidence

interval (CI)= -0.41- -0.21) and nominally associated with lower FEV_1/FVC ($p=1.1 \times 10^{-3}$; β_z= -0.17, 95%

CI= -0.28- -0.07). Fine-mapping analysis at this locus (chr14:103.7-104.3Mb) revealed one 95% credible

set of effect variables consisting of 59 SNPs, with rs10220464 having the highest individual posterior

inclusion probability among them (**S4 Fig**). We did not detect any significant differences in rs10220464

association effect size by ancestry or asthma status or study for FEV_1 (**Fig 4**). Furthermore, the *TDRD9*

locus remained the only genome-wide significant association when the two GWAS were performed

without adjustment for asthma status (**S5 Fig**). The overall effect size correlations between asthma-

adjusted and unadjusted GWAS results were r=0.981 for FEV_1 and r=0.954 for FEV_1/FVC.

160
161 **Fig 2. Genome-wide association results.** GWAS Manhattan plots for **A)** FEV_1 and **B)** FEV_1/FVC ratio.
162 The horizontal red line indicates genome-wide significance ($p \leq 2.5 \times 10^{-8}$). The dotted horizontal blue line
163 indicates $p=1 \times 10^{-5}$. Variants colored in green are in previously identified GWAS loci [23]. FEV_1, forced
164 expiratory volume in one second; FEV_1/FVC, ratio of FEV_1 to forced vital capacity.

165
166
Fig 3. FEV_1-associated variants on chromosome 14q32.33. FEV_1 association results are shown at the
TDRD9 gene locus. Each variant is plotted according to its position and $-\log_{10}$ p-value, colored by linkage
disequilibrium to the lead variant, rs10220464, within the sample. Candidate cis-Regulatory Elements
(cCREs) from ENCODE [59] are also shown for the region. The inset panel in the upper right shows the
distribution of adjusted FEV_1 values by rs10220464 genotype. FEV_1, forced expiratory volume in one
second; MAF, minor allele frequency; EnhD, distal enhancer-like signature; CTCF, CCCTC-binding factor
sites; enhP, proximal enhancer-like signature; prom, promoter-like signature; K4m3, trimethylation of
histone H3 at lysine 4.

175
176
Fig 4. Rs10220464 effect size heterogeneity. A forest plot of the associations between rs10220464 and
FEV_1 (% predicted) are shown for distinct sub-cohorts distinguished by self-identified race/ethnicity, study,
and asthma status. β_z, the association effect size between the rs10220464 allele count and the adjusted
and normalized FEV_1 (% predicted) values. FEV_1, forced expiratory volume in one second; N, total
number of individuals included in the association test; MAF, minor allele frequency within the sub-cohort;
P, the association p-value.

We examined association results for the previously identified FEV_1 and FEV_1/FVC loci reported

in the meta-analysis of the UK Biobank and SpiroMeta Consortium by Shrine and colleagues (n=400,102)

[23], which included 70 loci for FEV_1 and 117 for FEV_1/FVC. Of these, 64 of the lead SNPs for FEV_1

and 112 for FEV_1/FVC were genotyped in the APIC and URECA sample. Only one SNP, for FEV_1,

replicated with false discovery rate (FDR) q<0.05 (rs9610955; $p=1.0 \times 10^{-4}$; β_z= -0.38, 95% CI= -0.58- -

0.19; **S6 Fig, S7 Fig**). Cumulatively, 56% (n=36) and 54% (n=60) of these SNPs demonstrated consistent

directions of effect for FEV_1 and FEV_1/FVC, respectively, with effect size correlations of 0.29 (95% CI=

0.05- 0.50; p=0.020) for FEV_1 and 0.42 (95% CI=0.25- 0.56; $p=4.2 \times 10^{-6}$) for FEV_1/FVC.

Lung function risk alleles are associated with DNA methylation at the *TDRD9* locus in airway

epithelial cells

The majority of complex trait-associated variants exert effects by altering gene regulatory networks [60-

62]. These changes are often marked by quantitative differences in DNA methylation levels [63-65]. We

therefore investigated correlations between the FEV_1-associated allele at *TDRD9* and DNA methylation at

the locus in upper airway (nasal) epithelial cells (NECs) from URECA children at age 11 (n=286). We

tested for associations between the FEV_1 genotype, as tagged by rs10220464, and DNA methylation

levels at 796 CpG sites within 10 kb of any *TDRD9* locus variants associated with FEV_1 at $p<1 \times 10^{-5}$

(n=82 variants). The rs10220464 genotype was an meQTL for 5 CpG sites at an FDR <0.05 (**S3 Table**).

DNA methylation levels at only one of these CpG sites, cg03306306 ($p=2.3 \times 10^{-4}$; $\beta=0.07$, 95% CI=0.03-

0.10; **Fig 5a**), was also significantly associated with FEV_1 at age 10 in URECA ($p=0.011$; $\beta= -11.48$, 95%

CI= -20.27- -2.69; **Fig 5b**). The rs10220464 genotype accounted for 4.7% of residual variation in

cg03306306 methylation, and cg03306306 methylation explained 2.4% of residual variation in FEV_1.

Fig 5. Genotype and FEV_1 associations with DNA methylation. DNA methylation levels at
cg03306306 are shown by rs10220464 genotype and FEV_1 measures are plotted against cg03306306
methylation levels in NECs at age 11 (A, B), and PBMCs at age 7 (C, D) from URECA. FEV_1, forced
expiratory volume in one second; NECs, nasal epithelial cells; PBMCs, peripheral blood mononuclear
cells; URECA, Urban Environment and Childhood Asthma study.

We then analyzed cg03306306 methylation in PBMCs collected at age 7 (n=169) [66] from

URECA children to evaluate whether the genotype and lung function associations observed in NECs were

shared with blood cells. In PBMCs, we observed no correlation between the rs10220464 risk allele and

cg03306306 methylation (**Fig 5c**), nor was there an association between cg03306306 methylation and

FEV_1 (**Fig 5d**). These results indicate that cg03306306 methylation dynamics in the airway epithelium are

not present in peripheral blood cells.

Smoking exposure is associated with DNA methylation at the *TDRD9* locus

DNA methylation at the *TDRD9* locus had previously been associated with maternal smoking during

pregnancy [67,68]. Therefore, we tested for associations between environmental tobacco smoke exposure

(**S8 Fig**) and DNA methylation at this locus in the URECA children. Methylation at cg03306306 in NECs

was significantly associated with nicotine metabolite (cotinine) levels in urine collected at ages 7-10 years

(p=0.015; β=0.03, 95% CI=0.01-0.05; **Fig 6**). Methylation at cg03306306 in PBMCs from age 7 was not

associated with urine cotinine levels.

227
228 **Fig 6. DNA methylation association with smoking exposure.** DNA methylation at cg03306306 in nasal
229 epithelial cells at age 11 are plotted against urine cotinine levels from URECA at ages 8-10 as measured
230 using the NicAlert assay (n=285). URECA, Urban Environment and Childhood Asthma study.
231

To determine if there was an interaction effect between genotype and smoking exposure on DNA

methylation and/or lung function, we repeated the cotinine association tests in URECA with the addition

of an interaction term to assess if the genotype effect differed between individuals with low and high

exposures to smoking. There were no significant genotype-by-smoking exposure interaction effects on

methylation levels in NECs or PBMCs in URECA, nor were there any significant methylation-by-

smoking effects on FEV_1 (**S9 Fig**). There was modest evidence for a genotype-by-smoking exposure

interaction effect on FEV_1 in the combined APIC and URECA sample, but this did not reach statistical

significance (p=0.06, **S10 Fig**). Considering the ages of the participants in APIC and URECA, most

tobacco exposures were likely due to secondhand smoke.

Genetic effects on lung function are mediated by DNA methylation

To determine if DNA methylation at the *TDRD9* locus had a causal effect on lung function, we performed

a Mendelian randomization analysis using two-stage least squares (2SLS) regression. In the first stage,

cg03306306 methylation levels in NECs were regressed on an instrument composed of four meQTLs for

cg03306306 (rs11160777, rs137961671, rs7143936, rs11160776; **Materials and Methods**). In the

second stage, FEV_1 was regressed on the predicted DNA methylation values generated from the first stage

regression, thereby yielding a causal effect estimate of cg03306306 methylation on FEV_1. Urine cotinine

levels were included as a covariate in both stages. The variance explained in the first stage regression was

r^2=0.11. The causal effect of cg03306306 methylation on FEV_1 was statistically significant (p=0.020). We

also tested a single, unweighted allele score of the instrumental variables and observed a causal effect

association of p=0.045 (stage-one r^2=0.10). We further performed a bootstrapped mediation analysis to

test whether the rs10220464 risk allele effect on FEV_1 was mediated by DNA methylation. The indirect

effect of rs10220464 on FEV_1 via cg03306306 methylation was significant, both when including asthma

as a covariate (β_z=-0.04, 95% CI=-0.10- -0.003, percent mediated=14.4%) and when asthma was not

considered (β_z=-0.04, 95% CI=-0.10- -0.002, percent mediated=15.0%). These results indicate that the

effect of the FEV_1-associated genotype at the *TDRD9* locus is partially mediated through its impact on

nearby DNA methylation levels.

Gene expression and promoter-enhancer interactions implicate *PPP1R13B*

Trait-associated variants and DNA methylation often affect the transcriptome by influencing the

expression of one or more neighboring genes [69,70]. Identifying these correlations can help infer causal

mechanisms [71]. Therefore, we next explored the relationship between the genotype for the lead FEV_1

variant rs10220464 and the expression of genes within 1 Mb in NECs and PBMCs from URECA

children. Notably, the rs10220464 genotype was not associated with *TDRD9* expression levels in these

cells (NECs: p=0.60, β= 0.116; PBMCs: p=0.91, β= 0.014). Of the 27 genes that were evaluated (**S4**

Table), rs10220464 was significantly associated with the expression of only one gene, *PPP1R13B*

(Protein Phosphatase 1 Regulatory Subunit 13B; FDR q=$2.77.\times10^{-4}$; p=1.26×10^{-5}; β=0.12, 95% CI=0.06-

017; **Fig 7a**), in NECs. *PPP1R13B* expression levels were also the most strongly associated of the 27

genes with methylation at cg03306306 in NECs (p=0.018; β=0.10, 95% CI=0.02- 018; **Fig 7b**).

PPP1R13B expression in NECs, however, was not associated with FEV_1 or smoking exposure (**S11 Fig**).

Fig 7. *PPP1R13B* gene expression in NECs. *PPP1R13B* gene expression in NECs at age 11 are
plotted against A) rs10220464 genotype (n=324) and B) DNA methylation at cg03306306 in NECs at age
11 (n=254). NECs, nasal epithelial cells; CPM, counts per million.

The transcription start site of *PPP1R13B* resides 87 kb from rs10220464 and 152 kb from

cg03306306, suggesting long-range interactions between the FEV_1-associated genotype and the promoter

of *PPP1R13B*. To determine whether any of the FEV_1-associated GWAS variants at the *TDRD9* locus

resided in regions that physically interacted with the promoters of *cis*-genes, we evaluated chromatin

interactions in lower airway (bronchial) epithelial cells (BECs) [72], assessed by promoter-capture Hi-C.

Forty-two of the GWAS variants resided in regions that interacted with the promoters of 9 different genes

expressed in NECs (**Fig 8**; **S5 Table**). The gene most frequently mapped to these variants was

PPP1R13B, with 15 variants located in 3 different interaction loops. Moreover, the strongest observed

interaction was between a region containing 4 FEV_1-associated variants and the *PPP1R13B* promoter

(CHiCAGO score = 9.38; **S5 Table**), suggesting that this region is an enhancer for *PPP1R13B*

expression. This putative enhancer region is located just 2.21 kb from cg03306306.

288
Fig 8. Promoter-enhancer interactions at *TDRD9* locus in nasal epithelial cells. Promoter-to-
enhancer chromatin interactions captured by Hi-C in nasal epithelial cells from URECA at age 11 are
displayed as grey arcs. SNPs associated with FEV_1 ($p<1x10^{-5}$) are marked by blue lines in the top row
according to their genomic position on chromosome 14. The lead FEV_1 SNP, rs1022464, is highlighted in
yellow. CpG sites associated with rs1022464 (FDR<0.05) are displayed as green markers below the
genes, with cg03306306 highlighted in green. Chromatin Interactions containing SNPs associated with
FEV_1 ($p<1x10^{-5}$) are highlighted in blue. Magenta arcs highlight interactions between the *PPP1R13B*
promoter and regions containing FEV_1 SNPs and/or rs1022464-associated CpG sites. FEV_1, forced
expiratory volume in one second; SNPs, single nucleotide polymorphisms; meQTL, methylation
quantitative trait locus; pcHi-C, promoter capture Hi-C.

Summary of study associations

The associations between the *TDRD9* risk allele, cg03306306 DNA methylation in NECs, smoking

exposure, *PPP1R13B* gene expression, and FEV_1 (% predicted) reported in this study are summarized in

Fig 9.

305
Fig 9. Summary of study associations. The *TDRD9* locus was significantly associated with FEV_1 (%
predicted) in the APIC and URECA cohorts. This association was partially mediated by DNA methylation
at the cg03306306 CpG site in *TDRD9* in NECs, which was also significantly associated with
environmental tobacco smoke exposure. The TDRD9 risk allele and DNA methylation were both
significantly associated with *PPP1R13B* gene expression, but *PPP1R13B* gene expression was not
significantly correlated with FEV_1 itself. Unidirectional arrows represent inferred causality.
312
313

DISCUSSION

Using whole-genome sequence variant calls in an asthma-enriched cohort of predominantly African-

American children raised in urban environments, we identified a genotype at the *TDRD9* locus associated

with lower FEV_1 % predicted. This genotype effect was partially mediated by DNA methylation in airway

epithelial cells, which were also correlated with smoking exposure. Data from RNA-sequencing and

promoter-capture Hi-C in airway epithelial cells suggested that these FEV_1-associated genetic and

epigenetic variations influence the expression of the *PPP1R13B* gene through long-range interactions.

The *PPP1R13B* gene encodes a protein that promotes apoptosis, a form of programmed cell

death, via its interaction with the tumor suppressor p53 and is often referred to by its alias ASPP1

(apoptosis-stimulating protein of p53 1) [73]. In response to oncogenic stress, PPP1R13B translocates to

the nucleus, where it enhances the transcriptional activity of p53 on specific target genes relevant to

apoptosis [74,75]. Exposure to smoking and fine particulate matter induces epithelial apoptosis in the

lung via p53 [76-78]. PPP1R13B may also promote apoptosis in a p53-independent manner by inhibiting

autophagy in response to upregulation by EGR-1 (early growth response protein 1) [79]. EGR-1 mediates

stress-induced proinflammatory responses in the airway epithelium and contributes to the pathogenesis of

COPD [80-85]. Within the lung, *PPP1R13B* is indeed predominantly expressed in epithelial cells,

particularly in alveolar type 2 cells, and less so in immune cells and fibroblasts [86,87]. However, Cheng

and colleagues studied PPP1R13B function in lung fibroblasts and found that it was upregulated

following SiO_2 exposure, where it promoted fibroblast proliferation and migration through endoplasmic

reticulum stress and autophagy pathways [88]. Overall, these studies suggest that *PPP1R13B* plays a key

role in maintaining tissue homeostasis by regulating apoptosis and autophagy in response to

environmental stimuli [74,89,90]. The specific function(s) of this gene in the airway epithelium and its

potential impact on the development of airway obstruction remain to be elucidated. *PPP1R13B*

expression in airway epithelial cells at age 11 was not directly associated with lung function or urine

cotinine levels in the URECA children, but the cofactors of this gene [79,91] have been found previously

to be upregulated in smokers with COPD [81,92]. Given its association with lung function alleles in our

study, its expression in the airway epithelium, and its purported functions in autophagy and apoptosis

pathways, additional study of *PPP1R13B* in lung and airway development is warranted, particularly in the

context of adverse environmental stimuli, many of which are enriched in low-income urban

environments.

In NECs, *PPP1R13B* gene expression was significantly associated with DNA methylation levels

at the cg03306306 CpG site in *TDRD9*. Methylation at the *TDRD9* locus was previously reported to

correlate with specific environmental exposures [67,68,93] and with *TDRD9* expression in blood [67,94].

TDRD9 is lowly expressed in the lung but is detected in alveolar macrophages and in monocytes [86,87].

Interestingly, the gene was among the most differentially expressed genes in alveolar macrophages in

smokers relative to non-smokers [95], and its knockdown in TDRD9-expressing lung carcinomas resulted

in increased apoptosis [96]. Its expression was not correlated with the rs10220464 genotype in URECA

NECs or PBMCs, but rs10220464 is an eQTL for *TDRD9* expression in whole blood in GTEx data [97],

with the minor allele associated with lower *TDRD9* expression. Although evidence from this study points

to *PPP1R13B* in the airway epithelium, we can't exclude the possibility that *TDRD9* or other genes could

contribute to the locus' influence on lung function via other tissues.

The correlations of FEV_1 and rs10220464 with cg03306306 methylation and *PPP1R13B*

expression in NECs were absent in PBMCs. Although global DNA methylation patterns between tissues

are highly correlated [98], tissue-specific differentially methylated regions are more likely to be

functional, particularly if they are positively correlated with gene expression [99]. The *TDRD9* locus has

not been identified in epigenome-wide association studies of lung function [44,100-104], but these

measured DNA methylation from blood, which may be an insufficient proxy for methylation in the lungs

[105]. Indeed, previous studies have found that DNA methylation profiles in NECs are significantly more

predictive of pediatric asthma than those in PBMCs [106,107]. Furthermore, epigenetic biomarkers can

change with age. For example, epigenetic markers for lung function in adults do not replicate in children

[101].

The FEV_1 association signal at the *TDRD9* locus included many variants in high LD across a 200

kb region that could be independently contributing to function. Some of the variants lie in different long-

range enhancers [59]. It is also possible that one or more correlated variants were not included because

they failed quality control standards. In addition, due to the limited sample size of the WGS cohort, we

excluded rare variants (MAF<0.01) from consideration, which could contribute to the signal at this locus.

Additional functional studies are needed to identify the causal variant(s) and full mechanism of action.

If increased methylation at cg03306306 leads to lower FEV_1 in response to environmental

stressors, one might expect to see interaction effects with smoking exposure on FEV_1. We did not detect

any such interactions, although our analyses in that regard were likely underpowered given our observed

effects and sample sizes [36]. Furthermore, because this study was limited to children living in low-

income urban neighborhoods, environmental risk factors are likely to be more prevalent than in the

general population [55-57]. Additionally, such exposures are not necessarily ubiquitous across all the

different neighborhoods and communities represented in this sample, and although environmental tobacco

smoke exposure was examined and the socioeconomic range represented in this study is relatively

narrow, there could be relevant environmental factors that were not considered.

To infer causality, Mendelian randomization and mediation analyses rely on assumptions that are

often difficult to empirically verify. For the Mendelian randomization analysis, we identified instrumental

variants associated with the intermediate cg03306306 that were not independently associated with the

outcome, FEV_1. However, because these variants were selected from the same dataset that the outcome

testing was performed in, they were susceptible to bias from winner's curse [108]. To mitigate the

potential impact from this effect and from weak instruments, we performed a secondary analysis in which

we combined the instrumental variants into a single, unweighted score. For the mediation analysis,

unmeasured confounding can invalidate direct and indirect effect estimates [109]. To protect against such

bias, we systematically tested for confounding associations with additional environmental measures

available in APIC and URECA (**Materials and Methods**). Nonetheless, there may still exist unknown

confounding factors that were not measured. Ultimately the results of the Mendelian randomization and

mediation analyses indicate that methylation at cg03306306 in NECs mediated the rs10220464 genotype

effect on FEV_1, but there was residual correlation between rs10220464 and FEV_1, signifying that the

genotype effect was only partially mediated by cg03306306.

Another limitation of our study was the relatively small size for a GWAS. This likely contributed

to the lack of statistically significant replication for previously identified lung function loci [23],

considering that the observed effects were correlated with results of prior GWAS. However, the APIC

and URECA cohorts represent understudied, high-risk, pediatric populations that likely harbor distinct

genetic and environmental risk factors compared to older, primarily European ancestry cohorts included

in previous GWAS of lung function [14-20,23]. The findings of this study have yet to be replicated in an

independent cohort, and should therefore be considered preliminary; however, it is possible that these

associations would differ in populations with dissimilar ancestry, age, exposures, and/or asthma risk.

There are additional caveats to consider when interpreting our findings. First, this study

integrated data from two cohorts with different recruitment criteria, asthma definitions, and ancestral

compositions. Furthermore, most of the analyses beyond the GWAS were limited to subsets of the

URECA participants. However, we did not observe significant genetic effect heterogeneity for

rs10220464 by study, asthma status, or ancestry. To control for potential population stratification, we

used the first ten PCs of ancestry to adjust lung function values and then included the ancestry PCs as

fixed effects in the GWAS models (**Materials and Methods**). The linear mixed models also included a

genetic relatedness matrix as a random effect to account for residual population structure. Because

children with asthma have lower lung function overall (**Table 1**) and their lung function may be more

affected by environmental exposures [110-112], we adjusted for asthma status in the GWAS, as in

previous GWAS [113-116]. The likelihood of discovering lung function variants with consistent effects in

asthmatics and non-asthmatics was thereby increased, although genetic determinants of lung function may

differ by asthma status [117]. Furthermore, adjusting for disease status could potentially introduce

collider bias [118]. The significant genotype effect at the *TDRD9* locus, however, remained the only

genome-wide-significant association when asthma was excluded as a covariate, and adjustment for

asthma did not substantively alter the mediation results. Second, some of the analyses used data collected

at different timepoints. For example, most of the urine cotinine and spirometry measures were collected at

age 10, but the samples used for the NEC DNA methylation and RNA-seq analyses were collected at age

11. Because DNA methylation and gene expression can change over time [40,119-121], their values at

age 11 may not be fully representative of exposures at age 10. Finally, the promoter-capture Hi-C data

were from lower airway (bronchial) epithelial cells, whereas the DNA methylation and RNA-seq data

were generated from upper airway (nasal) epithelial cells. Although there are transcriptomic differences

between epithelial cells from each compartment, their respective profiles are highly correlated [122-126],

and the use of NECs as a proxy for the lower airway epithelium has been validated for both gene

expression and epigenetic studies [124-127].

Our study identified a novel avenue through which genetic risk and environmental exposures

could affect the airways of children raised in low-income urban neighborhoods. Further research into this

pathway may yield mechanistic insights into the early development of impaired lung function, perhaps

leading to interventions that can help reduce the high incidence and morbidity of chronic respiratory

diseases in socioeconomically disadvantaged children.

MATERIALS AND METHODS

Study Population and Phenotypes

We analyzed samples and phenotypes from two National Institutes of Allergy and Infectious Diseases (NIAID)-funded asthma studies conducted by the Inner-City Asthma Consortium (ICAC)[128]: the Asthma Phenotypes in the Inner City (APIC) study[55,56] and the Urban Environment and Childhood Asthma (URECA) birth cohort study [57]. The APIC study was a 1-year, prospective, epidemiological investigation of children and adolescents with asthma (ages 6-17) living in low-income areas ($\geq$20% of residents below poverty level) in nine U.S. cities (Baltimore, MD; Boston, MA; Chicago, IL; Cincinnati, OH; Dallas, TX; Denver, CO; Detroit, MI; New York, NY; Washington, DC). The APIC participants were required to have a diagnosis of asthma by a physician and to have had at least two episodes requiring bronchodilator administration within the past year [55]. The URECA study enrolled pregnant women living in low-income areas of four U.S. cities (Baltimore, MD; Boston, MA; New York, NY; St. Louis, MO) who reported that either or both parents of the index pregnancy had a history of asthma or allergic diseases [57]. This prospective, longitudinal study followed each child through adolescence, periodically collecting samples and clinical and environmental exposure data. Institutional review board approval was received from all participating sites, and written informed consent was obtained from legal guardians of all participating children, who also assented.

Lung function was assessed using spirometry. Lung function measures used in this study for APIC participants were taken at the study entry visit (V0). For URECA, measurements from age 10 were used when available; otherwise, the most recent measurement after age 5 was used (**S6 Table**). Asthma status was assigned according to study-specific criteria. For APIC, asthma was defined by a doctor's diagnosis of asthma and short-acting beta-agonist use in the year prior [55]. For URECA, asthma status was determined either by doctor diagnosis, lung function reversibility, or symptom recurrence [129]. The 2012 Global Lung Initiative reference equations [130] were applied to generate percent predicted estimates for FEV_1 and Z-scores for FEV_1/FVC ratio. Urine cotinine levels were measured using NicAlert immunochromatographic assays, which report results on a scale of 0-6 according to different cotinine

concentration ranges [131]. For URECA, urine cotinine results were available at age 10 for most

participants (n=391); otherwise, assays from age 8 (n=29) or age 7 (n=2) were used. This study utilized

DNA methylation and RNA-seq data generated for other URECA studies; therefore, the number of

samples included in each analysis varied and was limited by data availability (**S7 Table**, **S12 Fig**).

Whole-Genome Sequencing and Data Processing

DNA was extracted from peripheral blood (APIC, URECA) or cord blood (URECA) and quantified using

an Invitrogen Qubit 3 Fluorometer. DNA quality was assessed using the Thermo Scientific NanoDrop

One spectrophotometer and confirmed using an Agilent TapeStation system. DNA was processed in

batches of 60 using the Illumina Nextera DNA Flex library prep kit with unique dual adaptors. Each set of

60 libraries was sequenced over two NovaSEQ S4 flowcells. Whole-genome sequencing was performed

by the University of Chicago Genomics Facility using the Illumina NovaSEQ6000, which generated 150

bp paired-end reads. Sequencing data processing followed the Broad Institute's Genome Analysis Toolkit

(GATK) best practices for germline short variant discovery, as implemented in the harmonized pipeline

used by the New York Genome Center for TOPMed [132,133]. Reads were aligned to the GRCh38

human reference genome (including alternate loci and decoy contigs) using BWA-MEM (Burrows-

Wheeler Aligner; v0.7.17). Aligned reads further underwent duplicate removal (Picard MarkDuplicates;

v2.8.1) and base quality score recalibration (GATK BaseRecalibrator; v3.8) against known sites

(dbSNP138, known indels, and Mills and 1KG gold standard indels) provided in the GATK resource

bundle. Read alignment metrics were calculated using Picard CollectWgsMetrics (v2.8.1) for all aligned

reads and for aligned reads with base quality and mapping quality $\geq$ 20. DNA contamination levels were

estimated using VerifyBamID2 (v1.0.6) [134]. Samples with estimated DNA contamination >0.05 were

removed from consideration. Samples with poor coverage (<50% of the genome with $\geq$20x depth) were

also removed from further consideration. To identify potential sample swaps, WGS samples were

validated using independent genotyping arrays.

QC Array for Sample Validation

To identify potential WGS sample swaps, we independently genotyped the APIC and URECA

participants using the Illumina QC Array-24 BeadChip. SNPs were tested for Hardy-Weinberg

Equilibrium (HWE) within each self-identified ancestry group using the chi-square test and removed if

they deviated from HWE (Bonferroni-adjusted $p<0.05$) within at least one ancestry. SNPs with call rates

<0.98 were also removed. Samples with total variant call rates <0.95 were not used. Array data with

incorrect or indeterminate sex according to X-chromosome heterozygosity rates (Plink v1.90) were also

not used [135]. For fourteen of the sequenced URECA samples, we used results from the Illumina

Infinium CoreExome+Custom array for sample validation, which were generated and controlled for

quality as described by McKennan and colleagues [136]. WGS and array genotypes were tested for

concordance using VerifyBamID (v1.1.3) [137]. WGS samples that were not validated with array data

were not included in genetic analyses (n=2).

Variant Calling and Quality Control

Variant calls were generated using GATK HaplotypeCaller (v4.1.3.0), accounting for contamination

estimates, for single nucleotide variants and short insertions, deletions, and substitutions. Sample

genotypes were joined using GATK GenomicsDBImport and GenotypeGVCFs over the genomic

intervals defined in the GATK WGS calling region interval list provided in the GATK resource bundle.

Genotypes with read depth (DP) <10 or quality scores (GQ) <20 were set as missing. Sites with ≥ 0.1

missingness were then removed from consideration. Variants with minor allele frequencies >0.05 were

tested for accordance with HWE, accounting for population structure [138]. Sites with common variants

that deviated from structural HWE ($P<1\times10^{-6}$) were removed from consideration. Sites with quality by

depth ratios (QD) <4 or >34 were also removed, as we observed declines in variant transition/transversion

(TS/TV) ratios beyond these bounds (**S13 Fig**). Variant site quality was further evaluated using machine-

learning-based Variant Quality Score Recalibration (VQSR). First, SNPs were modeled using GATK

VariantRecalibrator (v4.1.3.0) with Hapmap 3 and with Omni 2.5M SNP chip array as truth resources,

1000G as a training resource, and dbSNP138 as a known sites resource. InDels were likewise trained with

the Mills and 1KG gold standard indels dataset as a truth resource and dbSNP138 as a known sites

resource. SNPs and InDels with resultant predicted true positive probabilities below 0.997 and 0.990,

respectively, were removed from consideration. Variant call accuracy was assessed by comparing call

concordance between three replicate sequencing samples using VCFtools (v0.1.14) vcf-compare [139].

Variant call file manipulation was conducted using BCFtools (v1.10.2) [140].

Ancestry Estimation

Ancestry principal components (PCs) were calculated on the intersect of high quality single-nucleotide

variants (SNVs) genotyped in the WGS data and several reference panels from the 1000 Genomes Project

(1KG; n=156) [141] and the Human Genome Diversity Project (HGDP; n=52) [142]. Native American

reference samples consisted of 52 samples from the HGDP with <5% non-native ancestry, according to

an analysis of roughly 2 million markers using the program ADMIXTURE (v1.3.0) [143]. These samples

were filtered for site quality (missingness 5%; ExcHet<60; VQSLOD$\geq$8.3929), genotype quality

(GQ$\geq$20) and depth (DP$\geq$10), MAF >0.02, and HWE (p>0.001) [142]. European, West African, and East

Asian reference samples were randomly selected from CEU (n=52), YRI (n=52), JPT (n=26), and CHB

(n=26) samples in the phase 3 1KG reference panel [141]. The combined genotypes were pruned for

linkage disequilibrium (LD) $\leq$0.1 within 1Mb intervals. Ancestry PCs were calculated, accounting for

subject relatedness, using PC-Air [144] and PC-Relate [145]. Initial kinship estimates were produced

using KING[146]. Kinship and PCs were iteratively estimated using PC-Relate and PC-Air, respectively,

until estimates for the top 5 PCs stabilized (n=3). Reference population admixture estimates were

estimated for each WGS sample with ADMIXTURE (v1.3.0), using the 1KG and HGDP reference

samples for supervised analysis [143]. Because sample relatedness can lead to biased admixture estimates

[144,147], admixture was estimated for each WGS sample separately.

Quantitative Trait Association Testing

Quantitative traits were adjusted for covariates and normalized using a two-stage approach [148,149].

First, each trait was regressed on age, sex, asthma status, and the first 10 PCs of ancestry. The residuals

were then rank-normalized using an inverse normal transformation. In the second stage, the normalized

residuals were considered outcome variables in the GWAS, adjusting for the same covariates as in the

first stage. Genome-wide association testing was performed for all high-quality common variant calls

(MAF$\geq$0.01) using a linear mixed model, as implemented in GEMMA [58], with subject relatedness

included as a random effect. Individuals who were not evaluated for asthma at ages 7 or 10 (n=127) were

excluded from trait association testing. The threshold we applied for genome-wide significance was

P$\leq$2.5x10^{-8}, based on a 5x10^{-8} GWAS threshold and further accounting for two tests. To identify potential

collider bias introduced by adjusting for asthma status, we repeated the GWAS without accounting for

asthma status in either covariate-adjustment stage.

Fine-mapping analysis was conducted using SuSiE (SusieR R package v0.12.27) [150]. SuSiE

applies a form of Bayesian variable selection in regression using iterative Bayesian stepwise selection to

identify "credible sets" of variables. Each credible set has a 95% probability of containing at least one

causal effect SNP. Prior to running SuSiE, we regressed asthma, age, sex, and ancestry PCs 1-10 from the

genotype matrix and outcome vector (the normalized FEV_1 residuals).

To explore whether there was lead-SNP effect heterogeneity by ancestry, study, or asthma status,

we performed additional single-SNP quantitative trait association tests within several different sub-

cohorts and introduced interaction effects into our models. For ancestry, we performed separate

association tests in each of the Black, Hispanic, and white populations, according to self-identified

race/ethnicity. We then tested for genotype-by-ancestry interaction effects across APIC and URECA by

using admixture proportions as covariates in our models, in lieu of ancestry PCs, and including an

interaction term with the lead SNP for each continental ancestry group in turn. We tested these interaction

effects using the --gxe argument in GEMMA in four separate models (one for each ancestry). To

determine whether there was effect heterogeneity by study (APIC vs. URECA), we performed separate

association tests in each study and also tested the association across APIC and URECA with the addition

of a study covariate and a genotype-by-study interaction term. For asthma status, we performed separate

association tests in the asthmatics and non-asthmatics and tested a genotype interaction term with asthma

status.

DNA Methylation Analysis

DNA from NECs was collected at age 11 from 287 URECA participants and assessed for genome-wide

methylation patterns using the Illumina Infinium Human Methylation EPIC Beadchip. DNA methylation

levels from PBMCs at age 7 in URECA were collected and processed as previously described [66].

MeQTL analysis was performed using Matrix eQTL [151]. NEC DNA methylation levels were adjusted

globally for sex, array, plate, collection site, DNA concentration, percent ciliated epithelial cells, percent

squamous cells, and ancestry PCs 1-3. Principal components analysis was then performed on the residual

methylation levels, and the first three PCs were included as covariates in the meQTL association

tests. Additional methylation PCs were not included in association tests, as they were significantly

correlated with asthma phenotypes. Associations with FDR-adjusted $P<0.05$ were considered significant.

MeQTL analysis with the PBMC data included sex, collection site, plate, ancestry PCs 1-3, and eight

latent factors [152] (protecting for FEV_1 at age 7) as covariates.

To test CpG site methylation associations with lung function in NECs, we performed linear

regressions on the most recent FEV_1 measures, with age, sex, ancestry PCs 1-3, and methylation PCs 1-3

as covariates. For the PBMC analysis, we set FEV_1 at age 7 as the dependent variable, with sex, collection

site, plate, ancestry PCs 1-3, and latent factors included as covariates.

For association testing with smoking exposures, we ran linear regressions for DNA methylation

and lung function in NECs and PBMCs, as described above, with the addition of cotinine concentrations

as a predictor. We further tested for smoking-by-genotype interaction effects on DNA methylation and

lung function using these models by adding an interaction term (cotinine concentration : rs10220464

genotype). Proportions of explained variance were calculated by squaring partial correlation coefficients

of regression model predictors [153]. One sample from one sibling pair was removed from all methylation

analyses to prevent confounding due to relatedness.

Mendelian Randomization and Mediation Analysis

To assess the causal effects of DNA methylation on lung function, we performed one-sample Mendelian

randomization analysis. We applied a 2SLS regression to URECA samples with WGS and DNA

methylation data (n=285) using ivreg [154]. DNA methylation levels in NECs at the cg03306306 CpG

site were first adjusted for methylation PCs 1-3 and used as the endogenous, exposure variable. The

adjusted and normalized FEV_1 values from the GWAS were set as the dependent outcome variable. Urine

cotinine levels were included as an exogenous covariate (included in both stages). The instrumental

variables were chosen from a set of candidate SNPs that were at least nominally associated with

cg03306306 methylation with $p<0.15$. Clustering of pairwise linkage disequilibrium values between these

SNPs revealed six distinct haplotypes (S**14 Fig**). To ensure instrument exogeneity, each candidate SNP

was tested for association with FEV_1 after conditioning on cg03306306 methylation and urine cotinine,

and SNPs associated with $p<0.05$ were removed from consideration. Of the remaining candidate SNPs,

one was chosen from each haplotype, resulting in an instrument composed of 4 SNPs (rs11160777,

rs137961671, rs7143936, rs11160776). Instrument relevance was validated using the F test, endogeneity

using the Wu-Hausman test, and instrument exogeneity using the Sargon test. We tested two 2SLS

models: one where the instrumental variables were included as individual predictors, and another

featuring an unweighted allele score of the four instrumental variants to reduce potential bias from weak

instruments and/or winner's curse [155,156].

Mediation analysis was conducted with ROBMED [157]. The adjusted and normalized FEV_1

residuals were set as the dependent variable, adjusted cg03306306 methylation as the mediator, and

rs10220464 as the independent variable. Age at FEV1 measurement, sex, asthma status, ancestry PCs 1-3,

and urine cotinine levels were included as covariates. We also performed a secondary mediation analysis

without adjusting for asthma status. To identify additional, potential confounders that could invalidate our

mediation model, we systematically tested for associations with 2 socioeconomic variables and 11

environmental exposures available in APIC and URECA (**S8 Table**, **S15 Fig**). For each environmental

exposure, we tested whether the variable was associated with the mediator (cg03306306) and whether the

variable was associated with the outcome (FEV_1) conditional on the mediator. To ensure no exposure-

mediator interactions, we repeated the cg03306306 association test with FEV_1 with rs10220464 included

as a predictor with a rs10220464: cg03306306 interaction term. The indirect effect of rs10220464 on

FEV_1 via cg03306306 methylation was estimated using 100,000 bootstrap resamples.

Gene Expression Analysis

We analyzed gene expression in NECs and PBMCs from the URECA birth cohort using RNA-seq. The

NEC data were derived from 323 children (155 females, 168 males) at age 11 years at the time of sample

collection, and the PBMC data were derived from 130 (53 females, 77 males) PBMC children aged 2

years at the time of collection. Sequencing reads were mapped and quantified using STAR (v2.6.1) [158]

and samples underwent trimmed means of M-value (TMM) normalization and voom transformation

[159]. Genes with <1 count per million mapped reads (CPM) were removed from analysis. For eQTL

association testing in NECs we corrected for sex, the first three ancestral PCs, collection site, epithelial

cell proportion, sequencing batch, and 14 latent factors [152] using limma [160]. In PBMCs, we corrected

for sex, the first three ancestral PCs, collection site, and 19 latent factors.

Chromatin Interaction Analysis

Chromatin interactions were assessed using promoter capture Hi-C [161,162] in ex vivo human BECs

from 8 adult lung donors, including 4 with asthma. The data was processed and analyzed as previously

described [72,163]. Chromosomal interactions were evaluated using the CHiCAGO algorithm [164].

Interactions with CHiCAGO scores ≥5 were considered significant [164]. Genetic variants within 1 kb of

a given interacting fragment were considered part of the chromatin loop. Genes that were not expressed in

NECs were not included in the analysis.

DATA AVAILABILITY

Phenotype, genotype, GWAS summary statistics, and whole-genome sequencing files are available in dbGaP under accession **phs002921.v1.p1**. The URECA gene expression data are available on the Gene Expression Omnibus (GEO) under the accession **GSE145505** (NECs) and **GSE96783** (PBMCs). The pcHi-C data are available in GEO under the reference series **GSE152550**. The DNA methylation data will be deposited to GEO (Morin, et al. A Functional Genomics Pipeline to Identify High-Value Asthma and Allergy CpGs in the Human Methylome. medRxiv. 2022:2022.05.19.22275204).

ACKNOWLEDGEMENTS

We are grateful to all participants and their families who took part in these studies. We would like to extend special thanks to Pieter Faber and the University of Chicago Genomics Facility, as well as to Petra LeBeau and Rebecca Z. Krouse, formerly of Rho Inc., for their respective contributions.

FUNDING

This work was supported by NIH grants U19 AI62310, HHSN272200900052C, HHSN272201000052I, UM1 AI114271, UG3 OD023282, and UM1 AI160040. Site data collection was supported by the following NIH grants: RR00052, UL1 TR001079 (Baltimore); M01 RR00533, UL1 RR025771, 1 UL1 TR001430 (Boston); UL1 TR000150 (Chicago); UL1 TR000451, UL1 TR001105 (Dallas); Ul1 RR025780 (Denver); UL1 TR000040, M01 RR00071, UL1 RR024156 (New York); UL1 TR000075 (D.C.); UL1 TR000077 (Cincinnati). M.D. was supported by TL1 TR002388 and T32 HL007605.

AUTHOR CONTRIBUTIONS

M.D., W.W.B., P.B., D.N., G.T.O., J.E.G., D.J.J., C.O. made contributions to the conception and/or design of the study. M.D., E.T., and S.C. performed computational analyses. E.T. prepared samples for 'omic analyses. W.W-S. oversaw 'omic data management. C.M.V. and A.C. oversaw clinical data management. J.E.S., D.G., G.T.O., J.E.G., D.J.J., C.O. contributed to interpretation of results. R.A.W.,

667   M.M., G.N.K.H., M.G.S., R.S.G, M.A.G., A.H.L., H.K., M.K., L.B.B., D.S., M.C.A. contributed patient

668   samples and/or data. M.D. wrote the manuscript. All authors read and approved the manuscript.

670   **DISCLOSURES**

REFERENCES

1. Hole DJ, Watt GC, Davey-Smith G, Hart CL, Gillis CR, Hawthorne VM. Impaired lung function and mortality risk in men and women: findings from the Renfrew and Paisley prospective population study. BMJ. 1996;313(7059):711-5; discussion 5-6.

2. Schunemann HJ, Dorn J, Grant BJ, Winkelstein W, Jr., Trevisan M. Pulmonary function is a long-term predictor of mortality in the general population: 29-year follow-up of the Buffalo Health Study. Chest. 2000;118(3):656-64.

3. Mannino DM, Holguin F, Pavlin BI, Ferdinands JM. Risk factors for prevalence of and mortality related to restriction on spirometry: findings from the First National Health and Nutrition Examination Survey and follow-up. Int J Tuberc Lung Dis. 2005;9(6):613-21.

4. Chinn S, Gislason T, Aspelund T, Gudnason V. Optimum expression of adult lung function based on all-cause mortality: results from the Reykjavik study. Respir Med. 2007;101(3):601-9.

5. Miller MR, Pedersen OF, Lange P, Vestbo J. Improved survival prediction from lung function data in a large population sample. Respir Med. 2009;103(3):442-8.

6. Agusti A, Noell G, Brugada J, Faner R. Lung function in early adulthood and health in later life: a transgenerational cohort analysis. Lancet Respir Med. 2017;5(12):935-45.

7. Stern DA, Morgan WJ, Wright AL, Guerra S, Martinez FD. Poor airway function in early infancy and lung function by age 22 years: a non-selective longitudinal cohort study. Lancet. 2007;370(9589):758-64.

8. Stocks J, Hislop A, Sonnappa S. Early lung development: lifelong effect on respiratory health and disease. Lancet Respir Med. 2013;1(9):728-42.

9. McGeachie MJ, Yates KP, Zhou X, Guo F, Sternberg AL, Van Natta ML, et al. Patterns of Growth and Decline in Lung Function in Persistent Childhood Asthma. N Engl J Med. 2016;374(19):1842-52.

10. Agusti A, Faner R. Lung function trajectories in health and disease. Lancet Respir Med. 2019;7(4):358-64.

11. Bisgaard H, Jensen SM, Bonnelykke K. Interaction between asthma and lung function growth in early life. Am J Respir Crit Care Med. 2012;185(11):1183-9.

12. Duijts L, Reiss IK, Brusselle G, de Jongste JC. Early origins of chronic obstructive lung diseases across the life course. Eur J Epidemiol. 2014;29(12):871-85.

13. Klimentidis YC, Vazquez AI, de Los Campos G, Allison DB, Dransfield MT, Thannickal VJ. Heritability of pulmonary function estimated from pedigree and whole-genome markers. Front Genet. 2013;4:174.

14. Wilk JB, Chen TH, Gottlieb DJ, Walter RE, Nagle MW, Brandler BJ, et al. A genome-
wide association study of pulmonary function measures in the Framingham Heart Study.
PLoS Genet. 2009;5(3):e1000429.

15. Repapi E, Sayers I, Wain LV, Burton PR, Johnson T, Obeidat M, et al. Genome-wide
association study identifies five loci associated with lung function. Nat Genet.
2010;42(1):36-44.

16. Hancock DB, Eijgelsheim M, Wilk JB, Gharib SA, Loehr LR, Marciante KD, et al. Meta-
analyses of genome-wide association studies identify multiple loci associated with
pulmonary function. Nat Genet. 2010;42(1):45-52.

17. Soler Artigas M, Loth DW, Wain LV, Gharib SA, Obeidat M, Tang W, et al. Genome-wide
association and large-scale follow up identifies 16 new loci influencing lung function. Nat
Genet. 2011;43(11):1082-90.

18. Yao TC, Du G, Han L, Sun Y, Hu D, Yang JJ, et al. Genome-wide association study of
lung function phenotypes in a founder population. J Allergy Clin Immunol.
2014;133(1):248-55 e1-10.

19. Loth DW, Soler Artigas M, Gharib SA, Wain LV, Franceschini N, Koch B, et al. Genome-
wide association analysis identifies six new loci associated with forced vital capacity. Nat
Genet. 2014;46(7):669-77.

20. Wain LV, Shrine N, Artigas MS, Erzurumluoglu AM, Noyvert B, Bossini-Castillo L, et al.
Genome-wide association analyses for lung function and chronic obstructive pulmonary
disease identify new loci and potential druggable targets. Nat Genet. 2017;49(3):416-25.

21. Burkart KM, Sofer T, London SJ, Manichaikul A, Hartwig FP, Yan Q, et al. A Genome-
Wide Association Study in Hispanics/Latinos Identifies Novel Signals for Lung Function.
The Hispanic Community Health Study/Study of Latinos. Am J Respir Crit Care Med.
2018;198(2):208-19.

22. Wyss AB, Sofer T, Lee MK, Terzikhan N, Nguyen JN, Lahousse L, et al. Multiethnic
meta-analysis identifies ancestry-specific and cross-ancestry loci for pulmonary function.
Nat Commun. 2018;9(1):2976.

23. Shrine N, Guyatt AL, Erzurumluoglu AM, Jackson VE, Hobbs BD, Melbourne CA, et al.
New genetic signals for lung function highlight pathways and chronic obstructive
pulmonary disease associations across multiple ancestries. Nat Genet. 2019;51(3):481-93.

24. Akenroye AT, Brunetti T, Romero K, Daya M, Kanchan K, Shankar G, et al. Genome-wide
association study of asthma, total IgE, and lung function in a cohort of Peruvian children. J
Allergy Clin Immunol. 2021. doi: 10.1016/j.jaci.2021.02.035.

25. Zhu Z, Li J, Si J, Ma B, Shi H, Lv J, et al. A large-scale genome-wide association analysis
of lung function in the Chinese population identifies novel loci and highlights shared
genetic aetiology with obesity. Eur Respir J. 2021;58(4).

26.   Obeidat M, Hao K, Bosse Y, Nickle DC, Nie Y, Postma DS, et al. Molecular mechanisms
underlying variations in lung function: a systems genetics analysis. Lancet Respir Med.
2015;3(10):782-95.

27.   Gharib SA, Loth DW, Soler Artigas M, Birkland TP, Wilk JB, Wain LV, et al. Integrative
pathway genomics of lung function and airflow obstruction. Hum Mol Genet.
2015;24(23):6836-48.

28.   Portas L, Pereira M, Shaheen SO, Wyss AB, London SJ, Burney PGJ, et al. Lung
Development Genes and Adult Lung Function. Am J Respir Crit Care Med.
2020;202(6):853-65.

29.   Kheirallah AK, Miller S, Hall IP, Sayers I. Translating Lung Function Genome-Wide
Association Study (GWAS) Findings: New Insights for Lung Biology. Adv Genet.
2016;93:57-145.

30.   Aschard H, Tobin MD, Hancock DB, Skurnik D, Sood A, James A, et al. Evidence for
large-scale gene-by-smoking interaction effects on pulmonary function. Int J Epidemiol.
2017;46(3):894-904.

31.   Park B, An J, Kim W, Kang HY, Koh SB, Oh B, et al. Effect of 6p21 region on lung
function is modified by smoking: a genome-wide interaction study. Sci Rep.
2020;10(1):13075.

32.   Kim W, Moll M, Qiao D, Hobbs BD, Shrine N, Sakornsakolpat P, et al. Smoking
Interaction with a Polygenic Risk Score for Reduced Lung Function. medRxiv. 2021. doi:
10.1101/2021.03.26.21254415:2021.03.26.21254415.

33.   Melbourne CA, Erzurumluoglu AM, Shrine N, Chen J, Tobin MD, Hansell A, et al.
Genome-wide gene-air pollution interaction analysis of lung function in 300,000
individuals. medRxiv. 2021. doi: 10.1101/2021.06.03.21256376:2021.06.03.21256376.

34.   Miller MD, Marty MA. Impact of environmental chemicals on lung development. Environ
Health Perspect. 2010;118(8):1155-64.

35.   Decrue F, Gorlanova O, Usemann J, Frey U. Lung functional development and asthma
trajectories. Semin Immunopathol. 2020;42(1):17-27.

36.   He Z, Wu H, Zhang S, Lin Y, Li R, Xie L, et al. The association between secondhand
smoke and childhood asthma: A systematic review and meta-analysis. Pediatr Pulmonol.
2020;55(10):2518-31.

37.   Thacher JD, Schultz ES, Hallberg J, Hellberg U, Kull I, Thunqvist P, et al. Tobacco smoke
exposure in early life and adolescence in relation to lung function. Eur Respir J.
2018;51(6).

38. Dratva J, Zemp E, Dharmage SC, Accordini S, Burdet L, Gislason T, et al. Early Life
Origins of Lung Ageing: Early Life Exposures and Lung Function Decline in Adulthood in
Two European Cohorts Aged 28-73 Years. PLoS One. 2016;11(1):e0145127.

39. Savran O, Ulrik CS. Early life insults as determinants of chronic obstructive pulmonary
disease in adult life. Int J Chron Obstruct Pulmon Dis. 2018;13:683-93.

40. Zeilinger S, Kuhnel B, Klopp N, Baurecht H, Kleinschmidt A, Gieger C, et al. Tobacco
smoking leads to extensive genome-wide changes in DNA methylation. PLoS One.
2013;8(5):e63812.

41. Rider CF, Carlsten C. Air pollution and DNA methylation: effects of exposure in humans.
Clin Epigenetics. 2019;11(1):131.

42. Jamieson E, Korologou-Linden R, Wootton RE, Guyatt AL, Battram T, Burrows K, et al.
Smoking, DNA Methylation, and Lung Function: a Mendelian Randomization Analysis to
Investigate Causal Pathways. Am J Hum Genet. 2020;106(3):315-26.

43. Kwak SY, Park CY, Shin MJ. Smoking May Affect Pulmonary Function through DNA
Methylation: an Epigenome-Wide Association Study in Korean Men. Clin Nutr Res.
2020;9(2):134-44.

44. Sunny SK, Zhang H, Relton CL, Ring S, Kadalayil L, Mzayek F, et al. Sex-specific
longitudinal association of DNA methylation with lung function. ERJ Open Res. 2021;7(3).

45. Mukherjee N, Lockett GA, Merid SK, Melen E, Pershagen G, Holloway JW, et al. DNA
methylation and genetic polymorphisms of the Leptin gene interact to influence lung
function outcomes and asthma at 18 years of age. Int J Mol Epidemiol Genet. 2016;7(1):1-
17.

46. Zhang H, Tong X, Holloway JW, Rezwan FI, Lockett GA, Patil V, et al. The interplay of
DNA methylation over time with Th2 pathway genetic variants on asthma risk and
temporal asthma transition. Clin Epigenetics. 2014;6(1):8.

47. Munoz-Pizza DM, Villada-Canela M, Reyna MA, Texcalac-Sangrador JL, Osornio-Vargas
AR. Air pollution and children's respiratory health: a scoping review of socioeconomic
status as an effect modifier. Int J Public Health. 2020;65(5):649-60.

48. Hajizadeh M, Nandi A. The socioeconomic gradient of secondhand smoke exposure in
children: evidence from 26 low-income and middle-income countries. Tob Control.
2016;25(e2):e146-e55.

49. Martinez CH, Mannino DM, Curtis JL, Han MK, Diaz AA. Socioeconomic Characteristics
Are Major Contributors to Ethnic Differences in Health Status in Obstructive Lung
Disease: An Analysis of the National Health and Nutrition Examination Survey 2007-2010.
Chest. 2015;148(1):151-8.

50. Thakur N, Oh SS, Nguyen EA, Martin M, Roth LA, Galanter J, et al. Socioeconomic status
and childhood asthma in urban minority youths. The GALA II and SAGE II studies. Am J
Respir Crit Care Med. 2013;188(10):1202-9.

51. Oraka E, Iqbal S, Flanders WD, Brinker K, Garbe P. Racial and ethnic disparities in current
asthma and emergency department visits: findings from the National Health Interview
Survey, 2001-2010. J Asthma. 2013;50(5):488-96.

52. Keet CA, Matsui EC, McCormack MC, Peng RD. Urban residence, neighborhood poverty,
race/ethnicity, and asthma morbidity among children on Medicaid. J Allergy Clin
Immunol. 2017;140(3):822-7.

53. Manrai AK, Funke BH, Rehm HL, Olesen MS, Maron BA, Szolovits P, et al. Genetic
Misdiagnoses and the Potential for Health Disparities. N Engl J Med. 2016;375(7):655-65.

54. Landry LG, Ali N, Williams DR, Rehm HL, Bonham VL. Lack Of Diversity In Genomic
Databases Is A Barrier To Translating Precision Medicine Research Into Practice. Health
Aff (Millwood). 2018;37(5):780-5.

55. Pongracic JA, Krouse RZ, Babineau DC, Zoratti EM, Cohen RT, Wood RA, et al.
Distinguishing characteristics of difficult-to-control asthma in inner-city children and
adolescents. J Allergy Clin Immunol. 2016;138(4):1030-41.

56. Zoratti EM, Krouse RZ, Babineau DC, Pongracic JA, O'Connor GT, Wood RA, et al.
Asthma phenotypes in inner-city children. J Allergy Clin Immunol. 2016;138(4):1016-29.

57. Gern JE, Visness CM, Gergen PJ, Wood RA, Bloomberg GR, O'Connor GT, et al. The
Urban Environment and Childhood Asthma (URECA) birth cohort study: design, methods,
and study population. BMC Pulm Med. 2009;9:17.

58. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association studies.
Nat Genet. 2012;44(7):821-4.

59. Consortium EP, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N, et al. Expanded
encyclopaedias of DNA elements in the human and mouse genomes. Nature.
2020;583(7818):699-710.

60. Ma M, Ru Y, Chuang LS, Hsu NY, Shi LS, Hakenberg J, et al. Disease-associated variants
in different categories of disease located in distinct regulatory elements. BMC Genomics.
2015;16 Suppl 8:S3.

61. Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From Polygenic to
Omnigenic. Cell. 2017;169(7):1177-86.

62. Watanabe K, Stringer S, Frei O, Umicevic Mirkov M, de Leeuw C, Polderman TJC, et al.
A global overview of pleiotropy and genetic architecture in complex traits. Nat Genet.
2019;51(9):1339-48.

63. Schulz H, Ruppert AK, Herms S, Wolf C, Mirza-Schreiber N, Stegle O, et al. Genome-
wide mapping of genetic determinants influencing DNA methylation and gene expression
in human hippocampus. Nat Commun. 2017;8(1):1511.

64. Huan T, Joehanes R, Song C, Peng F, Guo Y, Mendelson M, et al. Genome-wide
identification of DNA methylation QTLs in whole blood highlights pathways for
cardiovascular disease. Nat Commun. 2019;10(1):4267.

65. Hannon E, Gorrie-Stone TJ, Smart MC, Burrage J, Hughes A, Bao Y, et al. Leveraging
DNA-Methylation Quantitative-Trait Loci to Characterize the Relationship between
Methylomic Variation, Gene Expression, and Complex Traits. Am J Hum Genet.
2018;103(5):654-65.

66. McKennan C, Naughton K, Stanhope C, Kattan M, O'Connor GT, Sandel MT, et al.
Longitudinal data reveal strong genetic and weak non-genetic components of ethnicity-
dependent blood DNA methylation levels. Epigenetics. 2021;16(6):662-76.

67. Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, et al. DNA Methylation
in Newborns and Maternal Smoking in Pregnancy: Genome-wide Consortium Meta-
analysis. Am J Hum Genet. 2016;98(4):680-96.

68. Sikdar S, Joehanes R, Joubert BR, Xu CJ, Vives-Usano M, Rezwan FI, et al. Comparison
of smoking-related DNA methylation between newborns from prenatal exposure and adults
from personal smoking. Epigenomics. 2019;11(13):1487-500.

69. Nicolae DL, Gamazon E, Zhang W, Duan S, Dolan ME, Cox NJ. Trait-associated SNPs are
more likely to be eQTLs: annotation to enhance discovery from GWAS. PLoS Genet.
2010;6(4):e1000888.

70. Yao DW, O'Connor LJ, Price AL, Gusev A. Quantifying genetic effects on disease
mediated by assayed gene expression levels. Nat Genet. 2020;52(6):626-33.

71. Hormozdiari F, van de Bunt M, Segre AV, Li X, Joo JWJ, Bilow M, et al. Colocalization
of GWAS and eQTL Signals Detects Target Genes. Am J Hum Genet. 2016;99(6):1245-60.

72. Helling BA, Sobreira DR, Hansen GT, Sakabe NJ, Luo K, Billstrand C, et al. Altered
transcriptional and chromatin responses to rhinovirus in bronchial epithelial cells from
adults with asthma. Commun Biol. 2020;3(1):678.

73. Samuels-Lev Y, O'Connor DJ, Bergamaschi D, Trigiante G, Hsieh JK, Zhong S, et al.
ASPP proteins specifically stimulate the apoptotic function of p53. Mol Cell.
2001;8(4):781-94.

74. Aylon Y, Ofir-Rosenfeld Y, Yabuta N, Lapi E, Nojima H, Lu X, et al. The Lats2 tumor
suppressor augments p53-mediated apoptosis by promoting the nuclear proapoptotic
function of ASPP1. Genes Dev. 2010;24(21):2420-9.

75. Wang Y, Godin-Heymann N, Dan Wang X, Bergamaschi D, Llanos S, Lu X. ASPP1 and
ASPP2 bind active RAS, potentiate RAS signalling and enhance p53 activity in cancer
cells. Cell Death Differ. 2013;20(4):525-34.

76. Xue H, Li MX. MicroRNA-150 protects against cigarette smoke-induced lung
inflammation and airway epithelial cell apoptosis through repressing p53: MicroRNA-150
in CS-induced lung inflammation. Hum Exp Toxicol. 2018;37(9):920-8.

77. Xu F, Xu A, Guo Y, Bai Q, Wu X, Ji SP, et al. PM2.5 exposure induces alveolar epithelial
cell apoptosis and causes emphysema through p53/Siva-1. Eur Rev Med Pharmacol Sci.
2020;24(7):3943-50.

78. Song Q, Zhou ZJ, Cai S, Chen Y, Chen P. Oxidative stress links the tumour suppressor p53
with cell apoptosis induced by cigarette smoke. Int J Environ Health Res. 2021. doi:
10.1080/09603123.2021.1910211:1-11.

79. Zhao K, Yu M, Zhu Y, Liu D, Wu Q, Hu Y. EGR-1/ASPP1 inter-regulatory loop promotes
apoptosis by inhibiting cyto-protective autophagy. Cell Death Dis. 2017;8(6):e2869.

80. Reynolds PR, Cosio MG, Hoidal JR. Cigarette smoke-induced Egr-1 upregulates
proinflammatory cytokines in pulmonary epithelial cells. Am J Respir Cell Mol Biol.
2006;35(3):314-9.

81. Chen ZH, Kim HP, Sciurba FC, Lee SJ, Feghali-Bostwick C, Stolz DB, et al. Egr-1
regulates autophagy in cigarette smoke-induced chronic obstructive pulmonary disease.
PLoS One. 2008;3(10):e3316.

82. Shen N, Gong T, Wang JD, Meng FL, Qiao L, Yang RL, et al. Cigarette smoke-induced
pulmonary inflammatory responses are mediated by EGR-1/GGPPS/MAPK signaling. Am
J Pathol. 2011;178(1):110-8.

83. Wang SB, Zhang C, Xu XC, Xu F, Zhou JS, Wu YP, et al. Early growth response factor 1
is essential for cigarette smoke-induced MUC5AC expression in human bronchial
epithelial cells. Biochem Biophys Res Commun. 2017;490(2):147-54.

84. Xu F, Cao J, Luo M, Che L, Li W, Ying S, et al. Early growth response gene 1 is essential
for urban particulate matter-induced inflammation and mucus hyperproduction in airway
epithelium. Toxicol Lett. 2018;294:145-55.

85. Golebski K, Gorenjak M, Kabesch M, Maitland-Van Der Zee A-H, Melén E, Potočnik U,
et al. EGR-1 as a potential biomarker in asthma and proinflammatory responses in airway
epithelium. European Respiratory Journal. 2021;58(suppl 65):PA2041.

86. Wang A, Chiou J, Poirion OB, Buchanan J, Valdez MJ, Verheyden JM, et al. Single-cell
multiomic profiling of human lungs reveals cell-type-specific and age-dynamic control of
SARS-CoV2 host genes. Elife. 2020;9.

87. Karlsson M, Zhang C, Mear L, Zhong W, Digre A, Katona B, et al. A single-cell type
transcriptomics map of human tissues. Sci Adv. 2021;7(31).

88. Cheng Y, Luo W, Li Z, Cao M, Zhu Z, Han C, et al. CircRNA-012091/PPP1R13B-
mediated Lung Fibrotic Response in Silicosis via Endoplasmic Reticulum Stress and
Autophagy. Am J Respir Cell Mol Biol. 2019;61(3):380-91.

89. Vigneron AM, Ludwig RL, Vousden KH. Cytoplasmic ASPP1 inhibits apoptosis through
the control of YAP. Genes Dev. 2010;24(21):2430-9.

90. Manfredi JJ. An identity crisis for a cancer gene: subcellular location determines ASPP1
function. Cancer Cell. 2010;18(5):409-10.

91. Fogal V, Kartasheva NN, Trigiante G, Llanos S, Yap D, Vousden KH, et al. ASPP1 and
ASPP2 are new transcriptional targets of E2F. Cell Death Differ. 2005;12(4):369-76.

92. Zhou SJ, Li M, Zeng DX, Zhu ZM, Hu XW, Li YH, et al. Expression variations of
connective tissue growth factor in pulmonary arteries from smokers with and without
chronic obstructive pulmonary disease. Sci Rep. 2015;5:8564.

93. Eguchi A, Nishizawa-Jotaki S, Tanabe H, Rahmutulla B, Watanabe M, Miyaso H, et al. An
Altered DNA Methylation Status in the Human Umbilical Cord Is Correlated with
Maternal Exposure to Polychlorinated Biphenyls. Int J Environ Res Public Health.
2019;16(15).

94. Pierce BL, Tong L, Argos M, Demanelis K, Jasmine F, Rakibuz-Zaman M, et al. Co-
occurring expression and methylation QTLs allow detection of common causal variants and
shared biological mechanisms. Nat Commun. 2018;9(1):804.

95. Philibert RA, Sears RA, Powers LS, Nash E, Bair T, Gerke AK, et al. Coordinated DNA
methylation and gene expression changes in smoker alveolar macrophages: specific effects
on VEGF receptor 1 expression. J Leukoc Biol. 2012;92(3):621-31.

96. Guijo M, Ceballos-Chavez M, Gomez-Marin E, Basurto-Cayuela L, Reyes JC. Expression
of TDRD9 in a subset of lung carcinomas by CpG island hypomethylation protects from
DNA damage. Oncotarget. 2018;9(11):9618-31.

97. Consortium GT. The Genotype-Tissue Expression (GTEx) project. Nat Genet.
2013;45(6):580-5.

98. Lokk K, Modhukur V, Rajashekar B, Martens K, Magi R, Kolde R, et al. DNA methylome
profiling of human tissues identifies global and tissue-specific methylation patterns.
Genome Biol. 2014;15(4):r54.

99. Wan J, Oliver VF, Wang G, Zhu H, Zack DJ, Merbs SL, et al. Characterization of tissue-
specific differential DNA methylation suggests distinct modes of positive and negative
gene expression regulation. BMC Genomics. 2015;16:49.

100. den Dekker HT, Burrows K, Felix JF, Salas LA, Nedeljkovic I, Yao J, et al. Newborn DNA-methylation, childhood lung function, and the risks of asthma and COPD across the life course. Eur Respir J. 2019;53(4).

101. Imboden M, Wielscher M, Rezwan FI, Amaral AFS, Schaffner E, Jeong A, et al. Epigenome-wide association study of lung function level and its change. Eur Respir J. 2019;54(1).

102. Mukherjee N, Arathimos R, Chen S, Kheirkhah Rahimabad P, Han L, Zhang H, et al. DNA methylation at birth is associated with lung function development until age 26 years. Eur Respir J. 2021;57(4).

103. Wang T, Wang W, Li W, Duan H, Xu C, Tian X, et al. Genome-wide DNA methylation analysis of pulmonary function in middle and old-aged Chinese monozygotic twins. Respir Res. 2021;22(1):300.

104. Herrera-Luis E, Li A, Mak ACY, Perez-Garcia J, Elhawary JR, Oh SS, et al. Epigenome-wide association study of lung function in Latino children and youth with asthma. Clin Epigenetics. 2022;14(1):9.

105. Cosin-Tomas M, Bustamante M, Sunyer J. Epigenetic association studies at birth and the origin of lung function development. Eur Respir J. 2021;57(4).

106. Yang IV, Lozupone CA, Schwartz DA. The environment, epigenome, and asthma. J Allergy Clin Immunol. 2017;140(1):14-23.

107. Lin PI, Shu H, Mersha TB. Comparing DNA methylation profiles across different tissues associated with the diagnosis of pediatric asthma. Sci Rep. 2020;10(1):151.

108. Richmond RC, Davey Smith G. Mendelian Randomization: Concepts and Scope. Cold Spring Harb Perspect Med. 2022;12(1).

109. Robins JM, Greenland S. Identifiability and exchangeability for direct and indirect effects. Epidemiology. 1992;3(2):143-55.

110. Li YF, Gilliland FD, Berhane K, McConnell R, Gauderman WJ, Rappaport EB, et al. Effects of in utero and environmental tobacco smoke exposure on lung function in boys and girls with and without asthma. Am J Respir Crit Care Med. 2000;162(6):2097-104.

111. Gilliland FD, Berhane K, Li YF, Rappaport EB, Peters JM. Effects of early onset asthma and in utero exposure to maternal smoking on childhood lung function. Am J Respir Crit Care Med. 2003;167(6):917-24.

112. Schultz ES, Litonjua AA, Melen E. Effects of Long-Term Exposure to Traffic-Related Air Pollution on Lung Function in Children. Curr Allergy Asthma Rep. 2017;17(6):41.

113. Kreiner-Moller E, Bisgaard H, Bonnelykke K. Prenatal and postnatal genetic influence on lung function development. J Allergy Clin Immunol. 2014;134(5):1036-42 e15.

114. Lee EY, Mak ACY, Hu D, Sajuthi S, White MJ, Keys KL, et al. Whole-Genome
Sequencing Identifies Novel Functional Loci Associated with Lung Function in Puerto
Rican Youth. Am J Respir Crit Care Med. 2020;202(7):962-72.

115. Goddard PC, Keys KL, Mak ACY, Lee EY, Liu AK, Samedy-Bates LA, et al. Integrative
genomic analysis in African American children with asthma finds three novel loci
associated with lung function. Genet Epidemiol. 2021;45(2):190-208.

116. Akenroye AT, Brunetti T, Romero K, Daya M, Kanchan K, Shankar G, et al. Genome-wide
association study of asthma, total IgE, and lung function in a cohort of Peruvian children. J
Allergy Clin Immunol. 2021;148(6):1493-504.

117. Imboden M, Bouzigon E, Curjuric I, Ramasamy A, Kumar A, Hancock DB, et al. Genome-
wide association study of lung function decline in adults with and without asthma. J
Allergy Clin Immunol. 2012;129(5):1218-28.

118. Aschard H, Vilhjalmsson BJ, Joshi AD, Price AL, Kraft P. Adjusting for heritable
covariates can bias effect estimates in genome-wide association studies. Am J Hum Genet.
2015;96(2):329-39.

119. Jones MJ, Goodman SJ, Kobor MS. DNA methylation and healthy human aging. Aging
Cell. 2015;14(6):924-32.

120. Ambatipudi S, Cuenin C, Hernandez-Vargas H, Ghantous A, Le Calvez-Kelm F, Kaaks R,
et al. Tobacco smoking-associated genome-wide DNA methylation changes in the EPIC
study. Epigenomics. 2016;8(5):599-618.

121. Tommasi S, Zheng A, Besaratinia A. Exposure of mice to secondhand smoke elicits both
transient and long-lasting transcriptional changes in cancer-related functional networks. Int
J Cancer. 2015;136(10):2253-63.

122. Sridhar S, Schembri F, Zeskind J, Shah V, Gustafson AM, Steiling K, et al. Smoking-
induced gene expression changes in the bronchial airway are reflected in nasal and buccal
epithelium. BMC Genomics. 2008;9:259.

123. Zhang X, Sebastiani P, Liu G, Schembri F, Zhang X, Dumas YM, et al. Similarities and
differences between smoking-related gene expression in nasal and bronchial epithelium.
Physiol Genomics. 2010;41(1):1-8.

124. Brugha R, Lowe R, Henderson AJ, Holloway JW, Rakyan V, Wozniak E, et al. DNA
methylation profiles between airway epithelium and proxy tissues in children. Acta
Paediatr. 2017;106(12):2011-6.

125. Imkamp K, Berg M, Vermeulen CJ, Heijink IH, Guryev V, Kerstjens HAM, et al. Nasal
epithelium as a proxy for bronchial epithelium for smoking-induced gene expression and
expression Quantitative Trait Loci. J Allergy Clin Immunol. 2018;142(1):314-7 e15.

126. Kicic A, de Jong E, Ling KM, Nichol K, Anderson D, Wark PAB, et al. Assessing the
unified airway hypothesis in children via transcriptional profiling of the airway epithelium.
J Allergy Clin Immunol. 2020;145(6):1562-73.

127. Bergougnoux A, Claustres M, De Sario A. Nasal epithelial cells: a tool to study DNA
methylation in airway diseases. Epigenomics. 2015;7(1):119-26.

128. Gergen PJ, Teach SJ, Togias A, Busse WW. Reducing Exacerbations in the Inner City:
Lessons from the Inner-City Asthma Consortium (ICAC). J Allergy Clin Immunol Pract.
2016;4(1):22-6.

129. O'Connor GT, Lynch SV, Bloomberg GR, Kattan M, Wood RA, Gergen PJ, et al. Early-
life home environment and risk of asthma among inner-city children. J Allergy Clin
Immunol. 2018;141(4):1468-75.

130. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic
reference values for spirometry for the 3-95-yr age range: the global lung function 2012
equations. Eur Respir J. 2012;40(6):1324-43.

131. Bernert JT, Harmon TL, Sosnoff CS, McGuffey JE. Use of continine immunoassay test
strips for preclassifying urine samples from smokers and nonsmokers prior to analysis by
LC-MS-MS. J Anal Toxicol. 2005;29(8):814-8.

132. Regier AA, Farjoun Y, Larson DE, Krasheninina O, Kang HM, Howrigan DP, et al.
Functional equivalence of genome sequencing analysis pipelines enables harmonized
variant calling across human genetics projects. Nat Commun. 2018;9(1):4038.

133. Kowalski MH, Qian H, Hou Z, Rosen JD, Tapia AL, Shan Y, et al. Use of >100,000
NHLBI Trans-Omics for Precision Medicine (TOPMed) Consortium whole genome
sequences improves imputation quality and detection of rare variant associations in
admixed African and Hispanic/Latino populations. PLoS Genet. 2019;15(12):e1008500.

134. Zhang F, Flickinger M, Taliun SAG, In PPGC, Abecasis GR, Scott LJ, et al. Ancestry-
agnostic estimation of DNA sample contamination from sequence reads. Genome Res.
2020;30(2):185-94.

135. Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. Second-generation
PLINK: rising to the challenge of larger and richer datasets. Gigascience. 2015;4:7.

136. McKennan C, Naughton K, Stanhope C, Kattan M, O'Connor GT, Sandel MT, et al.
Longitudinal data reveal strong genetic and weak non-genetic components of ethnicity-
dependent blood DNA methylation levels. Epigenetics. 2020. doi:
10.1080/15592294.2020.1817290:1-15.

137. Jun G, Flickinger M, Hetrick KN, Romm JM, Doheny KF, Abecasis GR, et al. Detecting
and estimating contamination of human DNA samples in sequencing and array-based
genotype data. Am J Hum Genet. 2012;91(5):839-48.

138. Hao W, Storey JD. Extending Tests of Hardy-Weinberg Equilibrium to Structured
Populations. Genetics. 2019;213(3):759-70.

139. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011;27(15):2156-8.

140. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
SAMtools and BCFtools. Gigascience. 2021;10(2).

141. Genomes Project C, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, et al. A
global reference for human genetic variation. Nature. 2015;526(7571):68-74.

142. Bergstrom A, McCarthy SA, Hui R, Almarri MA, Ayub Q, Danecek P, et al. Insights into
human genetic variation and population history from 929 diverse genomes. Science.
2020;367(6484).

143. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res. 2009;19(9):1655-64.

144. Conomos MP, Miller MB, Thornton TA. Robust inference of population structure for
ancestry prediction and correction of stratification in the presence of relatedness. Genet
Epidemiol. 2015;39(4):276-93.

145. Conomos MP, Reiner AP, Weir BS, Thornton TA. Model-free Estimation of Recent
Genetic Relatedness. Am J Hum Genet. 2016;98(1):127-48.

146. Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. Robust relationship
inference in genome-wide association studies. Bioinformatics. 2010;26(22):2867-73.

147. Shringarpure SS, Bustamante CD, Lange K, Alexander DH. Efficient analysis of large
datasets and sex bias with ADMIXTURE. BMC Bioinformatics. 2016;17:218.

148. Sofer T, Zheng X, Gogarten SM, Laurie CA, Grinde K, Shaffer JR, et al. A fully adjusted
two-stage procedure for rank-normalization in genetic association studies. Genet
Epidemiol. 2019;43(3):263-75.

149. McCaw ZR, Lane JM, Saxena R, Redline S, Lin X. Operating characteristics of the rank-
based inverse normal transformation for quantitative trait analysis in genome-wide
association studies. Biometrics. 2020;76(4):1262-72.

150. Wang G, Sarkar A, Carbonetto P, Stephens M. A simple new approach to variable selection
in regression, with application to genetic fine mapping. Journal of the Royal Statistical
Society: Series B (Statistical Methodology). 2020;82(5):1273-300.

151. Shabalin AA. Matrix eQTL: ultra fast eQTL analysis via large matrix operations.
Bioinformatics. 2012;28(10):1353-8.

152. McKennan C, Nicolae D. Accounting for unobserved covariates with varying degrees of
1121   estimability in high-dimensional biological data. Biometrika. 2019;106(4):823-40.

1122   153. Zhang D. A Coefficient of Determination for Generalized Linear Models. The American
1123   Statistician. 2017;71(4):310-6.

1124   154. Fox J, Kleiber C, Zeileis A. ivreg: Two-Stage Least-Squares Regression with Diagnostics.
1125   https://john-d-fox.github.io/ivreg/2021.

1126   155. Burgess S, Small DS, Thompson SG. A review of instrumental variable estimators for
1127   Mendelian randomization. Stat Methods Med Res. 2017;26(5):2333-55.

1128   156. Burgess S, Thompson SG. Use of allele scores as instrumental variables for Mendelian
1129   randomization. International Journal of Epidemiology. 2013;42(4):1134-44.

1130   157. Alfons A, Ateş NY, Groenen PJF. A Robust Bootstrap Test for Mediation Analysis.
1131   Organizational Research Methods.0(0):1094428121999096.

1132   158. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
1133   universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.

1134   159. Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model
1135   analysis tools for RNA-seq read counts. Genome Biol. 2014;15(2):R29.

1136   160. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential
1137   expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.
1138   2015;43(7):e47.

1139   161. Jager R, Migliorini G, Henrion M, Kandaswamy R, Speedy HE, Heindl A, et al. Capture
1140   Hi-C identifies the chromatin interactome of colorectal cancer risk loci. Nat Commun.
1141   2015;6:6178.

1142   162. Mifsud B, Tavares-Cadete F, Young AN, Sugar R, Schoenfelder S, Ferreira L, et al.
1143   Mapping long-range promoter contacts in human cells with high-resolution capture Hi-C.
1144   Nat Genet. 2015;47(6):598-606.

1145   163. Montefiori LE, Sobreira DR, Sakabe NJ, Aneas I, Joslin AC, Hansen GT, et al. A promoter
1146   interaction map for cardiovascular disease genetics. Elife. 2018;7.

1147   164. Cairns J, Freire-Pritchett P, Wingett SW, Varnai C, Dimond A, Plagnol V, et al.
1148   CHiCAGO: robust detection of DNA looping interactions in Capture Hi-C data. Genome
1149   Biol. 2016;17(1):127.
1150

1151 **SUPPORTING INFORMATION**

1152

1153 **S1 Fig. Whole-genome sequencing depth and coverage. A)** Histogram of 1,035 whole-
1154 genome sequencing (WGS) samples from APIC and URECA by mean depth of coverage. **B)**
1155 Histogram of WGS samples based on proportion of genome covered at 20x, 25x, and 30x
1156 depth. APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment and
1157 Childhood Asthma study.

1158

1159

1160 **S1 Table. Post-QC sequencing call concordance between replicates.** Variant call
1161 concordance between three pairs of replicate samples, by variant type and cohort allele
1162 frequency. SNPs, single nucleotide polymorphisms; MAF, minor allele frequency; InDels,
1163 insertions and deletions.

1164

1165

S2 Table. FEV_1-associated variants in chr14q32.33. All variants in chr14q32.33 associated
with FEV_1 (% predicted) with $p<1\times10^{-5}$ (n=82) in GWAS of 896 participants from APIC &
URECA. N, number of genotyped individuals. MAF, minor allele frequency; 95% CI, 95%
confidence interval; SE, standard error; P, P-value (Wald); FEV_1, forced expiratory volume in
one second; APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment
and Childhood Asthma study.

1172

1173

1174 **S2 Fig**. **Distribution of lung function measures by study. A)** Distribution of FEV_1 (%
1175 predicted) in APIC and URECA. **B)** Distribution of FEV_1/FVC in APIC and URECA. APIC,
1176 Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma
1177 study. FEV_1, forced expiratory volume in one second; FVC, forced vital capacity.

1178

1179

1180 **S3 Fig. P-value distributions of GWAS results.** Quantile-quantile plots of the GWAS results
1181 with corresponding genomic control factors (lambda) are shown for A) FEV_1 (% predicted) and
1182 B) FEV_1/FVC. FEV_1, forced expiratory volume in one second; FVC, forced vital capacity.

1183

1184

1185 **S4 Fig. Fine-mapping results for FEV_1 (% predicted) at the *TDRD9* locus.** The X-axis shows
1186 the chromosome position on chromosome 14. The Y-axis is the posterior inclusion probability
1187 (PIP). Variants highlighted in red represent a credible set, in which there is a 95% probability
1188 that at least one of the variants is causal. FEV_1, forced expiratory volume in one second.

1189

1190

1191 **S5 Fig. Genome-wide association results without adjustment for asthma.** GWAS
1192 Manhattan plots for **A**) FEV_1 and **B**) FEV_1/FVC ratio, without adjustment for asthma status. The
1193 horizontal red line indicates genome-wide significance ($p \leq 2.5\times10^{-8}$). The dotted horizontal blue

1194   line indicates $p=1 \times 10^{-5}$. Variants colored in grey are the GWAS results with asthma adjustment.
1195 FEV_1, forced expiratory volume in one second; FEV_1/FVC, ratio of FEV_1 to forced vital capacity.
1196
1197
1198   **S6 Fig. Replication of FEV_1 GWAS SNPs.** Association statistics for previously identified FEV_1
1199   GWAS SNPs [23]. 64 out of 70 previously identified SNPs were genotyped in APIC & URECA.
1200   GWAS, genome-wide association study; SNP, single nucleotide polymorphism; APIC, Asthma
1201   Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma study.
1202 FEV_1, forced expiratory volume in one second.
1203
1204
S7 Fig. Replication of FEV_1/FVC GWAS SNPs. Association statistics for previously identified
FEV_1/FVC GWAS SNPs [23]. 112 out of 117 previously identified SNPs were genotyped in
APIC & URECA. GWAS, genome-wide association study; SNP, single nucleotide
polymorphism; APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment
and Childhood Asthma study. FEV_1, forced expiratory volume in one second; FVC, forced vital
capacity.
1211
1212
S3 Table. MeQTL analysis results and associations with FEV_1. All CpG sites where DNA
methylation levels in NECs at age 11 in URECA were associated with rs10220464 at FDR<0.05
are shown with their corresponding associations with FEV_1. The FDR-adjusted P-values (FDR
Q) correspond to a 5% false-discovery rate. FDR, false discovery rate; 95% CI, 95% confidence
interval; FEV_1, forced expiratory volume in one second; URECA, Urban Environment and
Childhood Asthma study.
1219
1220
1221   **S8 Fig. NicAlert Results by Study.** Distribution of urine cotinine levels, as measured using
1222   NicAlert immunochromatographic assays, which report results on a scale of 0-6 according to the
1223   labeled concentration ranges. Proportions were calculated relative to the number of samples
1224   with available NicAlert results. APIC, Asthma Phenotypes in the Inner City study; URECA,
1225   Urban Environment and Childhood Asthma study.
1226
1227
S9 Fig. DNA methylation at cg03306306 by smoking exposure. DNA methylation levels at
cg03306306 are shown by rs10220464 genotype in URECA participants with low and high
smoking exposures in (**A**) NECs at age 11 and (**B**) PBMCs at age 7. FEV_1 (% predicted) are
also shown by cg03306306 DNA methylation levels in URECA participants with low and high
smoking exposures in (**C**) NECs at age 11 and (**D**) PBMCs at age 7. NECs, nasal epithelial
cells; PBMCs, peripheral blood mononuclear cells; FEV_1, forced expiratory volume in one
second; URECA, Urban Environment and Childhood Asthma study.
1235
1236

1237 **S10 Fig. Genotype associations with FEV$_1$ by smoking exposure.** FEV$_1$ (% predicted) are
1238 shown by rs10220464 genotype in APIC & URECA participants with low and high smoking
1239 exposures according to urine cotinine levels. FEV$_1$, forced expiratory volume in one second;
1240 APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood
1241 Asthma study.
1242
1243
S4 Table. **rs10220464 eQTL analysis results.** Results of eQTL analyses in NECs and PBMCs
with rs10220464 for all genes within 1 Mb in URECA. Gene expression was measured in counts
per million mapped reads. The FDR-adjusted P-values (FDR Q) correspond to a 5% false-
discovery rate. FDR, false discovery rate; 95% CI, 95% confidence interval; NECs, nasal
epithelial cells; PBMCs, peripheral blood mononuclear cells; URECA, Urban Environment and
Childhood Asthma study.
1250
1251
1252 **S11 Fig. *PPP1R13B* expression in NECs vs. smoking exposure, FEV$_1$.** *PPP1R13B*
1253 expression in NECs at age 11 was not associated with smoking exposure at age 10 (**A**) nor with
1254 FEV$_1$ (% predicted) at age 10 (**B**) in URECA. NECs, nasal epithelial cells; FEV$_1$, forced
1255 expiratory volume in one second; URECA, Urban Environment and Childhood Asthma.
1256
1257
1258 **S5 Table. Chromatin interactions with FEV$_1$-associated SNPs.** Bait and target fragments
1259 refer to mapped Hi-C restriction fragments on chr14 (hg38) for gene promoters and putative
1260 enhancers, respectively. FEV$_1$ SNPs refer to number of FEV$_1$-associated variants (p<1x10^{-5})
1261 within 1kb of target fragment. SNPs, single nucleotide polymorphisms; FEV$_1$, forced expiratory
1262 volume in one second.
1263
1264
1265 **S6 Table. Age at used lung function measure in URECA.** URECA, Urban Environment and
1266 Childhood Asthma study; FEV$_1$, forced expiratory volume in one second; FVC, forced vital
1267 capacity.
1268
1269
1270 **S7 Table. Study samples.** APIC, Asthma Phenotypes in the Inner City study; URECA, Urban
1271 Environment and Childhood Asthma study; WGS, whole-genome sequencing; NECs, nasal
1272 epithelial cells; PBMCs, peripheral blood mononuclear cells.
1273
1274
1275 **S12 Fig. Data availability across APIC and URECA.** Data availability for measures used in
1276 this study are shown for all sequenced samples. Each row represents a pattern of available and
1277 missing data, with green squares indicating available data and grey squares indicating missing
1278 data. Total counts of available data points for each variable are listed across the top of the
1279 figure. Total counts for each data availability pattern are listed along the right.
1280
1281

S13 Fig. Transitions/transversions vs. quality/depth in WGS variant calls. The
transition/transversion ratio (TS/TV) is plotted against the variant call quality/depth metric (QD)
across all WGS SNP calls in APIC & URECA. Sites with QD less than 4 or greater than 34 were
removed from consideration in this study. SNPs, single nucleotide polymorphisms; WGS, whole-
genome sequencing; APIC, Asthma Phenotypes in the Inner City study; URECA, Urban
Environment and Childhood Asthma study.
1288
1289
1290 **S14 Fig. Intercorrelation of Mendelian randomization candidate instrument SNPs in**
1291 **URECA.** Instrumental variables were chosen from a set of candidate SNPs that were at least
1292 nominally associated with cg03306306 methylation with $p<0.15$. The correlation values between
1293 these SNPs are shown, clustered using Ward's method. The four SNPs used for the instrument
1294 are highlighted. URECA, Urban Environment and Childhood Asthma.
1295
1296
1297 **S8 Table. Additional phenotypic, socioeconomic, and environmental data.** Additional
1298 variables examined for potential confounding in mediation analyses for APIC & URECA. APIC,
1299 Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma
1300 study.
1301
1302
S15 Fig. Intercorrelation of phenotypes and environmental variables in APIC & URECA.
The correlations are shown between FEV_1 (% predicted), smoking exposure (NicAlert), the
primary the lead FEV_1 SNP rs10220464, DNA methylation at cg03306306, 11 environmental
exposures, and 2 socioeconomic indicators, clustered using Ward's method. APIC, Asthma
Phenotypes in the Inner City study; exp., exposure; URECA, Urban Environment and Childhood
Asthma.

Cohort	rs10220464 vs. FEV$_1$	N	MAF	β$_z$ [95% CI]	P
All		896	0.295	-0.31 [-0.41- -0.21]	2.42×10^{-9}
Black (non-Hispanic)		606	0.275	-0.27 [-0.40- -0.15]	1.88×10^{-5}
Hispanic		224	0.331	-0.34 [-0.53- -0.15]	7.15×10^{-4}
White (non-Hispanic)		9	0.464	-0.68 [-1.72- 0.37]	4.26×10^{-1}
APIC		504	0.331	-0.30 [-0.43- -0.16]	2.23×10^{-5}
URECA		392	0.260	-0.23 [-0.37- -0.09]	1.46×10^{-3}
Asthmatic		677	0.305	-0.31 [-0.43- -0.20]	1.62×10^{-7}
Non-asthmatic		219	0.275	-0.23 [-0.41- -0.05]	1.38×10^{-2}

β$_z$

NECs

PBMCs

S1 Table. Post-QC sequencing call concordance between replicates.

Variant Type	Variants (mean)	Call concordance
All Variants	4,284,228	0.9985
SNPs	3,793,694	0.9997
Common SNPs (MAF > 0.05)	3,244,661	0.9997
Low Frequency SNPs (0.01 > MAF ≤ 0.05)	383,918	0.9995
Rare SNPs (MAF ≤0.01)	165,115	0.9995
InDels	490,534	0.9892
Common InDels (MAF > 0.05)	428,621	0.9896
Low Frequency InDels (0.01 > MAF ≤ 0.05)	39,572	0.9845
Rare InDels (MAF ≤0.01)	22,340	0.9911

Variant call concordance between three pairs of replicate samples, by variant type and cohort allele frequency. SNPs, single nucleotide polymorphisms; MAF, minor allele frequency; InDels, insertions and deletions.

S2 Table. FEV$_1$-associated variants in chr14q32.33

Pos	rsID	N	Ref	Alt	MAF	Beta	SE	P	Type	Gene
103936818	**rs10220464**	**896**	**A**	**G**	**0.30**	**-0.311**	**0.052**	**2.42 x10^{-9}**	**intronic**	***TDRD9***
103926863	rs11850186	896	G	A	0.29	-0.307	0.052	5.34 x10^{-9}	intergenic	*ATP5MPL; TDRD9*
103934654	rs4906387	896	C	T	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103935736	rs9285602	896	C	T	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103937635	rs9324069	896	G	A	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103942966	.	896	GA	G	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103960917	rs4900604	896	G	A	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103963565	rs28574832	896	G	T	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103967867	rs4900605	896	G	T	0.29	-0.305	0.052	6.20 x10^{-9}	intronic	*TDRD9*
103997188	rs72712900	896	G	A	0.29	-0.304	0.052	6.51 x10^{-9}	intronic	*TDRD9*
103984806	rs11847797	896	G	A	0.29	-0.303	0.052	7.73 x10^{-9}	intronic	*TDRD9*
103980079	rs58289480	896	A	G	0.29	-0.303	0.052	8.28 x10^{-9}	intronic	*TDRD9*
103985095	rs4906396	896	A	C	0.29	-0.303	0.052	8.28 x10^{-9}	intronic	*TDRD9*
103953683	rs72712884	896	C	G	0.29	-0.303	0.052	8.28 x10^{-9}	intronic	*TDRD9*
104046654	rs143594477	896	C	CT	0.29	-0.302	0.052	9.12 x10^{-9}	intronic	*TDRD9*
104028226	rs10137997	896	T	C	0.29	-0.301	0.052	9.73 x10^{-9}	intronic	*TDRD9*
104034493	rs9944163	896	G	T	0.29	-0.301	0.052	9.73 x10^{-9}	intronic	*TDRD9*
104069368	rs72714937	896	C	T	0.29	-0.302	0.052	9.80 x10^{-9}	intergenic	*TDRD9; ASPG*
104059833	rs11160785	896	T	A	0.29	-0.300	0.052	1.03 x10^{-8}	intergenic	*TDRD9; ASPG*
103934222	rs28461376	896	A	G	0.29	-0.301	0.052	1.06 x10^{-8}	intronic	*TDRD9*
103963484	.	896	TC	T	0.29	-0.301	0.052	1.06 x10^{-8}	intronic	*TDRD9*
104037291	rs4906408	896	G	T	0.29	-0.299	0.052	1.14 x10^{-8}	intronic	*TDRD9*
104030861	rs10135338	896	G	A	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104035861	rs10134394	896	G	A	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104036696	rs11160780	896	G	A	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104037342	rs4900608	896	G	A	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104038178	rs4906409	896	T	C	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104039412	rs10144682	896	G	A	0.29	-0.299	0.052	1.18 x10^{-8}	intronic	*TDRD9*
104012687	rs11851441	896	A	G	0.29	-0.299	0.052	1.23 x10^{-8}	intronic	*TDRD9*
104031206	rs10135507	896	C	T	0.29	-0.298	0.052	1.25 x10^{-8}	exonic	*TDRD9*
103998027	rs55938939	896	G	A	0.29	-0.299	0.052	1.26 x10^{-8}	intronic	*TDRD9*
104048481	rs61244535	896	C	G	0.29	-0.299	0.052	1.29 x10^{-8}	intronic	*TDRD9*
104051127	rs28644198	896	C	A	0.29	-0.299	0.052	1.29 x10^{-8}	intronic	*TDRD9*
104054541	rs28377615	896	G	A	0.29	-0.299	0.052	1.29 x10^{-8}	intergenic	*TDRD9; ASPG*
103997438	rs4906397	896	C	T	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
103999081	rs4906399	896	C	T	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
104005628	rs12147936	896	C	T	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
104010259	rs145867870	896	C	CT	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
104013496	rs11851723	896	C	T	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
104014746	rs11160779	896	T	C	0.29	-0.298	0.052	1.31 x10^{-8}	exonic	*TDRD9*
104016537	rs72714909	896	C	A	0.29	-0.298	0.052	1.31 x10^{-8}	intronic	*TDRD9*
103986068	rs10143030	896	C	G	0.29	-0.300	0.052	1.41 x10^{-8}	intronic	*TDRD9*
103986303	rs10143389	896	G	A	0.29	-0.300	0.052	1.41 x10^{-8}	exonic	*TDRD9*
103986825	rs61248168	896	T	A	0.29	-0.300	0.052	1.41 x10^{-8}	intronic	*TDRD9*
103990858	rs1957518	896	G	A	0.29	-0.300	0.052	1.41 x10^{-8}	intronic	*TDRD9*
104072329	rs111576189	896	G	A	0.29	-0.298	0.052	1.54 x10^{-8}	intergenic	*TDRD9; ASPG*
104011541	rs28522352	896	T	A	0.29	-0.296	0.052	1.79 x10^{-8}	intronic	*TDRD9*
104003975	rs8010286	896	T	C	0.29	-0.293	0.052	2.06 x10^{-8}	intronic	*TDRD9*
104004349	rs28391043	896	G	A	0.29	-0.294	0.052	2.22 x10^{-8}	intronic	*TDRD9*
104027784	rs28725314	896	C	T	0.29	-0.290	0.052	2.96 x10^{-8}	intronic	*TDRD9*
103970594	rs11851097	896	C	T	0.36	-0.267	0.048	3.28 x10^{-8}	exonic	*TDRD9*

Pos	rsID	N	Ref	Alt	MAF	Beta	SE	P	Type	Gene
103987132	rs200735321	880	A	G	0.26	-0.311	0.057	6.98×10^{-8}	intronic	*TDRD9*
104067149	rs10782497	896	T	C	0.39	-0.265	0.049	1.08×10^{-7}	intergenic	*TDRD9; ASPG*
104019632	rs74089113	896	G	C	0.44	-0.246	0.046	1.25×10^{-7}	intronic	*TDRD9*
104071795	rs35775205	896	A	G	0.37	-0.264	0.050	1.72×10^{-7}	intergenic	*TDRD9; ASPG*
104038358	rs877009	896	T	C	0.43	-0.240	0.046	3.06×10^{-7}	intronic	*TDRD9*
104007407	rs12100528	896	G	A	0.43	-0.240	0.047	3.32×10^{-7}	intronic	*TDRD9*
104016337	rs7158223	896	G	T	0.42	-0.238	0.046	3.83×10^{-7}	intronic	*TDRD9*
103947548	rs7144813	896	G	C	0.35	-0.244	0.049	8.48×10^{-7}	intronic	*TDRD9*
103937851	rs10141985	896	G	A	0.34	-0.242	0.050	1.53×10^{-6}	intronic	*TDRD9*
103939314	rs7148877	896	A	G	0.34	-0.241	0.050	1.73×10^{-6}	intronic	*TDRD9*
103945800	rs12147460	896	T	G	0.34	-0.241	0.050	1.73×10^{-6}	intronic	*TDRD9*
104032500	rs9944058	896	C	T	0.42	-0.235	0.049	2.22×10^{-6}	intronic	*TDRD9*
103981884	rs4906395	896	T	C	0.42	-0.220	0.047	2.71×10^{-6}	intronic	*TDRD9*
103958246	rs10149734	896	A	C	0.41	-0.221	0.047	3.04×10^{-6}	intronic	*TDRD9*
103962412	rs72712889	896	T	C	0.41	-0.221	0.047	3.04×10^{-6}	intronic	*TDRD9*
103967964	rs4906392	896	T	C	0.41	-0.221	0.047	3.04×10^{-6}	intronic	*TDRD9*
103995217	rs10132153	896	C	A	0.41	-0.220	0.047	3.16×10^{-6}	intronic	*TDRD9*
104036896	rs7156122	896	G	A	0.41	-0.217	0.046	3.28×10^{-6}	intronic	*TDRD9*
104014863	rs397949797	896	A	AT	0.41	-0.216	0.046	3.75×10^{-6}	intronic	*TDRD9*
103979872	rs8022833	896	A	G	0.41	-0.219	0.047	3.87×10^{-6}	intronic	*TDRD9*
103981060	rs1957517	896	A	G	0.41	-0.219	0.047	3.87×10^{-6}	intronic	*TDRD9*
103995157	rs10132127	896	C	G	0.41	-0.218	0.047	4.02×10^{-6}	intronic	*TDRD9*
104004783	rs9671921	896	A	G	0.49	-0.214	0.046	4.98×10^{-6}	intronic	*TDRD9*
104008697	rs4906401	896	A	G	0.41	-0.215	0.047	5.23×10^{-6}	intronic	*TDRD9*
104001342	rs61319604	896	C	T	0.41	-0.215	0.047	5.35×10^{-6}	intronic	*TDRD9*
104002208	rs72714904	896	T	C	0.41	-0.215	0.047	5.35×10^{-6}	intronic	*TDRD9*
104003206	rs10135296	896	G	T	0.41	-0.215	0.047	5.35×10^{-6}	intronic	*TDRD9*
104006916	rs4900607	896	A	G	0.49	-0.212	0.046	5.90×10^{-6}	intronic	*TDRD9*
104010388	rs4906402	896	G	A	0.41	-0.213	0.047	6.22×10^{-6}	intronic	*TDRD9*
104000834	rs7160557	896	G	A	0.48	-0.207	0.046	8.97×10^{-6}	intronic	*TDRD9*
104078948	rs4906415	896	G	A	0.19	-0.268	0.060	9.00×10^{-6}	intergenic	*TDRD9; ASPG*

All variants in chr14q32.33 associated with FEV$_1$ (% predicted) with $p<1\times10^{-5}$ (n=82) in GWAS of 896 participants from APIC & URECA. N, number of genotyped individuals. MAF, minor allele frequency; 95% CI, 95% confidence interval; SE, standard error; P, P-value (Wald); FEV$_1$, forced expiratory volume in one second; APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma study.

S3 Table. **MeQTL analysis results and associations with FEV$_1$**

CpG Site	Position (strand), hg38 (chr14)	rs10220464			FEV$_1$	
		Beta [95% CI]	P	FDR Q	Beta [95% CI]	P
cg21567958	103909862 (-)	0.10 [0.05, 0.14]	1.79 x10^{-5}	0.01	-1.5 [-8.8, 5.7]	0.68
cg12183467	103885907 (-)	0.10 [0.05, 0.14]	1.31 x10^{-4}	0.04	5.3 [-1.5, 12.1]	0.13
cg16820107	103970549 (-)	0.08 [0.04, 0.12]	1.62 x10^{-4}	0.04	-5.4 [-13.4, 2.5]	0.18
cg03306306	**104001397 (+)**	**0.07 [0.03, 0.11]**	**2.28 x10^{-4}**	**0.04**	**-11.5 [-20.3, -2.7]**	**0.01**
cg17298714	103938355 (+)	0.14 [0.07, 0.21]	2.36x10^{-4}	0.04	-4.1 [-8.7, 0.4]	0.07

All CpG sites where DNA methylation levels in NECs at age 11 in URECA were associated with rs10220464 at FDR<0.05 are shown with their corresponding associations with FEV$_1$. The FDR-adjusted P-values (FDR Q) correspond to a 5% false-discovery rate. FDR, false discovery rate; 95% CI, 95% confidence interval; FEV$_1$, forced expiratory volume in one second; URECA, Urban Environment and Childhood Asthma study.

S4 Table. **rs10220464 eQTL analysis results**

Gene	NECs			PBMCs	
	Beta [95% CI]	P	FDR Q	Beta [95% CI]	P
PPP1R13B	**0.12 [0.06, 0.17]**	**1.26×10^{-5}**	**2.77×10^{-4}**	0.01 [-0.08, 0.09]	0.87
BAG5	-0.06 [-0.02, -0.10]	0.0054	0.059	0.03 [-0.03, 0.08]	0.31
ASPG	0.12 [-0.02, 0.26]	0.089	0.48	NOT EXPRESSED	
CKB	-0.04 [0.01, -0.09]	0.10	0.48	0.13 [-0.03, 0.30]	0.12
KLC1	-0.04 [0.01, -0.08]	0.12	0.48	0.02 [-0.04, 0.07]	0.53
TNFAIP2	-0.05 [0.02, -0.12]	0.15	0.48	-0.11 [0.06, -0.28]	0.22
MARK3	-0.03 [0.01, -0.07]	0.15	0.48	-0.02 [0.02, -0.06]	0.40
C14orf2	0.01 [-0.01, 0.03]	0.23	0.61	-0.03 [0.03, -0.09]	0.35
XRCC3	0.08 [-0.06, 0.22]	0.25	0.61	0.00 [0.06, -0.07]	0.92
ZFYVE21	0.02 [-0.01, 0.04]	0.28	0.61	-0.02 [0.03, -0.07]	0.49
ADSSL1	0.04 [-0.04, 0.11]	0.33	0.61	0.04 [-0.09, 0.17]	0.54
PLD4	0.07 [-0.07, 0.20]	0.33	0.61	0.07 [-0.12, 0.26]	0.47
SIVA1	-0.01 [0.01, -0.03]	0.36	0.61	0.01 [-0.06, 0.09]	0.77
EXOC3L4	-0.08 [0.14, -0.29]	0.47	0.69	NOT EXPRESSED	
TRMT61A	-0.02 [0.04, -0.09]	0.48	0.69	-0.04 [0.04, -0.12]	0.32
CEP170B	-0.02 [0.05, -0.09]	0.51	0.69	0.02 [-0.11, 0.14]	0.78
APOPT1	-0.02 [0.04, -0.08]	0.56	0.69	-0.03 [0.03, -0.09]	0.37
EIF5	-0.01 [0.02, -0.04]	0.60	0.69	-0.03 [0.05, -0.12]	0.44
TDRD9	-0.07 [0.21, -0.36]	0.60	0.69	0.01 [-0.23, 0.26]	0.91
ZBTB42	-0.02 [0.05, -0.08]	0.63	0.69	-0.09 [0.01, -0.19]	0.089
INF2	-0.01 [0.08, -0.10]	0.84	0.85	0.04 [-0.06, 0.13]	0.45
AKT1	0.00 [0.03, -0.03]	0.85	0.85	-0.02 [0.04, -0.08]	0.57
AL049840.1	NOT EXPRESSED			0.00 [0.08, -0.09]	0.93
AL049840.4	NOT EXPRESSED			0.01 [-0.05, 0.08]	0.65
AL049840.5	NOT EXPRESSED			0.02 [-0.08, 0.13]	0.68
AL133367.1	NOT EXPRESSED			0.09 [-0.02, 0.20]	0.12
LINC00638	NOT EXPRESSED			0.03 [-0.11, 0.18]	0.66

Results of eQTL analyses in NECs and PBMCs with rs10220464 for all genes within 1 Mb in URECA. Gene expression was measured in counts per million mapped reads. The FDR-adjusted P-values (FDR Q) correspond to a 5% false-discovery rate. FDR, false discovery rate; 95% CI, 95% confidence interval; NECs, nasal epithelial cells; PBMCs, peripheral blood mononuclear cells; URECA, Urban Environment and Childhood Asthma study.

S5 Table. Chromatin interactions with FEV$_1$-associated SNPs

Promoter	Bait Fragment	Target Fragment	Strand	Distance	FEV$_1$ SNPs	CHiCAGO Score
PPP1R13B	103845355-103845955	104003512-104004312	-	157557	4	9.38
CDCA4	105020395-105021827	104010864-104011532	-	1008863	3	7.65
PPP1R13B	103845355-103845955	104004616-104004922	-	158661	4	7.16
ZFYVE21	103716777-103717467	103948470-103949743	+	231003	1	6.57
ZFYVE21	103716777-103717467	103933835-103934479	+	216368	2	6.49
C14orf79	104985639-104987255	103935625-103936182	+	1049457	3	6.10
MARK3	103385080-103385712	103937160-103938502	+	551448	4	5.89
ZFYVE21	103715624-103716777	103981359-103981909	+	264582	2	5.89
PLD4	104927266-104927937	103963380-103964018	+	963248	3	5.69
ZFYVE21	103715624-103716777	103933397-103933835	+	216620	2	5.62
ADSSL1	104720007-104724747	104017055-104017482	+	702525	2	5.52
APOPT1	103562490-103563614	103999031-103999343	+	435417	1	5.48
CKB	103521551-103522109	103997308-103998030	-	475199	3	5.26
MARK3	103385881-103386341	103958731-103959336	+	572390	1	5.19
PPP1R13B	103845355-103845955	104036454-104037811	-	190499	7	5.00

Bait and target fragments refer to mapped Hi-C restriction fragments on chr14 (hg38) for gene promoters and putative enhancers, respectively. FEV$_1$ SNPs refer to number of FEV$_1$-associated variants (p<1x10^{-5}) within 1kb of target fragment. SNPs, single nucleotide polymorphisms; FEV$_1$, forced expiratory volume in one second.

S6 Table. Age at used lung function measure in URECA

Age (years)	Count	
	FEV$_1$	FEV$_1$/FVC
10	382	372
9	26	30
8	10	11
7	8	5
6	6	10
5	9	10

URECA, Urban Environment and Childhood Asthma study; FEV$_1$, forced expiratory volume in one second; FVC, forced vital capacity.

S7 Table. Study samples

Cohort	Sample	Cell Type	Age Collected	N
APIC	WGS	-	-	508
URECA	WGS	-	-	527
URECA	DNA Methylation	NECs	11	286
URECA	DNA Methylation	PBMCs	7	169
URECA	RNA-Seq	NECs	11	324
URECA	RNA-Seq	PBMCs	2	132

APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma study; WGS, whole-genome sequencing; NECs, nasal epithelial cells; PBMCs, peripheral blood mononuclear cells.

S8 Table. Additional phenotypic, socioeconomic, and environmental data

Variable	Description	Units	N
Socioeconomic factors			
Income<$15K	Family yearly income <$15,000	yes/no	1018
Caretaker education	Caretaker completed high school	yes/no	1031
Environmental exposures			
Gas stove at home	Gas stove in home	yes/no	982
Dampness at home	Water problems in home	yes/no	981
AC at home	AC unit in child's bedroom	yes/no	982
Airvents at home	Forced air for heat in home	yes/no	983
Rodent allergen exposure	Mus m 1 (mouse) bedroom allergen levels	log(μg/g)	847
Roach allergen exposure	Bla g 1 (German cockroach) bedroom allergen levels	log(μg/g)	850
Dust mite allergen exposure	Combined Der f 1 (*Dermatophagoides farina*) and Der p 1 (*Dermatophagoides pteronyssinus)* bedroom allergen levels	log(μg/g)	756
Dog allergen exposure	Can f 1 bedroom allergen levels	log(μg/g)	839
Cat allergen exposure	Fel d 1 bedroom allergen levels	log(μg/g)	839
NO$_2$ at home	Indoor nitrogen dioxide levels	log(ppb)	900
Caloric intake	Total caloric intake	log(calories/day)	787

Additional variables examined for potential confounding in mediation analyses for APIC & URECA. APIC, Asthma Phenotypes in the Inner City study; URECA, Urban Environment and Childhood Asthma study.

A

B

≥20x ≥25x ≥30x