**Predicting Hand, Foot, and Mouth Disease in Japan Using Google Trends: Infodemiology**

**Study**

Qian Niu,[1*] MS; Junyu Liu,[2*] ME; Miyu Onishi,[1] BS; Asuka Kawaguchi,[1] BS; Anuradhi Bandara,[1]

BS; Keiko Harada,[1] MS; Tomoki Aoyama,[1] MD, PhD; Momoko Nagai-Tanima,[1] PhD

[1] Department of Human Health Sciences, Graduate School of Medicine, Kyoto University, Kyoto,

Japan

[2] Department of Intelligence Science and Technology, Graduate School of Informatics, Kyoto

University, Kyoto, Japan

* Qian Niu and Junyu Liu contributed equally to this article.

Corresponding Author: Momoko Nagai-Tanima, PhD

Department of Human Health Sciences, Graduate School of Medicine,

Kyoto University, 53, Kawahara-Cho, Shogoin, Sakyo-Ku

Kyoto, 606-8507 Japan

Phone: 075-751-3964

Email: tanima.momoko.8s@kyoto-u.ac.jp

**Abstract**

**Objectives:** We aimed to predict the number of hand, foot, and mouth disease (HFMD) infections

before and during the COVID-19 pandemic using Internet search data.

**Methods:** We obtained actual HFMD cases from the National Institute of Infectious Disease and

Internet search data using Google Trends between 2004 and 2021 in Japan. We calculated the

Pearson correlation coefficients between actual HFMD cases and the search topic "HFMD" from

2004 to 2021. We conducted a cross-correlation analysis between the actual HFMD cases and 43

HFMD-related search terms before and during the pandemic. We identified the most significant

predictors of HFMD infection using stepwise multiple linear regression.

**Results:** We found that actual HFMD cases and Internet search volume peaked around July in most

years, except for 2020 and 2021. The search topic "HFMD" presented a strong correlation with

actual HFMD cases, but the correlation was weaker in 2004, 2008, and 2020. Results from stepwise

multiple linear regression exhibited that the search terms "infect," "daycare," "vomit," "HFMD,"

"eczema," "pain," and "high fever" were the most significant predictors before the pandemic, while

"infect," "enterovirus," "herpangina," "kindergarten," "myocarditis," "HFMD," "contact infection,"

"blister," "high fever" "dermatology," and "plantar" were the most significant predictors during the

pandemic.

**Conclusions:** The predictors for HFMD infections before and during the COVID-19 pandemic were

different. The awareness of HFMD infection in Japan may improved during the COVID-19

pandemic. Continuous monitoring is important to promote public health and prevent resurgence.

Public interest reflected in information-seeking behavior can be helpful for public health surveillance.

**Keywords**: HFMD, Infodemiology, Google Trends, Infection, Prediction

**Introduction**

Hand, foot, and mouth disease (HFMD) is an infectious disease that results in a blistering rash on the mouth, hands, and feet. Most infected individuals recover from HFMD within a few days. Various comorbidities, including myocarditis, neurogenic pulmonary edema, acute flaccid paralysis, and central nervous system complications, such as meningitis, cerebellar ataxia, and encephalitis, can also occur (Huang et al., 2018; Wang et al., 2017). HFMD has been reported in many countries, particularly in the Asia-Pacific region (Puenpa et al., 2019). Corresponding infectious consequences have been reported over the past decade, especially in Japan (Sumi et al., 2017), China (Zhuang et al., 2015; Zheng et al., 2017), and Singapore (Chew et al., 2015). During the summer of 2011, Japan had the largest epidemic of HFMD on record, with 347,362 cases reported (Japan IDWR, 2009). Coxsackievirus A6 (CV-A6) infection was responsible for most cases, with co-circulation of coxsackievirus A16 (CV-A16) and enterovirus A71 (EV-A71) (Fujimoto et al., 2012). EV-A71 has been sporadically detected from October 2014 onward. It became the predominant serotype in 2018, with approximately 70,000 reported cases, following an increased spread from the end of 2017 (Kabele et al., 2022). Since June 2019, a severe outbreak of HFMD has occurred in multiple regions of Japan, attracting the public attention again (IDWR, 2019). As enteroviruses can spread rapidly by droplet and fomite transmission among children in daycare and kindergartens, predicting HFMD outbreaks is vital to public health (Sun et al., 2018).

Rapid recognition and reporting of HFMD infection is essential, and several studies have constructed models for predicting HFMD infection (Rui et al., 2021; Yu et al., 2021; Zhang et al., 2021; Gao et al., 2021). Rui et al. explored epidemiological characteristics and calculated the early warning signals of HFMD using a logistic differential equation (LDE) model in seven regions of China (Rui et al., 2021). Yu et al. forecasted the number of HFMD cases with wavelet-based hybrid models in Zhengzhou, China (Yu et al., 2021). Zhang et al. proposed a landscape dynamic network marker (L-DNM) to detect pre-outbreak signals of HFMD in Tokyo, Hokkaido, and Osaka, Japan (Zhang et al.,

2021). Gao et al. used monthly HFMD infection cases and meteorological data to construct a

weather-based early warning model with a generalized additive model across China (Gao et al.,

2021). The above studies used a range of data, including monthly or weekly HFMD infectious cases

(Rui et al., 2021; Yu et al., 2021; Gao et al., 2021), dynamic information from city networks,

horizontal high-dimensional data, records of clinic visits (Zhang et al., 2021), and meteorological

data (Gao et al., 2021). However, no studies have used Internet search data to predict HFMD

infection. Traditional surveillance and reporting systems lag an outbreak by one to two weeks

because of the reporting and verification process. In Japan, the National Institute of Infectious

Disease (NIID) has monitored the outbreak of various infectious diseases and issued weekly reports

since 1999. However, this traditional type of surveillance uses data from several weeks prior (IDWR

Surveillance Data Table 2022 week, 2022). In contrast, Internet search data shows the information

that the public is searching for in a more real-time manner, which may be valuable for infection

surveillance.

The science of distribution and determinants of information in an electronic medium, specifically the

Internet, or in a population, to inform public health and policy is defined as "infodemiology"

(Eysenbach, 2009). Google Trends is frequently used in infodemiology research to gauge public

interest (Mavragani et al., 2018). Google Trends reflects public information-seeking behavior and

allows users to analyze Internet search data for specific search terms in any country or region over a

selected period (Nuti et al., 2014; Mavragani and Ochoa, 2019). Studies have shown that online

query trends correlate with real-life epidemiologic phenomena such as the flu (Pervaiz et al., 2012),

sinusitis (Sharma et al., 2020), lifestyle-related disease (Memon et al., 2020), asthma (Mavragani et

al., 2018), and pruritus (Tizek et al., 2019). Researchers have also investigated public interest and

information-seeking behavior in chronic obstructive pulmonary disease (COPD) (Boehm et al.,

2019), cancer (Schootman et al., 2015; Phillips et al., 2018), bariatric surgery (Linkov et al., 2014),

kidney stone surgery (Dreher et al., 2018), and suicide (Taira et al., 2021). During the COVID-19

pandemic, similar studies using Internet search data were conducted to predict COVID-19 infectious

cases (Husnayain et al., 2021; Higgins et al., 2020), explore public attitudes toward vaccination

(Pullan and Dey, 2021; Diaz et al., 2021; An et al., 2021), identify symptoms caused by pandemics

(Han et al., 2022; Kardeş et al., 2022; Knipe et al., 2021), and assess affected medical services

(Cohen et al., 2021; Akpan et al., 2021; Adelhoefer et al., 2021). The above studies indicate that

Google Trends could assist in gaining a better understanding and analysis of health information-

seeking behavior. Information from Google Trends could potentially be used to supplement the

current infection reports that have a lag time.

This study aimed to predict HFMD infection using Internet search data in Japan before and during

the COVID-19 pandemic.

**Methods**

**Data**

We obtained actual HFMD cases from the weekly reports issued by NIID, which included new

infectious cases and sentinel cases by prefecture and updated them from 1999 to the present (IDWR

Surveillance Data Table 2022 week, 2022). As for Internet search data, we downloaded the relative

search volume (RSV) of the "HFMD" search topic using Google Trends from January 1, 2004, to

December 31, 2021. The normalized RSV data represented the search interest relative to the highest

point for a given region and time. The scales of normalized RSV varied from 0 to 100, where 0

meant there were insufficient data for a term, while 100 was the peak popularity. We selected a

search topic instead of the search term "HFMD" for comprehensive search information and limited

the period from 2004 to 2021 because Google Trends provides data after 2004. In this study, we used

the search topic "HFMD" and the search term "HFMD". The weekly RSV of the search topic

"HFMD" from 2004 to 2021 was gathered for further analysis. To identify significant predictors of

HFMD infection, we selected 43 search terms including "HFMD" that reflected HFMD-related

symptoms and risk factors. We downloaded the weekly RSV of these 43 search terms in two periods:

before and during the pandemic. As Google Trends can provide data for up to five years on a weekly

basis, we selected the five years before the outbreak (the first case of COVID-19 in Japan was

confirmed on 16 January 2020) as period 1 (from 2015 to 2019) and the two years after the outbreak

as period 2 (2020 and 2021). We set the geographic location to Japan and the category to health to

limit irrelevant results. The word "child" has three common expressions in Japanese, so we divided it

into three search terms "child (kanji)," "child (kana)," and "child (mixed)."

**Statistical Analysis**

SPSS for MAC (version 26.0, IBM Corp.) was used for statistical analysis. Initially, we calculated

the Pearson correlation coefficient between the actual HFMD cases and RSV of the search topic

"HFMD" each year from 2004 to 2021 instead of the whole period due to the periodic characteristic.

Second, we conducted a cross-correlation analysis between the actual HFMD cases and the RSV of

related search terms. Cross-correlation is a measure of the similarity between two series as a function

of the displacement of one relative to the other and was used to objectively estimate the time lag

between cases of HFMD infection and related search terms (Bourke, 1996). We set the maximum lag

to ±20 weeks due to the periodic characteristics of HFMD infection. We obtained 40 cross-

correlation coefficients for each HFMD-related search term before and during the COVID-19

pandemic. Next, we selected the coefficients with the greatest absolute values and exhibited their

true values. Finally, we compared the coefficients with the greatest absolute value in the periods

before and during the COVID-19 pandemic. Regarding these coefficients, we assumed that negative,

zero, or positive values with the greatest absolute value represented the search terms that occurred

earlier, coincided with, or later than the actual HFMD cases, respectively.

Third, we conducted multiple linear regression to identify the most important Internet search terms

for predicting HFMD infection before and during the COVID-19 pandemic. We included HFMD-

related search terms for multiple linear regression as independent variables, with actual HFMD cases

as potential dependent variables. To identify the most significant search terms and limit the number

of independent variables, several common methods were used for independent variable selection,

including forward selection, backward elimination, and stepwise regression (Tranmer et al., 2020).

Stepwise regression alternates between forward and backward, bringing in and removing variables

that meet the criteria for entry or removal until a stable set of variables is attained. We utilized the

stepwise procedure to select the most significant search terms and reduce collinearity with

intercorrelations. Collinearity is the correlation between independent variables that expresses a linear

relationship in a regression model. When the independent variables are correlated in the same

regression model, they cannot independently predict the dependent variable. The variance inflation

factor provides a measure of the degree of collinearity, which is typically less than ten in the medical

field. We excluded terms with a variance inflation factor of > 10. A two-tailed P-value of <.05 was

considered statistically significant.

To assess the performance of the linear regression model, the coefficients of determination $R^2$ or

adjusted $R^2$, which indicates how much variation in response is explained by the model, are often

used (Akossou et al., 2013). The higher the $R^2$, the better the model fits to the data. To avoid

overfitting the model, we selected the adjusted $R^2$ value for model evaluation.

**Results**

**Basic description of the actual HFMD cases and search topic "HFMD"**

In **Figure 1**, we present the actual HFMD and RSV cases from 2004 to 2021. Visual inspection of

the figure indicated that both the actual HFMD cases and RSV of Google Trends peaked around July

in most years except for 2020 and 2021. The number of HFMD infections surged after 2011, peaking

every two years before 2020. The RSV coincided with this trend. In 2020, no periodic peak of

infection was observed, whereas, in 2021, the peak was delayed to November.

**Figure 1.** Monthly number and RSV of HFMD from 2004 to 2021

As shown in **Figure 2**, we calculated the correlation between the actual HFMD cases and RSV each

year from 2004 to 2021. The mean (standard error) of these correlations was 0.820 (0.052). Most of

the correlation values were greater than 0.7, except for 0.562, 0.238, and 0.338 in 2004, 2008, and

2020, respectively.

**Figure 2.** Correlation between actual HFMD cases and HFMD's Interest search RSV from 2004 to
2021

**Temporal correlation between the actual HFMD cases and Internet search terms before and during the pandemic**

We performed a cross-correlation analysis to determine the temporal relationship between the actual HFMD cases and HFMD-related search terms. Cross-correlation results before and during the pandemic are presented in **Table 1**. The cross-correlation coefficient with the greatest absolute value was selected, and its actual value and corresponding lag were represented.

In period 1, search terms including "infect," "coxsackie," "nursery," "mouth," "child (mixed)," "cerebellar ataxia," "acute flaccid paralysis," "infection control," "palm," "hand-foot," "excrement," "eczema," "pain," "group infection," "instep," "fecal-oral transmission," and "rash" presented one or more weeks earlier than actual HFMD cases. Search terms including "herpangina," "summer infection," "HFMD," and "blisters" coincided with the actual HFMD cases. Search terms including "virus," "entero," "enterovirus," "child (kana)," "infants," "daycare," "vomit," "child (kanji)," "pediatrics," "kindergarten," "myocarditis," "handwashing," "contact infection," "high fever," "meningitis," "droplet infection," "headache," "plantar," "encephalitis," "neurogenic pulmonary edema," "dermatology," and "fever" presented one or more weeks later than actual HFMD cases.

In period 2, search terms including "infect," "coxsackie," "nursery," "child (mixed)," "acute flaccid paralysis," "palm," "excrement," "eczema," "pain," "instep," "fecal-oral transmission," "rash," "herpangina," "blister," "virus," "entero," "enterovirus," "child (kana)," "daycare," "child (kanji)" "pediatrics," "myocarditis," "handwashing," "high fever," "meningitis," "headache" "plantar," "encephalitis," "neurogenic pulmonary edema," "dermatology," and "fever" presented one or more weeks earlier than actual HFMD cases. Search terms including "mouth," "hand-foot," and "HFMD" coincided with the actual HFMD cases. The search terms "cerebellar ataxia," "infection control," "group infection," "summer infection," "infants," "vomit," "kindergarten," "contact infection," and "droplet infection" presented one or more weeks later than the actual HFMD cases.

Compared with period 1, the temporal correlation between the actual HFMD cases and Internet

search terms changed in period 2. The search terms "herpangina," "summer infection," "HFMD,"

and "blisters" coincided with actual HFMD cases in period 1. However, in period 2, the search terms

"HFMD," "mouth," and "hand-foot" coincided, while "herpangina" and "blister" were used earlier

and "summer infection" later. In period 1, 17 (37.8%) search terms presented earlier than the HFMD

cases, and 22 (48.9%) search terms presented later. In period 2, 31 (68.9%) search terms presented

earlier than the HFMD cases, and only nine (20%) search terms presented later. Furthermore, we

compared the coefficients with the greatest absolute value in the periods before and during the

COVID-19 pandemic. 26 (67.4%) search terms were searched for earlier in period 2 than in period 1,

might indicating higher public awareness of HFMD infections as well as COVID-19.

**Table 1.** Cross-correlation analysis of the actual hand, foot, and mouth disease cases and Internet

search terms

| Search terms | Period 1 (2015-2019) | | Period 2 (2020-2021) | |
|---|---|---|---|---|
| | Lags (week) | Cross-correlation (r) | Lags (week) | Cross-correlation (r) |
| **Earlier** | | | | |
| infect | -14 | -0.171 | -12 | 0.303 |
| coxsackie* | -1 | 0.349 | -5 | 0.290 |
| nursery* | -5 | 0.256 | -15 | 0.407 |
| mouth | -1 | 0.971 | 0 | 0.833 |
| child (mixed) | -16 | -0.221 | -12 | 0.631 |
| cerebellar ataxia | -6 | 0.124 | 4 | 0.209 |
| acute flaccid paralysis | -15 | 0.070 | -13 | 0.144 |
| infection control | -17 | 0.256 | 2(5)[†] | 0.115 |

| | | | | |
|---|---|---|---|---|
| palm* | -7 | 0.425 | -17 | 0.403 |
| hand-foot | -1 | 0.978 | 0 | 0.835 |
| excrement* | -12 | -0.124 | -13 | 0.100 |
| eczema* | -1 | 0.488 | -13 | 0.505 |
| pain* | -9 | 0.232 | -10 | 0.277 |
| group infection | -6(-7) [†] | -0.232 | 1 | -0.301 |
| instep | -7 | 0.261 | -2 | 0.254 |
| fecal-oral transmission* | -2 | 0.197 | -20 | -0.128 |
| rash* | -1 | 0.797 | -14 | 0.462 |
| **Coincide** | | | | |
| herpangina* | 0 | 0.641 | -4 | 0.717 |
| summer infection | 0 | 0.534 | 6 | -0.173 |
| hand, foot, and mouth disease | 0 | 0.977 | 0 | 0.922 |
| blister* | 0 | 0.695 | -6(-7) [†] | 0.689 |
| **Later** | | | | |
| virus* | 2 | -0.187 | -1(-2) [†] | 0.287 |
| entero* | 11 | 0.217 | -3 | 0.302 |
| enterovirus* | 14 | 0.201 | -7 | 0.088 |
| child (kana)* | 14 | 0.375 | -6 | 0.669 |
| infants* | 7 | -0.154 | 5 | 0.259 |
| daycare* | 13 | -0.088 | -16 | 0.277 |
| vomit | 2 | -0.399 | 9 | 0.229 |
| child(kanji)* | 4 | -0.142 | -15 | 0.555 |
| pediatrics* | 15 | 0.389 | -7 | 0.640 |

| | | | | |
|---|---|---|---|---|
| kindergarten | 2 | -0.252 | 6 | -0.214 |
| myocarditis* | 17 | -0.188 | -6 | 0.408 |
| handwashing* | 15 | 0.300 | -8 | 0.248 |
| contact infection* | 20 | 0.256 | 2(3)[†] | 0.372 |
| high fever* | 12(13)[†] | -0.308 | -12 | 0.640 |
| meningitis* | 1 | 0.426 | -8 | 0.084 |
| droplet infection* | 20 | 0.360 | 1 | -0.336 |
| headache* | 3 | -0.127 | -9(-10)[†] | 0.534 |
| plantar* | 5 | 0.397 | -18 | 0.372 |
| encephalitis* | 20 | -0.229 | -13 | 0.456 |
| neurogenic pulmonary edema* | 17 | 0.092 | -2(-3)[†] | 0.231 |
| dermatology* | 1 | 0.363 | -16 | 0.501 |
| fever* | 12 | -0.217 | -11 | 0.885 |

a. The number with the greatest absolute value of cross-correlation coefficient is selected, and its actual value and corresponding lag are represented.

b. *(Asterisk) labeled the cross-correlation lags of search terms earlier in 2020-2021 than 2015-2019.

c. [†]Different lags with the same cross-correlation coefficient were represented as a lag number (lag number).

**Predictors for actual HFMD cases before and during the pandemic**

We identified the most significant predictors of Internet search terms for HFMD infection using multiple linear regression with a stepwise procedure. In **Table 2**, the search terms "infect," "daycare," "vomit," "HFMD," "eczema," "pain," and "high fever" were found to be significant predictors in period 1 and accounted for an adjusted $R^2$=96.1% of the variation in HFMD infection. The search terms "infect," "enterovirus," "herpangina," "kindergarten," "myocarditis," "HFMD," "contact infection," "blister," "high fever" "dermatology" and "plantar" were significant predictors in period 2 and accounted for an adjusted $R^2$=90.6% of the variation in HFMD infection (**Table 3**). More Internet search terms were identified for predicting HFMD infection in period 2, especially "enterovirus" and "contact infection", which might indicate that the public attention affected by the COVID-19 pandemic.

**Table 2.** Stepwise regression model of the hand, foot, and mouth disease cases and correlated interest search trends from 2015 to 2019

| Search term | Model summary | | |
|---|---|---|---|
| | Standardized β | Std.Error | P-value |
| infect | .094 | 14.347 | <.001 |
| daycare | -.026 | 5.207 | .041 |
| vomit | -.054 | 10.903 | .001 |
| HFMD | .940 | 6.453 | <.001 |
| eczema | .039 | 13.841 | .013 |
| pain | -.062 | 19.304 | <.001 |
| high fever | -.067 | 14.683 | .001 |
| a. Dependent variable: hand, foot, and mouth disease cases | | | |
| b. Predictors: infect, child-rearing facility, vomit, HFMD, eczema, pain, high fever | | | |
| c. Adjust $R^2$ = 96.1% | | | |

**Table 3.** Stepwise regression model of the hand, foot, and mouth disease cases and correlated

interest search trends in 2020 and 2021

| Search term | Model summary | | |
|---|---|---|---|
| | Standardized β | Std.Error | P-value |
| infect | .159 | 6.661 | .011 |
| enterovirus | -.091 | 2.022 | .009 |
| herpangina | .252 | 3.11 | <.001 |
| kindergarten | .084 | 4.327 | .019 |
| myocarditis | .081 | 4.024 | .018 |
| HFMD | .696 | 4.514 | <.001 |
| contact infection | -.121 | 2.308 | .002 |
| blister | .143 | 4.725 | .004 |
| high fever | -.174 | 5.41 | .007 |
| dermatology | -.130 | 6.984 | .012 |
| plantar | -.104 | 4.726 | .028 |
| a. Dependent variable: hand, foot, and mouth disease cases  b. Predictors: infect, enterovirus, herpangina, kindergarten, myocarditis, HFMD, contact infection, blister, high fever, dermatology, plantar  c. Adjust $R^2$ = 90.6 % | | | |

243
244

**Discussion**

This study has presented the trends in actual HFMD cases and RSV of the search topic "HFMD" from 2004 to 2021. We found that the actual HFMD cases and RSV peaked around July in most years, except in 2020 and 2021, and surged after 2011 with peaks every two years before 2020. The search topic "HFMD" presented a strong correlation with actual HFMD cases, but the correlation was weak in 2004, 2008, and 2020. We conducted a cross-correlation analysis between the actual HFMD cases and related search terms before and during the pandemic. This indicated that the public might have improved awareness of HFMD infection during the pandemic. We used multiple linear regression to identify the most significant predictors of Internet search terms for actual HFMD cases before and during the pandemic. We identified the search terms "infect," "daycare," "vomit," "HFMD," "eczema," "pain," and "high fever" as the most significant predictors before the pandemic, and the search terms "infect," "enterovirus," "herpangina," "kindergarten," "myocarditis," "HFMD," "contact infection," "blister," "high fever" "dermatology" and "plantar" were the most significant predictors during the pandemic. Our research indicated that the Japanese public was more aware of HFMD infection during the COVID-19 pandemic. To our knowledge, this study is the first to predict HFMD cases using related Internet search terms and identify different predictors before and during the COVID-19 pandemic. This study also shows that Internet search data could supplement public health surveillance and help authorities respond to potential outbreaks rapidly.

From 2004 to 2021, the RSV of "HFMD" coincided with the actual HFMD cases in most years, except for 2004, 2008, and 2020. The data showed similar trends, with peaks occurring around July. However, in 2021, the peak was delayed until November. In 2004, 2008, and 2020, Google Trends search data did not match the actual "HFMD" cases. The search topic "HFMD" still presented a peak around July 2004, but the number of cases was low, which could result from the lower penetration of the Google search engine at the time (ComScore on top search engines for December 2004, 2005). In 2008, the search topic "HFMD" peaked on May 2-10 and July 20-26. The peak on July 20-26 was

consistent with previous years in Japan, whereas the large outbreak in China drove the peak on May

2-10, which could be why the search topic "HFMD" presented a weak correlation with the actual

HFMD cases. The latest epidemic occurred in China in 2008. More than 600,000 HFMD cases and

126 deaths in children were reported from March 2008 to June 2009 (Zhang et al., 2010), and Japan

reported that this infection came from China (National Institute of Public Health, 2008). In 2020, the

Japanese government implemented several measures to control COVID-19. Subsequently, the spread

of HFMD did not occur as in previous years, and the search topic "HFMD" presented a relatively

small peak in July. This may explain why the search topic "HFMD" did not correlate strongly with

the actual HFMD cases in the first year of the pandemic. Despite a small peak in the search topic

"HFMD", the RSV was much lower than in previous years. Many researchers have shown that the

Internet search data from Google Trends represents the public interest in a specific topic. However,

Internet search data should be used cautiously as a surveillance system because large events can

easily interfere with it. Obtaining fine-grained data could help develop surveillance systems that can

effectively exclude biased or irrelevant information to predict infection outbreaks.

Our results indicate that the global pandemic might have enhanced public awareness of HFMD in

addition to COVID-19. Firstly, the cross-correlation results showed that 31 search terms presented

earlier than the actual HFMD cases, and only nine search terms presented later during the pandemic.

Of the HFMD-related search terms, 67.4% showed an earlier lag than before the pandemic. Secondly,

significant predictors for HFMD infections during the pandemic comprised more comprehensive

information. Before the pandemic, search terms mainly including vulnerable area and symptoms of

HFMD infection. Additional predictors were identified during the pandemic containing the

information with disease-causing virus, infection pathways, susceptible body parts, complications,

and visiting departments. Public might desire more comprehensive information of HFMD by Internet

searching affected by the COVID-19 pandemic. According to previous studies, the prevalence of

respiratory infectious diseases, such as influenza, varicella, herpes zoster, rubella, and measles, was

reduced during the COVID-19 pandemic (Sakamoto et al., 2020; Wu et al., 2020; Wu et al., 2020;

Kies et al., 2021; Veiga et al., 2021; Li et al., 2021; Wan et al., 2021). This might have been due to

adherence to non-pharmaceutical interventions and lower non-polio enterovirus activity during the

COVID-19 pandemic compared with 2014-2019 (Kuo et al., 2021). Switzerland had an

unprecedented complete absence of pediatric enteroviral meningitis in 2020 (Stoffel et al., 2021). In

Japan, community-acquired pneumonia (Yan et al., 2022) and influenza (Hirose et al., 2021)

admissions have been reduced during the COVID-19 pandemic. COVID-19 preventative actions and

better personal hygiene are beneficial for preventing the spread of disease. In the upcoming season,

the prevalence of common diseases may rise as the public gradually complies less with infection

control measures (Yan et al., 2022). Our results show that there was a peak in HFMD infection and

public interest in November 2021. Continuous monitoring of HFMD is required, and public

information-seeking behavior may be useful for public health surveillance.

**Limitations**

This study had several limitations. First, our findings are limited to those who used Google to search

for health-related information, which may not represent the entire community. Second, the specific

HFMD-related search terms we selected did not represent all search terms in public use. Third,

search interest analysis is complementary to infection prediction and cannot replace traditional

research methods because of its hypersensitivity to large events. Fourth, we used search data from

2015 to 2019 to represent the period before the COVID-19 pandemic due to restrictions of Google

Trends, which may not represent the entire period. Finally, our study included only Japan; therefore,

our findings might not apply to other countries.

**Conclusion**

We identified different predictors for HFMD infections before and during the COVID-19 pandemic.

Our results indicate that the public may have had enhanced awareness of HFMD infection and paid

more attention to viruses during the COVID-19 pandemic. It is critical to continuously monitor

resurgent common infections as the public gradually reduces compliance with infection control

measures. Public information-seeking behavior using Internet search query data may be useful for

public health surveillance.

**Declarations**

**Acknowledgments:** Not applicable**.**

**Funding:** This work was supported by the Japan Science and Technology for pioneering research

initiated by the next generation (SPRING; grant number JPMJSP2110).

**Ethics Statement:** No ethical approval was required.

**Availability of Data and Materials:** We used publicly available data published by Google Trends

and the National Institute of Infectious Diseases in Japan. All data generated or analyzed during this

study are included in this published article (and its supplementary information files).

**Competing Interests:** The authors declare that they have no competing interests.

**Authors' Contributions:** QN, JYL, MNT, and TA contributed to the study conception and design.

QN, AB, KH, MO, and AK collected data. QN and JYL participated in data analysis and

interpretation and drafted the manuscript. All authors contributed to the manuscript revision and

approved the final version of the manuscript.

Abbreviations**: HFMD: h**and, f**oot, and** m**outh** d**isease; EV-A71: enterovirus A71; CV-A6:**

**coxsackievirus A6; CV-A16: coxsackievirus A16; NIID: National Institute of Infectious Disease;**

**LDE: logistic differential equation; L-DNM: landscape dynamic network marker; COPD:**

**chronic obstructive pulmonary disease; RSV: relative search volume**

**References**

Adelhoefer S, Berning P, Solomon SB, Maybody M, Whelton SP, Blaha MJ, Dzaye O. Decreased

public pursuit of cancer-related information during the COVID-19 pandemic in the United

States. Cancer Causes Control. 2021;32(6):577–85.

Akossou P. Impact of data structure on the estimators R-square and adjusted R-square in linear

regression. Int J Math Comput. Available from: https://www.researchgate.net/profile/Arcadius-

Akossou/publication/289526309_Impact_of_data_structure_on_the_estimators_R-

square_and_adjusted_R-square_in_linear_regression/links/586a90ce08ae329d621114f2/Impact-

of-data-structure-on-the-estimators-R-square-and-adjusted-R-square-in-linear-regression.pdf.

Akpan IJ, Aguolu OG, Kobara YM, Razavi R, Akpan AA, Shanker M. Association between what

people learned about COVID-19 using web searches and their behavior toward public health

guidelines: Empirical Infodemiology study. J Med Internet Res. 2021;23(9):e28975.

An L, Russell DM, Mihalcea R, Bacon E, Huffman S, Resnicow K. Online search behavior related to

COVID-19 vaccines: Infodemiology study. JMIR Infodemiology. 2021;1(1):e32127.

Boehm A, Pizzini A, Sonnweber T, Loeffler-Ragg J, Lamina C, Weiss G, Tancevski I. Assessing

global COPD awareness with Google Trends. Eur Respir J. 2019;53(6).

doi:10.1183/13993003.00351-2019.

Bourke P. Cross Correlation. Auto correlation—2D pattern. Available from:

https://www.researchgate.net/profile/Paul-Bourke-

2/publication/245817141_Cross_correlation_autocorrelation_2d_pattern_identification/links/00

46352c36f114ecbf000000/Cross-correlation-autocorrelation-2d-pattern-identification.pdf.

Chew SP, Chong SL, Barbier S, Matthew A, Lee JH, Chan YH. Risk factors for severe hand foot

mouth disease in Singapore: A case control study. BMC Infect Dis. 2015;15:486.

Cohen SA, Ebrahimian S, Cohen LE, Tijerina JD. Online public interest in common malignancies

and cancer screening during the COVID-19 pandemic in the United States. J Clin Transl Res.

2021;7(6):723–32.

Diaz P, Reddy P, Ramasahayam R, Kuchakulla M, Ramasamy R. COVID☐19 vaccine hesitancy

linked to increased internet search queries for side effects on fertility potential in the initial

rollout phase following Emergency Use Authorization. Andrologia. 2021;53(9):e14156.

doi:10.1111/and.14156.

Dreher PC, Tong C, Ghiraldi E, Friedlander JI. Use of google trends to track online behavior and

interest in kidney stone surgery. Urology. 2018;121:74–8.

Eysenbach G. Infodemiology and Infoveillance: Framework for an emerging set of public health

informatics methods to analyze search, communication and publication behavior on the Internet.

J Med Internet Res. 2009;11(1):e11. doi:10.2196/jmir.1157.

Fujimoto T, Iizuka S, Enomoto M, Abe K, Yamashita K, Hanaoka N, Okabe N, Yoshida H, Yasui Y,

Kobayashi M, Fujii Y, Tanaka H, Yamamoto M, Shimizu H. Hand, foot, and mouth disease

caused by coxsackievirus A6, Japan, 2011. Emerg Infect Dis. 2012;18(2):337–9.

Gao Q, Liu Z, Xiang J, Tong M, Zhang Y, Wang S, Zhang Y, Lu L, Jiang B, Bi P. Forecast and early

warning of hand, foot, and mouth disease based on meteorological factors: Evidence from a

multicity study of 11 meteorological geographical divisions in mainland China. Environ Res.

2021;192:110301.

National Institute of Public Health [Overseas Infectious Disease Information] China Enterovirus(EV-

71) 2008/05/07 – H・CRISIS. [cited 30 May 2022]. Available: https://www.niph.go.jp/h-

crisis/archives/21996/.

Han J, Kamat S, Agarwal A, O'Hagan R, Tukel C, Owji S, Ghalili S, Luu Y, Dautriche Svidzinski C,

Abittan BJ, Ungar J, Gulati N. Correlation between interest in COVID-19 hair loss and COVID-

19 surges: Analysis of google trends. JMIR Dermatol. 2022;5(2):e37271.

Higgins TS, Wu AW, Sharma D, Illing EA, Rubel K, Ting JY, Snot Force Alliance. Correlations of

online search engine trends with coronavirus disease (COVID-19) incidence: Infodemiology

study. JMIR Public Health Surveill. 2020;6(2):e19702.

Hirose T, Katayama Y, Tanaka K, Kitamura T, Nakao S, Tachino J, Nakao S, Nitta M, Iwami T,

Fujimi S, Uejima T, Miyamoto Y, Baba T, Mizobata Y, Kuwagata Y, Shimazu T, Matsuoka T.

Reduction of influenza in Osaka, Japan during the COVID-19 outbreak: A population-based

ORION registry study. IJID Reg. 2021;1:79–81.

Huang J, Liao Q, Ooi MH, Cowling BJ, Chang Z, Wu P, Liu F, Li Y, Luo L, Yu S, Yu H, Wei S.

Epidemiology of recurrent hand, foot and mouth disease, China, 2008–2015. Emerg Infect Dis.

2018;24(3). doi:10.3201/eid2403.171303.

Husnayain A, Shim E, Fuad A, Su EC-Y. Predicting new daily COVID-19 cases and deaths using

search engine query data in South Korea from 2020 to 2021: Infodemiology study. J Med

Internet Res. 2021;23(12):e34178.

IASR. Summer Diseases (herpangina/foot, and mouth disease). [Cited 23 May 2022]. Available

from: https://www.niid.go.jp/niid/ja/iasr/510-surveillance/iasr/graphs/4892-iasrgnatus.html.

IDWR; Year 2019 Issue 29 <Infectious Diseases Requiring Attention> Hand, Foot, and Mouth

Disease. [Cited 16 Jun 2022]. Available from: https://www.niid.go.jp/niid/ja/hfmd-m/hfmd-

idwrc/9017-idwrc-1929.html.

IDWR Surveillance Data Table 2022 week;18. [Cited 23 May 2022]. Available from:

https://www.niid.go.jp/niid/en/survaillance-data-table-english/11133-idwr-sokuho-data-e-

2218.html; May 17 2022.

Kabele Pavel, Mojhová Martina, Smíšková Dita. Hand-foot-mouth disease in puerperium. Ceska

Gynekol. 2022;87(1):47–9.

Kardeş S, Erdem A, Gürdal H. Public interest in musculoskeletal symptoms and disorders during the

COVID-19 pandemic: Infodemiology study. Z Rheumatol. 2022;81(3):247–52.

Kies KD, Thomas AS, Binnicker MJ, Bashynski KL, Patel R. Decrease in enteroviral meningitis: An

unexpected benefit of coronavirus disease 2019 (COVID-19) Mitigation? Clin Infect Dis

2021;73(9):e2807–9–e2809.

Knipe D, Gunnell D, Evans H, John A, Fancourt D. Is Google Trends a useful tool for tracking

mental and social distress during a public health emergency? A time-series analysis. J Affect

Disord. 2021;294:737–44.

Kuo SC, Tsou HH, Wu HY, Hsu YT, Lee FJ, Shih SM, Hsiung CA, Chen WJ. Nonpolio enterovirus

activity during the COVID-19 pandemic, Taiwan, 2020. Emerg Infect Dis. 2021;27(1).

doi:10.3201/eid2701.203394.

Li Q, Wang J, Lv H, Lu H. Impact of China's COVID-19 prevention and control efforts on outbreaks

of influenza. BioSci Trends. 2021;15(3):192–5.

Linkov F, Bovbjerg DH, Freese KE, Ramanathan R, Eid GM, Gourash W. Bariatric surgery interest

around the world: What Google Trends can teach us. Surg Obes Relat Dis. 2014;10(3):533–8.

Mavragani A, Ochoa G. Google trends in Infodemiology and Infoveillance: Methodology

framework. JMIR Public Health Surveill. 2019;5(2):e13439.

Mavragani A, Ochoa G, Tsagarakis KP. Assessing the methods, tools, and statistical approaches in

google trends research: Systematic review. J Med Internet Res. 2018;20(11):e270.

Mavragani A, Sampri A, Sypsa K, Tsagarakis KP. Integrating smart health in the US health care

system: Infodemiology study of asthma monitoring in the google era. JMIR Public Health

Surveill. 2018;4(1):e24.

Memon SA, Razak S, Weber I. Lifestyle disease surveillance using population search behavior:

Feasibility study. J Med Internet Res. 2020;22(1):e13347.

Nuti SV, Wayda B, Ranasinghe I, Wang S, Dreyer RP, Chen SI, Murugiah K. The use of google

trends in health care research: A systematic review. PLOS ONE. 2014;9(10):e109583.

Pervaiz F, Pervaiz M, Abdur Rehman N, Saif U. FluBreaks: Early epidemic detection from Google

flu trends. J Med Internet Res. 2012;14(5):e125.

Phillips CA, Barz Leahy A, Li Y, Schapira MM, Bailey LC, Merchant RM. Relationship between

state-level google online search volume and cancer incidence in the United States: Retrospective

study. J Med Internet Res. 2018;20(1):e6.

Puenpa J, Wanlapakorn N, Vongpunsawad S, Poovorawan Y. The history of enterovirus A71

outbreaks and molecular epidemiology in the Asia-Pacific region. J Biomed Sci. 2019;26(1):75.

Pullan S, Dey M. Vaccine hesitancy and anti-vaccination in the time of COVID-19: A Google

Trends analysis. Vaccine. 2021;39(14):1877–81.

Rui J, Luo K, Chen Q, Zhang D, Zhao Q, Zhang Y, Zhai X, Zhao Z, Zhang S, Liao Y, Hu S, Gao L,

Lei Z, Wang M, Wang Y, Liu X, Yu S, Xie F, Li J, Liu R, Chiang YC, Zhao B, Su Y, Zhang

XS, Chen T. Early warning of hand, foot, and mouth disease transmission: A modeling study in

mainland, China. PLOS Negl Trop Dis. 2021;15(3):e0009233.

Sakamoto H, Ishikane M, Ueda P. Seasonal influenza activity during the SARS-CoV-2 outbreak in

Japan. JAMA. 2020;323(19):1969–71.

Schootman M, Toor A, Cavazos-Rehg P, Jeffe DB, McQueen A, Eberth J, Davidson NO. The utility

of Google Trends data to examine interest in cancer screening. BMJ Open. 2015;5(6):e006678.

Sharma D, Sandelski MM, Ting J, Higgins TS. Correlations in trends of sinusitis-related online

google search queries in the United States. Am J Rhinol Allergy. 2020;34(4):482–6.

Stoffel L, Agyeman PKA, Keitel K, Barbani MT, Duppenthaler A, Kopp MV, Aebi C. Striking

decrease of enteroviral meningitis in children during the COVID-19 pandemic. Open Forum

Infect Dis. 2021;8(6):ofab115.

Sumi A, Toyoda S, Kanou K, Fujimoto T, Mise K, Kohei Y, Koyama A, Kobayashi N. Association

between meteorological factors and reported cases of hand, foot, and mouth disease from 2000

to 2015 in Japan. Epidemiol Infect. 2017;145(14):2896–911.

Sun BJ, Chen HJ, Chen Y, An XD, Zhou BS. The risk factors of acquiring severe hand, foot, and

mouth disease: A meta-analysis. Can J Infect Dis Med Microbiol. 2018;2018:2751457.

Taira K, Hosokawa R, Itatani T, Fujita S. Predicting the number of suicides in Japan using Internet

search queries: Vector autoregression time series model. JMIR Public Health Surveill.

2021;7(12):e34016.

Tizek L, Schielein M, Rüth M, Ständer S, Pereira MP, Eberlein B, Biedermann T, Zink A. Influence

of climate on google Internet searches for pruritus across 16 German cities: Retrospective

analysis. J Med Internet Res. 2019;21(7):e13739.

Tranmer, M., Murphy, J., Elliot, M., and Pampaka, M. (2020) Multiple Linear Regression (2nd

Edition); Cathie Marsh Institute Working Paper 2020-01.

https://hummedia.manchester.ac.uk/institutes/cmist/a rchive-publications/working-

papers/2020/2020-1- multiple-linear-regression.pdf

Veiga ABGD, Martins LG, Riediger I, Mazetto A, Debur MDC, Gregianini TS. More than just a

common cold: Endemic coronaviruses OC43, HKU1, NL63, and 229E associated with severe

acute respiratory infection and fatality cases among healthy adults. J Med Virol.

2021;93(2):1002–7.

Wan WY, Thoon KC, Loo LH, Chan KS, Oon LLE, Ramasamy A, Maiwald M. Trends in

respiratory virus infections during the COVID-19 pandemic in Singapore, 2020. JAMA Netw

Open. 2021;4(6):e2115973.

Wang J, Hu T, Sun D, Ding S, Carr MJ, Xing W, Li S, Wang X, Shi W. Epidemiological

characteristics of hand, foot, and mouth disease in Shandong, China, 2009–2016. Sci. Rep.. Sci

Rep. 2017;7(1):8900.

Wu D, Liu Q, Wu T, Wang D, Lu J. The impact of COVID-19 control measures on the morbidity of

varicella, herpes zoster, rubella and measles in Guangzhou, China. Immun Inflam Dis.

2020;8(4):844–6.

Wu D, Lu J, Liu Y, Zhang Z, Luo L. Positive effects of COVID-19 control measures on influenza

prevention. Int J Infect Dis. 2020;95:345–6.

Yan Y, Tomooka K, Naito T, Tanigawa T. Decreased number of inpatients with community-

acquired pneumonia during the COVID-19 pandemic: A large multicenter study in Japan. J

Infect Chemother. 2022;28(5):709–13.

Yu G, Feng H, Feng S, Zhao J, Xu J. Forecasting hand-foot-and-mouth disease cases using wavelet-

based SARIMA-NNAR hybrid model. PLOS ONE. 2021;16(2):e0246673.

ComScore on top search engines for December 2004: Google –35%, Yah00 –32%; Feb 13 2005. In:

ZDNet. [Cited 15 Jun 2022]. Available from: https://www.zdnet.com/article/comscore-on-top-

search-engines-for-december-2004-google-35-yah00-32/.

Zhang X, Xie R, Liu Z, Pan Y, Liu R, Chen P. Identifying pre-outbreak signals of hand, foot and

mouth disease based on landscape dynamic network marker. BMC Infect Dis. 2021;21(Suppl

1):6.

Zhang Y, Zhu Z, Yang W, Ren J, Tan X, Wang Y, Mao N, Xu S, Zhu S, Cui A, Zhang Y, Yan D, Li

Q, Dong X, Zhang J, Zhao Y, Wan J, Feng Z, Sun J, Wang S, Li D, Xu W. An emerging

recombinant human enterovirus 71 responsible for the 2008 outbreak of hand foot and mouth

disease in Fuyang city of China. Virol J. 2010;7:94.

Zheng YM, Yang J, Liao QH. Direct medical costs and influencing factors in severe hand, foot and

mouth disease in children aged between six months and five years old. Zhonghua Yu Fang Yi

Xue Za Zhi. 2017;51(1):87–92.

Zhuang ZC, Kou ZQ, Bai YJ, Cong X, Wang LH, Li C, Zhao L, Yu XJ, Wang ZY, Wen HL.

Epidemiological research on hand, foot, and mouth disease in Mainland China. Viruses.

2015;7(12):6400–11.

Ministry of Health, Labour and Welfare. National Institute of Infectious Diseases. Infectious Disease

Weekly Report (Japan IDWR);11. Available from: https://ci.nii.ac.jp/naid/10027747966/; 2009.

**Figure Captions**

**Figure 1.** Monthly number and relative search volume of hand, foot, and mouth disease from 2004 to

2021.

**Figure 2.** Correlation between the actual hand, foot, and mouth disease cases and the relative search

volume from 2004 to 2021.