Digitalization impacts the COVID-19 pandemic and the stringency of government measures

Helen Heinrichs[1], Florian Mueller[1,2], Lucia Rohfleisch[1], Volkmar Schulz[1,2,3], Steven R. Talbot[4,†,*] & Fabian Kiessling[1,3,5,†,*]

[1] Institute for Experimental Molecular Imaging, Center for Biohybrid Medical Systems, RWTH Aachen University Clinic, Forckenbeckstr. 55, 52074 Aachen, Germany

[2] Department of Physics of Molecular Imaging Systems, Institute for Experimental Molecular Imaging, Center for Biohybrid Medical Systems, RWTH Aachen University Clinic, Forckenbeckstr. 55, 52074 Aachen, Germany

[3] Fraunhofer Institute for Digital Medicine MEVIS, Bremen, Germany

[4] Institute for Laboratory Animal Science, Hannover Medical School, Carl-Neuberg-Str. 1, 30625 Hannover, Germany

[5] Comprehensive Diagnostic Center Aachen, RWTH Aachen University Clinic, Pauwelstr. 30, 52074 Aachen, Germany

[†] these authors contributed equally

[*] Correspondences: fkiessling@ukaachen.de & talbot.steven@mh-hannover.de

Abstract

COVID-19 poses a significant burden on populations worldwide. Although the pandemic has accelerated digital transformation, little is known about digitalization's influence on pandemic developments. Using the Digital Adoption Index (DAI), we examined the effects of digitalization on early COVID-19 cases, deaths, and stringency indices (SI) of government measures. Gradient Tree Boosting pinpointed essential features, such as populations' smoker fraction, age, poverty, and DAI. Then, regression analyses indicated that higher DAI was associated with significant declines in new cases (β=-362.25/pm; p<0.001) and deaths (β=-5.53/pm; p<0.001) months after the peak. Over the entire range, higher DAI was associated with higher mean SI (β=4.86; p<0.01). Interestingly, however, in countries with high DAI >0.68, the correlation inverted (β=-55.26; p<0.05). Finally, a scoping review identified 70 publications providing valuable arguments for our findings. Thus, digitalization is evident for handling current and future pandemics but can also facilitate stronger government actions restricting individual freedom.

Introduction

The global coronavirus disease 2019 (COVID-19) outbreak has kept the world on hold for some time. Countries have suffered large numbers of infections and deaths [1,2], which tremendously affected economies and societies [3,4]. As a response, governments have implemented coordinated containment and mitigation strategies [5]. Although authorities worldwide have employed similar policies, such as work and school closures, border restrictions, quarantine, or lockdown, their impact on the pandemic and the stringency of implementation have varied across countries [6-8]. Recent research suggests that the effectiveness of policies depends on many factors, including socioeconomic and demographic conditions as well as cultural and political structures [5,9]. In addition, digital information technology may be a distinguishing factor determining the course of the pandemic [10].

The pandemic has accelerated digital transformation in many areas of life. The health care sector is rapidly adopting telemedicine [10], schools are working with collaborative online platforms [11], and business models are becoming more hybrid [12]. However, little is known about how digitalization has shaped the crisis and how it has been managed. Asian countries have introduced many innovative technologies to attenuate the spread of COVID-19. South Korea and Singapore strengthened their digital contact tracing tools by promptly implementing existing surveillance technologies [13,14]. Hong Kong and Taiwan used government-issued mobile phones and wristbands as quarantine compliance measures [10,15]. Such measures are potentially relevant factors in controlling the pandemic, leading to less stringent control measures [8,13]. The effectiveness of digital contact tracing in controlling outbreaks has been discussed in much earlier work by Fraser et al. [16] and has recently been adapted to the case of COVID-19 [17]. However, it is unclear how digital tracing and surveillance are implementable in regions with diverse digital infrastructure. Evidence in existing research focuses on the promises of digitalization. Nevertheless, as described in previous papers, contact tracing requires extensive digital infrastructure [13-15]. The pandemic has exposed fundamental gaps in the universality of digital access, resulting in challenges in the implementation and use of digital technologies [18].

This study aimed to explore the effect of digitalization on COVID-19 developments in a larger context. To investigate whether digital technologies and infrastructures may have influenced pandemic trends, we examined the role of countries' digital adoption, expressed by the World Bank's Digital Adoption Index (DAI), in changes in COVID-19 cases, deaths, and the stringency index (SI) of government actions.

Our approach allows for evaluating the effect of digitalization on pandemic propagation and government activity beyond factors of individual digital health applications. Our findings highlight the need for efficient and technologically advanced infrastructures to combat future potential health crises.

Results

We retrieved data from the open-access COVID-19 dataset, available via GitHub and maintained by *Our World in Data* [19]. These data are updated daily and facilitate an easy-to-use and comparative framework for this transnational study. It also entails the SI from the Oxford COVID-19 Government Response Tracker, which we included as a standardized method to compare countries' stringency of government measures [20]. We used the reported data until March 10, 2021. This study focused on the early phase of COVID-19 when countries took initial actions, and new technological tools were not yet

available or being developed to assess countries' digital preparedness facing COVID-19. The World Bank's DAI from 2016 expresses countries' levels of digital adoption across three dimensions of the economy: government, business, and people. [21,22]. Each of the three sub-indices comprises various equally weighted and normalized indicators on a 0 to 1 scale, while the DAI represents the simple average. The "government" pillar combines online public services, digital identification, and core administrative systems. The "business" pillar includes the number of secure servers (in millions), internet speed (kbps), percentage of businesses with websites, and 3G coverage. Finally, the "people" pillar indicates mobile and internet access at home.

The more advanced countries' digitalization, the lower the number of cases and the faster new cases decline

Gradient Tree Boosting (GTB) was first applied to compare DAI, demographic and health-related variables with our outcome variables of interest. At the peak of COVID-19 cases in 2020, GTB indicated a strong dependence on digitalization across countries (0.23, SD±0.12) next to the proportion of female smokers (0.32, SD±0.22) and older people (0.22, SD±0.14). Male smokers (0.14, SD±0.04) and per capita gross domestic (0.09, SD±0.05) were also essential characteristics (Fig.2a). Values in parentheses indicate the relative feature importance and standard deviation.

We plotted the change in new cases per million people (pm) by countries' peaks against DAI using linear regression. We scrutinized the slope to determine how DAI potentially counteracted pandemic trends within months after reaching maximum new cases. Figure 2b shows a significant decline in new COVID-19 cases with increasing DAI in most countries, trending toward 362 fewer new cases per unit DAI increase (β=-362.25/pm, p<0.001). This trend analysis was repeated with a regression weighted by the difference in caseload, also showing a greater ability to reduce new cases pm with increasing DAI (β=-446.54/pm, p<0.001) (Table 1).

Table 1 Linear regression analyses of the association between DAI and changes in COVID-19 new cases and deaths months after their peaks.

	β	Std. Error of β	t	*p-value*
Effect of DAI on changes in new cases (smoothed) pm				
Constant	92.9	30.46	3.049	0.003*
DAI	-362.25	53.79	-6.734	0.001**
Effect of DAI on changes in new cases (smoothed) pm (weighted)				
Constant	35.43	68.46	0.518	0.606
DAI	-446.54	101.06	-4.419	0.001**
Effect of DAI on changes in new deaths (smoothed) pm				
Constant	1.02	0.73	1.398	0.164
DAI	-5.53	1.29	-4.293	0.001**
Effect of DAI on changes in new deaths (smoothed) pm (weighted)				
Constant	-3.42	1.97	-1.738	0.084
DAI	-3.79	2.99	-1.269	0.207

* p≤0.05; ** p≤0.001;
Notes: The regression coefficient (β) is the degree of change in the outcome variable for every one-unit change in the predictor variable. The t-statistic is the regression coefficient divided by its standard error. Digital adoption is expressed in arbitrary units of DAI.

Of the 5012 articles retrieved by our search, we removed 1454 duplicates and another 3450 records during the title and abstract screening. Finally, of the 108 records eligible for the full-text screening, 70 met the inclusion criteria and were included in the scoping review (Supplementary Fig. 5). We grouped

keywords into the following themes: i) health care system, ii) government responses, and iii) determinants facilitating or hindering public health outcomes (Fig. 1). In this results section, we discuss the most prominent findings related to the first two themes, and in the Supplementary Results section, we discuss the determinants of public health outcomes (Supplementary Fig. 3). Many articles on digital technologies covered cases and deaths, summarized under "health care system". Supplementary Fig. 6 and Supplementary Fig. 7 provide some descriptive findings on the geographical mapping of included records.

Fig. 1 Funnel model of included themes and keywords in the scoping review. The funnel model illustrates the scoping review process. We began our literature review with a broad search strategy that included COVID-19-related terms (e.g., "Covid-19" OR "sars-cov-2" OR "pandemic") and digit*, which represents the digital domain with different word endings (as indicated by an asterisk (*)). After an extensive selection process, 70 papers were included in our review. From these papers, we identified keywords reflecting recurring aspects of the literature. These were grouped into the following themes: i) health care system, ii) government responses, and iii) determinants facilitating or hindering public health outcomes. Eventually, the scoping review offered essential arguments for our statistical findings. (Note: NPIs = non-pharmaceutical interventions; NPIs are public health measures to prevent and/or mitigate transmission of SARS-CoV-2 that do not rely on medications or vaccination. These include, but are not limited to, surveillance measures, travel restrictions, and recommended use of face masks [23]).

Digital technologies within the health care system.

The scoping review's first theme (n=26) addressed health care technological features that may affect care-delivery processes during outbreaks or that were present before the pandemic (Fig. 2c). Telehealth involves various technical methods (e.g., virtual consultations or remote patient monitoring) rapidly adopted in the US, UK, Australia, and elsewhere [24-26]. Such electronic communication tools improve access to care by eliminating the additional risk of exposure [24]. With resource shortages looming, Scott

et al. introduced the National Emergency Tele-Critical Care Network to deliver care and expertise via mobile devices across the US [27]. Chatterjee et al. provided an overview of integrated communication systems in South America [25]. Ecuador, Bolivia, and Peru, which score medium in DAI, introduced telemedicine platforms providing access to specialists virtually. Chile, achieving a higher DAI, enforced a new model of digital hospitals with upcoming features like electronic health records (EHRs). Additionally, they propose using IoT or AI-based devices to assess health parameters more accurately [25]. Wood et al. investigated the state EHRs in the UK [28]. While researchers initially lacked access to critical population health data, the British Heart Foundation Data Science Centre developed a digital environment with secure access to crucial COVID-19 data for researchers. During the COVID-19 outbreak, China rapidly expanded the establishment of internet hospitals. These combine various web-based services for the public, such as telemedicine, drug delivery, and psychological counseling [29,30]. An early adopter of robots, Guangdong Provincial People's Hospital assisted the medical staff by disseminating health information or delivering medicines to isolation wards [31]. While telecommunication facilitates medical treatment in First Nations or sparsely populated islands [26], illiterate or low-income populations are severely affected by the digital divide and face unequal access to telehealth [25]. Despite a high mobile penetration in India, the introduction of eHealth services proved difficult for similar reasons [26].

Fig. 2 The impact of digital adoption on the development of COVID-19 cases. a The gradient tree boosting results show the five most important characteristics for peak total COVID-19 cases in 2020 out of 15 features. Relative feature importance was normalized, and the sum was set to 1.0. Data are expressed as mean ± standard deviation. **b** Change in new cases pm after the peak in 2020-2021. I) The result of the simple linear regression illustrates the effect of the DAI on the reduction in new COVID-19 cases (7-day-smoothed) pm in the post-peak

period. Countries with higher DAI were more efficient in significantly reducing the number of new cases. II) The second regression is weighted by the delta of minimum and maximum new cases to show the impact of more successful countries in lowering the new case numbers. The blue (I) and red (II) lines indicate the linear fits of regression analyses. The areas of the 95% confidence intervals are shaded gray. DAI is expressed as an arbitrary unit. **c** The radar chart displays the keywords addressed in the "health care system" theme of the scoping review. The scale for the radial axis diagrams shows how frequently the keywords occurred in the included studies. Papers may contain different numbers of keywords.

The more advanced countries' digitalization, the lower the number of deaths and the faster new deaths decline

We employed GTB to identify essential features of the peak numbers of deaths (Fig. 3a). We found that the proportion of female smokers (0.49, SD±0.12) held the highest feature importance, followed by extreme poverty (0.17, SD±0.02) and the population aged 70 or older (0.11, SD±0.07). Additionally, GTB indicated feature importance for DAI (0.13, SD±0.05) and the proportion of male smokers (0.1, SD±0.03).

Regression analysis shows the impact of digitalization on post-peak COVID-19 deaths (Table 1; Fig. 53b). Higher DAI was associated with a significant decrease in new deaths pm, resulting in approximately 5.53 (p<0.001) fewer deaths pm after the peak in 2020/2021. The slope expresses the decline in cases after reaching the peak for each country. We performed another regression analysis weighted by the decrease in the number of deaths, showing the same trend (β=-3.79/pm; p=0.207).

Fig. 3 The impact of digital adoption on the development of COVID-19 deaths. **a** Gradient tree boosting results show the top five important features for peak COVID-19 deaths in 2020. Relative feature importance was normalized, and the sum was set to 1.0. Data are expressed as mean ± standard deviation. **b** Regression results

show the effect of the DAI on reducing new deaths pm during the post-peak period in 2020-2021. I) Countries with higher DAI could significantly reduce their number of new COVID-19 deaths (7-day-smoothed) months after the peak. II) In the weighted regression analysis, the significant effect of the DAI on reducing deaths is no longer visible. The blue (I) and red (II) lines indicate the linear fits of regression analyses. The area of the 95% confidence interval is shaded gray. DAI is expressed as an arbitrary unit.

The influence of digitalization on the stringency of government measures

The most important features identified with GTB and associated with SI were hospital beds per thousand people (0.24, SD±0.03), followed by DAI (0.22, SD±0.03). Other informative features included extreme poverty (0.19, SD±0.04), male smokers (0.17, SD±0.02), and diabetes prevalence (0.17, SD±0.02) (Fig. 4a).

The final regression model shows that, on average, countries with a higher DAI adopted slightly stricter policy measures during the first year of the pandemic (β=4.86/pm, $p<0.01$) (Fig. 4bI). After plotting DAI against mean SI, we identified three trends in data and grouped the DAI accordingly. Then, the estimated effect of each group (low, medium, and high DAI) was analyzed by including the DAI:class interaction term in the model. Subsequent regression analyses of all three classes showed that the interaction of DAI and its high class (>0.68) could significantly reduce the mean SI compared to the interaction of DAI and low class (β=-55.26, $p<0.05$) (Fig. 4bII, Table 2). See Supplementary Table 1 and Supplementary Figure 2 for more information on the classification of DAI classes.

Table 2 Linear regression models of the overall effect of DAI on SI and its regression per class (low, medium, high).

	β	Std. Error of β	t	*p-value*
Overall effect of DAI on SI$_{norm}$				
Constant	-0.40	1.00	-0.398	0.691
DAI	4.86	1.75	2.780	0.006**
Effect of low, medium, and high DAI on the trend in SI				
Constant	38.61	12.10	3.191	0.003**
low [0;0.36]	54.18	42.01	1.290	0.205
Constant	71.18	9.98	7.133	0.001**
medium [0.37;0.68]	-10.61	18.23	-0.582	0.562
Constant	102.36	18.75	5.458	0.001**
high [0.69;1]	-55.26	24.32	-2.272	0.028*

* $p\leq0.05$; ** $p\leq0.001$;
Notes: The regression coefficient (β) is the degree of change in the outcome variable for every one-unit change in the predictor variable. The t-statistic is the regression coefficient divided by its standard error. Digital adoption is expressed in arbitrary units of DAI. The numbers in squared brackets indicate the intervals of the classes.

Digital technologies in government responses.

46 of 70 review articles offered insights into government responses (Fig. 4c). Throughout the pandemic, contact tracing and surveillance management were utilized to support government decision-making [32,33]. Southeast Asia led the way in leveraging digital technologies to track transmission routes early in the pandemic. Digital contact tracing in South Korea, Singapore, and Hong Kong [34,35] is attributed to the successful containment of COVID-19 in its early stages [15]. The countries' populations are particularly obedient to following government guidelines and using their cell phones' GPS. Israel also collected cell phone location data to aid contact tracing [36], and Taiwan ordered staying-at-home measures immediately after detecting a case [34]. Some Asian countries have been exceptionally proactive and

diligent in integrating digital governance. A case study in Wuhan [32] exhibited community-based digital contact tracing, such as the Health QR Code for smartphones, a few weeks into the pandemic. This QR code held personal and health information and was regularly checked by contact tracers to monitor people's mobility. The government used this to divide Wuhan's inhabitants into community grids according to their risk status. Real-time data enabled governments to identify clusters of viral spreading, eventually reducing total infections and hospitalizations shortly after implementing the app. In Bangladesh, the government initiated mobile awareness initiatives where people received preventive information on their cell phones if they were subscribed to a phone operator [37]. While some digitalized countries have been more proactive, Germany and Austria were relatively slow to respond to the pandemic and eventually introduced stringent measures [35]. The initiation of lockdowns or travel restrictions in the US appeared less organized and varied between states [15]. Other studies show great potential for implementing transmission models and providing evidence-based guidance to governments [38]. For example, Leung et al. established a framework to track Hong Kong's replication number of COVID-19 in near real-time, without multi-day delays, using age-specific digital mobility data [39]. Cencetti et al. introduced a model for contact tracing combined with direct quarantine orders for infected individuals [40].

Fig. 4 The influence of digital adoption on the stringency of government measures. a GTB shows the relative importance of characteristics on government stringency indices. Relative feature importance was normalized, and the sum was set to 1.0. Data are expressed as mean ± standard deviation. **b** I) The regression model shows that, on average, countries with a higher DAI adopted slightly more stringent policy measures during the first year of the pandemic. To allow for a suitable comparison of the stringency index across countries, we adjusted for the different onsets of the SI ($SI_{norm} = SI_{mean}/SI_{initial}$). The area of the 95% confidence interval is shaded gray. II) Plotting DAI

against SI$_{mean}$ revealed three trends in data. Accordingly, we split the DAI into three groups (low, medium, and high). The thresholds of the groups were determined using a thresholded interaction analysis (lower threshold=0.36 and upper threshold=0.68). An increase in SI was visible in the low DAI group, while countries in the high DAI group decreased their stringency in government measures. **c** Findings related to government responses within the scoping review: contact tracing, non-pharmaceutical interventions, and evidence-based guidance. The scale for the radial axis diagrams shows how frequently the keywords occurred in the included studies.

To identify potential confounding characteristics for DAI, we performed an ANOVA with countries' income levels. ANOVA followed by a Games-Howell post-hoc test revealed significant differences among all four income levels, $F(3,160)=154.87$, $p<0.001$, and showed that DAI was proportionally higher in high-income countries ($p<0.001$) (Fig. 5).

Fig. 5 Global level of digital adoption by income. Digital adoption by country is represented by the digital adoption index (DAI), chosen as the one representative score covering numerous countries. The DAI is based on three sub-indices implying governments, businesses, and people. Regions include economies at all income levels [41]. Adoption of digital technologies varies across countries, with higher- and upper-middle-income countries scoring a higher DAI.

Discussion

COVID-19 prompted rapid and enthusiastic adoption of numerous digital technologies and processes. Given the ongoing devastating situation in many countries and the risk of further SARS-CoV-2 transmission, optimization of digital transformation is needed. Our findings assert that digitalization will not act as a single silver bullet to prevent outbreaks but significantly reduces cases and deaths and influences governments' SI. Therefore, digitalization may be an effective measure for mitigating COVID-19 and future pandemics.

GTB results indicate associations between several variables, including DAI and pandemic trends. The proportion of smokers and elderly are relevant characteristics in COVID-19 cases and deaths. Some studies attribute smoking to a worse outcome after contracting the virus [42-44], similar to how it was a risk factor for mortality in MERS-CoV cases in South Korea in 2015 [45]. The risk of death rises with age [43,46] and is highest in countries with elderly populations [47]. Also, the risk of comorbidities, e.g., diabetes, increases with age, which increases the likelihood of a severe infection course [48]. COVID-19 cases were associated with lower per capita gross domestic [49] and deaths with poverty [50]. The association between hospital bed capacity and SI could be explained due to increased pressure on hospital beds, resulting in governments imposing new policies [51].

Linear regression analyses show the relationship between DAI and the evolution of the pandemic. Countries with higher DAI proved more resilient in managing cases and deaths. Jurzik et al. note that higher levels of digitalization likely contributed to the ability of some Asian governments to respond effectively, highlighting the digital divide [52]. The COVID-19 pandemic has exacerbated the digital divide, disproportionally affecting minorities [53] and people living in relatively poorer regions [54]. WHO's "Global Strategy for Digital Health 2020-2025" advocates for accelerated implementation and universal and equitable access to digital health and health policies [55]. Digital health, including competencies in communication and information technologies, internet, and broadband access, is classified as a determinant of health [56,57]. A lack of digital literacy and skills can put people at risk for adverse health outcomes [58].

Review papers explored digital technologies and tools used during the pandemic. Additionally, some studies underline that sociocultural and political contexts, and public attitudes can strongly influence the effectiveness of government measures. Although countries with higher DAI have better prerequisites to cope with pandemics, countries like the US [36] have failed to promptly adopt contact tracing mainly because of privacy concerns [40]. Knowledge of previous respiratory syndrome outbreaks was invaluable for effective early response in Singapore, Taiwan, and Hong Kong [59]. After the MERS-CoV outbreak, South Korea implemented several tracing methods allowing authorities to collect data on peoples' mobility via credit cards and cell phones [60]. These approaches require numerous contact tracers, but affluent Western countries lacked such positions when COVID-19 struck [61]. However, the success of surveillance management in Asian countries comes at a price: a lack of privacy and freedom [62]. As seen in our findings, some of the countries scoring high in DAI, like England, France, Germany, and the US, were more reluctant to use digital measures and introduced restrictive non-pharmaceutical interventions (NPIs), e.g., lockdowns. These had farer-reaching negative personal, social, and economic impacts [62]. For NPIs to contribute meaningfully to the containment of COVID-19, they should be used effectively and targeted [23]. A high DAI (>0.68) facilitates preparedness and eases the stringency of government measures applied. Countries that deployed digital surveillance and contact tracing remained pioneers in managing disease burden [10], ultimately strengthening people's autonomy to maintain near-normal lives [62]. However, not all digital innovations are transferable to countries struggling with massive outbreaks. Internet use has spread globally but is less well utilized in poorer countries, posing a barrier to basic requirements for using digital technologies in those countries [22].

This study has some limitations: The GTB models included constant demographic and health-related parameters and cannot be used for predictive purposes. Furthermore, the lower fits of regression models suggest inherent variation likely caused by other covariates or the heterogeneous country-level nature. Future research could consider expanding the covariates to include those identified during GTB, countries' social, economic, environmental, and health characteristics, and additional information on testing and vaccination. In addition, disaggregating these data would allow for a more in-depth analysis of the determinants of COVID-19 trends and a more targeted approach to mitigating the pandemic's effects. Finally, the choice of timing may have influenced statistical analyses. Regardless of the time point, values change and have a high variance. Therefore, we took dates at their maximum growth rate and computed the post-peak period for regression analyses, allowing practical and efficient monitoring of pandemic evolution.

To our knowledge, this is the first study examining the influence of digitalization on pandemic trends worldwide. The use of digital technologies has had a positive impact on the pandemic, resulting in a faster decline in new cases and deaths. Countries with higher DAI scores were generally able to take stronger government measures. To reap the benefits of digitalization in preserving individual freedom, certain higher thresholds for digital technology adoption are a prerequisite. This paper aims to encourage readers to recognize and appreciate the major impact of digitalization on societies and health systems worldwide. Our findings underscore the need for efficient and technologically advanced infrastructures to combat future potential health crises.

Methods

Feature identification and ranking

GTB is a supervised machine learning algorithm building classification and regression models for labeled data sets (training data) [63,64]. GTB models are organized as an ensemble of sequential decision trees, evaluated as simple comparisons with binary outcomes. The tree topology allows following the inference process and identifying the most relevant features based on the frequency and position of the used features in the decision trees, referred to as feature importance [65]. Thus, GTB offers better data interpretability. A sequential elimination method was applied to identify the most relevant input features. The XGBoost implementation handled missing input features intrinsically [66]. For the identification of important features, we took countries' DAIs' and country-level demographics and health-related parameters adapted from the COVID-19 dataset (population density, median age, aged 65+, aged 70+, gross domestic product per capita, extreme poverty, cardiovascular-related death rates, diabetes prevalence, share of female and male smokers, handwashing facilities, hospital beds, life expectancy at birth, and human development index) [19]. We identified peaks of total confirmed cases and deaths of COVID-19 pm and SIs in 2020 and performed three analyses separately. We conducted a hyperparameter search to limit the complexity of GTB models [66]. Therefore, a random split into training and validation data was applied. The found set of hyperparameters was fixed for the following steps. We trained GTB models following a 10-fold cross-validation scheme to probe the data repeatably. For every GTB model of the ten training rounds, feature importance values were extracted for all input features and averaged. We discarded the least important feature and repeated the analysis until only one remained. If multiple features showed equal importance, the discarded one was randomly chosen.

Linear regression analyses

Simple linear regression showed the relationship between DAI and the change in COVID-19 cases and deaths. To express the change in COVID-19 new cases pm and new deaths pm (7-day-smoothed), we determined countries' post-peak case and death reductions. Starting at peak values for each country of log-transformed data, we fitted linear regressions to the subsequent minimum of monthly maxima between 2020 and 2021. Then, we extracted slope parameters (change of log¬10 (new cases and deaths 7-day smoothed pm) over time) for each country. Results were visualized by plotting DAI against slopes using ggplot2 [67]. Supplementary Fig. 1 shows how we estimated the decline in new deaths pm in the post-peak period, using a few country examples. The same method was applied to determine the post-peak period in new cases pm. We excluded countries that a) were unable to reduce their death

rates, b) experienced an increase in death rates that exceeded half of the peak, or c) reached the peak at the end of our dataset. Based on the defined criteria, the following countries were excluded from assessing the post-peak change in new cases (n=121) and deaths (n=123): Cambodia, Dominica, Laos, Marshall Islands, Saint Kitts and Nevis, Samoa, Solomon Islands, Timor, Vanuatu (n=9). In addition, Hong Kong and Macao were excluded from regression analyses due to missing DAI data. In another step, we added a weight to the post-peak regression analysis by measuring the difference between the previously determined monthly minimum and the maximum of new cases, respectively, deaths (Δ=max-min).

Following this, we analyzed the association between DAI and SI. To allow appropriate comparisons of SIs between countries, we normalized the actual SI by the initial SI ($SI_{initial}$) per country in the dataset ($SI_{norm}= SI_{mean}/SI_{initial}$). We then performed linear regression between DAI and SI_{norm}. Further, when plotting the SI_{mean} against DAI, we noticed three micro-trends in data. To characterize these trends with linear models, DAI was thresholded in the range [0·2: 0·8] in 0·01 steps. The range was chosen so that sufficient data remained to the left of the lower and right of the upper threshold to perform regression. The value of each DAI step was compared to the actual DAI value in each country and classified with a binary function as either "smaller" or "larger" than the current DAI-threshold. This resulted in n=61 labeled data sets with SI_{mean}, DAI, and binary class information for each of the 155 countries. We hypothesized that a local change in SI trend would show as a significant change in the statistical interaction of DAI with the binary class. Therefore, each data set was analyzed with a linear regression (SI_{mean} ~ DAI:class), and the p-value of the interaction was monitored. We observed two breaks in the inferential interaction analysis, where the p-value changed from $p \leq 0·05$ to $p \geq 0·05$ (Supplementary Table 1, Supplementary Fig. 2). These thresholds were used to classify the DAI into three classes (low, medium, and high), which were subsequently analyzed with linear regressions for micro-trend analysis. ANOVA was performed to detect mean differences in DAI estimates within income groups: low (LIE), lower-middle (LMIE), upper-middle (UMIE), and high-income economies (HIE). For post-hoc analysis, we applied a Games-Howell test to adjust for variances and corrected the p-value via Holm adjustment (Supplementary Table 2, Supplementary Fig. 4). We performed regression analyses and ANOVA using R software [68].

Explanation of findings

The review contextualized preceding findings with current literature and sought explanations for the relationship between DAI and COVID-19 values, guided by the PRISMA Extension for Scoping Reviews [69] (Supplementary Table 6). The search strategy combined alternative spellings of COVID-19 terms and pandemic-related keywords with "digit*" (Supplementary Note). This review covers English-language studies published in peer-reviewed scientific journals, including original research, review articles, and perspectives. Our search excluded news reports, comments, letters, policy briefs, and reports not fully accessible.

On May 3, 2021, we conducted a comprehensive literature search in MEDLINE/PubMed and Web of Science databases. We searched for publications published the day after the WHO China County Office was informed about case clusters in Wuhan, January 1, 2020 [1], to the day of screening. We first selected studies based on their titles and abstracts and then evaluated their full texts for inclusion. Duplicates

were removed using EndNote. Two authors (HH and LR) independently screened articles at both stages using Rayyan [70]. Literature that did not meet the predefined criteria was removed. Disagreements between reviewers were resolved through discussion until inter-rater agreement was high (Cohen's kappa coefficient >0.9). Our study team developed a screening manual with inclusion and exclusion criteria and a data extraction form (Supplementary Tables 3 and 4).

Eligible studies were thematically analyzed following Mays et al.'s narrative review approach [71]. We identified and juxtaposed common keywords to capture knowledge from different sources. Recurrent keywords were identified by repeatedly examining the reviewed articles' key pieces to ensure consistent interpretation and later merged them into themes. We mapped the keywords' occurrences in a matrix, revealing considerable overlap between themes (Supplementary Table 5). The three themes are described as follows:

The first theme encompasses the importance of digital health tools integrated into the *health care system*. These include various electronic methods such as telemedicine to support clinical staff in managing patients or electronic medical records to facilitate the digital infrastructure of medical facilities and reduce massive pressures on hospitals. By strengthening health systems, hospital management, and clinical workflows, these technologies, in one way or another, may have a direct impact on decreasing morbidity and mortality.

The second theme refers to studies on *government responses* aiming at containing and mitigating the spread of COVID-19. For example, some studies focused on tracing and detecting suspected cases through comprehensive surveillance systems, digital contact tracing, and exposure notifications. Others addressed non-pharmaceutical interventions, including stay-at-home orders, restrictions of gatherings, and mask-wearing requirements. In addition, some studies have developed evidence-based guidance for policymakers and scientists in the form of epidemiologic models and empirical frameworks.

The third theme focused on literature involving papers that provide information on *determinants facilitating or hindering public health outcomes* beyond the themes mentioned above. Those papers revolve around risk communication and "infodemic" management on national, regional, and individual levels that are crucial in shaping individual health-seeking behavior. Next to communication, we found other factors that influenced pandemic management due to complex, interacting factors such as socioeconomic, organizational, political, and privacy concerns.

Data Availability

The data that support the findings of this study are publicly available. We have compiled COVID-19 related data (i.e., number of cases and deaths and SI and demographic and epidemiological country-specific parameters) from *Our World in Data* at https://covid.ourworldindata.org/. The DAI data can be found at https://www.worldbank.org/en/publication/wdr2016/Digital-Adoption-Index and the World Bank data about countries' income groups at https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups. All reports selected for the scoping review and from which data were generated are included in the Supplementary Information.

References

Zhu, N. et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

Zhou, P. et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).

Cutler, D. M. & Summers, L. H. The COVID-19 Pandemic and the $16 Trillion Virus. *JAMA* **324**, 1495-1496, doi:10.1001/jama.2020.19759 (2020).

Chakraborty, I. & Maity, P. COVID-19 outbreak: Migration, effects on society, global environment and prevention. *Sci. Total Environ.* **728**, 138882, doi:10.1016/j.scitotenv.2020.138882 (2020).

Haug, N. et al. Ranking the effectiveness of worldwide COVID-19 government interventions. *Nat. Hum. Behav.* **4**, 1303-1312, doi:10.1038/s41562-020-01009-0 (2020).

Flaxman, S. et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature* **584**, 257-261, doi:10.1038/s41586-020-2405-7 (2020).

Hsiang, S. et al. The effect of large-scale anti-contagion policies on the COVID-19 pandemic. *Nature* **584**, 262-267, doi:10.1038/s41586-020-2404-8 (2020).

Bendavid, E., Oh, C., Bhattacharya, J. & Ioannidis, J. P. A. Assessing mandatory stay-at-home and business closure effects on the spread of COVID-19. *Eur. J. Clin. Invest.* **51**, e13484, doi:10.1111/eci.13484 (2021).

Bavel, J. J. V. et al. Using social and behavioural science to support COVID-19 pandemic response. *Nat. Hum. Behav.* **4**, 460-471, doi:10.1038/s41562-020-0884-z (2020).

Whitelaw, S., Mamas, M. A., Topol, E. & Van Spall, H. G. C. Applications of digital technology in COVID-19 pandemic planning and response. *Lancet Digit. Health* **2**, e435-e440, doi:10.1016/S2589-7500(20)30142-4 (2020).

Cone, L. et al. Pandemic Acceleration: Covid-19 and the emergency digitalization of European education. *Eur. Educ. Res. J.*, doi:10.1177/14749041211041793 (2021).

Amankwah-Amoah, J., Khan, Z., Wood, G. & Knight, G. COVID-19 and digitalization: The great acceleration. *J. Bus. Res.* **136**, 602-611, doi:10.1016/j.jbusres.2021.08.011 (2021).

Dighe, A. et al. Response to COVID-19 in South Korea and implications for lifting stringent interventions. *BMC Med.* **18**, 321, doi:10.1186/s12916-020-01791-8 (2020).

Chen, H., Shi, L., Zhang, Y., Wang, X. & Sun, G. A cross-country core strategy comparison in China, Japan, Singapore and South Korea during the early COVID-19 pandemic. *Glob. Health* **17**, 22, doi:10.1186/s12992-021-00672-w (2021).

Zeng, K., Bernardo, S. N. & Havins, W. E. The Use of Digital Tools to Mitigate the COVID-19 Pandemic: Comparative Retrospective Study of Six Countries. *JMIR Public Health Surveill.* **6**, e24598, doi:10.2196/24598 (2020).

Fraser, C., Riley, S., Anderson, R. M. & Ferguson, N. M. Factors that make an infectious disease outbreak controllable. *PNAS* **101**, 6146-6151, doi:10.1073/pnas.0307506101 (2004).

Ferretti, L. et al. Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science* **368**, doi:10.1126/science.abb6936 (2020).

Faraj, S., Renno, W. & Bhardwaj, A. Unto the breach: What the COVID-19 pandemic exposes about digitalization. *Information and Organization* **31**, 100337, doi:10.1016/j.infoandorg.2021.100337 (2021).

Ritchie, H. et al. Coronavirus pandemic (COVID-19). *Our world in data* (2020).

Hale, T. et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat. Hum. Behav.* **5**, 529-538, doi:10.1038/s41562-021-01079-8 (2021).

The World Bank, *Digital Adoption Index*, https://www.worldbank.org/en/publication/wdr2016/Digital-Adoption-Index (2016).

The World Bank. *World development report 2016: Digital dividends*. (World Bank Publications, Washington DC, 2016).

ECDC. Guidelines for non-pharmaceutical interventions to reduce the impact of COVID-19 in the EU/EEA and the UK. . (2020).

Anthony Jnr, B. Implications of telehealth and digital care solutions during COVID-19 pandemic: a qualitative literature review. *Inform. Health. Soc. Care* **46**, 68-83 (2021).

Chatterjee, P., Tesis, A., Cymberknop, L. J. & Armentano, R. L. Internet of Things and Artificial Intelligence in Healthcare During COVID-19 Pandemic-A South American Perspective. *Front. Public Health* **8**, 600213 (2020).

Liaw, S. T. et al. Primary Care Informatics Response to Covid-19 Pandemic: Adaptation, Progress, and Lessons from Four Countries with High ICT Development. *Yearb. Med. Inform.* **30**, 44-55 (2021).

Scott, B. K. et al. Advanced Digital Health Technologies for COVID-19 and Future Emergencies. *Telemed. J. E. Health* **26**, 1226-1233, doi:10.1089/tmj.2020.0140 (2020).

Wood, A. et al. Linked electronic health records for research on a nationwide cohort of more than 54 million people in England: data resource. *Bmj* **373**, n826 (2021).

Xu, T. & Yang, R. COVID-19 Epidemic and Public Health Measures in China. *J. Epidemiol. Glob. Health* **10**, 118-123, doi:10.2991/jegh.k.200421.001 (2020).

Ye, Q., Zhou, J. & Wu, H. Using Information Technology to Manage the COVID-19 Pandemic: Development of a Technical Framework Based on Practical Experience in China. *JMIR Med. Inform.* **8**, e19515 (2020).

Yan, A., Zou, Y. & Mirchandani, D. A. How hospitals in mainland China responded to the outbreak of COVID-19 using information technology-enabled services: An analysis of hospital news webpages. *J. Am. Med. Inform. Assoc.* **27**, 991-999 (2020).

Boeing, P. & Wang, Y. H. Decoding China's COVID-19 'virus exceptionalism': Community-based digital contact tracing in Wuhan. *R & D Management* **51**, 339-351 (2021).

Storeng, K. T. & de Bengy Puyvallée, A. The Smartphone Pandemic: How Big Tech and public health authorities partner in the digital response to Covid-19. *Glob. Public Health* **16**, 1482-1498 (2021).

Nageshwaran, G., Harris, R. C. & Guerche-Seblain, C. E. Review of the role of big data and digital technologies in controlling COVID-19 in Asia: Public health interest vs. privacy. *Digit. Health* **7**, 20552076211002953 (2021).

Anttiroiko, A. V. Successful Government Responses to the Pandemic: Contextualizing National and Urban Responses to the COVID-19 Outbreak in East and West. *International Journal of E-Planning Research* **10**, 1-17 (2021).

Skoll, D., Miller, J. C. & Saxon, L. A. COVID-19 testing and infection surveillance: Is a combined digital contact-tracing and mass-testing solution feasible in the United States? *Cardiovasc. Digit. Health J.* **1**, 149-159 (2020).

Islam, M. N. & Islam, A. A Systematic Review of the Digital Interventions for Fighting COVID-19: The Bangladesh Perspective. *Ieee Access* **8**, 114078-114087 (2020).

Martin, T. et al. Demystifying COVID-19 Digital Contact Tracing: A Survey on Frameworks and Mobile Apps. *Wireless Communications & Mobile Computing* **2020** (2020).

Leung, K., Wu, J. T. & Leung, G. M. Real-time tracking and prediction of COVID-19 infection using digital proxies of population mobility and mixing. *Nat. Commun.* **12**, 1501 (2021).

Cencetti, G. et al. Digital proximity tracing on empirical contact networks for pandemic control. *Nat. Commun.* **12**, 1655, doi:10.1038/s41467-021-21809-w (2021).

The World Bank, *World Bank Country and Lending Groups*; [retrieved 2021 Sep]. Available from: https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups.

Guan, W.-J. et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N. Engl. J. Med.* **382**. 1708-1720 (2020).

Zhou, F. et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054-1062 (2020).

Huang, C. et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506 (2020).

Nam, H.-S. et al. High fatality rates and associated factors in two hospital outbreaks of MERS in Daejeon, the Republic of Korea. *Int. J. Infect Dis.* **58**, 37-42, doi:10.1016/j.ijid.2017.02.008 (2017).

Tan, T. et al. Association between high serum total cortisol concentrations and mortality from COVID-19. *Lancet Diabetes Endocrinol.* **8**, 659-660, doi:10.1016/S2213-8587(20)30216-3 (2020).

Clark, A. et al. Global, regional, and national estimates of the population at increased risk of severe COVID-19 due to underlying health conditions in 2020: a modelling study. *Lancet Glob. Health* **8**, e1003-e1017, doi:10.1016/s2214-109x(20)30264-3 (2020).

Wang, D. et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA* **323**, 1061-1069 (2020).

Pardhan, S. & Drydakis, N. Associating the Change in New COVID-19 Cases to GDP per Capita in 38 European Countries in the First Wave of the Pandemic. *Front. Public Health* **8** (2020).

Finch, W. H. & Hernández Finch, M. E. Poverty and Covid-19: rates of incidence and deaths in the United States during the first 10 weeks of the pandemic. *Front. Sociol.* **5**, 47 (2020).

OECD. *The territorial impact of COVID-19: Managing the crisis across levels of government*; [retrieved 2022 March]. Available from: https://www.oecd.org/coronavirus/policy-responses/the-

territorial-impact-of-covid-19-managing-the-crisis-across-levels-of-government-d3e314e1/#boxsection-d1e10825 (2020).

Jurzik, E. et al. COVID-19 and Inequality in Asia: Breaking the Vicious Cycle. *Available at SSRN 3744684* (2020).

Rimmer, A. Covid-19: Disproportionate impact on ethnic minority healthcare workers will be explored by government. *BMJ* **369**, m1562, doi:10.1136/bmj.m1562 (2020).

Budd, J. et al. Digital technologies in the public-health response to COVID-19. *Nat. Med.* **26**, 1183-1192 (2020).

World Health Organization. *Global Strategy on digital health 2020-2025.* (World Health Organization, Geneva, 2021).

Sieck, C. J. et al. Digital inclusion as a social determinant of health. *NPJ Digit. Med.* **4**, 52, doi:10.1038/s41746-021-00413-8 (2021).

Reddick, C. G., Enriquez, R., Harris, R. J. & Sharma, B. Determinants of broadband access and affordability: An analysis of a community survey on the digital divide. *Cities* **106**, 102904 (2020).

Meskó, B., Drobni, Z., Bényei, É., Gergely, B. & Győrffy, Z. Digital health is a cultural transformation of traditional healthcare. *Mhealth* **3**, 38-38, doi:10.21037/mhealth.2017.08.07 (2017).

An, B. Y. & Tang, S.-Y. Lessons From COVID-19 Responses in East Asia: Institutional Infrastructure and Enduring Policy Instruments. *The American Review of Public Administration* **50**, 790-800, doi:10.1177/0275074020943707 (2020).

Lin, L. & Hou, Z. Y. Combat COVID-19 with artificial intelligence and big data. *J. Travel. Med.* **27** (2020).

Lewis, D. Why many countries failed at COVID contact-tracing - but some got it right. *Nature* **588**, 384-388 (2020).

Parker, M. J., Fraser, C., Abeler-Dörner, L. & Bonsall, D. Ethics of instantaneous contact tracing using mobile phone apps in the control of the COVID-19 pandemic. *J. Med. Ethics* **46**, 427-431, doi:10.1136/medethics-2020-106314 (2020).

Friedman, J. H. Greedy function approximation: A gradient boosting machine. *Ann. Stat.* **29**, 1189-1232, 1144 (2001).

Friedman, J. H. Stochastic gradient boosting. *Computational Statistics & Data Analysis* **38**, 367-378, doi:10.1016/S0167-9473(01)00065-2 (2002).

Natekin, A. & Knoll, A. Gradient boosting machines, a tutorial. *Front. Neurorobot.* **7**, doi:10.3389/fnbot.2013.00021 (2013).

Chen, T. & Guestrin, C. XGBoost: A Scalable Tree Bossting System. *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 785-794, doi:10.1145/2939672.2939785 (Association for Computing Machinery, San Francisco, 2016).

Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*, (Springer-Verlag, New York, 2016).

R Core Team. *R: A language and environment for statistical computing.* (R Foundation for Statistical Computing, Vienna, 2020).

Tricco, A. et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann. Intern. Med.* **169**, 467-473, doi:10.7326/m18-0850 %m 30178033 (2018).

Ouzzani, M., Hammady, H., Fedorowicz, Z. & Elmagarmid, A. Rayyan—a web and mobile app for systematic reviews. *Systematic reviews* **5**, 1-10 (2016).

Mays, N., Pope, C. & Popay, J. Systematically reviewing qualitative and quantitative evidence to inform management and policy-making in the health field. *J. Health Serv. Res. Policy* **10**, 6-20, doi:10.1258/1355819054308576 (2005).

Acknowledgements

This work is funded by the German Federal Ministry of Education and Research (BMBF, grant no.: 01GP1910A).

Author contributions

Conception and design: H.H. and F.K.; acquisition of data: H.H., F.M., and S.R.T.; analysis and interpretation of data: H.H., L.R., F.M., V.S., S.R.T., and F.K.; writing of the manuscript: H.H.; study supervision: F.K. All authors reviewed the manuscript and gave their final approval.

Competing interests

All authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to S.R.T. or F.K.