Tradeoff between speed and reproductive number in pathogen evolution

Andreas Eilersen,[1,2,*] Bjarke Frost Nielsen,[1,2] and Kim Sneppen[2]

[1]*PandemiX Center, Roskilde University*
[2]*Niels Bohr Institute, University of Copenhagen*
(Dated: January 8, 2023)

The rapid succession of new variants of SARS-CoV-2 emphasizes the need to understand the factors driving pathogen evolution. Here, we investigate a possible tradeoff between the speed at which a disease progresses and its reproductive number. Using an SEIR framework, we show that in the exponential growth phase of an epidemic, there is an optimal disease duration that balances the advantage of a fast disease progression with that of causing many secondary infections. This result offers one possible explanation for the ever shorter generation times of novel variants of SARS-CoV-2, as we progressed from the original strain to the Alpha, Delta, and finally Omicron variants. In the endemic state, the optimum disappears and longer disease duration becomes advantageous for the pathogen. However, selection pressures depend on context: mitigation strategies such as quarantine of infected individuals may slow down the evolution towards longer-lasting, more infectious variants. This work then suggests that, in the future, the trend towards shorter generation times may reverse, and SARS-CoV-2 may instead evolve towards longer-lasting variants.

Keywords: epidemiology, evolution, latency, computational, model

INTRODUCTION

Since the emergence of SARS-CoV-2, multiple variants of the virus with faster transmission dynamics have arisen. The variants have supplanted each other in successive waves, with variants with ever higher transmission rates and/or shorter generation times replacing older, slower variants [1, 2]. This unfolding evolutionary race suggests a dynamic that can be explored through modeling.

Some work has already been done on modeling the evolution of the infection profile of SARS-CoV-2 and other pathogens with similar generation times. Saad-Roy *et al.* studied the evolution of a presymptomatic infectious state under the assumption that such a state is less infectious [3], and in the context of superinfection and within-host competition [4]. In addition, the relationship between the duration of a disease or parasitic infection and the infection rate has been studied under the assumption of a tradeoff between the two given by some functional relationship [5, 6]. Porco *et al.* [7] investigated the effects of treatment and other interventions on disease evolution under the assumption of a similar tradeoff. Analogous studies have been done on other ecological relationships, such as predation [8, 9]. Finally, Park *et al.* [10] have studied the interplay between disease infectivity and speed with a focus on mitigation rather than evolution. However, the possibility that a longer infectious period might be an evolutionary advantage for a disease only up to a certain threshold has not been studied in detail.

* andreaseilersen@nbi.ku.dk

In this article, we focus on the tradeoff between the duration of the individual infections and the number of secondary cases that each infected individual generates. We assume that infected individuals transmit the disease at a constant rate during the infectious period. This means that a long disease duration should lead to a higher effective reproductive number, R_e, that is, to more secondary infections. On the other hand, a long disease duration might also be a disadvantage to the disease, as it may be associated with a long latency and thereby a slow epidemic progression. This is particularly the case if one assumes direct proportionality between the duration of the latency time of a disease and its infectious period. Table 1 suggests that across diseases, longer infectious periods indeed correlate with longer latent periods. In a susceptible-exposed-infectious-recovered (SEIR) type compartmental model, which we will consider in this study, the latent period corresponds to the E-state. We will derive relations and carry out epidemic simulations based on systems of ordinary differential equations (ODEs) to investigate the condition for an optimum disease duration.

MODEL SETUP

We assume that the transmission rate β of a disease is constant throughout the duration of the infectious period T, giving a linear relationship between disease duration and number of secondary cases. While strict proportionality does not necessarily hold, [39, 40] a positive, monotonic relation between the two is expected since since a longer infectious period leads to more opportunities for passing on the infection.

Throughout this work, we will distinguish between the latent and incubation periods of a disease. The latent period is the time from the initial infection until the patient becomes infectious. In contrast, the incubation period is the time from infection until the onset of symptoms.

Disease	Latency time	Infectious period	c	k	R_0
COVID-19 (wild-type)	≤ 4.0 [11]	$6.5 - 9.5$ [12]	$0.4 - 0.6$	5 [13]	2.9 ± 0.5 [14]
Influenza H1N1	2.6 [15]	3.4 [15]	0.8	2 [16]	1.6 [17]
Smallpox	14.6 [18]	8.6 [18]	1.7	35 [19]	$3.5 - 6$ [18]
Measles	~ 8.5 [20] [a]	~ 8 [21]	~ 1	15 [22]	13 [23]
Ebola	12.7 [24]	7.2 [25] [b]	1.8	10 [24]	$1.4 - 1.8$ [26]
SARS	~ 10 [27]	~ 12 [27] [c]	~ 0.8	2 [28]	$2 - 4$ [29]
Yellow fever	4.3 [30]	$1 - 4$ [31]	$1.1 - 4.3$	8 [30]	2.4 [31]
Cholera	1.7 [32]	2.0 [33]	0.9	2 [32]	$1.1 - 2.6$ [34]
Syphilis	$9 - 90$ [35]	~ 365 [36]	$0.02 - 0.2$	-	-
Rabies	$10 - 700$ [37]	< 7 [38]	$1.4 - 100$	-	-

TABLE I. Duration (days) of the latent and infectious periods for some infectious diseases. c is the ratio of latency time to the infectious period of the disease, while $k = k_{incubation}$ is the shape factor for a Gamma distribution which we fit approximately to the measured distributions of incubation times found in the cited literature (the coefficient of variation of these distributions is given by $CV^2 = 1/k$). Note that we show the k-values for the incubation periods as opposed to the latent periods, which we assume are similarly distributed. The incubation period is the time from infection to symptom onset, while the latent period is the time from infection to onset of infectiousness. The last column shows the estimated reproductive number.

[a] As infectiousness begins four days before the onset of a rash, latency time is calculated as incubation period minus four days.

[b] The cited study reports separate infectious periods for survivors and deceased patients. We have here indicated the average.

[c] Incubation period is reported as 4.6 days. Here, we define the infectious period by the requirement that at least 50 % of patients secrete measurable quantities of the virus. In that case, infectiousness begins on day five after the onset of symptoms and ends on day 17, yielding a latency time of 9.6 days.

The SEIR model reads:

$$\frac{\mathrm{d}S}{\mathrm{d}t} = -\beta\, S\, I \tag{1}$$

$$\frac{\mathrm{d}E}{\mathrm{d}t} = \beta\, S\, I \,-\, \frac{1}{\tau}\, E \tag{2}$$

$$\frac{\mathrm{d}I}{\mathrm{d}t} = \frac{1}{\tau}\, E \,-\, \frac{1}{T}\, I \tag{3}$$

$$\frac{\mathrm{d}R}{\mathrm{d}t} = \frac{1}{T}\, I. \tag{4}$$

where S, E, I, and R are susceptible, exposed (but non-infectious), infectious, and recovered compartments respectively. β is the transmission rate per unit time, τ is the average duration of the pre-infectious exposed period, and T is the duration of the infectious period. We take the total population of the system to be fixed at $N = 1$ and assume that the time-scale of the entire scenario is short enough that vital dynamics - births and deaths - can be neglected.

We will also investigate the effects of variability in the durations of the E and I states. Usually, an exponential distribution is assumed in SEIR models. However, by subdividing each of the compartments E and I into k equally long sub-compartments we instead obtain a Gamma distribution of latency times and infectious periods with a shape parameter k. The shape parameter is related to the coefficient of variation CV by $k = 1/\sqrt{CV}$. Thus, a greater k corresponds to lower person-to-person variation in the durations of the E and I states. Since

we assume integer values of k, this distribution coincides with an Erlang distribution. The above equations for the simple SEIR model correspond to letting the shape parameter take on the value $k = 1$, yielding the familiar exponential case.

When considering the evolution of the disease in an endemic state with a high degree of existing immunity in the population, we slightly modify the above SEIR model. To make the endemic state possible, we allow individuals to lose immunity at a rate ω, corresponding to an SEIRS model of disease progression. This model is then solved numerically, including multiple co-circulating variants with different disease durations T. Thereby we emulate the natural competition between variants.

Finally, mitigation by isolation of infected individuals is implemented by the addition of a quarantine rate q and corresponding noninfectious quarantine compartment Q to the SEIRS model. This represents how individuals have some chance of becoming symptomatic, being contact traced, or otherwise being diagnosed and isolated for each day of illness. We expect that individuals suffering from a very long-lasting infectious disease will eventually self-quarantine. The resulting equations thus become

FIG. 1. Direct simulation of the initial growth of an epidemic for different disease durations T. When disease duration is short, the epidemic grows faster as T increases because each infected individual can infect more people. However, for larger values of T, the cost of the longer latent period increases and leads to slower exponential growth. Here, $\beta = 1$ and $c = \tau/T = 1$, corresponding to assuming equal duration of the E and I states. The initial exposed and infected fractions of the population are $E(0) = 10^{-12}$ and $I(0) = 10^{-12}$.

$$\frac{\mathrm{d}S}{\mathrm{d}t} = \omega\, R - \sum_i^{n_{var}} \beta\, S\, I_i \qquad (5)$$

$$\frac{\mathrm{d}E_i}{\mathrm{d}t} = \beta\, S\, I_i - \frac{1}{\tau_i}\, E_i \qquad (6)$$

$$\frac{\mathrm{d}I_i}{\mathrm{d}t} = \frac{1}{\tau_i}\, E_i - \left(\frac{1}{T_i} + q\right) I_i \qquad (7)$$

$$\frac{\mathrm{d}Q_i}{\mathrm{d}t} = q\, I_i - \frac{1}{T_i}\, Q_i \qquad (8)$$

$$\frac{\mathrm{d}R}{\mathrm{d}t} = \sum_i^{n_{var}} \frac{1}{T_i}\, (I_i + Q_i) - \omega\, R. \qquad (9)$$

Here we have introduced an index i to indicate the possibility of including n_{var} different variants, all of which we assume to have perfect cross immunity with respect to the others. This will become important when simulating competition between strains with different disease durations and latency times.

RESULTS

Optimum disease duration for exponential growth

In the exponential growth phase of an epidemic, the growth rate may be determined by linearizing the system of equations (1-4) around the disease-free equilibrium. The epidemic growth rate r is then the largest eigenvalue of the Jacobian [41]:

$$r = \frac{-(1+c) + \sqrt{(1+c)^2 - 4(1-\beta T)c}}{2cT}, \qquad (10)$$

where $c \equiv \tau/T$, and we have chosen the physically allowable positive branch. The above function has a maximum for the duration T given by

$$R_{0,fastest} = \beta T_{fastest} = 2 + \sqrt{c} + \frac{1}{\sqrt{c}}, \qquad (11)$$

In the limit of $k \to \infty$, the durations of the exposed and infectious stages are deterministic. If we further neglect the variation in the timing of disease transmission, disease generations thus last exactly $\tau + T$. Under these approximations, the expression for the number of infected in the exponential growth phase approaches

$$I(t) = R_0^{t/(T+\tau)} = (\beta T)^{t/((1+T)c)}. \qquad (12)$$

Maximising the exponential growth rate as a function of T in this limit, we obtain the finite value

$$T_{fastest} = e/\beta, \qquad (13)$$

where e denotes Euler's number.

In the exponential growth phase, the variant with the highest growth rate will quickly come to dominate. An illustration of this phase for different disease durations can be seen in Fig. 1, while a plot of the growth rate as a function of T for various values of c is shown in Fig. 2(a). As shown, the exponential growth rate is much higher for lower values of c. Furthermore, there is good agreement between our analytical and numerical calculations. When increasing c, the maximum growth rate decreases strongly, and the optimum with respect to T becomes less clear.

In the equation for r (Eq. (10)), it is assumed that the probability distribution for the duration of the latency times and infectious periods of each individual is an exponential distribution, corresponding to a shape parameter $k = 1$. We wish to explore how the degree of variability in the latent and infectious period affects the optimal value disease duration and the resulting growth rate. We do this numerically by solving the SEIR equations with a Gamma distributed latent and infectious period and varying the shape parameter.

In Fig. 2(b), the effects of varying k are shown. At $c = 1$, the effect of an increase in k is modest. However, when the latent period is much longer than the infectious period ($c = 10$), a more definite duration ($k = 10$) is associated with a clearly lower resultant growth rate. A more complete overview of growth rates and maxima as functions of c for different k is given in Fig. 3.

Overall, Fig. 3 illustrates that the maximal daily growth rate decreases monotonically as a function of c, reflecting a longer latency period. Increasing k, i.e., making the distribution of latency times more sharply peaked,

(a)

(b)

(a)

(b)

FIG. 2. Growth rate as a function of disease duration T as simulated, compared with the growth rate derived in Eq. (10). Here, we set $\beta = 1/\text{day}$. One observes that the growth rate has a maximum at the duration given by Eq. (11) (giving $T = 4$ for $c = 1$ and $T = 5.48$ for $c = 10$). Panel (b) shows the corresponding plot when considering a larger shape factor k for the duration of latent and infectious periods.

also leads to a slight decrease in maximal growth rate for $c > 1$.

Fig. 3 (b) shows that the value of T which maximises growth rate r, denoted $T_{fastest}$, is highest in the limit of $c \to 0$. At low values of k, $T_{fastest}$ exhibits a dependence on c, with a local minimum around $c = 1$ beyond which it grows as c increases. $T_{fastest}(c)$ for $k = 1$ reproduces Eq. (11). As k is increased, $T_{fastest}$ becomes increasingly independent of c and approaches a value of approximately e, as predicted by Eq. (13) for $\beta = 1$. The good agreement is remarkable given the simplifications contained in that equation.

In a situation where the disease is growing exponentially, e.g., when an epidemic is breaking out or control measures are failing, the variant with the fastest exponential growth rate will win as illustrated in Figs. 1 and 2. This, however, only holds transiently and we will see

FIG. 3. Growth rates and optimal disease duration as a function of c for various k. (a) shows the maximal exponential growth rate r of an epidemic pathogen given different values of c and k. We see that growth rates are highest for low c and also decrease very slightly with k. (b) shows the disease duration $T_{fastest}$ that maximizes the growth rate. For $\beta = 1$ this is also the value of R_0 that maximizes the growth rate. This is equal to the predictions from Eq. (10) for $k = 1$ and drops as k increases. At higher k, $T_{fastest}$ approaches e as predicted in the limit of $k \to \infty$.

that the situation is reversed when considering the endemic state in a model with immunity loss.

The endemic state

In the endemic state, the disease prevalence is sustained at an approximately constant level. In our model, this is done by introducing a small rate of loss from the recovered (R) state, corresponding to the waning of acquired immunity. Furthermore, we will consider a quarantine rate q that quantifies the probability per day that an individual in the I state goes into isolation. As previously described, this is represented by a noninfectious

quarantine compartment Q which infected persons may leave upon recovery.

The results are illustrated in Fig. 4. The figure shows that in this case, longer-lasting variants always outcompete shorter-lasting ones. If the system starts out with an equal fraction of the population infected with each variant, we thus expect to see a succession of variant takeovers. Longer-lasting variants take longer to grow, but eventually always end up taking over due to their higher basic reproductive number. However, by increasing the quarantine rate q (Fig. 4(b)), we see that this development may be slowed significantly. We have also examined the sensitivity to variations in the immunity loss rate ω, but varying this rate has little effect, except on the magnitude of outbreaks.

DISCUSSION

Our analysis illustrates that being fast-acting can be an evolutionary advantage for a pathogen, even if it comes at the cost of a lower reproductive number. However, this is only the case in the initial exponential growth phase of the epidemic. In the endemic phase the longer-lasting variants which have a higher R_0 will always eventually come to dominate. This is the case regardless of interventions such as quarantine, though evolution towards long-lasting, more infectious variants may be slowed by quarantining infectious individuals.

We expect the two scenarios modelled here to be applicable across most of the trajectory of a real-world epidemic. With regard to quarantine, we would expect the onset of symptoms to increase the chance that individuals stay home or are bedridden, effectively self-quarantining. This is likely the case regardless of large-scale mitigation policies.

In addition, the exponential growth scenario is not necessarily limited to the short initial stage of the epidemic. In the case of the COVID-19 pandemic, mitigation efforts in various locations often kept the local R_e at or below 1. When such efforts failed or were relaxed, local epidemics entered a new exponential growth phase. Our results may thus help understand the successive shifts in dominance from the Wuhan strain, to the Alpha and Delta variants, and finally the various Omicron subvariants. The Delta variant has been shown to have a somewhat shorter incubation period and significantly shorter generation time than the ancestral strain [42–44]. Hart et al. [45] measure a generation time of 5.5 days for the Alpha variant and 4.6 days for the Delta variant. Omicron was even faster, with a reported serial interval of only 2.2 days [46]. The analysis in Abbott et al. further supports the tendency of faster disease progression for the latter SARS-CoV-2 variant, although Pung et al. disputes whether generation times of Delta were in fact significantly lower than for the Alpha variant [2, 47].

Each new SARS-CoV-2 variant has been accompanied by changes in transmission rates as well as generation

(a)

(b)

FIG. 4. The figure shows the competition between variants of different duration T as the disease progresses to the endemc phase. The simulations use fixed $\beta = 1$, $c = 1$, and $\omega = 10^{-3}$. The plots show the total exposed, infected, and quarantined infected population $E + I + Q$. (a) An ODE-based simulation of six different variants of duration $T = 10, 50, 100, 200, 300, 400$ days without quarantine. We see that the longer-lasting variants replace the shorter-lived ones. (b) The same simulation with a quarantine rate of $q = 0.1$ per day of infectious illness. The longer-lasting variants still win, but quarantine slows down the evolution.

times. The analysis presented here focuses on the time aspect while ignoring the obvious evolutionary gain a pathogen may obtain by increasing the infection rate β. We found that the growth rates in the exponential growth phase of an epidemic indeed seem to be optimal for rather short generation times, which agrees with the obervation that new SARS-CoV-2 variants tend to be faster than older variants. This overall tendency during 2020-2021 with ever faster virus variants may later be broken, however. Our simulations demonstrate that this is likely if the pandemic reaches a more endemic state where slower variants of the disease gain in fitness, as mitigation and

quarantine efforts are dropped. Even under the assumption of quarantine measures, the optimal strategy should shift towards a longer disease duration in the endemic state, albeit more slowly.

Our analysis focuses on pathogens like SARS-CoV-2, which are transmitted through social contact and act on a relatively short timescale. There are of course also pathogens that act on timescales longer than those predicted here. This is for example true of sexually transmitted infections such as syphilis and HIV which cause lifelong infection ($c << 1$). These infections violate the assumption of latency times being roughly proportional to disease duration, and thus are not captured by our model. In the case of these STIs, the two aspects of pathogen dynamics are essentially decoupled in the body.

Alternative approaches have been used to investigate the reasons for the variation in disease duration observed in the real world. These have predicted the existence of several "regimes" of disease duration [48]: from fast-acting childhood infections in situations with high contact rates to lifelong infections in low-contact situations.

It is of course both idealized and highly simplified to assume that latency times scale proportionally with infectious periods. As mentioned in the introduction, there is however some support for our assumption of a relationship between the two quantities. In Table 1, we show examples of latency times and infectious period durations for various diseases, along with the shape parameters k of the associated distributions of incubation period duration. We here assume these k values to be the same as for the latency time distributions of the diseases. The values of c are indeed often on the order of 1, except for zoonotic diseases like rabies or sexually transmitted infections like syphilis. Interestingly, the data show a large variation in how sharply peaked the incubation periods are, with k varying from < 5 in respiratory diseases such as SARS, COVID-19, and influenza, to an estimated 35 in smallpox.

The findings of this article highlight the context-dependence of evolutionary fitness as it pertains to disease duration and latency. Our results may help explain some of observed dynamics of emerging SARS-CoV-2 variants. In a wider perspective, our work also sheds some light on the apparent division of infectious diseases into a group of quite fast diseases characterized by epidemic outbreaks, and another group which are slow with long latent periods and an endemic pattern of infection.

ACKNOWLEDGMENTS

We wish to thank Lone Simonsen, Viggo Andreasen, and Nils Christian Stenseth for enlightening discussions.

KS and AE have received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement no. 740704). BFN received funding from the Carlsberg Foundation under its Semper Ardens programme (grant no. CF20-0046), and AE received funding from NordForsk under the Nordic Programme for Interdisciplinary Research (grant no. 104910).

DATA AVAILABILITY

The code used to generate the plots shown in this article is available on GitHub under the URL `https://github.com/gks973/Evolution_of_speed` or in a permanent archived version at `https://doi.org/10.5281/zenodo.7512664`.

AUTHOR CONTRIBUTIONS STATEMENT

KS proposed the model. AE, KS and BFN derived the expressions, and BFN and AE wrote the code for simulations. KS, AE, and BFN wrote the manuscript.

[1] W. S. Hart, E. Miller, N. J. Andrews, P. Waight, P. K. Maini, S. Funk, and R. N. Thompson, Generation time of the alpha and delta sars-cov-2 variants: an epidemiological analysis, The Lancet Infectious Diseases **22**, 603 (2022).

[2] S. Abbott, K. Sherratt, M. Gerstung, and S. Funk, Estimation of the test to test distribution as a proxy for generation interval distribution for the omicron variant in england, medRxiv (2022).

[3] C. M. Saad-Roy, N. S. Wingreen, S. A. Levin, and B. T. Grenfell, Dynamics in a simple evolutionary-epidemiological model for the evolution of an initial asymptomatic infection stage, Proceedings of the national academy of sciences **117**, 11541 (2020).

[4] C. M. Saad-Roy, B. T. Grenfell, S. A. Levin, L. Pellis, H. B. Stage, P. Van Den Driessche, and N. S. Wingreen, Superinfection and the evolution of an initial asymptomatic stage, Royal Society open science **8**, 202212 (2021).

[5] R. M. Anderson and R. M. May, Coevolution of hosts and parasites, Parasitology **85**, 411 (1982).

[6] S. Alizon, A. Hurford, N. Mideo, and M. Van Baalen, Virulence evolution and the trade-off hypothesis: history, current state of affairs and the future, Journal of evolutionary biology **22**, 245 (2009).

[7] T. C. Porco, J. O. Lloyd-Smith, K. L. Gross, and A. P. Galvani, The effect of treatment on pathogen virulence, Journal of Theoretical Biology **233**, 91 (2005).

[8] J. Reed and N. C. Stenseth, On evolutionarily stable strategies, Journal of theoretical biology **108**, 491 (1984).

[9] K. Grunert, H. Holden, E. R. Jakobsen, and N. C. Stenseth, Evolutionarily stable strategies in stable and periodically fluctuating populations: The rosenzweig–macarthur predator–prey model, Proceedings of the National Academy of Sciences **118** (2021).

[10] S. W. Park, B. M. Bolker, S. Funk, C. J. E. Metcalf, J. S. Weitz, B. T. Grenfell, and J. Dushoff, The importance of the generation interval in investigating dynamics and control of new sars-cov-2 variants, Journal of the Royal Society Interface **19**, 20220173 (2022).

[11] H. Nishiura, N. M. Linton, and A. R. Akhmetzhanov, Serial interval of novel coronavirus (covid-19) infections, International journal of infectious diseases **93**, 284 (2020).

[12] A. W. Byrne, D. McEvoy, A. B. Collins, K. Hunt, M. Casey, A. Barber, F. Butler, J. Griffin, E. A. Lane, C. McAloon, *et al.*, Inferred duration of infectious period of sars-cov-2: rapid scoping review and analysis of available evidence for asymptomatic and symptomatic covid-19 cases, BMJ open **10**, e039856 (2020).

[13] Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. Leung, E. H. Lau, J. Y. Wong, *et al.*, Early transmission dynamics in wuhan, china, of novel coronavirus–infected pneumonia, New England journal of medicine (2020).

[14] M. A. Billah, M. M. Miah, and M. N. Khan, Reproductive number of coronavirus: A systematic review and meta-analysis based on global level evidence, PloS one **15**, e0242128 (2020).

[15] A. R. Tuite, A. L. Greer, M. Whelan, A.-L. Winter, B. Lee, P. Yan, J. Wu, S. Moghadas, D. Buckeridge, B. Pourbohloul, *et al.*, Estimated epidemiologic parameters and morbidity associated with pandemic h1n1 influenza, Cmaj **182**, 131 (2010).

[16] H. Nishiura and H. Inaba, Estimation of the incubation period of influenza a (h1n1-2009) among imported cases: addressing censoring using outbreak data at the origin of importation, Journal of theoretical biology **272**, 123 (2011).

[17] C. Fraser, C. A. Donnelly, S. Cauchemez, W. P. Hanage, M. D. Van Kerkhove, T. D. Hollingsworth, J. Griffin, R. F. Baggaley, H. E. Jenkins, E. J. Lyons, *et al.*, Pandemic potential of a strain of influenza a (h1n1): early findings, science **324**, 1557 (2009).

[18] R. Gani and S. Leach, Transmission potential of smallpox in contemporary populations, Nature **414**, 748 (2001).

[19] H. Nishiura, Determination of the appropriate quarantine period following smallpox exposure: an objective approach using the incubation period distribution, International journal of hygiene and environmental health **212**, 97 (2009).

[20] J. Lessler, N. G. Reich, R. Brookmeyer, T. M. Perl, K. E. Nelson, and D. A. Cummings, Incubation periods of acute respiratory viral infections: a systematic review, The Lancet infectious diseases **9**, 291 (2009).

[21] CDC, *CDC yellow book 2020* (Oxford University Press, 2019) Chap. Measles (Rubeola), pp. 287–291.

[22] D. Klinkenberg and H. Nishiura, The correlation between infectivity and incubation period of measles, estimated from households with two cases, Journal of theoretical biology **284**, 52 (2011).

[23] F. M. Guerra, S. Bolotin, G. Lim, J. Heffernan, S. L. Deeks, Y. Li, and N. S. Crowcroft, The basic reproduction number (r0) of measles: a systematic review, The Lancet Infectious Diseases **17**, e420 (2017).

[24] M. Eichner, S. F. Dowell, and N. Firese, Incubation period of ebola hemorrhagic virus subtype zaire, Osong Public Health and Research Perspectives **2**, 3 (2011).

[25] G. E. Velásquez, O. Aibana, E. J. Ling, I. Diakite, E. Q. Mooring, and M. B. Murray, Time from infection to disease and infectiousness for ebola virus disease, a systematic review, Clinical Infectious Diseases **61**, 1135 (2015).

[26] Z. Wong, C. Bui, A. Chughtai, and C. Macintyre, A systematic review of early modelling studies of ebola virus disease in west africa, Epidemiology & Infection **145**, 1069 (2017).

[27] R. M. Anderson, C. Fraser, A. C. Ghani, C. A. Donnelly, S. Riley, N. M. Ferguson, G. M. Leung, T. H. Lam, and A. J. Hedley, Epidemiology, transmission dynamics and control of sars: the 2002–2003 epidemic, Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences **359**, 1091 (2004).

[28] B. J. Cowling, M. P. Muller, I. O. Wong, L.-M. Ho, M. Louie, A. McGeer, and G. M. Leung, Alternative methods of estimating an incubation distribution: examples from severe acute respiratory syndrome, Epidemiology , 253 (2007).

[29] WHO *et al.*, *Consensus document on the epidemiology of severe acute respiratory syndrome (SARS)*, Tech. Rep. (World Health Organization, 2003).

[30] M. A. Johansson, N. Arana-Vizcarrondo, B. J. Biggerstaff, and J. E. Staples, Incubation periods of yellow fever virus, The American journal of tropical medicine and hygiene **83**, 183 (2010).

[31] A. Curtis, J. W. Mills, and J. K. Blackburn, A spatial variant of the basic reproduction number for the new orleans yellow fever epidemic of 1878, The Professional Geographer **59**, 492 (2007).

[32] J. Eberhart-Phillips, R. Besser, M. Tormey, D. Feikin, M. Araneta, J. Wells, L. Kilman, G. Rutherford, P. Griffin, R. Baron, *et al.*, An outbreak of cholera from food served on an international aircraft, Epidemiology & Infection **116**, 9 (1996).

[33] A. A. Weil, Y. Begum, F. Chowdhury, A. I. Khan, D. T. Leung, R. C. LaRocque, R. C. Charles, E. T. Ryan, S. B. Calderwood, F. Qadri, *et al.*, Bacterial shedding in household contacts of cholera patients in dhaka, bangladesh, The American journal of tropical medicine and hygiene **91**, 738 (2014).

[34] Z. Mukandavire, D. L. Smith, and J. G. Morris Jr, Cholera in haiti: reproductive numbers and vaccination coverage estimates, Scientific reports **3**, 1 (2013).

[35] B. T. Goh, Syphilis in adults, Sexually transmitted infections **81**, 448 (2005).

[36] P. O'Byrne and P. MacPherson, Syphilis, British Medical Journal **365** (2019).

[37] C. E. Rupprecht, Rhabdoviruses: rabies virus, Medical microbiology **4** (1996).

[38] C. R. Fisher, D. G. Streicker, and M. J. Schnell, The spread and evolution of rabies virus: conquering new frontiers, Nature Reviews Microbiology **16**, 241 (2018).

[39] S. Lehtinen, P. Ashcroft, and S. Bonhoeffer, On the relationship between serial interval, infectiousness profile and generation time, Journal of the Royal Society Interface **18**, 20200756 (2021).

[40] J. J. van Kampen, D. A. van de Vijver, P. L. Fraaij, B. L. Haagmans, M. M. Lamers, N. Okba, J. P. van den Akker,

H. Endeman, D. A. Gommers, J. J. Cornelissen, *et al.*, Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (covid-19), Nature communications **12**, 1 (2021).

[41] M. Lipsitch, T. Cohen, B. Cooper, J. M. Robins, S. Ma, L. James, G. Gopalakrishna, S. K. Chew, C. C. Tan, M. H. Samore, *et al.*, Transmission dynamics and control of severe acute respiratory syndrome, science **300**, 1966 (2003).

[42] N. M. Linton, T. Kobayashi, Y. Yang, K. Hayashi, A. R. Akhmetzhanov, S.-m. Jung, B. Yuan, R. Kinoshita, and H. Nishiura, Incubation period and other epidemiological characteristics of 2019 novel coronavirus infections with right truncation: a statistical analysis of publicly available case data, Journal of clinical medicine **9**, 538 (2020).

[43] J. Griffin, M. Casey, Á. Collins, K. Hunt, D. McEvoy, A. Byrne, C. McAloon, A. Barber, E. A. Lane, and S. More, Rapid review of available evidence on the serial interval and generation time of covid-19, BMJ open **10**, e040263 (2020).

[44] M. Zhang, J. Xiao, A. Deng, Y. Zhang, Y. Zhuang, T. Hu, J. Li, H. Tu, B. Li, Y. Zhou, *et al.*, Transmission dynamics of an outbreak of the covid-19 delta variant b. 1.617. 2guangdong province, china, may–june 2021, China CDC weekly **3**, 584 (2021).

[45] W. S. Hart, E. Miller, N. J. Andrews, P. Waight, P. K. Maini, S. Funk, and R. N. Thompson, Generation time of the alpha and delta sars-cov-2 variants, medRxiv (2021).

[46] D. Kim, J. Jo, J.-S. Lim, and S. Ryu, Serial interval and basic reproduction number of sars-cov-2 omicron variant in south korea, medRxiv (2021).

[47] R. Pung, T. M. Mak, A. J. Kucharski, and V. J. Lee, Serial intervals in sars-cov-2 b. 1.617. 2 variant cases, The Lancet **398**, 837 (2021).

[48] A. Lange and N. M. Ferguson, Antigenic diversity, transmission mechanisms, and the evolution of pathogens, PLoS Computational Biology **5**, e1000536 (2009).