

1 **Impact of dementia, living in a long-term care facility, and physical activity status**
2 **on COVID-19 severity in older adults**

3

4 Shinya Tsuzuki^{1,2,3}, Takayuki Akiyama², Nobuaki Matsunaga², Norio Ohmagari^{1,2}

5

6 ¹Disease Control and Prevention Center and ²AMR Clinical Reference Center, National
7 Center for Global Health and Medicine, Tokyo, Japan

8 ³Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium

9

10 Corresponding Author: Shinya Tsuzuki

11 Disease Control and Prevention Center

12 National Center for Global Health and Medicine, Tokyo, Japan

13 1-21-1 Toyama, Shinjuku-ku, Tokyo 162-8655, Japan

14 Tel: +81-3-3202-7181

15 Fax: +81-3-3207-1038

16 E-mail: stsuzuki@hosp.ncgm.go.jp

17 Running title: The impact of physical activity on COVID-19

18 Keywords: COVID-19, dementia, long-term care facility, physical activity, propensity

19 score

20

21

22 **Abstract**

23 **Background:** Japan is fast becoming an extremely aged society and older adults are
24 known to be at risk of severe COVID-19. However, the impact of risk factors specific to
25 this population for severe COVID-19 caused by the Omicron variant of concern (VOC)
26 are not yet clear.

27 **Methods:** We performed an exploratory analysis using logistic regression to identify
28 risk factors for severe COVID-19 illness among 4,868 older adults with a positive
29 SARS-CoV-2 test result who were admitted to a healthcare facility between 1 January
30 2022 and 16 May 2022. We then conducted one-to-one propensity score (PS) matching
31 for three factors—dementia, admission from a long-term care facility, and poor physical
32 activity status—and used Fisher's exact test to compare the proportion of severe
33 COVID-19 cases in the matched data. We also estimated the average treatment effect on
34 treated (ATT) in each PS matching analysis.

35 **Results:** Of the 4,868 cases analyzed, 1,380 were severe. Logistic regression analysis
36 showed that age, male sex, cardiovascular disease, cerebrovascular disease, chronic lung
37 disease, renal failure and/or dialysis, physician-diagnosed obesity, admission from a
38 long-term care facility, and poor physical activity status were risk factors for severe
39 disease. Vaccination and dementia were identified as factors associated with non-severe
40 illness. The ATT for dementia, admission from a long-term care facility, and poor

41 physical activity status was -0.04 (95% confidence interval $-0.07, -0.01$), 0.09 ($0.06,$
42 0.12), and 0.17 ($0.14, 0.19$), respectively.

43 **Conclusions:** Our results suggest that poor physical activity status and living in a
44 long-term care facility have a substantial impact on the risk of severe COVID-19 caused
45 by the Omicron VOC, while dementia might be associated with non-severe illness.

46

47 **Introduction**

48 Coronavirus disease 2019 (COVID-19), caused by severe acute respiratory coronavirus
49 2 (SARS-CoV-2), has become a global health threat [1–3]. Since the first case of
50 COVID-19 was identified in Wuhan, China [4], there have been various changes in the
51 approaches to clinical management of COVID-19. The risk factors for severe disease
52 now seem to have been identified [5,6] and the pharmaceutical treatments for
53 COVID-19 cases are at least somewhat established [7–9]. However, the SARS-CoV-2
54 B.1.1.529 variant of concern (VOC) that causes COVID-19 has somewhat different
55 clinical characteristics from the pre-existing variants. This variant, which was first
56 identified in South Africa on 24 November 2021 [10] and was subsequently named
57 Omicron, has now spread worldwide. Fewer patients develop serious illness with this
58 variant and vaccines are less effective [11–13].

59 In Japan, the Omicron VOC has spread rapidly, as in many countries, although
60 there have been fewer severe cases than when the Delta variant was dominant [14].
61 Nevertheless, the higher infectivity of the Omicron VOC than for previous variants led
62 to a huge number of new confirmed cases and a corresponding increase in the number
63 of severe cases. This has burdened our health systems and society further [15].

64 Although previous studies have identified the risk factors associated with

65 severe illness caused by other variants [5,16–18], the same factors may not be
66 applicable to the disease caused by Omicron VOC. Examining the risk factors
67 associated with severe COVID-19 caused by the Omicron VOC is therefore desirable
68 especially since, throughout the entire pandemic period, older Japanese adults have
69 accounted for a large proportion of the severe COVID-19 cases that required
70 hospitalization [19,20]. Because Japan is fast becoming an extremely aged society, an
71 evaluation of the risk factors specific to the elderly population would be extremely
72 valuable. For instance, the US Centers for Disease Control and Prevention (CDC)
73 reported that living in a long-term care facility was an independent risk factor for
74 mortality [21]. Moreover, dementia or pre-existing Alzheimer’s disease was reported to
75 be associated with late mortality due to COVID-19 [22,23]. According to Steenkamp et
76 al., a moderate or high level of physical activity had a preventive effect on severe
77 COVID-19 [24]. However, the impact of these factors on the severity of COVID-19
78 caused by the Omicron VOC is not clear.

79 The main objectives of this study were to identify the risk factors for severe
80 COVID-19 caused by the Omicron VOC and to assess the impact of three factors
81 specific to the elderly population—dementia, living in a long-term care facility, and
82 physical activity status—on the severity of the disease.

83

84 **Methods**

85 **Study population and data**

86 Healthcare facilities that voluntarily participated in the COVID-19 Registry Japan
87 (COVIREGI-JP) [19,25], which is managed by the REBIND (Repository of Data and
88 Biospecimen of Infectious Disease) project, enrolled the patients. Research
89 collaborators in each facility manually entered the data into the registry. The inclusion
90 criteria for enrolment were (i) a positive SARS-CoV-2 test result and (ii) admittance to a
91 healthcare facility between 1 January 2022 and 16 May 2022. The exclusion criteria
92 were positive test results for any of the N501Y, E484K, E484Q, and L452R mutants in
93 SARS-CoV-2 genome tests.

94

95 **Statistical analysis**

96 As a descriptive analysis of the whole data, continuous variables are presented as
97 median and interquartile range (IQR) and categorical variables as number of cases and
98 percentages. We then performed an exploratory analysis using logistic regression to
99 identify risk factors for severe illness. In this study, we defined severe illness as a need
100 for supplementary oxygen during admission. The following variables were included in

101 the regression model: age, sex, vaccinated at least twice, current smoking habit,
102 cardiovascular disease, cerebrovascular disease, chronic lung disease, asthma, liver
103 disease, renal failure and/or dialysis, diabetes mellitus, solid tumor, blood cancer,
104 collagen disease, physician-diagnosed obesity, dementia, admission from a long-term
105 care facility, and physical activity. We defined physical activity as a dichotomous
106 variable and patients were considered to have good physical activity status if they could
107 (i) eat a normal diet, (ii) walk independently, and (iii) take care of themselves; otherwise,
108 they were considered to have poor physical activity status. Each of the three criteria was
109 assessed by a physician.

110 Next, we conducted one-to-one propensity score (PS) matching [26] using
111 three categorizations: presence/absence of dementia, admission from a long-term care
112 facility/home or an acute healthcare facility; and good/poor physical activity status. The
113 PS was calculated by logistic regression analysis with the same variables included in the
114 exploratory analysis. All matching processes were based on nearest neighbor matching
115 with a caliper width of 0.05 and no replacement was allowed. The standardized
116 difference was used to measure the covariate balance, and an absolute standardized
117 difference above 0.1 was interpreted as a meaningful imbalance. Fisher's exact test was
118 used to compare the proportion of severe COVID-19 cases in matched data. In addition,

119 we estimated the average treatment effect on treated (ATT) in each PS matching
120 analysis.

121 Two-sided p values of < 0.05 were considered to show statistical significance.

122 All analyses were conducted with R version 4.1.3 [27].

123

124 **Results**

125 The characteristics of participants are shown in Table 1. We included 4,868 patients in
126 the analysis, 1,380 of whom had severe COVID-19. The age distribution of the severe
127 group was about 20 years older than that of the non-severe group. The severe group
128 showed a higher proportion of past medical history and comorbidity.

129

130 (Table 1)

131

132 Table 2 and Figure 1 show the results of the logistic regression analysis. Age,
133 male sex, cardiovascular disease, cerebrovascular disease, chronic lung disease, renal
134 failure and/or dialysis, physician-diagnosed obesity, admission from a long-term care
135 facility, and poor physical activity status were identified as risk factors for severe illness
136 (i.e., need for supplementary oxygen during admission). Vaccination and dementia were

137 identified as factors associated with non-severe illness.

138

139 (Table 2)

140

141 (Figure 1)

142

143 Figures 2, 3, and 4 show the standardized mean difference before and after the
144 matching procedure for dementia, admission from a long-term care facility, and physical
145 activity status, respectively. For all categorizations, neither group showed significant
146 differences in each item included in the model. The details of the three datasets after
147 matching are available in Tables S1, S2, and S3 in the supplementary file.

148

149 (Figure 2)

150

151 (Figure 3)

152

153 (Figure 4)

154

155 Table 3 shows the estimated ATT of each cohort. All three factors were
156 significantly associated with disease severity. Dementia showed a negative impact on
157 illness severity. Both admission from a long-term care facility and poor physical activity
158 status were associated with severe illness, with the latter showing a larger ATT. Fisher's
159 exact test determined similar results for admission from a long-term care facility and
160 poor physical activity status ($p = 0.014$ and < 0.001 , respectively), whereas dementia
161 did not show a significant difference ($p = 0.435$).

162

163 **Discussion**

164 This study identified the risk factors for severe illness due to COVID-19 caused by the
165 Omicron VOC in older adults. There is little evidence on these factors from Japan until
166 now, and our results show that the risk factors for severe illnesses due to the Omicron
167 VOC are very similar to those of other variants (e.g., Alpha and Delta variants). As
168 suggested previously, older age, male sex, cardiovascular disease, chronic lung disease,
169 and obesity are among the factors associated with severe COVID-19 [5,9,12,17,18]. In
170 addition, these risk factors are similar to those specific to the elderly population [6].
171 These facts may suggest that the original pathology of SARS-CoV-2 infection is
172 inflammation of the respiratory tract, regardless of the strain or variant of the causative

173 organism.

174 At least two vaccine doses is strongly associated with non-severe illness for the
175 Omicron VOC, as well as for other strains [13,28–30]. This result supports the need to
176 promote vaccination at the population level, even though there has been concern about
177 the efficacy of the currently available COVID-19 vaccines against the Omicron VOC
178 [13].

179 The results pertaining to dementia should be carefully interpreted. Although
180 Fisher’s exact test did not show significant differences in the risk of severe disease in
181 the data after PS matching, logistic regression analysis identified dementia as one of the
182 risk factors for severe illness. In addition, the ATT for dementia showed a negative
183 impact on its severity. These results are not in agreement with those of previous studies
184 [22,23,31,32], and dementia itself would not intuitively seem to have a positive effect
185 on the severity of COVID-19. However, two reports from Japan stated that dementia
186 was not a significant risk factor for severe illness due to COVID-19 [6,33]. Examination
187 of the reason for this discrepancy is a future challenge.

188 According to our results, both living in a long-term care facility and poor
189 physical activity status were associated with severe COVID-19, but the ATT for poor
190 physical activity status was the larger of the two. This may suggest that poor physical

191 activity status has a greater impact on the severity of COVID-19 than living in a
192 long-term care facility; in other words, older adults who live in such facilities may also
193 have different risk. Those who have a good physical activity status may have a lower
194 risk of severe illness even if they live in a long-term care facility. This finding may be
195 beneficial when we consider how long-term care facilities manage COVID-19,
196 particularly in super-aged societies like Japan.

197 Several limitations of this study should be noted. First, this was a retrospective
198 cohort study and not a randomized controlled trial. Although we adjusted for various
199 factors using PS matching, not all factors were included in the model. Next, we
200 excluded a large number of patients from the final analyses because if any two groups
201 (i.e., presence/absence of dementia, living in a long-term care facility/elsewhere,
202 good/poor physical activity status) had substantially different demographic
203 characteristics, then numerous patients had to be removed to adjust the background of
204 both groups. Although this would strengthen the internal validity of our results, the
205 external validity and generalizability were sacrificed to some extent. Furthermore,
206 because our registry data did not include information directly assessing the activities of
207 daily living of each patient (e.g., Barthel Index), we had to adopt a new indicator
208 comprising several factors associated with physical activity. These factors—normal diet,

209 independent walking, and ability to take care of oneself—were assessed by each
210 physician in each facility participating in our registry, and subjectivity might have
211 affected the result.

212

213 **Conclusions**

214 Our results suggest that physical activity and living in a long-term care facility have a
215 substantial impact on severe illness caused by COVID-19 owing to the Omicron-19
216 variant, whereas dementia might be associated with non-severe illness. We should take
217 these factors into consideration in the management of older adults with COVID-19.

218

219 **Acknowledgments**

220 The authors thank all of the participating facilities for their care of patients with
221 COVID-19 and their cooperation with data entry into the registry. The data used for this
222 research were provided by the REBIND (Repository of Data and Biospecimen of
223 Infectious Disease) project, which was commissioned for the National Center for Global
224 Health and Medicine by the Ministry of Health, Labour and Welfare of Japan.

225

226 **Ethics approval**

227 Our study data were provided by Research Electronic Data Capture, a secure,
228 Web-based data capture application hosted at the JCRAC Data Center of the National
229 Center for Global Health and Medicine. The opt-out recruitment method was applied,
230 and informed consents for individuals were waived, as approved by the National Center
231 for Global Health and Medicine Ethics Review Board. Information about the entire
232 research is available through the COVID-19 Registry Japan website
233 (<https://covid-registry.ncgm.go.jp/>). This study was approved by the National Center for
234 Global Health and Medicine Ethics Review Board (approval number:
235 NCGM-G-003494-0).

236

237 **Conflict of interest statement**

238 All authors have no conflicts of interest to be disclosed.

239

240 **Authors' contributions**

241 ST and NO conceived the study. TA curated the data. ST and TA analyzed and

242 interpreted the data. ST wrote the first draft of the manuscript. All authors critically

243 reviewed the manuscript and approved the final version.

244

245 **Funding**

246 This research was supported by the Health and Labour Sciences Research Grant,
247 “Research on Emerging and Re-emerging Infectious Diseases and Immunization” (grant
248 number 19HA1003) and JSPS KAKENHI (grant number 20K10546).

249

250 **Availability of data**

251 The data supporting the findings of this study are not publicly available due to the
252 privacy of research participants and sites but are available upon reasonable request.

253 Data on an individual level are shared with limitations to participating healthcare
254 facilities through application to the REBIND project.

255

256

References

- 257 1 Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of
258 Coronavirus Disease 2019 in China. *N Engl J Med*. 2020 Apr;382(18):1708–20.
- 259 2 Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical
260 characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China:
261 a descriptive study. *The Lancet*. 2020 Feb;395(10223):507–13.
- 262 3 Nozaki I, Miyano S. The necessity of continuous international cooperation for
263 establishing the coronavirus disease 2019 diagnostic capacity despite the challenges
264 of fighting the outbreak in home countries. *GHM*. 2020 Apr;2(2):145–7.
- 265 4 Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients
266 infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020
267 Feb;395(10223):497–506.
- 268 5 Terada M, Ohtsu H, Saito S, Hayakawa K, Tsuzuki S, Asai Y, et al. Risk factors for
269 severity on admission and the disease progression during hospitalisation in a large
270 cohort of patients with COVID-19 in Japan. *BMJ Open*. 2021 Jun;11(6):e047007.
- 271 6 Asai Y, Nomoto H, Hayakawa K, Matsunaga N, Tsuzuki S, Terada M, et al.
272 Comorbidities as Risk Factors for Severe Disease in Hospitalized Elderly

- 273 COVID-19 Patients by Different Age-Groups in Japan. GER. 2022 Jan;1–11.
- 274 7 Ministry of Health, Labour and Welfare. Clinical Guideline for COVID-19
275 [Internet]. [cited 2021 Aug 12]. Available from:
276 https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000121431_00111.html
- 277 8 National Institutes of Health. Information on COVID-19 Treatment, Prevention and
278 Research [Internet]. COVID-19 Treatment Guidelines. 2021 [cited 2021 Jan 27].
279 Available from: <https://www.covid19treatmentguidelines.nih.gov/>
- 280 9 World Health Organization. Therapeutics and COVID-19: living guideline [Internet].
281 [cited 2021 Jan 27]. Available from:
282 [https://www.who.int/publications-detail-redirect/therapeutics-and-covid-19-living-g](https://www.who.int/publications-detail-redirect/therapeutics-and-covid-19-living-guideline)
283 [uideline](https://www.who.int/publications-detail-redirect/therapeutics-and-covid-19-living-guideline)
- 284 10 World Health Organization,. Classification of Omicron (B.1.1.529): SARS-CoV-2
285 Variant of Concern [Internet]. [cited 2022 Mar 20]. Available from:
286 [https://www.who.int/news/item/26-11-2021-classification-of-omicron-\(b.1.1.529\)-sa](https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern)
287 [rs-cov-2-variant-of-concern](https://www.who.int/news/item/26-11-2021-classification-of-omicron-(b.1.1.529)-sars-cov-2-variant-of-concern)
- 288 11 Modes ME, Directo MP, Melgar M, Johnson LR, Yang H, Chaudhary P, et al.
289 Clinical Characteristics and Outcomes Among Adults Hospitalized with

290 Laboratory-Confirmed SARS-CoV-2 Infection During Periods of B.1.617.2 (Delta)
291 and B.1.1.529 (Omicron) Variant Predominance — One Hospital, California, July
292 15–September 23, 2021, and Dec. MMWR Morbidity and Mortality Weekly Report.
293 2022;71(6):217–23.

294 12 Bouzid D, Visseaux B, Kassasseya C, Daoud A, Fémy F, Hermand C, et al.
295 Comparison of Patients Infected With Delta Versus Omicron COVID-19 Variants
296 Presenting to Paris Emergency Departments. *Annals of Internal Medicine*. 2022
297 DOI: 10.7326/m22-0308

298 13 Luring AS, Tenforde MW, Chappell JD, Gaglani M, Ginde AA, Mcneal T, et al.
299 Clinical severity of, and effectiveness of mRNA vaccines against, covid-19 from
300 omicron, delta, and alpha SARS-CoV-2 variants in the United States: prospective
301 observational study. *BMJ*. 2022:e069761.

302 14 Ministry of Health, Labour and Welfare. The number of patients and other
303 information about COVID-19 in Japan [Internet]. 2022 [cited 2021 Jan 27].
304 Available from: <https://www.mhlw.go.jp/stf/covid-19/kokunainohasseijoukyou.html>

305 15 Ministry of Health, Labour and Welfare. Investigation about treatment conditions
306 and number of beds: COVID-19 [Internet]. [cited 2021 Aug 18]. Available from:

- 307 https://www.mhlw.go.jp/stf/seisakunitsuite/newpage_00023.html
- 308 16 Chen J, Bai H, Liu J, Chen G, Liao Q, Yang J, et al. Distinct Clinical Characteristics
309 and Risk Factors for Mortality in Female Inpatients With Coronavirus Disease 2019
310 (COVID-19): A Sex-stratified, Large-scale Cohort Study in Wuhan, China. *Clin*
311 *Infect Dis.* 2020 Dec;71(12):3188–95.
- 312 17 Booth A, Reed AB, Ponzo S, Yassaee A, Aral M, Plans D, et al. Population risk
313 factors for severe disease and mortality in COVID-19: A global systematic review
314 and meta-analysis. *PLOS ONE.* 2021 Mar;16(3):e0247461.
- 315 18 CDC. Coronavirus Disease 2019 (COVID-19) [Internet]. Centers for Disease
316 Control and Prevention. 2020 Feb [cited 2021 Aug 12]. Available from:
317 <https://www.cdc.gov/coronavirus/2019-ncov/index.html>
- 318 19 Matsunaga N, Hayakawa K, Asai Y, Tsuzuki S, Terada M, Suzuki S, et al. Clinical
319 characteristics of the first three waves of hospitalised patients with COVID-19 in
320 Japan prior to the widespread use of vaccination: a nationwide observational study.
321 *The Lancet Regional Health - Western Pacific.* 2022;22:100421.
- 322 20 Coronavirus Disease (COVID-19) Situation Report in Japan [Internet]. [cited 2020
323 Oct 9]. Available from: <https://toyokeizai.net/sp/visual/tko/covid19/>

- 324 21 François Watkins LK, Mitruka K, Dorough L, Bressler SS, Kugeler KJ, Sadigh KS,
325 et al. Characteristics of Reported Deaths Among Fully Vaccinated Persons With
326 Coronavirus Disease 2019—United States, January–April 2021. *Clinical Infectious*
327 *Diseases*. 2022 DOI: 10.1093/cid/ciac066
- 328 22 Vlachogiannis NI, Baker KF, Georgiopoulos G, Lazaridis C, Loeff IS, Hanrath AT,
329 et al. Clinical frailty, and not features of acute infection, is associated with late
330 mortality in COVID-19: a retrospective cohort study. *Journal of Cachexia,*
331 *Sarcopenia and Muscle*. 2022 DOI: 10.1002/jcsm.12966
- 332 23 Kim JH, Chang IB, Kim YH, Min CY, Yoo DM, Choi HG. The Association of
333 Pre-existing Diagnoses of Alzheimer’s Disease and Parkinson’s Disease and
334 Coronavirus Disease 2019 Infection, Severity and Mortality: Results From the
335 Korean National Health Insurance Database. *Front Aging Neurosci*.
336 2022;14:821235.
- 337 24 Steenkamp L, Saggars RT, Bandini R, Stranges S, Choi Y-H, Thornton JS, et al.
338 Small steps, strong shield: directly measured, moderate physical activity in 65 361
339 adults is associated with significant protective effects from severe COVID-19
340 outcomes. *British Journal of Sports Medicine*. 2022;bjsports-2021-1.

- 341 25 Matsunaga N, Hayakawa K, Terada M, Ohtsu H, Asai Y, Tsuzuki S, et al. Clinical
342 epidemiology of hospitalized patients with COVID-19 in Japan: Report of the
343 COVID-19 REGISTRY JAPAN. Clin Infect Dis. 2020 Sep DOI:
344 10.1093/cid/ciaa1470
- 345 26 Rosenbaum PR, Rubin DB. The central role of the propensity score in observational
346 studies for causal effects. 15.
- 347 27 R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
348 Austria: R Foundation for Statistical Computing; 2018.
- 349 28 Dagan N, Barda N, Kepten E, Miron O, Perchik S, Katz MA, et al. BNT162b2
350 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. New England
351 Journal of Medicine. 2021 Apr;384(15):1412–23.
- 352 29 Gier B de, Andeweg S, Joosten R, Schegget R ter, Smorenburg N, Kasstele J van
353 de, et al. Vaccine effectiveness against SARS-CoV-2 transmission and infections
354 among household and other close contacts of confirmed cases, the Netherlands,
355 February to May 2021. Eurosurveillance. 2021 Aug;26(31):2100640.
- 356 30 Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety
357 and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. New England Journal of

- 358 Medicine. 2020 Dec DOI: 10.1056/NEJMoa2034577
- 359 31 July J, Pranata R. Prevalence of dementia and its impact on mortality in patients
360 with coronavirus disease 2019: A systematic review and meta-analysis. *Geriatrics*
361 & *Gerontology International*. 2021;21(2):172–7.
- 362 32 Wang Q, Davis PB, Gurney ME, Xu R. COVID-19 and dementia: Analyses of risk,
363 disparity, and outcomes from electronic health records in the US. *Alzheimer’s &*
364 *Dementia*. 2021;17(8):1297–306.
- 365 33 Miyashita S, Yamada T, Mikami T, Miyashita H, Chopra N, Rizk D. Impact of
366 dementia on clinical outcomes in elderly patients with coronavirus 2019 (COVID
367 -19): an experience in New York. *Geriatrics & Gerontology International*.
368 2020;20(7):732–4.
- 369
- 370

371 **Figure legends**

372 **Figure 1. Results of multivariable logistic regression analysis**

373 Black circles indicate median. Whiskers indicate 95% confidence intervals. LTCF,

374 long-term care facility.

375

376 **Figure 2. Balance of demographic characteristics of older COVID-19 inpatients**

377 **before and after propensity score matching (with dementia/without dementia)**

378 LTCF, long-term care facility.

379

380 **Figure 3. Balance of patients' demographic characteristics before and after**

381 **propensity score matching (admission from a long-term care facility/elsewhere)**

382


383 **Figure 4. Balance of patients' demographic characteristics before and after**

384 **propensity score matching (poor/good physical activity status)**


385 LTCF, long-term care facility.

386


Odds ratio



Covariate Balance



Covariate Balance



Covariate Balance

