Modelling the epidemiological implications for SARS-CoV-2 of Christmas household bubbles in England

Edward M. Hill[1,2*]

1 The Zeeman Institute for Systems Biology & Infectious Disease Epidemiology Research, School of Life Sciences and Mathematics Institute, University of Warwick, Coventry, United Kingdom.
2 Joint UNIversities Pandemic and Epidemiological Research, https://maths.org/juniper/.

* Corresponding Author. Email: Edward.Hill@warwick.ac.uk

Abstract

The emergence of SARS-CoV-2 saw severe detriments to public health being inflicted by COVID-19 disease throughout 2020. In the lead up to Christmas 2020, the UK Government sought an easement of social restrictions that would permit spending time with others over the Christmas period, whilst limiting the risk of spreading SARS-CoV-2. In November 2020, plans were published to allow individuals to socialise within 'Christmas bubbles' with friends and family. This policy involved a planned easing of restrictions in England between 23–27 December 2020, with Christmas bubbles allowing people from up to three households to meet throughout the holiday period. We estimated the epidemiological impact of both this and alternative bubble strategies that allowed extending contacts beyond the immediate household. We used a stochastic individual-based model for a synthetic population of 100,000 households, with demographic and SARS-CoV-2 epidemiological characteristics comparable to England as of November 2020. We evaluated five Christmas bubble scenarios for the period 23-27 December 2020, assuming our populations of households did not have symptomatic infection present and were not in isolation as the eased social restrictions began. Assessment comprised incidence and cumulative infection metrics. We tested the sensitivity of the results to a situation where it was possible for households to be in isolation at the beginning of the Christmas bubble period and also when there was lower adherence to testing, contact tracing and isolation interventions. We found that visiting family and friends over the holiday period for a shorter duration and in smaller groups was less risky than spending the entire five days together. The increases in infection from greater amounts of social mixing disproportionately impacted the eldest. We provide this account as an illustration of a real-time contribution of modelling insights to a scientific advisory group, the Scientific Pandemic Influenza Group on Modelling, Operational sub-group (SPI-M-O) for the Scientific Advisory Group for Emergencies (SAGE) in the UK, during the COVID-19 pandemic.

Introduction

The COVID-19 pandemic has had grave consequences for global public health. In the initial absence of known effective therapeutics and vaccinations against SARS-CoV-2, the strategies implemented by many nations during 2020 to attempt to curb SARS-CoV-2 outbreaks revolved around nonpharmaceutical interventions (NPIs), including stringent lockdown rules and social distancing measures [1]. Despite the use of such measures in England, by 30 November 2020 there had been 52,881 people who died within 28 days of being identified as a COVID-19 case by a positive test [2].

The vacation period around Christmas is an important time for many people of all faiths and none who come together over the holidays. In light of the hardships suffered during 2020, the UK government was considering actions to balance the desire to spend time with others over the Christmas period, whilst limiting the risk of spreading SARS-CoV-2. On that basis, on 24 November 2020 plans were published to allow individuals to socialise within 'Christmas bubbles' with friends and family [3]. This policy involved a planned easing of restrictions in England between 23–27 December inclusive. The intent was that anyone who was not self-isolating could form a Christmas bubble according to three main rules: (i) only being in one Christmas bubble; (ii) not changing their Christmas bubble; and (iii) the Christmas bubble could not include people from more than three households. Christmas bubbles could meet, without socially distancing, in private homes, places of worship or public outdoor spaces.

Ultimately, these original plans for Christmas bubbles never came to fruition, being heavily revised following the emergence of the (later named) Alpha variant in Kent, England [4]. On 19th December 2020 [5], it was announced that Christmas bubbles in England would be permitted only on Christmas Day (25th December 2020) and only in locations not in the highest level of restrictions (Tier 4) [6].

Nonetheless, under the proviso of the original plan for Christmas bubbles in England throughout 23-27 December 2020, the expected increase in social interactions during that time meant there was interest in ascertaining the epidemiological implications of household bubbles. Contact clustering in social bubbles beyond the immediate household had already undergone consideration during the COVID-19 pandemic, as part of both modified lockdown policies and gradual lockdown exit strategies granting greater social interaction [7, 8]. It had been demonstrated that social network-based reduction of contact strongly enhanced the effectiveness of social distancing measures while keeping risks lower [7]. Prior work using an individual based model had estimated the impact of such bubble strategies on epidemic and mortality risk using the UK as a case study, finding that social bubbles could be an effective way of extending contacts beyond the household while limiting the increase in epidemic risk, if managed appropriately [8].

The data and science surrounding the SARS-Cov-2 infection is fast moving. In the context of Christmas bubbles in England during 2020, contributory tools to evaluate epidemiological questions on household bubbles and their ramifications included scenario assessments using mathematical models. We present here our associated analyses conducted in late November and early December 2020. We treat our manuscript as a record of the state of our modelling and the epidemiological context at that time. There have been subsequent additions to the literature relevant to this topic, which are remarked upon in the Discussion.

In this study, we describe a stochastic individual-based model that was used to compare a set of household bubble scenarios during 23-27 December 2020, with a synthetic population initialised to reflect the demography and epidemiological context of England. We considered a population of households initialised with no symptomatic infection present, with between-household mixing occurring through implementation of a household visit schedule. Purely focused on epidemiological outcomes, we found

conditional on there being any easing of rules on household gatherings then shorter and smaller gatherings were preferred. Furthermore, the model quantified the increase in incidence in the over 65s in circumstances where additional mixing occurred over more days.

We present the manuscript as an example of the real-time contribution mathematical epidemiological and modelling of infectious disease dynamics can make to the knowledge base informing policy in response to a public health emergency. The work was presented to the Scientific Pandemic Influenza Group on Modelling, Operational sub-group (SPI-M-O) for the Scientific Advisory Group for Emergencies (SAGE) in the UK on 2nd December 2020, contributing to paragraphs in the consensus statement on the impact of a festive break in relaxations [9].

Methods

We simulated an epidemic process amongst a population of households and assessed the epidemiological impact of temporarily expanding the permitted size of household bubbles during Christmas 2020.

In the Methods we detail: (i) the parameterisation of the households (subsection *Household demographics*), (ii) the epidemiological model for SARS-CoV-2 transmission (subsection *SARS-CoV-2 epidemiological model*), (iii) our implementation of test, trace and isolate guidance (subsection *Testing, contact tracing and isolation*), (iv) the Christmas bubble scenarios evaluated (subsection *Christmas bubble scenarios*), and (v) the simulation protocol used to assess the Christmas bubble scenarios (subsection *Simulation overview*).

Household demographics

In our model, each individual was assigned to a household and classified into one of three age groups: 0-19yrs, 20-64yrs, 65+yrs. The three-class age structure allowed for age-based heterogeneity in probabilities of asymptomatic infection and susceptibility to infection. We considered a population of 100,000 households, sampling household sizes and the proportion of households satisfying a given age composition (e.g.one 0-19yrs individual and two 65+ year olds) from 2011 census data for England and Wales [10].

Single-occupancy households and single parent households could form support bubbles with another household [11], which we refer to as an extended household. We modelled extended households as in essence having the same transmission rates as if they shared the same physical household over the simulated period.

In the absence of data on the proportion of single person households/single-parent households in a support bubble, for each simulation we sampled the propensity to form a support bubble from a Uniform(0.5,0.75) distribution (*i.e.* sampling on the basis of the majority of those households joining up with another).

SARS-CoV-2 epidemiological model

In this section we outline our assumptions regarding the natural course of infection, asymptomatic infection and transmission, susceptibility and the relationship between transmission risk and household size. For a listing of associated parameters, see Table 1.

Disease state structure

We ran a disease process on the households, highlighting that we only considered infection resulting from person-to-person interactions due to household mixing. We did not consider transmission aris-

ing from other settings based on a wish to evaluate the additional infections that would arise from increased between household mixing. The implications of this assumption is expanded upon in the Discussion.

We modelled the natural infection history using a Susceptible-Exposed-Infected-Recovered (SEIR) type structure. Namely, each individual was either in a susceptible, latent (infected but not infectious), infectious or recovered state. Once infected, we assumed infectiousness could start from the following day. We assumed an Erlang-distributed incubation period to symptom onset, with shape parameter 6 and scale parameter 0.88 [12].

The distribution of infectiousness relative to symptom onset was based on a Gamma distribution with shape parameter 97.2 and scale parameter 0.2689, shifted by 25.6 days [13, 14]. It had a fixed four day pre-symptomatic phase, followed by a fixed ten day symptomatic phase. This gave a total of 14 days of infectivity and a minimum 15 day infection duration. The infectiousness temporal profile weighted the transmission risk (see the subsequent subsection on *Transmission risk*) across the duration of the infectious period (for the full temporal profile, see Table 1). Following completion of the infectious period, the individual entered the recovered state. We provide a schematic representation of the epidemiological model in Fig. S1.

Probability of asymptomatic infection and relative infectiousness

Infected individuals could be either asymptomatic throughout infection or symptomatic (*i.e.* develop symptoms following the latent period), according to a specified asymptomatic probability. We allowed for the probability of asymptomatic infection to be age dependent (split into children and adolescents, 0-19yrs, and adults, 20+yrs. In each simulation run we sampled an asymptomatic infection probability for the 0-19yrs age group from a Uniform(0.2,0.35) and an asymptomatic infection probability for the 20+yrs age group from a Uniform(0.05,0.2) distribution [15]. The sampled asymptomatic probabilities remained fixed throughout that particular simulation run.

There was limited data available to provide a robust quantitative estimate of the relative infectiousness of asymptomatic and symptomatic individuals, with some support for the reduced infectiousness of asymptomatic individuals [15, 16]. Therefore, we assumed that asymptomatic individuals had a lower risk of transmitting infection compared to symptomatic individuals. To reflect the uncertainty in this area, for each simulation we sampled the relative infectiousness of asymptomatic individuals compared to symptomatic individuals from a Uniform(0.3, 0.7) distribution. This was sampled independently to the asymptomatic probability. The sampled value was applied as a scaling on transmission risk, applied evenly throughout the duration of infectiousness (*i.e.* with no time dependence, see the subsection *Probability of transmission per contact*).

Susceptibility

Based on available estimates at the time, we set the relative susceptibility to SARS-CoV-2 infection for the child and adolescent age group (0-19yrs) to be approximately half that of people aged 20+yrs [17]. Explicitly, for each simulation we sampled the relative susceptibility of the child and adolescent age group (0-19yrs) from a Uniform(0.4, 0.6) distribution.

Transmission risk

We used a data-driven approach to obtain the relative risk of transmission within households of differing composition.

To obtain the transmission risk from an individual, we modulated the infectivity profile using estimates of adjusted household secondary attack rates from a UK based surveillance study [18]. We attributed

a household secondary attack rate to each person based on the size of their household. We sampled the attack rates for each individual from a normal distribution with mean dependent on the household size [18]: 0.48 for a household size of two, 0.40 for three, 0.33 for four, and 0.22 for five or more. The standard deviation of the normal distribution for households of size two or three was 0.06, and for households of four or more was 0.05. It should be noted that these estimates were made using a sample of 379 confirmed COVID-19 cases, meaning the robustness of the central estimates could be low. As such, we sampled from the previously mentioned normal distributions to ensure this uncertainty was captured.

On a day where households visited one another it was assumed they would be gathered for the entire day. The overall transmission risk on a given day was dictated either by the total number of individuals in the household if there was no mixing with other households, or if multiple households were meeting as part of a support bubble or a Christmas bubble then the transmission risk was governed by the total number of individuals in the gathering. For example, if in a Christmas bubble there were three households of sizes 2, 1 and 1, then we assumed the transmission risk to be like a household of size 4.

Probability of transmission per contact

We outline here how the household transmission risk, relative infectiousness of the infector, relative susceptibility of the infectee and the infectiousness temporal profile were used to compute the probability of transmission across an infectious-susceptible contact pair.

For an infectious individual j on day t of their infectious state, the probability of transmission to each susceptible contact k in household bubble h, denoted $p_{j,k,h}(t)$, was given by the product of four components:

$$p_{j,k,h}(t) = r_h a_j s_k i_j(t),$$

where r_h is the sampled household attack rate in household h, a_j the relative infectiousness of individual j (taking either value 1 for cases that were symptomatic during the infection episode, or the sampled value for relative infectiousness of asymptomatics compared to symptomatics otherwise), s_k the relative susceptibility of individual k and $i_j(t)$ the value of the infectiousness temporal profile on day t for individual j (Table 1).

Table 1: Description of epidemiological parameters.

Description	Distribution	Source
Incubation period	Erlang(6, 0.88)	[12]
Infectiousness profile	Infectivity profile over 14 days: [0.0369, 0.0491, 0.0835, 0.1190, 0.1439, 0.1497, 0.1354, 0.1076, 0.0757, 0.0476, 0.0269, 0.0138, 0.0064, 0.0044]	[13, 14]
Proportion of cases asymptomatic (0-19yrs)	Uniform(0.20, 0.35)	[15]
Proportion of cases asymptomatic (20+yrs)	Uniform(0.05, 0.20)	[15]
Relative infectiousness of an asymptomatic	Uniform(0.30, 0.70)	[15, 16]
Relative susceptibility of 0-19yrs age class	Uniform(0.40, 0.60)	[17]
Household attack rates	Size 2: Normal(0.48,0.06) Size 3: Normal(0.40,0.06) Size 4: Normal(0.33,0.05) Size ≥5: Normal(0.22,0.05)	[18]

Testing, contact tracing and isolation

Other measures in active use during this period included symptomatic testing, contact tracing and isolation [19, 20]. We next expand on our implementation of the testing, contact tracing and isolation components within our household model. Associated parameters are listed in Table 2.

Adherence

We used an adherence parameter to capture the proportion of individuals that follow the recommended guidance. In our simulations all individuals within the same household (or extended household) had the same adherence status, representing whether an individual would both adhere to isolation guidance and engage with test and trace. We did not allow partial adherence to one measure and not the other. Adherence to isolation encompassed isolation for any reason: presenting with symptoms themselves, being in the same household as someone displaying symptoms, or being identified as a close contact of an infected individual via contact tracing.

We assumed a constant 70% adherence to isolation and test-and-trace measures, sampled using a Bernoulli trial. Though we acknowledge a subjective nature to this assumption, indicators from the Office for National Statistics (ONS) Opinions and Lifestyle Survey covering the period 5 to 8 November 2020 reported around 7 in 10 adults "finding it very easy or easy to follow the current lockdown measures where they live"[21]. To garner insights into the sensitivity to the adherence parameter we also performed simulations with a lower adherence, catering for a situation where the majority of the population were not partaking in other NPIs (see subsection on *Sensitivity analysis*).

Testing

An adherent individual household member was assumed to take a PCR test if they displayed symptoms. We assumed individuals would isolate between taking a PCR test and receiving the result. To enable a clearer assessment of the potential increase in infection resulting from the size and duration of Christmas bubbles, rather than possible confounding feedbacks that may be induced in the infection dynamics by lags in test result reporting, we also assumed test results were delivered on the same day as symptom onset.

We assumed the PCR test had 100% specificity, in line with data from the Office for National Statistics COVID-19 Infection survey suggesting the false-positive rate to be very low, under 0.005% [22]. PCR test sensitivity was a fixed constant of 87%, based on the reported clinical sensitivity in symptomatic individuals [23], which also lay within the estimated range of 85% and 98% sensitivity reported across other studies [22]. These statistics agreed with the available estimates in the UK as of November 2020. That being said, at a similar time we were seeing the first reports in the literature that provided information on the sensitivity of PCR testing for SARS-CoV-2 infections dependent upon time since infection (studies such as Hellewell *et al.* [24], Wikramaratna *et al.* [25], for example). When carrying out this analysis we had yet to incorporate these time-dependent sensitivity profiles into the model.

In instances where the symptomatic household member returned a negative test result, they remained in self-isolation if they had independently been identified via contact tracing as a contact of a known infected; otherwise, the symptomatic individual left self-isolation. This matches government guidance on contact tracing at the time - see the subsection below on *Contact tracing*.

Household members were also released from isolation upon a negative result, as long as no other symptomatic cases (that were confirmed positive) were present in the household and they had not been independently identified as requiring to self-isolate through contact tracing.

Contact tracing

Contacts of individuals returning a positive test result were retrieved as far back as two days prior to the emergence of symptoms for the index case. Contacts of a confirmed case were required to spend up to 14 days in self-isolation [20], matching the UK government guidance in November 2020. We set the isolation period to elapse 14 days after the index case became symptomatic.

Isolation

Within an adherent household, symptomatic infected individuals isolated for 10 days upon symptom onset. All other individuals in the extended household were set to self-isolate for 14 days from their infected household members data of symptom onset [19]. Individuals in isolation did not participate in planned household visits due to occur during the five-day bubble window.

Table 2: Description of testing and isolation related parameters.

Description	Value	Source
Adherence	70% (30% in adherence sensitivity analysis)	[21]
Test specificity	100%	[22]
Test sensitivity	87%	[23]
Duration of self-isolation if symptomatic	10 days	UK government guidance in November 2020 [19]
Household isolation period	14 days	UK government guidance in November 2020 [19]
Duration of isolation if contact traced	14 days (beginning from the day the index case first displays symptoms)	UK government guidance in November 2020 [20]

Christmas bubble scenarios

We evaluated five Christmas bubble scenarios, which could be in use during 23-27 December 2020. The five scenarios, ranging from least to greatest amount of unique additional contacts, were:

Scenario A - No change to household bubble guidance: Only support bubbles (extended households) permitted to meet each day.

Scenario B - Short duration fixed exclusive bubbles: The same three households meet each day between 25-26 December 2020, with no overlapping between bubbles.

Scenario C - Fixed exclusive bubbles: The same three households meet each day between 23-27 December 2020, with no overlapping between bubbles. This scenario most closely corresponds to the original Christmas bubble plan for England.

Scenario D - Fixed non-exclusive bubbles: Each household meets two other households each day between 23-27 December 2020. Bubbles may overlap.

Scenario E - Daily change in household triplets: Each day, all households are assigned to a household triplet. The household triplets change each day between 23-27 December 2020.

In all scenarios we assumed extended households (households part of a support bubble) mixed every day during 23-27 December 2020. Furthermore, in Scenarios B-E existing support bubbles counted as one household towards the three household limit, concurring with the original Christmas bubbles guidance [3]. For a visual illustration of the different scenarios, see Fig. 1.

Fig. 1: Illustration of the modelled Christmas bubble scenarios. We depict our five modelled scenarios. Scenario A: No change (support bubbles only); Scenario B: Short duration fixed exclusive bubbles, meet 25 and 26 December only; Scenario C: Fixed exclusive bubbles, meet every day 23-27 December; Scenario D: Fixed non-exclusive bubbles, meet every day 23-27 December; Scenario E: New household triplets meet each day between 23-27 December. Households contained within dashed ellipses signify an existing support bubble. Households within solid ellipses are households contacted as part of a Christmas bubble. We use markers to denote individuals and different marker types signify individuals in different age groups; 0-19yrs (red squares); 20-64yrs (blue circles); 65+yrs (grey stars).

Simulation overview

Model initialisation

For each scenario we performed 100 model runs on a system containing 100,000 households, with an approximate overall population of 310,000 people. These quantities were pragmatic choices, balancing between generating sufficient realisations spanning the plausible range of parameter values and having a large overall population size for which epidemic simulations could be completed in a confined time frame.

At the outset of each simulation we assigned household bubbles in a two-step process. Firstly, for each single-occupancy household we checked whether they would form a support bubble with another household (see the subsection *Household demographics*). Secondly, for Scenarios B-E, we created Christmas bubbles and visitation schedules according to the specifics of each scenario. The second step of the household bubble allocation was not applicable for Scenario A.

We then proceeded to initialise the epidemiological state of each individual. Accordingly, we assigned to each individual one of the following possible disease statuses; susceptible, latently infected, asymp- tomatically infected, presymptomatic infectious, recovered. There were intentionally no individuals initialised as symptomatically infected at the start of the simulated time horizon, meaning no house- holds began in isolation.

The probability of being attributed a given disease state was age-dependent, with age-stratified es- timates for our three age groups (0-19yrs, 20-64yrs, 65+yrs) informed by inferred values from the Warwick SARS-CoV-2 compartmental transmission model in late November 2020 [26]. We set the status of individuals in each household at random, with acknowledgement that in truth disease sta- tuses within a household may be correlated due to within household transmission (remarked upon further in the Discussion). We provide initial condition values in Table 3.

Table 3: Percentage of each age group initialised in each infection status.

	Age (years)		
	0-19	20-64	65+
Susceptible	73%	74%	84.5%
Latent infected	1%	0.5%	0.25%
Asymptomatic infected	0.3%	0.1%	0.05%
Presymptomatic infected	0.7%	0.4%	0.2%
Recovered	25%	25%	15%

Main analysis

For our main analysis we began simulations on 23 December 2020 and and ran each simulation for 15 days. During 23-27 December 2020 contacts were made according to the Christmas bubble scenario being studied. After 27 December 2020, we assumed extended households continued to interact each day for ten further days.

The described setup corresponded to a population of households that had no symptomatic infection present, and therefore no households being in isolation, at the beginning of the five-day Christmas bubble window on 23 December 2020.

Sensitivity analysis: Alternative time horizon

As a sensitivity analysis, we considered an alternative set of circumstances where it was possible for households to be in isolation as the Christmas bubble period began.

In this case we began the simulations ten days prior to 23 December 2020 (*i.e.* simulations began on 13 December 2022), with the same initial conditions. On each day during 13-22 December 2020 extended households would meet each day. For 23-27 December we ran the previously described set of five Christmas bubble scenarios. As before, after 27 December 2020 we assumed extended households continued to interact each day for ten further days.

The intent of this setup was to more closely reflect the likely epidemiological impacts averaged over the whole general population, rather than the epidemiological impacts amongst solely households that had no symptomatic infection present and requirement to isolate upon the temporary relaxation of social restrictions.

Sensitivity analysis: Adherence to test, trace and isolation measures

Lastly, for both the main and alternative analyses time horizons we investigated epidemiological outcomes given a lower adherence to isolation and test-and-trace measures of 30% (rather than 70%).

Epidemiological outcomes

Our assessment of the Christmas bubble and visitation scenarios primarily considered a pair of incidence and cumulative infection statistics. We grouped these epidemiological measures into: (i) Daily incidence each day between 23-27 December 2020; (ii) Cumulative infections over the entire modelled time horizon. For these quantities we were also interested in the share attributable to each age group.

Software and code

We performed the model simulations in Julia v1.5 and generated the figures in Matlab R2022a. The code repository for the study is available at: https://github.com/EdMHill/SARS-CoV-2_Christmas_bubbles_2020.

Results

Longer and larger visits alters the age distribution of cases

We visualise the daily incidences from 23 December 2020 (Fig. 2(a)) to 27 December 2020 (Fig. 2(e)), for all five Christmas bubble scenarios ordered from the lowest level of unique contacts (Scenario A, darkest colour) to the highest level (Scenario E, lightest colour), and sorted by age group (blue, left across to purple, right). We particularly note the dynamics exhibited by Scenario B that allowed exclusive bubbles of three households on 25 and 26 December 2020 only.

For 23 and 24 December 2020, Scenario B had the same rules as Scenario A, and thus similar daily incidence distributions (Figs. 2(a) and 2(b), Table S1). For the two days where Christmas bubbles were implemented (25 and 26 December 2020), the daily incidence distributions were similar to Scenarios C-E, where households were allowed to meet two other households on all five days between 23-27 December 2020 (Figs. 2(c) and 2(d), Table S1). On 27 December 2020, the incidence returned to levels comparable to Scenario A, though median estimates for Scenario B were now slightly larger (Fig. 2(e), Table S1); the overall estimates for all ages in the population were 0.067% (95% PI:0.0589%-0.079%) for Scenario A, compared to 0.069% (95% PI: 0.060%-0.079%) for Scenario B.

We next remark on an apparent age-dependence on the expected increase in incidence on days when Christmas bubbles were permitted to meet. Compared to maintaining support bubbles only (Scenario A), on days Christmas bubbles could meet we found the incidence approximately trebled for those aged 65+yrs. In contrast, for the 20-64yrs age group incidence roughly doubled, whereas the relative change was smallest for those aged 0-19yrs (Figs. 2(a) to 2(e)).

We observe a shift towards a greater proportion of infections occurring in the 65+yrs age group (Fig. 2(f)) as a result of the use of exclusive bubbles of three households for a short duration (Scenario B). Under Scenario A, the percentage of new infections during this five day window estimated to occur in those aged 65+yrs was 23% (95% PI: 19%-29%). Under Scenario B, the analogous median and uncertainty estimates were 30% (95% PI: 26%-34%) for Scenario B. The 0-19yrs and 20-64yrs age groups had similar declines of 3-4%, in contribution to new infections during that time period.

With an increase in duration of interaction (Scenarios C and D) or the number of contacts (Scenario E), we observed a larger shift in the age distribution of new infections that occurred during the period 23-27 December 2020. Specifically, there was an appreciable rise of roughly 10% in the proportion of cases attributed to those aged 65 and above, with a corresponding decrease in the region of 5% of the share of new infections attributable to those aged 0-19yrs and 20-64yrs (Fig. 2(f)).

Implications of Christmas bubbles on post-Christmas SARS-CoV-2 prevalence

All Christmas bubble scenarios that allowed for mixing beyond extended households resulted in an increase in the amount of new infections over the whole 15-day time horizon, 23 December 2020 to 06 January 2021 (Fig. 3). Notably, the total number of days spent mixing within that period could impact post-Christmas prevalence. Compared to having no additional household bubbling (Scenario A), for which cumulative infections for all ages reached 0.60% (95% PI: 0.53%-0.64%), the rises in cumulative infections were most stark for household visit schedules where three households could meet each day for the full five days (Scenarios C-E).

In detail, we found modelled cumulative infections across all ages of: exclusive bubbles (Scenario C) - 0.96% (95% PI: 0.88%-1.06%); non-exclusive fixed bubbles (Scenario D) - 0.96% (95% PI: 0.87%-1.06%); daily change in household triplets (Scenario E) - 0.99% (95% PI: 0.88%-1.07%). We remark that although Scenario E, with new household triplets meeting each day of 23-27 December, would result in a greater amount of distinct person-to-person contacts being made compared to having fixed

Fig. 2: Distributions for the daily incidence and age-stratification of new infections between 23-27 December 2020 under each Christmas bubble scenario. Estimates produced from 100 realisations per scenario for daily incidence on: **(a)** 23 December 2020; **(b)** 24 December 2020; **(c)** 25 December 2020; **(d)** 26 December 2020; **(e)** 27 December 2020. Panel **(f)** displays the share of new infections between 23-27 December 2020 attributable to each of the three age groups. In each violin plot the white squares represent the medians and solid black lines correspond to the interquartile range. See Table S1 for median values and 95% prediction intervals. Labels A-E correspond to the distributions associated with Scenarios A-E. The intensity of shading of the violin plots also distinguishes between the scenarios; darkest for Scenario A to lightest for Scenario A: No change (support bubbles only): Scenario B: Short duration fixed exclusive bubbles, meet 25 and 26 December only; Scenario C: Fixed exclusive bubbles, meet every day 23-27 December; Scenario D: Fixed non-exclusive bubbles, meet every day 23-27 December; Scenario E: New household triplets meet each day between 23-27 December. Relative to Scenario A, the increase in incidence for Scenarios C-E was greatest for the 65+yrs age group.

bubbles comprised of three households (Scenarios C and D), the modelled outcomes across these three scenarios were similar. On the other hand, restricting the duration of exclusive bubbles to 25-26 December 2020 (Scenario B) reduced the relative increase in cumulative infections over the 15-day time horizon by approximately half (Fig. 3, Table S1).

Stratifying by age, we found that the greatest relative increase in cumulative infection for Scenario E versus Scenario A was once more amongst the eldest age group (Scenario A - 0.32% (95% PI: 0.26%-0.39%), Scenario E - 0.79% (95% PI: 0.66%-0.92%), Fig. 3). The 0-19yrs child and adolescent age group had the lowest relative increase (Scenario A - 0.49% (95% PI: 0.38%-0.58%), Scenario E - 0.74% (95% PI: 0.57%-0.89%)).

Sensitivity analysis: Alternative time horizon

We next inspected our simulations that began on 13 December 2020, which included households with initially symptomatic individuals and those already in isolation on 23 December 2020, the start of the (up to) five-day temporary easement of social restrictions. One may consider such a setting to be more analogous to the likely state of the general population as Christmas bubbles began.

We observed qualitatively similar findings to the main analysis when inspecting epidemiological outcomes both constrained to the 23-27 December 2020 period (Fig. S2, Table S3) and over the entire time horizon through to 06 January 2021 (Fig. S3, Table S4). Shorter duration exclusive bubbles (Scenario B) roughly halved the additional infections accumulated compared to Scenarios C-E where meeting with two other households was allowed for all five days during 23-27 December 2020 (Fig. S3(a)). Relative to maintaining the status quo of support bubbles (Scenario A), allowing in the period 23-27 December fixed exclusive bubbles of three households to meet all five days (Scenario C), fixed non-exclusive bubbles of three households to meet all five days (Scenario D), or on each day different household triplets to meet (Scenario E) resulted in a higher proportion of cases occurring in the 65+yes age class (Figs. S2(f)&S3(b)). These three bubbling scenarios also caused similar levels of additional infections (Figs. S2&S3).

There were four quantitative differences between the outcomes for our alternative analysis compared to the main analysis (see Tables S3-S4 for summary statistics). First, in the alternative analysis the daily incidence during 23-27 December 2020 for each Christmas bubble scenario and age group combination was of the order of five-fold less compared to the corresponding scenario in the main analysis. Second, independent of the Christmas bubble strategy in use, the share of new infections was higher in the 0-19yrs age group. Third, the magnitude of the relative increase in cumulative infections compared to there being no temporary easement in social restrictions (Scenario A) was less for household visit schedules that resulted in interactions of longer duration (Scenarios C and D) or a greater amount of contacts (Scenario E). Finally, there was greater variability in all predicted distributions (Figs. S2&S3)

Sensitivity analysis: Adherence to test, trace and isolation measures

The last piece of analysis investigated sensitivity of the epidemiological outcomes to the adherence parameter. We considered a background where the majority of the population were not adherent to isolation and test-and-trace measures, setting the adherence probability to 30%.

As anticipated, irrespective of the Christmas bubble scenario and age group, lower adherence led to higher incidence and cumulative infections. However, we still found qualitatively similar findings regarding age distributions, chiefly a marked uplift in the proportion of infections occurring in those aged 65 and above (Figs. S4-S7, Tables S5-S8).

(a)

(b)

Fig. 3: Cumulative infection distributions for the entire 15 day time horizon, 23 December 2020 to 06 January 2021, under each Christmas bubble scenario. Estimates produced from 100 realisations per scenario. **(a)** The percentage of each age group newly infected during the time horizon. **(b)** The percentage of infections during the time horizon attributed to each age group. In each violin plot the white squares represent the medians and solid black lines correspond to the interquartile range. See Table S2 for median values and 95% prediction intervals. Labels A-E correspond to the distributions associated with Scenarios A-E. The intensity of shading of the violin plots also distinguishes between the scenarios; darkest for Scenario A to lightest for Scenario E. Scenario A: No change (support bubbles only); Scenario B: Short duration fixed exclusive bubbles, meet 25 and 26 December only; Scenario C: Fixed exclusive bubbles, meet every day 23-27 December; Scenario D: Fixed non-exclusive bubbles, meet every day 23-27 December; Scenario E: New household triplets meet each day between 23-27 December. All scenarios beyond extended household bubbles resulted in an increase in the amount of new infections amongst the population over the considered time horizon.

Discussion

In this study, we have developed a stochastic individual-based model to analyse the spread of SARS-CoV-2 in a population experiencing a temporary easement in social restrictions, permitting the creation of bubbles of three households. We have investigated the impact of different bubble strategies, akin to the 'Christmas bubbles' proposed for use in England between 23-27 December 2020, upon SARS-CoV-2 incidence and prevalence amongst a synthetic population of households.

We found incidence and cumulative infection metrics were similar for bubble policies allowing either longer duration meetings of fixed household bubbles or multiple one-day gatherings of unique household triplets. Further, it was likely that visiting family and friends over the holiday period for two days would be less risky than spending the entire time together; shorter and smaller gatherings reduced the relative amount of infection. We also observed the increases in infection from greater amounts of social mixing disproportionately impacted the eldest. This is of particular concern due to the worse clinical outcomes associated with increasing age, with higher hospitalisation and mortality rates of COVID-19 disease observed in older age groups ages [2, 27]. These conclusions were maintained for a population of households where households could begin the Christmas bubble time period in isolation, thus fragmenting the potential maximum connectivity that usage of Christmas bubbles could bring, and in a setting where the majority of the population would not be adherent to test, trace and isolate guidance.

Our findings have since been corroborated by subsequent additions to the literature on the topic of quantifying the effect of household bubbles on the transmission of SARS-CoV-2, with a variety of modelling approaches represented [28, 29]. In detail, Danon *et al.* [28] used the configuration model to link households and generate a network description of households in the UK. They found that whilst the creation of support bubbles between a single-person household and another household of any size had a small impact on transmission, scenarios where all households form a bubble would be highly likely to cause extensive transmission. Later work from Hilton *et al.* [29] provided an infectious disease modelling framework formulated in terms of tractable systems of ordinary differential equations that includes explicit representation of age/risk structure and household structure. Our results are in agreement with their findings on the usage of short-term social bubbles, which suggested that short-term relaxation in mixing restrictions would have a small but non-negligible impact on the epidemic dynamics, with larger temporary bubbles and longer mixing periods associated with higher prevalence.

The model framework described here is reasonably parsimonious compared to fine-detailed household models [29–32]. With regards to informing policy in response to a global public health emergency such as a pandemic, parsimonious models that can be promptly set up and that are able to generate results in extremely short time-scales over can be helpful tools for providing high-level insights, on the basis that a reasonable but imperfect answer is better than none [33]. The model also has ongoing prospective use in situations where substantial mixing between household groups is expected over a short duration of time, in relation to between household contact levels before and after said mixing period. Such situations are of interest with the period of increased mixing between household groups resulting in a new transmission network as new connections are made. In circumstances where the network then returns to normal work and school connections afterwards, infection can potentially be introduced into new groups. These effects, however, would be uncertain and remain challenging to resolve quantitatively in real-time.

Where possible, we have taken a data-driven approach to parameterise the system and instruct heterogeneities we expect to be present, such as in age-dependent susceptibility and household composition by age group. Nevertheless, this work has made simplifying assumptions and our results therefore have limitations.

We only considered infection resulting from person-to-person interactions due to household mixing. We did not consider transmission arising from other settings, with it also possible that the formation and ceasing of bubbles may have additional effects on transmission. As an example, infections could be seeded in households due to external contacts arising from the workplace (although there would typically be fewer people attending the workplace during the Christmas and New Year period). Such connectivity bridging households would naturally result in a greater number of contacts being made overall and likely hasten the spread of infection. Perhaps counterintuitively, it could also result in households entering isolation earlier and a more immediate curbing of the outbreak. These nonlinearities exhibited between isolation levels and infection burden merit further exploration.

Our findings may be sensitive to alternative epidemiological model structures and intervention assumptions. Of particular note is adherence to isolation and test-and-trace measures. By November 2020 there was growing evidence for nationally applied NPIs (such as social distancing, self-isolation upon symptom onset and household quarantine) having helped reduce the spread of SARS-CoV-2 [34–36]. Though we provide a sensitivity analysis to the adherence parameter (of 30% compared to the value of 70% used in the main analysis), conducting the evaluation of different Christmas bubble options within the time compressed needs of decision making in the policy arena was the priority. The true adherence of the population, and how this could change over time, warrants continued appraisal.

Our assumptions regarding bubble formation were also a simplified representation of the real-world social system. First, we assumed all households would join a Christmas bubble, which would not necessarily be the case. Second, we did not include the possibility of pre-existing household units splitting apart to join their own independent bubbles over the course of the five day window. Third, we did not investigate clustered initial infections (which were instead seeded randomly in the population). Finally, we did not consider childcare bubbles [37]. Childcare bubbles provided informal childcare (informal in the sense the childcare was unregistered) for anyone under 14 from friends and/or family from one other household. Stipulations included only having one childcare bubble with one other household at a time, meaning no household could be part of more than one childcare bubble. The inclusion of these other types of allowed household bubbles would alter the plausible combinations of households that may form a Christmas bubble. Our findings should be interpreted with due consideration of the pragmatic decisions made to ensure analyses could be delivered in a timely manner.

Models of infectious disease transmission are able to provide quantitative evidence for the potential impact of different intervention options on the future course of an outbreak. This study demonstrates the potential uses of a stochastic individual-based model representing a synthetic population of households, which could be developed swiftly and deployed in situations necessitating assessment of the epidemiological impact of extending contacts beyond the immediate household. Our findings offer general insights on how short term changes to household bubbles might influence the infectious disease dynamics, particularly alterations to the age distributions of epidemiological outcomes of interest.

Acknowledgements

With thanks to Louise Dyson, Matt Keeling, Mike Tildesley and Robin Thompson for helpful discussions and comments on the manuscript.

Author contributions

Edward M. Hill: Conceptualisation, Data curation, Formal analysis, Methodology, Software, Validation, Visualisation, Writing - Original Draft, Writing - Review & Editing.

Financial disclosure

EMH was supported by the Medical Research Council through the COVID-19 Rapid Response Rolling Call [grant number MR/V009761/1]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data availability

All data utilised in this study are publicly available, with relevant references and data repositories stated within the main manuscript and Supporting Information.

Code availability

The code repository for the study is available at https://github.com/EdMHill/SARS-CoV-2_Christmas_bubbles_2020.

Archived code: https://doi.org/10.5281/zenodo.6794136

Competing interests

The author declares that they have no competing interests.

References

[1] Thomas Hale, Noam Angrist, Emily Cameron- Blake, Laura Hallas, Beatriz Kira, Saptarshi Majumdar, Anna Petherick, Toby Phillips, Helen Tatlow, Samuel Webster. Oxford COVID-19 Government Response Tracker, Blavatnik School of Government. (2022). URL www.bsg.ox.ac.uk/covidtracker. [Online] (Accessed: 04 July 2022).

[2] UK Government. Coronavirus (COVID-19) in the UK dashboard: Deaths (2022). URL https://coronavirus.data.gov.uk/details/deaths. [Online] (Accessed: 04 July 2022).

[3] Cabinet Office. Making a Christmas bubble with friends and family (Published 24 November 2020. (2020). URL https://web.archive.org/web/20201129225012/https://www.gov.uk/government/publications/making-a-christmas-bubble-with-friends-and-family/making-a-christmas-bubble-with-friends-and-family#meeting-with-your-christmas-bubble-and-other-friends [Online] (Accessed: 04 July 2022).

[4] Public Health England. Investigation of novel SARS-CoV-2 variant: Variant of Concern 202012/01 (2020). URL https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/959438/Technical_Briefing_VOC_SH_NJL2_SH2.pdf. [Online] (Accessed: 04 July 2022).

[5] Prime Minister's Office, 10 Downing Street. Prime Minister's statement on coronavirus (COVID-19): 19 December 2020. (2020). URL https://www.gov.uk/government/speeches/prime-ministers-statement-on-coronavirus-covid-19-19-december-2020. [Online] (Accessed: 04 July 2022).

[6] Cabinet Office. Making a Christmas bubble with friends and family. (2020). URL https://www.gov.uk/government/publications/making-a-christmas-bubble-with-friends-and-family/making-a-christmas-bubble-with-friends-and-family#households-where-everybody-is-not-in-the-same-christ [Online] (Accessed: 04 July 2022).

[7] Block P, Hoffman M, Raabe IJ, Dowd JB, Rahal C, *et al.* Social network-based distancing strategies to flatten the COVID-19 curve in a post-lockdown world. *Nature Human Behaviour* **4**(6):588–596 (2020). doi:10.1038/s41562-020-0898-6.

[8] Leng T, White C, Hilton J, Kucharski A, Pellis L, *et al.* The effectiveness of social bubbles as part of a Covid-19 lockdown exit strategy, a modelling study. *Wellcome Open Research* **5**:213 (2020). doi:10.12688/wellcomeopenres.16164.1.

[9] Scientific Pandemic Influenza Group on Modelling, Operational sub-group (SPI-M-O) for the Scientific Advisory Group for Emergencies (SAGE). SPI-M-O: Consensus statement on COVID-19, 2 December 2020. (2020). URL https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/945801/S0940_201202_SPI-M-O_Consensus_Statement_FINAL_UPDATEDv2_for_release.pdf. [Online] (Accessed: 04 July 2022).

[10] Office for National Statistics. CT0820_2011 Census - household type, household size and age of usual residents (people) - England and Wales (2018). URL https://www.ons.gov.uk/peoplepopulationandcommunity/housing/adhocs/008635ct08202011censushouseholdtypehouseholdsizeandageofusualresidentspeopleenglandandwales?:uri=peoplepopulationandcommunity/housing/adhocs/008635ct08202011censushouseholdtypehouseholdsiz. [Online] (Accessed: 04 July 2022).

[11] Department of Health and Social Care. Making a support bubble with another household. (2021). URL https://www.gov.uk/guidance/making-a-support-bubble-with-another-household. [Online] (Accessed: 04 July 2022).

[12] Lauer SA, Grantz KH, Bi Q, Jones FK, Zheng Q, *et al.* The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Annals of Internal Medicine* **172**(9):577–582 (2020). doi:10.7326/M20-0504.

[13] He X, Lau EHY, Wu P, Deng X, Wang J, *et al.* Temporal dynamics in viral shedding and trans-

missibility of COVID-19. *Nature Medicine* **26**(5):672–675 (2020). doi:10.1038/s41591-020-0869-5.

[14] Ashcroft P, Huisman JS, Lehtinen S, Bouman JA, Althaus CL, *et al.* COVID-19 infectivity profile correction. *Swiss Medical Weekly* **150**:w20336 (2020). doi:10.4414/smw.2020.20336.

[15] Buitrago-Garcia D, Egli-Gany D, Counotte MJ, Hossmann S, Imeri H, *et al.* Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLOS Medicine* **17**(9):e1003346 (2020). doi:10.1371/journal.pmed.1003346.

[16] McEvoy D, McAloon CG, Collins AB, Hunt K, Butler F, *et al.* The relative infectiousness of asymptomatic SARS-CoV-2 infected persons compared with symptomatic individuals: A rapid scoping review. *medRxiv* page 2020.07.30.20165084 (2020). doi:10.1101/2020.07.30.20165084.

[17] Davies NG, Klepac P, Liu Y, Prem K, Jit M, *et al.* Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nature Medicine* **26**(8):1205–1211 (2020). doi:10.1038/s41591-020-0962-9.

[18] Lopez Bernal J, Panagiotopoulos N, Byers C, Garcia Vilaplana T, Boddington N, *et al.* Transmission dynamics of COVID-19 in household and community settings in the United Kingdom, January to March 2020. *Eurosurveillance* **27**(15):pii=2001551 (2022). doi:10.2807/1560-7917.ES.2022.27.15.2001551.

[19] Public Health England. Stay at home: guidance for households with possible or confirmed coronavirus (COVID-19) infection (05 November 2020 update). (2020). URL https://web.archive.org/web/20201130170108/https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance/stay-at-home-guidance-for-households-with-possible-coronavirus-covid-19-infection. [Online] (Accessed: 04 July 2022).

[20] Department of Health and Social Care. NHS test and trace: how it works (08 October 2020 update). (2020). URL https://web.archive.org/web/20201129190212/https://www.gov.uk/guidance/nhs-test-and-trace-how-it-works. [Online] (Accessed: 04 July 2022).

[21] Office for National Statistics. Coronavirus and the social impacts on Great Britain: 13 November 2020 (2020). URL https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/healthandwellbeing/bulletins/coronavirusandthesocialimpactsongreatbritain/13november2020#lockdown-and-preventative-measures. [Online] (Accessed: 04 July 2022).

[22] Office for National Statistics. Coronavirus (COVID-19) Infection Survey, UK: 26 November 2020 (2020). URL https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurveypilot/26november2020#test-sensitivity-and-specificity. [Online] (Accessed: 04 July 2022).

[23] Holborow A, Asad H, Porter L, Tidswell P, Johnston C, *et al.* The clinical sensitivity of a single SARS-CoV-2 upper respiratory tract RT-PCR test for diagnosing COVID-19 using convalescent antibody as a comparator. *Clinical medicine (London, England)* **20**(6):e209–e211 (2020). doi:10.7861/clinmed.2020-0555.

[24] Hellewell J, Russell TW, Beale R, Kelly G, Houlihan C, *et al.* Estimating the effectiveness of routine asymptomatic PCR testing at different frequencies for the detection of SARS-CoV-2 infections (version 1). *medRxiv* page 2020.11.24.20229948 (2020). doi:10.1101/2020.11.24.20229948version1.

[25] Wikramaratna PS, Paton RS, Ghafari M, Lourenço J. Estimating the false-negative test probability of SARS-CoV-2 by RT-PCR. *Eurosurveillance* **25**(50):2000568 (2020). doi:10.2807/1560-7917.ES.2020.25.50.2000568.

[26] Keeling MJ, Dyson L, Guyver-Fletcher G, Holmes A, Semple MG, *et al.* Fitting to the UK COVID-19 outbreak, short-term forecasts and estimating the reproductive number. *Statistical Methods in Medical Research* (2022). doi:10.1177/09622802211070257.

[27] UK Government. Coronavirus (COVID-19) in the UK dashboard: Healthcare (2022). URL

https://coronavirus.data.gov.uk/details/healthcare. [Online] (Accessed: 04 July 2022).

[28] Danon L, Lacasa L, Brooks-Pollock E. Household bubbles and COVID-19 transmission: insights from percolation theory. *Philosophical Transactions of the Royal Society B: Biological Sciences* **376**(1829):20200284 (2021). doi:10.1098/rstb.2020.0284.

[29] Hilton J, Riley H, Pellis L, Aziza R, Brand S, *et al.* A computational framework for modelling infectious disease policy based on age and household structure with applications to the COVID-19 pandemic. *ArXiv e-prints* (2022).

[30] House T, Keeling MJ. Deterministic epidemic models with explicit household structure. *Mathematical Biosciences* **213**(1):29–39 (2008). doi:10.1016/j.mbs.2008.01.011.

[31] Glass K, McCaw J, McVernon J. Incorporating population dynamics into household models of infectious disease transmission. *Epidemics* **3**(3-4):152–158 (2011).

[32] Hilton J, Keeling MJ. Incorporating household structure and demography into models of endemic disease. *Journal of The Royal Society Interface* **16**(157):20190317 (2019). doi:10.1098/rsif.2019.0317.

[33] Whitty CJM. What makes an academic paper useful for health policy? *BMC Medicine* **13**:301 (2015). doi:10.1186/s12916-015-0544-8.

[34] Ferguson NM, Laydon D, Nedjati-Gilani G, Imai N, Ainslie K, *et al.* Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand. *Imperial College London* (2020). doi:10.25561/77482.

[35] Davies NG, Kucharski AJ, Eggo RM, Gimma A, Edmunds WJ, *et al.* Effects of non-pharmaceutical interventions on COVID-19 cases, deaths, and demand for hospital services in the UK: a modelling study. *The Lancet Public Health* **5**(7):e375–e385 (2020). doi:10.1016/S2468-2667(20)30133-X.

[36] Keeling MJ, Hill EM, Gorsich EE, Penman B, Guyver-Fletcher G, *et al.* Predictions of COVID-19 dynamics in the UK: Short-term forecasting and analysis of potential exit strategies. *PLOS Computational Biology* **17**(1):e1008619 (2021). doi:10.1371/journal.pcbi.1008619.

[37] Department of Health and Social Care. Making a childcare bubble with another household. (2021). URL https://www.gov.uk/guidance/making-a-childcare-bubble-with-another-household. [Online] (Accessed: 04 July 2022).