Preoperative predictors of instrumental activities of daily living disability in older adults

six months after knee arthroplasty

<u>**Abstract**</u>

Objective: To investigate preoperative predictors of instrumental activities of daily living

(IADL) disability in older adults six months after knee arthroplasty (KA).

Design: Prospective study.

Setting: General hospital with an orthopedic surgery department.

Participants: Two hundred twenty (N = 220) patients $\geq$ 65 years old with total knee

arthroplasty (TKA) or unicompartmental knee arthroplasty (UKA).

Interventions: Not applicable.

Main Outcome Measure: IADL status was evaluated for six activities. Participants chose

"able," "need help," or "unable" according to their capacity of executing these IADL

activities. If they chose "need help" or "unable" for one or more items, they were defined as

"disabled." Their usual gait speed (UGS), range of motion for the knee, isometric knee

extension strength (IKES), pain status, depressive symptoms, pain catastrophizing, and self-

efficacy were evaluated as predictors. Baseline and follow-up assessments were conducted

one month before and six months after KA, respectively. Logistic regression analyses with

IADL status as the dependent variable were performed at follow-up. All models were adjusted

using age, sex, severity of the knee deformity, operation type (TKA or UKA) and

preoperative IADL status as covariates.

Results: In total, 166 patients completed the follow-up assessment, and 83 (50.0%) reported

IADL disability six months after KA. Preoperative UGS, IKES on the non-operated side and

self-efficacy were significantly different between those with a disability at follow-up and

those who did not and were therefore included in logistic regression models as independent

variables. UGS (odds ratio [OR] = 3.22, 95% confidence interval [95% CI] = 1.38–7.56, p =

0.007) was determined as a significant independent variable.

Conclusion: The present study demonstrated the importance of evaluating preoperative gait

speed to predict the presence of IADL disability in older adults six months after KA.

Therefore, those with poorer preoperative mobility should be provided careful postoperative

care and treatment.

Keywords: activities of daily living**,** disability, gait speed, knee arthroplasty, osteoarthritis

<u>Abbreviations:</u>

AUC, area under the curve;

BADL, basic activities of daily living;

CI, confidence interval**;**

GDS-15, fifteen-item Geriatric Depression Scale;

IADL, instrumental activities of daily living;

IKES, isometric knee extension strength;

KA, knee arthroplasty;

KL-score, Kellgren−Lawllence score;

KOOS-pain, pain subscale of the knee injury and osteoarthritis outcome score;

OA, osteoarthritis;

OR, odds ratio;

PCS-6, six-item short form of the pain catastrophizing scale;

POD, postoperative day;

PSEQ-4, four-item short form of the pain self-efficacy questionnaire;

ROC curve, receiver characteristic operating curve;

1   ROM, range of motion;

2   TKA, total knee arthroplasty;

3   UGS, usual gait speed;

4   UKA, unicompartmental knee arthroplasty

Introduction

Knee osteoarthritis (OA) is a prevalent chronic disease related to aging, and the number of

older persons with knee OA, who experience daily life disability, is increasing globally.[1, 2]

Knee arthroplasty (KA), including total knee arthroplasty (TKA) and unicompartmental knee

arthroplasty (UKA), is a well-established operative treatment for degenerative knee joint

disease in patients suffering from pain and disability. However, 10–20% of patients reportedly

have subjective functional limitations after KA.[3-6] Considering the increasing demand of KA

for older adults, it is necessary to strengthen countermeasures against postoperative

disabilities that limit activities of daily life.

For the aging population, independence in performing activities of daily living is crucial.

Basic activities of daily living (BADL) are fundamental skills required to independently care

for oneself, such as toileting and dressing;[7] whereas, instrumental activities of daily living

(IADL) are more complex activities related to the ability to live independently, such as

shopping, transportation, and performing household chores.[8] Thus, IADL disability usually

precedes that of the BADL.[9] Various cohort studies have shown that disability in IADL can

affect all-cause mortality,[10] the rate of cognitive functional decline,[11] and quality of life,[12]

having a substantial impact on geriatric mental and physical health.

Several systematic reviews have described various factors, such as functional, pain and

psychological statuses, as predictors of disability after KA.[13-15] However, most studies defined

disability as an impairment in both BADL and IADL. Considering the hierarchical

relationship between IADL and BADL,[9] determining the predictors of IADL disability in the

geriatric population after KA is potentially useful for both preoperative and postoperative

management. Multiple meta-analyses regarding postoperative rehabilitation revealed that

most intervention periods have been set within six months after KA,[16, 17] meaning that

disability should be addressed within a limited period. Additionally, it is desirable for patients

with an elevated risk for IADL disability to be screened early, preferably before surgery.

Studies regarding IADL disability in older persons undergoing KA are few,[18] and preoperative

predictors for disability in IADL are unclear. This study investigates the preoperative

predictors of IADL disability in older adults six months after KA. We hypothesized that

certain motor functions and psychological factors hold a predictive power of IADL disability.

Methods

Participants

Participants who underwent unilateral TKA or UKA between May 17, 2019 and June 30,

2021 at Shonan Kamakura General Hospital ere included in this study upon meeting the

following inclusion criteria: (1) being ≥ 65 years of age and (2) diagnosed with knee OA

based on the guidelines[19]: a radiologic Kellgren–Lawrence score (KL-score) of ≥ 2 (definite

osteophytes and narrowed joint space) and knee pain for > 3 months. Patients were excluded

by the following exclusion criteria: (1) being diagnosed with post-traumatic OA; (2)

rheumatoid arthritis; (3) cognitive decline (mini-mental state examination score < 23); (4)

serious pathologies such as cancer, and neurological fallout that could affect the test

performance such as muscle paralysis; (5) history of lower extremity operation except KA;

(6) preoperative BADL disability (Barthel index < 100); and (7) intention of having a TKA or

UKA on the other knee before our study's assessment. 321 patients were selected at the

beginning of the study. Eighty-six of them were excluded because of: a history of lower

extremity operation outside the knee joint region (n = 32), rheumatoid arthritis (n = 19),

revision TKA or UKA (n = 18), Barthel index <100 (n = 7), mini-mental state examination

score < 23 (n = 6), and post-traumatic knee OA (n = 4). Of the 235 patients who were invited

to take part in the current study, 15 refused, leaving 220 participants.

Study design

This was a prospective study conducted and reported using the STROBE guidelines.[20] The

institutional review board at Tokushukai Group Ethics Committee (NO. (No.

TGE01198-024)) approved the study design, and all participants gave written informed

consent. This study was conducted in accordance with the Declaration of Helsinki. All

participants were assessed by well-trained physical therapists a month before the operation as

a baseline measurement and six months after the operation as a follow-up measurement. They

were managed by using a set clinical pathway of our institution and permitted full weight

bearing on the postoperative day (POD) 1. Patients started gait exercises with a walker-cane

on POD1 and a single cane on POD2. Ascending and descending stairs were practiced on

POD5. Patients who could walk with a single cane independently were discharged by

POD10–16 for TKA and 5–12 for UKA. Baseline characteristics such as age, sex, body mass

index, bilateral KL-score, Charlson comorbidity index,[21] and intra-operative information were

obtained from clinical records and used as possible covariates.

Measurements

IADL status

The Lawton and Brody IADL scale[8] investigated independent living skills by examining six

activity items (shopping, preparing food, housekeeping, using transportation, doing laundry,

and handling finances). Using a telephone and taking medications were excluded because

these activities would not be completely affected by symptoms of knee OA. Participants were

questioned to score their capability of performing these tasks trichotomously, with either

"able," "need help," or "unable" as choices for answers. We defined "disabled" participants as

those who answered "need help" or "unable" to perform one or more items; they were defined

as "non-disabled" otherwise, on the basis of previous studies.[22, 23]

Motor functions

Their usual gait speed (UGS) was measured using a 5 m timed gait test. Participants were

instructed to walk straight on an 11 m flat surface at their usual pace. A gait speed of 5 m was

measured in the middle of the walkway.[24] A goniometer measured the range of motion

(ROM) for knee extension and flexion,[25] while a handheld dynamometer [a] assessed the

isometric knee extension strength (IKES) to quantify knee muscle power.[26] ROM and IKES

were measured on both the operated and unoperated knees.

Pain status

The pain was evaluated by using the pain subscale of the Japanese Knee Injury and

Osteoarthritis Outcome Score (KOOS-pain).[27] Participants were asked about their pain in

various kinds of situations one week before the evaluation date. Each question was scored in

the range 0–4. The normalized total score is converted to 100 points, with higher scores

indicating less pain. KOOS-pain has satisfactory internal consistency (Cronbach's alpha =

0.90) and correlates with the body pain subscale of the 36-item short form health survey ($r =$

$0.67, p < 0.01$).[27]

Psychological factors

Depressive symptoms, pain catastrophizing, and self-efficacy were evaluated as potential

psychological factors on the basis of a previous systematic review.[15] The Japanese version of

the fifteen-item Geriatric Depression Scale (GDS-15) [28] was used for the assessment of

depressive symptoms. The GDS-15 has satisfactory internal consistency (Cronbach's alpha =

0.83) and high detectability of depression (area under the curve [AUC] of the receiver

operating characteristic curve [ROC curve] = 0.96). [28] The participants were questioned with

"yes" or "no" answer choices. The total score of 0 to 15 was assigned with higher scores

indicating more depressive symptoms.

The Japanese version of the six-item short form of the Pain Catastrophizing Scale (PCS-6) [29]

was used for the assessment of pain catastrophizing. PCS-6 had satisfactory internal

consistency (Cronbach's alpha = 0.90)[29] and correlated to pain status ($r = 0.30$, $p < 0.001$).[30]

Participants were required to evaluate their experience of six emotions and thoughts during

pain. Each item is evaluated on a scale of 0 ("not at all") to 4 ("all the time"). The total score

of 0 to 24 was assigned with higher scores representing higher levels of catastrophe.

The Japanese version of the four-item short form of the Pain Self-Efficacy Questionnaire

(PSEQ-4) [31] was used for the assessment of self-efficacy. The PSEQ-4 had satisfactory

internal consistency (Cronbach's alpha = 0.90) and correlated to pain status ($r = -0.35$, $p <$

1 0.001).[30] The PSEQ-4 is comprised of four questions measuring the participant's confidence

2 when conducting specific activities despite the pain. Each item was evaluated on a scale of 0

3 ("not at all confident") to 6 ("completely confident"). The total score of 0 to 24 was assigned

4 with higher scores representing more self-efficacy.

6 ***Statistical analysis***

7 The participants were divided into two groups according to their IADL status six months after

8 KA: the disabled (answered "need help" or "unable" for one or more items) and non-disabled

9 (answered "able" for all items) groups. All preoperative baseline characteristics and

10 measurements were compared between the two groups using the Mann–Whitney U test, chi-

11 square, or student's t-test. In the disabled group, the impairment of each IADL tested at

12 follow-up was recorded. Baseline characteristics were also compared between those who were

13 followed up and dropped out.

15 Measurements with $p < 0.05$ in the two comparative groups were represented nominally based

16 on the cutoff value for determining the presence of IADL disability at six postoperative

17 months, calculated from the ROC curve. The ideal cutoff value for balancing the sensitivity

and specificity of a test was determined as the point on the curve closest to (0, 1).[32]

Thereafter, logistic regression was used to predict the presence of IADL disability after six

months of KA (dependent variable). The independent variables were included in each model

separately, using nominal scales. In model 1, age, sex, KL-score of both knees and operation

type (TKA or UKA) were used as covariates. In model 2, the presence of IADL disability

before the operation was added to model 1. KL-score on the non-operated side for participants

with a history of TKA or UKA was treated as score 0.

Additionally, participants were categorized into four subgroups based on pre- and

postoperative IADL status: group 1 without IADL disability; group 2 with IADL disability

before operation only; group 3 with IADL disability after operation only; and group 4 with

IADL disability before and after the operation. UGS and KOOS-pain were compared pre- and

postoperatively in each group to confirm the treatment effect of KA using the paired *t*-test.

The baseline characteristics in the four subgroups were also compared using the chi-square

goodness of fit test, Fisher's exact test, one-way analysis of variance followed by Tukey's test,

or Kruskal–Wallis test followed by the Steel–Dwass test. All data were analyzed using R

version 4.0.3.[b] A p-value < 0.05 was used to indicate statistical significance.

In order to decide the sample size for logistic regression analysis, the

$number\ of\ cases\ (N)\ =\ 10\,k/p$ are needed; k is the number of independent variables as

covariates and p is defined as a ratio of responders to non-responders at the follow-up

points.[33] We predicted that eight factors out of all the variables would be required to avoid

overfitting.[34] In a previous report (n = 407), 60.9% of older adults with joint pain had IADL

disability.[22] Based on this study, we hypothesized that the ratio p of non-disabled to disabled

was 1:1.5 and calculated the minimum number of participants required for our study as

$10 \times 8/0.6 = 134$.

Results

Of the 220 participants who joined the study, 166 (75.5%) completed the follow-up

assessment after TKA or UKA. Fifty-four (24.5%) participants dropped out, namely due to

incomplete follow-up (n = 21), reparative surgery on the other knee within six months (n =

14), cancelation of the operation (n = 9), onset of other diseases not related to KA (n = 4),

periprosthetic joint infection (n = 3), and periprosthetic fracture (n = 3). Baseline

characteristics were not significantly different between those who were followed up and

dropped out.

Of participants who were followed up, 83 (50.0%) had IADL disability six months after KA.

Sixty-one (69.3%) struggled with shopping, and 47 (53.4%) struggled using transportation

(Table 1). The non-disabled group was significantly younger ($p = 0.004$) with a larger

percentage of women (85.5% vs 69.9%; $p = 0.03$) when compared to the IADL disabled

group. The percentage of patients showing IADL disability before surgery was 20.5% and

72.3% ($p < 0.001$) in the non-disabled and disabled groups, respectively. In the IADL non-

disabled group, the values of UGS, IKES on the non-operated side and PSEQ-4 were

significantly higher. (Table 2)

Based on the two groups' comparison, the cutoff values of UGS, IKES on the non-operated

side, and PSEQ-4 were calculated by inspecting the ROC curve to discern the presence of

IADL disability at six months after KA. The optimal cutoff value was 1.0 m/sec for UGS

(AUC, 0.71; 95% confidential interval [CI], 0.63–0.79), 1.0 Nm/kg for IKES on the non-

operated side (AUC, 0.61; 95% CI, 0.53–0.70) and 13.0 for PSEQ-4 (AUC, 0.60; 95% CI

0.52–0.69) (Table 3). Table 4 shows the odds ratios (ORs) of independent variables that were

expressed nominally. UGS was a significant independent variable in models 1 and 2, and

IKES on the non-operated side and PSEQ-4 were significant in model 1 but not in model 2.

In the baseline characteristics comparisons among subgroups, significant differences in the

age and number of women were detected. Regarding KL-score on the non-operated side, there

was a cell with 0 frequency, making statistical analysis impossible (Table 5). The values of

KOOS-pain were significantly higher in all subgroups after operation versus before it. The

values of UGS after operation were significantly higher than before, group 3 being the

exception (Table 6).

Discussion

This study investigated preoperative predictors of IADL disability in older adults after six

months of TKA or UKA, where half of the participants reported disabilities. Gait speed, knee

muscle strength, and self-efficacy were potential predictors. However, the logistic regression

analysis adjusted for covariates showed that only gait speed was a true predictor of IADL

disability, six months after KA. Thus, the results of this study partially supported our

hypothesis.

Our study suggested that the cutoff value for preoperative UGS was 1.0 m/sec. The AUC

needed to be at least 0.7 to be considered satisfactory.[32] Thus, our study is significant, as the

cutoff value for preoperative UGS exceeded the minimum criterion. A previous prospective

study in community-dwelling older adults showed that participants with UGS below 1.0

m/sec had a sharper increase in the 2-year incidence of IADL disability than those above.[35]

Other cohort studies involving older adults have also revealed a 1.0 m/sec cutoff value for risk

prediction of health events.[36, 37] Considering the incidence of BADL disability, 0.8 m/sec has

been suggested as a more sensible cutoff value.[38] Therefore, considering the hierarchical

relationship between IADL and BADL,[9] our cutoff value for UGS was valid.

The predictive power of PSEQ-4 and IKES on the non-operated side was low based on the

AUCs of ROC curves, something we could not wholly explain. However, previous studies

showed that self-efficacy[15] and muscle strength[14] were predictors of decreased functional

status after KA. Thus, these factors should not be overlooked when considering disability

after surgery.

George et al. have shown that the number of older adults with preoperative IADL disability

improved after TKA.[18] In contrast, patients with BADL disability (100 < Barthel Index) were

excluded. Most participants did not have moderate to severe comorbidity preoperatively;

however, the proportion of older adults in our study who presented with IADL disability

before operation worsened after TKA or UKA. The aforementioned study assessed fewer

IADLs and included younger participants than those in our study, which we believe could

have contributed to the different results. In our participants, the main items of IADL disability

were outdoor activities, such as shopping (69.3%) and using transportation (53.4%), which

require gait proficiency. When comparing subgroups, only those presenting with new IADL

disability after operation (group 3) failed to improve UGS after KA. In group 4, UGS

significantly improved after KA despite having the lowest baseline and follow-up values of

the four groups. As a result, we believe that those who have a slow preoperative gait speed,

which does not increase to a certain value postoperatively, tend to present IADL disability at

six months after KA, even if their pain status has improved.

Some studies have shown that preoperative functional limitation affects postoperative

disability.[39-41] In contrast to our study, they assessed disability using patient-reported outcome

measures such as Western Ontario and McMaster University's Osteoarthritis Index[42] and

Oxford Knee Score[43]. These tools assess subjective difficulty in various activities and

situations, but specific modifiable factors affecting disability in daily life remained unclear.

The strength of our findings lies in the predictive power of preoperative gait speed as a

modifiable factor for IADL disability in older adults after six months of KA. Furthermore, the

specific value of UGS was suggested. A clinical practice guideline[44] recommended

preoperative physical therapy to improve postoperative outcome after KA; however, the

specific regimen is unclear. Our results could be useful for preoperative planning and

intervention. In particular, improving preoperative gait speed may enhance the IADL status.

Additionally, the systematic review by Sayah et al. revealed that functional limitations

including IADL disability improved drastically in six months and gradually keep improving 2

years after TKA.[45] Therefore, our study population, especially participants in group 3, could

potentially improve IADL disability six months postoperatively. However, considering the

relationship between IADL disability, adverse events,[10, 12] and general rehabilitative

intervention periods after KA,[16, 17] such patients are at high risk for adverse events and require

careful follow-up even beyond six months postoperatively.

Study limitations

The present study had some limitations. First, since IADL disability was treated as a

categorical value, we could not determine its severity. Hence, previous cohort studies have

shown that the presence of as few as one IADL disability affected all-cause mortality[10] and

the cognitive function decline rate.[11] Second, ours was a single-center analysis; the area

distribution where the participants lived was limited. Third, our study population included

those with a history of KA, and we could not exclude the effect of past KA on IADL

disability. However, the number of those with a history of TKA or UKA was similar between

IADL disabled and IADL non-disabled groups.

Conclusion

Our study revealed the importance of evaluating UGS before a TKA or UKA as a predictor of

six-month postoperative IADL disability in older adults, in addition to preoperative IADL

status. Those with poor mobility before surgery should be carefully rehabilitated after KA.

Future studies investigating IADL disability during long-term follow-up periods are needed to

further validate our findings.

Suppliers list:

a: μ-tus F-1, Anima Corporation, 3-65-1, Shimoishihara, Chofu, Tokyo, 182-0034, Japan

b: R version 4.0.3, R Foundation for Statistical Computing, Welthandelsplatz 1, 1020 Vienna,

Austria

References

1. Bijlsma JW, Berenbaum F, Lafeber FP. Osteoarthritis: an update with relevance for clinical practice. Lancet. 2011;377(9783):2115-26. doi: 10.1016/S0140-6736(11)60243-2.

2. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study. 2015. Lancet. 2016;388(10053):1545-602. doi: 10.1016/S0140-6736(16)31678-6.

3. Bourne RB, Chesworth BM, Davis AM, Mahomed NN, Charron KD. Patient satisfaction after total knee arthroplasty: who is satisfied and who is not? Clin Orthop Relat Res. 2010;468(1):57-63. doi: 10.1007/s11999-009-1119-9.

4. Noble PC, Conditt MA, Cook KF, Mathis KB. The John Insall Award: Patient expectations affect satisfaction with total knee arthroplasty. Clin Orthop Relat Res. 2006;452:35-43. doi: 10.1097/01.blo.0000238825.63648.1e.

5. Von Keudell A, Sodha S, Collins J, Minas T, Fitz W, Gomoll AH. Patient satisfaction after primary total and unicompartmental knee arthroplasty: an age-dependent analysis. Knee. 2014;21(1):180-4. doi: 10.1016/j.knee.2013.08.004. Epub 2013 Aug 15.

6. Lee M, Huang Y, Chong HC, Ning Y, Lo NN, Yeo SJ. Predicting Satisfaction for

Unicompartmental Knee Arthroplasty Patients in an Asian Population. J Arthroplasty.

2016;31(8):1706-10. doi: 10.1016/j.arth.2016.02.005. Epub 2016 Feb 8.

7. Kats S, Ford AB, Moskowitz RW, Jackson BA, Jafee MW. Studies of illness in the aged.

The index of ADL: A standardized measure of biological and psychosocial function

JAMA. 1963;185:914-9. doi: 10.1001/jama.1963.03060120024016.

8. Lawton MP, Brody EM. Assessment of older people: self-maintaining and instrumental

activities of daily living. Gerontologist. 1969;9(3):179-86.

9. Edjolo A, Proust-Lima C, Delva F, Dartigues JF, Pérès K. Natural History of Dependency

in the Elderly: A 24-Year Population-Based Study Using a Longitudinal Item Response

Theory Model. Am J Epidemiol. 2016;183(4):277-85. doi: 10.1093/aje/kwv223. Epub

2016 Jan 28.

10. Hennessy S, Kurichi JE, Pan Q, Streim JE, Bogner HR, Xie D et al. Disability Stage is an

Independent Risk Factor for Mortality in Medicare Beneficiaries Aged 65 Years and

Older. PM R. 2015;7(12):1215-25. doi: 10.1016/j.pmrj.2015.05.014. Epub 2015 May 21.

11. Rajan KB, Hebert LE, Scherr PA, Mendes de Leon CF, Evans DA. Disability in basic and

instrumental activities of daily living is associated with faster rate of decline in cognitive

function of older adults. J Gerontol A Biol Sci Med Sci. 2013;68(5):624-30. doi: 10.1093/gerona/gls208.

12. Williams JS, Egede LE. The Association Between Multimorbidity and Quality of Life, Health Status and Functional Disability. Am J Med Sci. 2016;352(1):45-52. doi: 10.1016/j.amjms.2016.03.004. Epub 2016 Mar 18.

13. Lungu E, Vendittoli PA, Desmeules F. Preoperative Determinants of Patient-reported Pain and Physical Function Levels Following Total Knee Arthroplasty: A Systematic Review. Open Orthop J. 2016;10:213-31. doi: 10.2174/1874325001610010213. eCollection 2016.

14. Devasenapathy N, Maddison R, Malhotra R, Zodepy S, Sharma S, Belavy DL. Preoperative Quadriceps Muscle Strength and Functional Ability Predict Performance-Based Outcomes 6 Months After Total Knee Arthroplasty: A Systematic Review. Phys Ther. 2019;99(1):46-61. doi: 10.1093/ptj/pzy118.

15. Vissers MM, Bussmann JB, Verhaar JA, Busschbach JJ, Bierma-Zeinstra SM, Reijman M. Psychological factors affecting the outcome of total hip and knee arthroplasty: a systematic review. Semin Arthritis Rheum. 2012;41(4):576-88. doi: 10.1016/j.semarthrit.2011.07.003. Epub 2011 Oct 28.

16. Dávila Castrodad IM, Recai TM, Abraham MM, et al. Rehabilitation protocols following

total knee arthroplasty: a review of study designs and outcome measures. Ann Transl

Med. 2019;7(Suppl 7):S255. doi: 10.21037/atm.2019.08.15.

17. Li D, Yang Z, Kang P, Xie X. Home-Based Compared with Hospital-Based

Rehabilitation Program for Patients Undergoing Total Knee Arthroplasty for

Osteoarthritis: A Systematic Review and Meta-analysis of Randomized Controlled Trials.

Am J Phys Med Rehabil. 2017;96(6):440-7. doi: 10.1097/PHM.0000000000000621.

18. George LK, Ruiz D, Sloan FA. The effects of total knee arthroplasty on physical

functioning in the older population. Arthritis Rheum. 2008;58(10):3166-71. doi:

10.1002/art.23888.

19. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K et al. Development of criteria

for the classification and reporting of osteoarthritis. Classification of osteoarthritis of the

knee. Diagnostic and Therapeutic Criteria Committee of the American Rheumatism

Association. Arthritis Rheum 1986;29(8):1039-49.

20. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of

Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting

observational studies. J Clin Epidemiol. 2008;61(4):344-9. doi:

10.1016/j.jclinepi.2007.11.008.

21. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying

prognostic comorbidity in longitudinal studies: development and validation. J Chronic

Dis. 1987;40(5):373-83. doi: 10.1016/0021-9681(87)90171-8.

22. Hermsen LA, Leone SS, Smalbrugge M, Dekker J, van der Horst HE. Frequency, severity

and determinants of functional limitations in older adults with joint pain and comorbidity:

results of a cross-sectional study. Arch Gerontol Geriatr. 2014;59(1):98-106. doi:

10.1016/j.archger.2014.02.006. Epub 2014 Feb 28.

23. Lee MC, Hsu CC, Tsai YF, Chen CY, Lin CC, Wang CY. Criterion-Referenced Values of

Grip Strength and Usual Gait Speed Using Instrumental Activities of Daily Living

Disability as the Criterion. J Geriatr Phys Ther. 2018;41(1):14-9. doi:

10.1519/JPT.0000000000000106.

24. Shinkai S, Watanabe S, Kumagai S, et al. Walking speed as a good predictor for the onset

of functional dependence in a Japanese rural community population. Age Ageing.

2000;29(5):441-6. doi: 10.1093/ageing/29.5.441.

25. Norkin CC, White DJ. Measurement of joint motion : a guide to goniometry. 2nd ed.

Philadelphia: F.A. Davis; 1995.

26. Katoh M, Yamasaki H. Comparison of Reliability of Isometric Leg Muscle Strength

Measurements Made Using a Hand-Held Dynamometer with and without a Restraining

Belt. J Phys Ther Sci. 2009;21(1):37-42.

27. Nakamura N, Takeuchi R, Sawaguchi T, Ishikawa H, Saito T, Goldhahn S. Cross-cultural

adaptation and validation of the Japanese Knee Injury and Osteoarthritis Outcome Score

(KOOS). J Orthop Sci. 2011;16(5):516-23. doi: 10.1007/s00776-011-0112-9. Epub 2011

Jul 16.

28. Sugishita K, Sugishita M, Hemmi I, Asada T, Tanigawa T. A Validity and Reliability

Study of the Japanese Version of the Geriatric Depression Scale 15 (GDS-15-J). Clin

Gerontol. 2017;40(4):233-40. doi: 10.1080/07317115.2016.1199452. Epub 2016 Jun 29.

29. Nishigami T, Mibu A, Tanaka K, Yamashita Y, Watanabe A, Tanabe A. Psychometric

properties of the Japanese version of short forms of the Pain Catastrophizing Scale in

participants with musculoskeletal pain: A cross-sectional study. J Orthop Sci.

2017;22(2):351-6. doi: 10.1016/j.jos.2016.11.015. Epub 2017 Jan 10.

30. McWilliams LA, Kowal J, Wilson KG. Development and evaluation of short forms of the

Pain Catastrophizing Scale and the Pain Self-efficacy Questionnaire. Eur J Pain.

2015;19(9):1342-9. doi: 10.1002/ejp.665. Epub 2015 Mar 11.

31. Adachi T, Enomoto K, Yamada K, et al. Evaluating the psychometric properties of two-

item and four-item short forms of the Japanese Pain Self-Efficacy Questionnaire: a cross-

sectional study. J Anesth. 2019;33(1):58-66. doi: 10.1007/s00540-018-2583-8. Epub 2018

Nov 16.

32. Akobeng AK. Understanding diagnostic tests 3: Receiver operating characteristic curves.

Acta Paediatr 2007;96(5):644-7. doi.org/10.1111/j.1651-2227.2006.00178.x.

33. Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the

number of events per variable in logistic regression analysis. J Clin Epidemiol.

1996;49(12):1373-9. doi: 10.1016/s0895-4356(96)00236-3.

34. Vittinghoff E, McCulloch CE. Relaxing the rule of ten events per variable in logistic and

Cox regression. Am J Epidemiol. 2007;165(6):710-8. doi: 10.1093/aje/kwk052. Epub

2006 Dec 20.

35. Donoghue OA, Savva GM, Cronin H, Kenny RA, Horgan NF. Using timed up and go and

usual gait speed to predict incident disability in daily activities among community-

dwelling adults aged 65 and older. Arch Phys Med Rehabil. 2014;95(10):1954-61. doi:

10.1016/j.apmr.2014.06.008. Epub 2014 Jun 28.

36. Cesari M, Kritchevsky SB, Penninx BW, et al. Prognostic value of usual gait speed in

well-functioning older people--results from the Health, Aging and Body Composition

Study. J Am Geriatr Soc. 2005;53(10):1675-80. doi: 10.1111/j.1532-5415.2005.53501. x.

37. Bohannon RW. Comfortable and maximum walking speed of adults aged 20-79 years:

reference values and determinants. Age Ageing. 1997;26(1):15-9. doi:

10.1093/ageing/26.1.15.

38. Abellan van Kan G, Rolland Y, Andrieu S, et al. Gait speed at usual pace as a predictor of

adverse outcomes in community-dwelling older people an International Academy on

Nutrition and Aging (IANA) Task Force. J Nutr Health Aging. 2009;13(10):881-9. doi:

10.1007/s12603-009-0246-z.

39. Lingard EA, Katz JN, Wright EA, Sledge CB, Kinemax Outcomes Group. Predicting the

outcome of total knee arthroplasty. J Bone Joint Surg Am. 2004;86(10):2179-86. doi:

10.2106/00004623-200410000-00008.

40. Judge A, Arden NK, Cooper C, et al. Predictors of outcomes of total knee replacement

surgery. Rheumatology (Oxford). 2012;51(10):1804-13. doi:

10.1093/rheumatology/kes075. Epub 2012 Apr 24.

41. Desmeules F, Dionne CE, Belzile É, Bourbonnais R, Champagne F, Frémont P.

Determinants of pain, functional limitations and health-related quality of life six months

after total knee arthroplasty: results from a prospective cohort study. BMC Sports Sci

Med Rehabil. 2013;5:2. doi: 10.1186/2052-1847-5-2.

42. Bellamy N, Buchanan WW, Goldsmith CH, Campbell J, Stitt LW. Validation study of

WOMAC: a health status instrument for measuring clinically important patient relevant

outcomes to antirheumatic drug therapy in patients with osteoarthritis of the hip or knee.

J Rheumatol. 1988;15(12):1833-40.

43. Dawson J, Fitzpatrick R, Murray D, Carr A. Questionnaire on the perceptions of patients

about total knee replacement. J Bone Joint Surg Br. 1998;80(1):63-9. doi: 10.1302/0301-

620x.80b1.7859.

44. Jette DU, Hunter SJ, Burkett L, et al. Physical Therapist Management of Total Knee

Arthroplasty. Phys Ther. 2020;100(9):1603-31. doi: 10.1093/ptj/pzaa099.

45. Sayah SM, Karunaratne S, Beckenkamp PR, et al. Clinical Course of Pain and Function

Following Total Knee Arthroplasty: A Systematic Review and Meta-Regression. J

Arthroplasty. 2021;36(12):3993-4002.e37. doi: 10.1016/j.arth.2021.06.019. Epub

2021Jun 29.

Legends of tables

Table 1. Content of disability in IADL disabled group

Table 2. Baseline characteristics of the study population and comparative preoperative

measurements between the IADL non-disabled and disabled groups

Table 3. Variable characteristics to discriminate those with or without IADL disability six

months after KA

Table 4. ORs of independent variables in a logistic regression model with IADL disability status

at six months after KA as dependent variable

Table 5. Demographic comparison in four subgroups

Table 6. Comparison of KOOS-pain and UGS between pre- and postoperative subgroups

Table 1. Content of disability in the IADL-disabled group

Items	IADL disabled 6-months after operation (n=83)*
Preparing food	33 (37.5)
Shopping	61 (69.3)
Housekeeping	40 (45.5)
Doing laundry	6 (6.8)
Using transportation	47 (53.4)
Handling finances	15 (17.0)

* Number (%)

Table 2. Baseline characteristics of the study population and comparative preoperative measurements between the IADL non-disabled and disabled groups

Variable	All (n=166)	IADL non-disabled (n=83)	IADL disabled (n=83)	p-value	Effect size*
Age; mean±SD	74.9± 6.5	73.4± 6.8	76.3 ± 6.2	0.004	0.22
Woman; number (%)	129 (77.7)	58 (69.9)	71 (85.5)	0.03	0.17
Body mass Index (kg/m^2); median (IQR)	24.4 ± 3.4	24.1± 3.0	24.7±3.8	0.2	0.11
Family status (living alone); number (%)	24 (17.5)	16 (19.2)	13 (15.7)	0.68	0.03
Charlson comorbidity index; median (IQR)	0 (0-0)	0 (0–0)	0 (0–0)	1	<0.001
UKA; number (%)	109 (65.7)	57 (68.7)	52 (62.6)	0.51	0.05
KL score on operated side				0.07	0.18
II; number (%)	23 (13.9)	16 (19.2)	7 (8.4)		
III; number (%)	91 (54.8)	46 (55.4)	45 (54.2)		
IV; number (%)/	52 (31.3)	21 (25.3)	31 (37.3)		
KL score on non-operated side				0.57	0.11
History of TKA or UKA on non-operated side; number (%)	35 (21.1)	18 (21.7)	17 (20.5)		
II; number (%)	51 (30.7)	28 (33.7)	23 (27.7)		
III; number (%)	56 (33.7)	28 (33.7)	28 (33.7)		
IV; number (%)	9 (14.5)	9 (10.8)	15 (18.1)		
IADL disabled before operation; number (%)	63 (46.3)	17 (20.5)	60 (72.3)	<0.001	0.51
Motor and knee joint function					

UGS (m/s); mean±SD	0.95 ± 0.27	1.05 ± 0.26	0.86 ± 0.24	<0.001	0.39
IKES on operated side (Nm/kg); median(IQR)	1.01 (0.77–1.28)	1.03 (0.83–1.28)	0.99 (0.69–1.27)	0.26	0.09
IKES on non-operated side (Nm/kg); median (IQR)	1.27 (0.98–1.60)	1.31 (1.09–1.75)	1.21 (0.82–1.49)	0.01	0.2
Knee flexion ROM on operated side (°); median (IQR)	125 (115–135)	125 (120–135)	125 (110–135)	0.19	0.1
Knee flexion ROM on non-operated side (°); median (IQR)	130 (120–140)	135 (125–140)	130 (115–140)	0.06	0.15
Knee extension ROM on operated side (°); median (IQR)	-5 (-10 to -5)	-5 (-10 to -5)	-5 (-10 to -5)	0.14	0.12
Knee extension ROM on non-operated side (°); median (IQR)	-5 (-5 to 0)	-5 (-5 to 0)	-5 (-10 to 0)	0.053	0.15
Pain status					
KOOS pain; mean±SD	45.4 ± 16.9	46.7 ± 18.7	44.1 ± 14.8	0.31	0.08
Psychological status					
GDS-15; median (IQR)	3.0 (1.0–6.0)	3.0 (1.0–6.0)	3.0 (1.0–6.0)	0.16	0.11
PCS-6; median (IQR)	14.0 (9.0-18.0)	14.0 (7.0-18.0)	15.0 (10.0-19.0)	0.16	0.11
PSEQ-4; median (IQR)	14.0 (12.0-18.0)	15.0 (12.5-18.5)	13.0 (11.0-17.5)	0.02	0.18

* Effect size was estimated using *r* for continuous variables and Cramér 's *V* for categorical variables

Table 3. Variable characteristics to discriminate those with or without IADL disability six months after KA

Variables	Cut off	AUC (95%CI)	Sensitivity (%)	Specificity (%)	OR (95% CI)
UGS (m/s)	1.0	0.71 (0.63–0.79)	57.8	79.5	5.32 (2.57–10.6)
IKES on the non-operated side (Nm/kg)	1.0	0.61 (0.53–0.70)	84.3	38.6	3.38 (1.61–7.07)
PSEQ–4 (points)	13	0.60 (0.52–0.69))	69.9	54.2	2.49 (1.29–4.81)

Table 4. ORs of independent variables in a logistic regression model with IADL disability status at six months after KA as dependent variable

	UGS		IKES non-operated side		PSEQ-4	
	Crude OR (95%CI)	Adjusted OR (95%CI)	Crude OR (95%CI)	Adjusted OR (95%CI)	Crude OR (95%CI)	Adjusted OR (95%CI)
	5.32 (2.67–10.6)		3.38 (1.61–7.07)		2.49 (1.29–4.81)	
Model 1*		4.96† (2.31–10.7)		2.85‡ (1.24–6.54)		2.28‡ (1.12–4.64)
Model 2§		3.22∣∣ (1.38–7.56)		1.70 (0.65–4.45)		1.32 (0.58–2.98)

* Adjusted for age(years), sex, KL score on both knees and operation type (TKA or UKA)

† $p<0.001$

‡ $p<0.05$

§ Adjusted for age(years), sex, KL score on both knees, operation type (TKA or UKA) and IADL disabled before operation or not.

∣∣ $p<0.01$

Table 5. Demographic comparison in four subgroups

Variable	Group1* (n=66)	Group 2† (n=17)	Group 3‡ (n=23)	Group 4§ (n=60)	*p*-value	Multiple comparison
Age; mean±SD	72.8±6.6	75.5±7.3	74.7±6.3	76.9±6.2	0.007	1 < 4
Women; number (%)	45 (68.2)	13 (76.5)	17 (73.9)	54 (90.0)	0.02	
Body mass Index (kg/m²); median (IQR)	23.6 (22.4–26.1)	23.4 (21.6–25.7)	24.8 (22.2–26.9)	23.9 (22.5–26.7)	0.8	
Family status (living alone); number (%)	15 (22.7)	1 (5.9)	2 (8.7)	11 (18.3)	0.3	
Charlson comorbidity index; median (IQR)	0 (0–0)	0 (0–1)	0 (0–0.5)	0 (0–0)	0.95	
UKA; number (%)	46 (76.7)	11 (64.7)	16 (69.6)	36 (60.0)	0.68	
KL score on operated side					0.09	
II; number (%)	10 (15.2)	6 (35.3)	2 (8.7)	5 (8.3)		
III; number (%)	39 (59.1)	7 (41.2)	11 (47.8)	34 (56.7)		
IV; number (%)	17 (25.8)	4 (23.5)	10 (43.5)	21 (35.0)		
KL score on non-operated side					–	
History of TKA or UKA on non-operated side; number (%)	15 (22.7)	3 (17.6)	4 (17.4)	13 (21.7)		
II; number (%)	18 (27.3)	10 (58.8)	9 (39.1)	14 (23.3)		
III; number (%)	18 (27.3)	4 (23.5)	7 (30.4)	21 (35.0)		
IV; number (%)	18 (27.3)	0 (0.0)	3 (13.0)	12 (20.0)		

* Those who did not have IADL disability before and after operation

† Those who have IADL disability only before operation

‡ Those who have IADL disability only after operation

§ Those who have IADL disability before and after operation.

Table 6. Comparison of KOOS-pain and UGS between pre- and postoperative subgroups

Variable		Group 1* (n=66)	Group 2† (n=17)	Group 3‡ (n=23)	Group 4§ (n=60)
UGS (m/s)	Baseline‖	1.09±0.23	0.87±0.24	0.95±0.16	0.81±0.26
	Follow-up‖	1.17±0.19	1.11±0.28	1.02±0.26	0.94±0.22
	p-value	0.01	<0.001	0.18	<0.001
	Effect size¶	0.33	0.84	0.29	0.49
KOOS pain (point)	Baseline‖	49.4±18.9	36.6±14.03	48.4±11.1	42.5±15.8
	Follow-up‖	82.2±12.9	81.7±10.7	74.2±12.1	71.3±14.6
	p-value	<0.001	<0.001	<0.001	<0.001
	Effect size¶	0.88	0.94	0.88	0.89

* Those who did not have IADL disability before and after operation

† Those who have IADL disability only before operation

‡ Those who have IADL disability only after operation

§ Those who have IADL disability before and after operation.

‖ Values of baseline and follow-up are shown on mean±SD.

¶ Effect size was estimated using *r*.