Risk factor modelling of antimicrobial resistance in cancer patients: A systematic review

Anders Skyrud Danielsen[1,2], Léa Franconeri[3,4], Samantha Page[3], Anders Eivind Myhre[5], Ragnhild Agathe Tornes[6], Oliver Kacelnik[3], Jørgen Vildershøj Bjørnholt[1,2]

[1]Department of Microbiology, Oslo University Hospital
[2]Institute of Clinical Medicine, Faculty of Medicine, University of Oslo
[3]Department of Infection Control and Preparedness, Norwegian Institute of Public Health
[4]ECDC Fellowship Programme, Field Epidemiology path (EPIET), European Centre for Disease Prevention and Control, (ECDC), Stockholm, Sweden
[5]Department of Haematology, Oslo University Hospital
[6]The Library for the Healthcare Administration, Norwegian Institute of Public Health

Corresponding author: Anders Skyrud Danielsen, andersskyrud.danielsen@fhi.no

Keywords: antimicrobial resistance, infection, mortality, risk factors, cancer, malignancy, systematic review

Abstract

Background

Infections are major causes of disease in cancer patients and pose a major obstacle to the success of cancer care. The global rise of antimicrobial resistance threatens to make these obstacles even greater and hinder continuing progress in cancer care. To prevent and handle such infections, a better understanding of the risk factors are needed. This internally funded systematic review (PROSPERO registration: CRD42021282769) aims to review what risk factors for resistant infections and deaths due to resistant infections have been investigated, and with what methodological approaches.

Methods

We employed two broad searches of antimicrobial resistance in cancer patients, using terms associated with antimicrobial resistant infections, in MEDLINE and Embase through Ovid, in addition to Cinahl through EBSCOhost and Web of Science Core Collection. Primary, observational studies in English from January 2015 to November 2021 on human cancer patients that explicitly modelled infection or mortality associated with antimicrobial resistance in a multivariable model were included. We extracted data on the study populations and their malignancies, risk factors, the microbial aetiology, methods for variable selection, and assessed the risk of bias using the NHLBI Study Quality Assessment Tools.

Results

Two searches yielded a total of 27151 unique records, of which 144 studies were included after screening and reading. Of the outcomes studied, mortality was the most common (68/144, 47%). Forty-five percent (65/144) of the studies focused on haemato-oncological patients, and 27% (39/144) studied several bacteria or fungi. Studies included a median of 200 patients and 46 events. One-hundred-and-three (72%) studies used a p-value-based variable selection. Studies included a median of seven variables in the final (and largest) model, which yielded a median of 6.9 events per variable. An in-depth example of vancomycin-resistant enterococci was reported.

Conclusions

We found the current research on this topic to be heterogeneous, in both the methodological and epidemiological approach. Methodological choices resulting in very diverse models made it difficult or even impossible to draw statistical inferences and summarise what risk factors were of clinical relevance. The development and adherence to more standardised protocols that build on existing literature is urgent.

Background

Cancer patients have higher risk and worse outcome of infectious disease complications, compared with healthy people (1). Autopsy studies have indicated that infections may play a role in more than half of all cancer patient fatalities (1). A recent study has also shown that cancer patients have a higher risk of contracting infections with antimicrobial resistant organisms (2). Importantly, infections often necessitate caregivers to postpone or withhold adequate cancer treatment, which may impair cancer outcome. Clinicians may use risk stratification tools and scoring systems to guide anti-infective therapy. Some examples widely used in the management of infectious complications in cancer patients are the Multinational Association for Supportive Care in Cancer (MASCC) risk indices for febrile neutropenic patients or the Pitt bacteremia score (3,4). These risk indices, however, were developed in relatively small patient cohorts in the early 1990s.

In recent years, there have been major changes in disease-causing microbial ecology, particularly in hospitals (5). Bacteria and fungi are becoming increasingly resistant to antimicrobial drugs, and in Europe alone it is estimated that more than 33,000 people die each year from resistant microbes (6,7). Not only are microbes acquiring antimicrobial resistance, but the microbial spectrum is changing, with an increasing proportion of intrinsically resistant species (8). Infections in cancer patients are increasingly often caused by resistant organisms, which threaten to slow, or even reverse, recent years' advances in the treatment of cancer (9). In other words, there is a pressing need to understand the changing epidemiology of bacterial and fungal infections among cancer patients, so as to design better preventive measures. Central to adapting to this new reality is to understand the mechanisms by which infections occur and cause disease. The research that leads to the discovery and description of these mechanisms, as well as testing them in preliminary models, has been called 'prognostic factor research' by the PROGRESS Group (10). Here, we use the term 'risk factors' as the patients studied only rarely had infections at the start of follow-up. Understanding these risk factors could allow us to update and expand on the currently used risk stratification tools, or even create new ones where needed. Such risk stratification tools or scoring systems are usually based on clinical predictions models, which

typically are regression models where individual-level clinical data are used to predict a clinical outcome of interest (11,12).

In this systematic review we aim to review what risk factors on resistant infections and their outcomes in cancer patients have been investigated, and with what methodological approaches, to guide this research in the future.

Methods

Protocol registration and reporting standards

To review what risk factors for resistant infections and corresponding mortality in cancer patients have been investigated in 2015-2021, we conducted a systematic review that used a broad and extensive search. This systematic review was registered at PROSPERO (ID: CRD42021282769) (13) and follows the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline (checklist can be found in Supplementary Material 6) (14).

Search strategy

Our search strategy was implemented in three steps. First, we performed preliminary searches in PubMed to identify some main keywords to be included. We then performed a search and, after screening and sorting the results of the search, we manually reviewed the records and references to discover keywords that were not covered by our initial search. We then performed a second search designed to expand the first search findings.

The first search was conducted by a research librarian June 22th to 24th, 2021. It consisted of terms covering all cancers and terms covering antibiotic resistance and infections. A spectrum of synonyms with appropriate truncations and proximity operators was used for searching title, abstract, and author keywords. In addition, controlled subject headings were searched when available. The search strategy was tailored to each database's search interface. The search was run in the OVID MEDLINE and OVID Embase , in addition to EBSCO Cinahl and Web of Science Core Collection (Science Citation Index Expanded, Social Sciences Citation Index, Arts & Humanities Citation Index and Emerging Sources Citation Index - searched simultaneously). The strategies were limited to Danish, English, Norwegian, Spanish, and Swedish. They also included a time limit for publications from 2015 onwards. A total of 25 881 records were retrieved. After removing duplicate records in EndNote, 14 153 references were identified.

The second search expanded on this by including specific antibiotics, resistance mechanisms, and bacteria and fungi often associated with either acquired resistance or high levels of intrinsic resistance. The terms covering antibiotics were variations of piperacillin/tazobactam, meticillin/methicillin, cephalosporin, carbapenem, aminoglycoside, gentamicin, amikacin, fluoroquinolone, linezolid, vancomycin, echinocandin, azole, colistin. The terms covering mechanisms of resistance or microbial properties associated with resistance were beta-lactamase/β-lactamase, extended-spectrum beta-lactamase/β-lactamase/betalactamase, carbapenemase, biofilm-producing, non-fermenting. The terms covering bacteria and fungi were *Pseudomonas aeruginosa,*

Acinetobacter spp., Acinetobacter baumannii, Stenotrophomonas maltophilia, Clostridium/Clostridioides difficile, Enterococcus faecium, Enterococcus faecalis, coagulase-negative staphylococci, Candida non-albicans, *Candida auris, Aspergillus fumigatus*. Those were selected from a wide range of terms during a multidisciplinary meeting.which led to a consensus. This search was run November 19th 2021 in the same databases (except for Cinahl) and with the same limitations as the June version.

Search strategies can be found in Supplementary material 1 and 2.

Study selection

The titles of all records were first screened by AD and LF to exclude any records that were not about antimicrobial resistance and cancer using the Rayyan tool (15)**.** We then screened all abstracts of the records where the subject was antimicrobial resistance and cancer to sort these into different study designs. The remaining original records with an observational study design were then read in full text by both AD and LF to see if they matched the eligibility criteria (Table 1). If there was uncertainty about inclusion of a record, both authors discussed until reaching a decision.

Table 1. Eligibility criteria.

Inclusion criteria	Exclusion criteria
- Primary study with original data - Observational study design (prospective/retrospective cohort study, case-control study, cross-sectional study) - The study population consists only of cancer patients, haematological orotherwise, of any age, sex, or ethnicity and in any institution. - The study includes a multivariable model. - A resistant bacterial or resistant fungal infection is either a risk factor or an outcome - The outcome is either infection (or colonisation) or death - Published after 2015. - Published in English. - Studies humans.	- Any other publication type than a primary observational study. - Cancer is a covariate or comparator, not everyone in the study population has cancer. - Resistance to antimicrobial drugs is not included in the model (neither as a risk factor, nor as an outcome). - The outcome is not infection or mortality, e.g. distribution of bacteria or bacterial strains. - Studies where the causal assumption is reversed, so that infection causes cancer (e.g. increased risk of gastric cancer after *Helicobacter pylori* infection)

Data extraction and analysis

The risk of bias in all included studies was assessed by AD and LF separately in a blinded process using the National Institutes of Health study quality assessment tools (16), and an arbitration meeting was held where OK/JB acted as an arbiter to come to a consensus about the final risk of bias assessment. The guidance for the use of these tools were followed, but the tool was modified to include an item with a question of whether the outcome was well defined in case-control studies.

Data extraction was then performed by AD and LF. We extracted the year of publication, the title of the study, the authors, the number of patients, the country of the study setting, the patient population statement, the risk factors included in the final model, the microbial aetiology, whether the studies employed a screening for statistical significance, and the

events per variable in the final model. This was included together with the risk of bias assessment and a short commentary in three different tables of all included studies, one for studies with an infection outcome, one for studies with a mortality outcome and one for studies with both outcomes. A table summarising the findings descriptively were created, presenting frequencies and medians. Analyses were performed using RStudio (17). The studies were checked against pre-specified criteria for a potential meta-analysis, being that models would have the same aetiology, outcome, and risk factors. These criteria were not met for any aetiology. In lieu of a quantitative analysis, a qualitative in-depth example of the investigation of risk factors was reported. We chose research that included vancomycin-resistant enterococci (VRE), as this research represented a typical cross-section of the studies selected. Furthermore, VRE is a typical hospital-associated microbe that may readily be prevented.

Results

Studies selected

After excluding 47048 duplicates, the two searches yielded a total of 27151 unique records from a total of three major databases - Ovid MEDLINE, Ovid Embase, and Web of Science - in addition to the Cinahl database (EBSCO) (Figure 1). Title screening for relevance to both antimicrobial resistance and cancer excluded 25341 records, where most excluded records were only about cancer, e.g. basic research on the drug resistance of the cancer disease. Abstract screening for a non-observational study design excluded a further 845 records, where 103 had a review design, 50 had an interventional design, 454 were case reports, and 238 were miscellaneous, mostly commentary articles and conference abstracts. Finally, after full text screening; 821 records were excluded, of which 165 included other diagnoses than cancer (including healthy individuals), 197 did not include an infection or mortality outcome (often microbial distribution or endpoints like length of stay), 130 did not explicitly include antimicrobial resistance in the model, 311 did not include a multivariable model, 16 were not in English, and we failed to gain access to two leaving 144 studies included.

Figure 1: PRISMA flowchart of the study selection (14).

Of the 144 studies, 55/144 (38%) had an infection outcome, 66/144 (46%) had a mortality outcome, and 23/144 (16%) had both outcomes, all of which are listed in detail and cited in tables with all extracted data in Supplementary Material S3, S4, and S5, respectively. In total, there were 23/144 (16%) studies of patients with solid cancers (18–40), 65/144 (45%) studies of patients with haematological cancers (41–105), and 56/144 (39%) studies with patients of both or unspecified cancer types (106–161). Most studies selected (39/144, 27%) reported and modelled several bacteria and/or fungi that were tested for resistances towards several antimicrobials (18–33,41–53,106–115). Eight of 144 (6%) reporting and modelling several microorganisms focused only on Gram-negative bacteria (54–58,116–118) and 1/144 (1%) focused only on fungi (59). Twelve of 144 (8%) studies studied the family *Enterobacterales* (especially *Escherichia coli* and *Klebsiella pneumoniae* combined), either ESBL- or carbapenemase-producing (34,60–64,119–124). The most commonly studied single organism was *Clostridioides difficile* with 27/144 (19%) studies (35–37,65–77,125–135). Also studied were three non-fermenters with 5/144 (3%) studies about *Acinetobacter baumannii* (78,136–139), 4/144 (3%) studies about *Pseudomonas aeruginosa* (79,80,140,141), and 5/144 (3%) studies about *Stenotrophomonas maltophilia*

(81–84,142), often specified as either multidrug- or extensively drug resistant. We also found several studies focusing on four well-known healthcare-associated bacteria, with 11/144 (8%) focusing on VRE (85–94,143,144), 7/144 (5%) focusing on carbapenemase-producing *K. pneumoniae* (KPC) (95–98,145–147), 2/144 (1%) focusing on methicillin-resistant *Staphylococcus aureus* (MRSA) (38,148), and 6/144 (4%) focusing on extended-spectrum beta-lactamase-producing *Escherichia coli* (ESBL-E) (99,100,149–152). We also found that fungi typically resistant to antifungals were studied, 3/144 (2%) studies about *Aspergillus* spp. (39,101,102) and 9/144 (6%) studies about *Candida* non-albicans (40,103,153–159), respectively. Finally, there was one of 144 (1%) studies for each of the microbes *Staphylococcus epidermidis* (with linezolid-resistance) (104), *Bacillus* spp. (160), *Streptococcus pneumoniae* (several resistance mechanisms) (161), and *Aeromonas sobria* (105).

Summary of findings and in-depth example

The most common microbial aetiologies were several bacteria/fungi (39, 25%), followed by *C. difficile* (27, 19%) and Enterobacteriaceae (12, 8%) (Table 2). The most common country for the study setting was the United States of America (USA) (40, 28%), followed by China (15, 10%), and Italy (11, 8%). The selected studies included a median of 200 patients (IQR 102-338) and 46 events (IQR 25-83.5). 103 (72%) studies used a p-value-based variable selection, either bivariable screening or stepwise regression. These studies screened a median of 16 variables (IQR 9-27), and included a median of 7 variables in the final (and largest) model. The median events per variable was close to 7.

Table 2. Summary of findings.

Characteristics	Frequencies
Outcome	**% (n)**
Infection	38 (55)
Death	45.8 (66)
Both	15.9 (23)
Country/setting	**% (n)**
USA	28 (40)
China	10 (15)
Italy	8 (11)
Spain	7 (10)
Germany	6 (9)
South Korea	6 (9)

Microbial aetiology	% (n)
Several bacterias and / or fungi	27 (39)
C. difficile	19 (27)
Enterobacteriaceae	8 (12)
Vancomycin-resistant enterococci	8 (11)
Carbapenemase-producing *K. pneumoniae*	5 (7)
Patients (median)	200 (IQR 102-338)
Events (median)	46 (IQR 25-84)
Events per variable (median)	6.9 (IQR 4.2-12.7)
Variables screened (median)	16 (IQR 9-27)
Variables in the final model (median)	7 (IQR 4-10)
Stepwise regression	**% (n)**
	72 (103)
Total	**144**

As a model organism, we explored how risk factors for and of VRE in cancer patients was described and modelled in the selected studies. *Enterococcus* is a genus of Gram-positive cocci, and often refer to the two commensal *E. faecium* and *E. faecalis*. These bacteria may acquire resistance towards the antibiotic vancomycin. Of our selection, 46/144 (32%) of the studies mention either *Enterococcus* spp., *E. faecium*, or *E. faecalis*, which is either described as vancomycin-resistant or "multidrug-resistant" (sometimes not further specified), or where the resistance mechanism is not described at all. Most of the studies (30/46, 65%) that mention the organism include it in a larger group, like Gram-positive bacteria or multidrug-resistant organisms, which hampers any specific focus on VRE (18–30,41–49,106–113).

Eight studies model the risk of being either colonised or infected with VRE. The number of patients included ranged from 72 to 342, and the variables screened ranged from 6 to 46. One study focused on patients undergoing liver transplantation for hilar cholangiocarcinoma, and seven focused on haematological cancer patients. What risk factors the authors

investigated for VRE infection or colonisation in a multivariable model, and what patient population they included, can be found in table 3. Although a plethora of risk factors were investigated, five of the studies selected variables based on their corresponding p-values in bivariable analyses, and as such dropped several variables from the analysis due to their failure to achieve the p-value criterion. Authors did not necessarily conclude that all risk factors investigated in the multivariable model were of importance. The risk of bias was rated low for three, medium for five, while none had a "fatal flaw". However, five of eight studies had less than ten events per variable in the final model, which may lead to a lack of statistical power. Among the common conclusions from the studies of VRE infection, four concluded that antibiotic exposure was a risk factor for VRE infection or colonisation, and two highlighted neutropenia. None of the antibiotic exposure risk factors was the same (one was vancomycin, one was carbapenem, one was general antibiotic exposure and one was daptomycin). When looking at the findings in detail, Aktürk et al. found that severe neutropenia and previous bacteraemia with another pathogen may increase the risk of progressing from VRE colonisation to VRE infection in paediatric haematological cancer patients (143). In 2015, Ford et al. found that severe neutropenia and number of stools per day was associated with VRE bloodstream infections in leukaemia patients, and in 2019 some of the same authors concluded that VRE colonisation rates fell when the hospital started using less carbapenems (85,86). Herc et al reported that only previous daptomycin exposure was associated with daptomycin-resistant VRE infections in patients with haematological malignancies (87). Hefazi et al found that VRE colonisation is associated with VRE infection in stem cell transplantation patients and Ramanan et al found that VRE colonisation pre-transplantation was associated with any infection post-transplantation (31,88). Heisel et al found that cephalosporin use and intravenous vancomycin were associated with VRE infections in patients with acute myeloid leukaemia or myelodysplastic syndrome undergoing intensive induction therapy, and finally Klein et al found that in multiple myeloma patients, granulocyte-colony stimulating factor was associated with fewer VRE cases than antibiotic prophylaxis (89,90).

Table 3. Risk factors for VRE infection or colonisation investigated in multivariable models.

Authors and publishing year	Patient population	Risk factors investigated
Akturk, Sutcu, Somer et al 2016	All patients admitted to the paediatric haematology/oncology ward with a documented Vancomycin-resistant enterococci colonisation 48-72 h after admission.	Severe neutropenia (<100/mm3), Previous bacteremia with another pathogen
Ford, Coombs, Stofer et al 2019	Patients admitted with newly diagnosed acute myelogenous leukaemia (AML), acute lymphoblastic leukaemia (ALL), biphenotypic leukaemia (BPL), or chronic myelogenous leukaemia in blast phase (CML-BP) between September 2011 and August 2017	Age, Days of empiric antibiotics, LOS, Cycling vs carbapenem period, Days of carbapenem, Days of cefepime or piperacillin/tazobactam
Ford, Lopansri, Haydoura et al 2015	Patients with newly diagnosed acute leukaemia between 2006 and 2012	Severe neutropenia, Number of stools/day
Hefazi, Damlaj, Alkhateeb et al 2016	All consecutive patients who underwent their first allogeneic HCT for AML at ta clinic between April 2004 and December 2014	Age (≥60 years vs <60), HCI-CI (≥3 vs 0- 2), KPS (>80 vs ≤80), Conditioning (RIC vs MAC), VRE

		colonisation (within 30 days of HCT)
Heisel, Sutton, Mascara et al 2017	Adult patients who received intensive induction chemotherapy with the standard 7+3 regimen of cytarabine and idarubicin for newly diagnosed AML or MDS from January 2012 to December 2015	Gender (male), Vancomycin IV, Cephalosporin
Herc, Kauffman, Marini et al 2017	Adult patients treated on an inpatient haematology service from January 2011 to December 2015 with a haematological malignancy and a blood culture positive for an enterococcus species.	Congestive heart failure, Typhlitis, Gastrointestinal bleeding, Daptomycin, Cefepime received within 90 days, Cumulative Daptomycin days within 90 days, Cumulative Fluoroquinolone days within 90 days, Prior enterococcal infection, Hospital LOS prior to positive culture, Stage of malignancy, Number induction chemotherapy regimens, Clofarabine within 30 days
Ramanan, Cummins, Wilhelm et al 2017	Adult patients who underwent liver transplantation for biopsy-proven hilar cholangiocarcinoma at a clinic between January 2004 and March 2013	VRE colonisation pre-transplant, Living donor transplant, Cold ischemia time, CMV donor seropositivity, Hepatic artery thrombosis post- transplant, Biliary stricture post- transplant, Intra-abdominal fluid collection post-transplant, Number of re-operations in 1st month post-transplant.
Klein, Sauer, Klein et al 2021	Multiple myeloma inpatients who received high-dose therapy with melphalan followed by autologous stem cell transplantation at a hospital between March 2016 and July 201	Antibiotic prophylaxis (vs. G-CSF support), No prophylaxis (vs. G-CSF support), Age (per ten years), ASCT at relapse (vs. first-line treatment), ≥VGPR before ASCT (vs. ≤PR), Stem cell amount ≥2.5 * (vs. <2.5)

Another eight studies analyse the deaths associated with VRE in cancer patients. The number of patients included ranged from 95 to 1424, and the variables screened ranged from 11 to 56, although the exact number was indeterminable for one of the studies. Six studies selected variables based on a specified p-value threshold, but for one study the method for variable selection could not be determined. A full list of the different patient populations and risk factors investigated can be found in table 4. We assessed the risk of bias among these studies and found four studies at low risk and four studies at a medium risk of bias. Of the eight studies, six had less than 10 events per variable in the final model. The studies had few conclusions about risk factors for mortality among cancer patients associated with VRE in common. However, three studies found that VRE bacteremia was a risk factor for death and two studies found no risk factors after running their model. When looking at the findings in detail, Akhtar et al. found that only shock (not further specified) was associated with difference in mortality between VRE and VSE bacteraemia in cancer patients (144). Kamboj et al did not find any factors that were associated with higher mortality in stem cell transplantation patients with a VRE bacteraemia (91). Kern et al modelled the mortality associated with an enterococcal bacteraemia in haematological cancer patients, but did not specify vancomycin-resistance (50). Kirkizlar et al found that in

leukaemia patients colonised with VRE, a low neutrophil count and coinfection were associated with increased mortality (92). Mendes et al included VRE in a bivariable screening, but discarded the factor as it did not meet the criterion of p<0.1 (51). Ornstein et al found that leukaemia patients with a VRE bacteraemia at the induction of chemotherapy had poorer survival than patients with other bloodstream infections (93). Papanicolaou et al found that VRE bacteraemia increased the mortality in patients receiving their first stem cell transplantation, but did not disclose how variables were selected for the multivariable model (94). Finally, Pugliese et al modelled the risk of mortality associated with several bacteria in leukaemia inpatients, among them *Enterococcus* spp. (no vancomycin-resistance mentioned), but did not find an association (52).

Table 4. Risk factors for death associated with VRE investigated in multivariable models.

Authors and publishing year	Patient population	Risk factors investigated
Akhtar, Sultan, Nizamuddin et al 2016	Cancer patients whose blood culture grew either vancomycin-sensitive enterococcus (VSE) or VRE from January 2012 to December 2014	Paediatric (Compared to adults), Male (Compared to females), Haematological malignancies (Compared to solid ones), Location in hospital at onset of bacteraemia (Compared to patients in ICU), Inpatient, Others(OPD/ ER), APACHE-II score at onset of bacteraemia, Length of hospital stay in days before onset of bacteraemia, Shock (Compared to patients not in shock at onset of bacteraemia), DIC score, Received vancomycin within 4 weeks prior to bacteraemia
Kamboj, Cohen, Huang et al 2019	Adults 18 years and older who underwent allo-HSCT and developed VREB between January 1, 2005 and December 31, 2014, at this NY cancer centre	Prior VRE bacteremia, Hypotension (<90/50), Persistent VREB >48 hours
Kern, Roth, Bertz et al 2019	Patients with BSI during neutropenia subsequent to high‑dose chemotherapy for acute leukaemia (HDC) or autologous or allogeneic hematopoietic stem cell transplantation (auto‑HSCT or allo‑HSCT) from the beginning of 2002 through April 2015	Age, Sex, Antineoplastic treatment, Admitted with neutropenia, Days of neutropenia before BSI onset, BSI pathogen
Kirkizlar, Akalin, Kirkizlar et al 2020	Adult acute leukaemia patients with VRE colonised who had febrile neutropenia	Age, Female, < 0,5x109/L neutrophil count while VRE (+), Coinfection
Mendes, da Silva, da Costa Bandeira de Melo et al 2021	Patients aged >=16 years with newly diagnosed AML who started any regimen of intensive treatment	Age >60 years, GNB colonisation, CRP levels >15 mg/dL, monocytic AML, AGR
Ornstein, Mukherjee, Keng et al 2015	Patients with de novo and secondary AML who received cytarabine-based induction chemotherapy at Cleveland Clinic between 1 January 2000 and 1 April 2008	Age (>60 vs. <60), Cytology (Poor vs. Favourable, Poor vs. Intermediate, Poor vs. Unknown), VRE bacteraemia, Other bloodstream infections, Aetiology (secondary vs. de novo), white

		blood cell at diagnosis, Year of diagnosis
Papanicolaou, Ustun, Young et al 2019	Patients who received their first allogeneic HCT for acute myelogenous leukaemia (AML), acute lymphoblastic leukaemia (ALL), or myelodysplastic syndrome (MDS) between January 2008 and December 2012	VRE BSI, Age (21-40, 41-50, 51-60, >60), AML/ALL intermediate, AML/ALL/MDS advanced, MDS advanced, Cord blood allograft, Human leukocyte antigen 7/8, Karnofsky performance scale <90, Cytomegalovirus donor or receptor positive, HCT during 2010-2012
Pugliese, Salvatore, Iula et al 2017	Adult patients with newly diagnosed AML hospitalised at a haematological department from 1 January 2002 to 31 December 2012 in order to receive cytotoxic agent induction courses for haematological remission	High-dose cytarabine-containing chemotherapy, Ultrasonography-driven NEC therapy with antibiotic regimens including tigecycline.

Discussion

In our systematic review of studies from the previous six years with multivariable models of risk factors for antimicrobial resistant infections in cancer patients, we selected 144 studies that were eligible for inclusion. Most studies focused on haematological cancer patients and explored a host of different microbes. Studies often had small sample sizes, screened a large number of variables, and used p-value-based methods for variable selection like bivariable screening or stepwise regression. In general, the models were highly heterogeneous, with nuances in the study populations and microbial aetiologies, and major differences in which risk factors were modelled, as we have exemplified through the research on VRE in cancer patients. The issues with heterogeneity when performing a systematic review of basic prognostic factor research have been described before (162). Although it is not possible to summarise factors that appear to be of importance, we find that known general risk factors for infections like immunosuppression and specific risk factors for resistant infections like previous antibiotic use are recurring, as other non-systematic reviews do (1,2,8,9). A comprehensive list of these factors can be found in the supplementary material S3 and S4. To the best of our knowledge, no previous systematic review has investigated the risk factors for all resistant infections in cancer patients. However, three systematic reviews have looked at risk factors for methicillin-resistant *S. aureus*, extended-spectrum β-lactamase-producing *Enterobacteriaceae*, and vancomycin-resistant enterococci, specifically (163–165). These studies were able to pool the prevalence of such infections and identify some risk factors of importance, providing a good basis for future research. Some systematic reviews also investigate the changing epidemiology of antimicrobial resistant infections in cancer patients without a particular focus on risk factors (8,166).

Some conservative choices were made in the selection process which may have reduced the final number of studies included. Some studies were excluded because study participants had other diagnoses than cancer, e.g. recipients of hematopoietic stem cell transplantation due to non-malignant haematological disorders. Several of these studies did

not describe the full diagnostic panorama, although comprehensive patient characteristics remain an important backbone in epidemiological research. Other studies did not explicitly describe or model resistant microbes, either by including resistance or resistant microbes as a variable or an outcome in the model. Given the rapid increase in antimicrobial resistance, current research in the epidemiology of infectious diseases in healthcare-settings should include detailed data on this topic. The most common reason for excluding observational studies during full-text reading was the lack of a multivariable model. These studies seemed to have a low number of patients, and a multivariable model may have been avoided due to low statistical power. As discussed by the PROGRESS Research Group in their recommendations for prognostic factor research, the discovery and investigation into new prognostic factors should rely on multivariable modelling to discern these from factors already known to be of importance, and to provide a basic adjustment of confounders (10). Studies simply testing whether a number of factors differ between groups often rely heavily on null-hypothesis statistical significance testing and are subject to the multiple comparisons problem (167). Although exploratory studies are important, such studies sometimes either test risk factors that are already known to be associated with the outcome, test factors in too small samples, or they test too many factors at once. This may lead to wrong inferences due to known issues such as sparse data bias or winner's curse inflating effect sizes (168,169). An alternative can be to establish larger research collaborations that can pool data into larger cohorts, like we found several examples of (59,140). Worth noting, we found two clinical prediction models. In one of them, the IRONIC group developed a scoring system for the risk of multidrug resistance in bloodstream infections by *P. aeruginosa*, and in another, Colombian researchers developed a scoring system for the risk of ESBL-producing *Enterobactericaea* (124,140).

Studies were assessed for their risk of bias using the NIH Quality Assessment tool, in which we found several recurring issues. Most studies did not include a sample size calculation or described a blinding of exposure assessors. Differences in the risk of bias assessments were often determined by allowing continuous variables to be treated continuously or the correct definition of the exposure. We also found that several studies lacked a reliable definition of the outcome, in particular the definition of mortality. As discussed by the NIH Quality Assessment reviewers, even though death as an outcome seems to be objective by nature (i.e. researchers rely on "face validity"), even this outcome should be defined. Such a definition should include information on particulars like from which register or medical record information about the death was collected, if the patient perished within or outside of the institution, and if the latter was the case, who reported the death. Most studies did have a sufficient follow-up time to capture infectious disease outcomes (in particular mortality), but these follow-up times varied greatly for all studies. However, the NIH Quality Assessment tool included items that were not relevant to all studies (e.g. the adjustment of confounders in a prediction context), and the issue with finding good quality assessment tools for prognostic factor research has been described earlier (170). Systematic reviews of clinical prediction models should use other more specialised tools like PROBAST (171).

We chose to summarise the research by describing an in-depth example of studies on VRE, which provided us with a representative cross-section in terms of both patient population, sample size, methodological approaches, and number of variables. This example showed how heterogeneity in how the subject is studied may hamper our ability to build on this basic research, either through the development of more comprehensive predictive models, or

through pooling or meta-analysis. Most studies studying resistant enterococci group the bacteria together with other bacteria and/or fungi. Furthermore, authors often study the bacteria in a highly selected patient population, in which it is difficult to infer how baseline characteristics pertain to the risk factors studied, either from a table of characteristics or the model. However, the most major source of heterogeneity that reduces our ability to build on the research is the arbitrary way in which variables are selected.

We find that throughout the entire material, there was a widespread use of either stepwise regression or a bivariable screening of variables as a method for variable selection in regression models, where variables that achieve some pre-specified p-value threshold are included in the final model. This method is, however, not recommended for either estimating the effect of an exposure or predicting an outcome (12,172–174). In short, the reason is its reliance on p-values as a criterion for variable selection, although the p-value is not an indicator of a causal relationship or of predictive power. A factor confounding a causal effect may not be statistically significant, and there may be situations in which a variable that is not statistically significant in a bivariable analysis may increase the predictive power of a multivariable model. Consequently, including all candidate factors in a multivariable model and testing them by means of statistical significance may not be a valid method of discovering new prognostic factors. Unfortunately, there is no alternative to these methods based on statistical significance that is as simple and practical, or automated and data-driven. The basis for all modelling is a theoretical understanding of covariates, outcomes and the relationship between them. All in all, it is difficult and maybe even impossible to summarise what risk factors are shown to be of relevance in this literature. Simply counting how many times a certain risk factor is found to be statistically significant tells us little of its relative importance.

The strengths of our study are the large scope of our searches, which has resulted in a comprehensive overview of the current state of this research topic. However, there are several limitations to our study. First, we only searched for studies where any type of antimicrobial resistance was mentioned in the title, abstract, controlled vocabulary, or keywords, which may have excluded some studies that only mention infections in broader terms, but still model the risk for contracting resistant infections or any potential outcomes of such infections. Furthermore, we narrowed our inclusion criteria to studies that model either an infectious or mortality outcome, but this does not fully cover the ways in which resistant microbes may increase the disease burden among cancer patients, like repeated hospitalisations and/or increased length of stays. Several of the studies that were excluded mainly had patients with a haematological malignancy, but also some patients with aplastic anaemia or other haematological disorders. Other studies did not explicitly include (acquired or intrinsic) resistance in the models, although they may have included resistance in other implicit ways, like the failure of an empiric carbapenem cure. Patients with other haematological disorders or other infectious etiologies may have similar risk profiles, and their exclusion may have unreasonably narrowed the scope of the review. We also searched for records in Swedish, Danish and Spanish, but records in these languages were not included to ensure this systematic review would be available and reproducible for all readers. As many of the studies we included were from countries where English is not the native language, the exclusion of other languages may represent a limitation of our review.

Conclusions

In this systematic review, we found a great level of heterogeneity in the approach to studying risk factors for resistant infections and mortality due to resistant infections among cancer patients. We argue that it's difficult or even impossible to use these models to summarise which risk factors are of particular importance. This is due to differences in the patient populations selected, and the different ways of grouping microbes. Furthermore, studies on this subject often have a small sample size and use p-value-based variable selection methods, which lead to very diverse models. The consequence of this heterogeneity is not only that the literature is unprepared for either a meta-analysis or a pooled analysis. It also means that it is difficult to use this research to understand the mechanisms by which resistant microbes cause disease in cancer patients, and thus that it is difficult for clinicians to use the research to guide their prevention of such conditions. This represents a serious shortcoming of this body of literature. There is a need to develop and adhere to more standardised protocols when investigating new risk factors. Such protocols should include a clear aim of what risk factors are to be explored and build on existing literature, e.g. by selecting similar patient populations and being careful to include factors previously shown to be of importance regardless of their p-value in a bivariable screening.

Declarations

Acknowledgements
We would like to thank bachelor's student Sofie Almark Jeppesen for assistance in the screening process. We would also like to thank the authors who shared their research with us where we did not otherwise have access.

Ethics approval and consent to participate
Not applicable

Consent for publication
Not applicable

Availability of data and materials
All data extracted or generated and then analysed are included in the article or the supplementary material.

Competing interests
The authors declare that they have no competing interests.

Funding
The study was internally funded. AD is funded by a grant from the South-Eastern Norway Regional Health Authority.

Authors' contributions
AD conceived of the idea, drafted the manuscript, screened the search, extracted and organised the data, and performed the risk of bias assessment. LF revised the manuscript, screened the search, extracted and organised the data, analysed the data, and performed

the risk of bias assessment. SP revised the manuscript. AM revised the manuscript. RT revised the manuscript, and performed the search. OK revised the manuscript, and performed the risk of bias assessment. JB conceived the idea, and revised the manuscript.

References

(1)   Zembower TR. Epidemiology of Infections in Cancer Patients. In: Stosor V, Zembower TR, editors. Infect. Complicat. Cancer Patients, Cham: Springer International Publishing; 2014. https://doi.org/10.1007/978-3-319-04220-6_2.

(2)   Nanayakkara AK, Boucher HW, Fowler Jr VG, Jezek A, Outterson K, Greenberg DE. Antibiotic resistance in the patient with cancer: Escalating challenges and paths forward. CA Cancer J Clin 2021;71. https://doi.org/10.3322/caac.21697.

(3)   Klastersky J, Paesmans M, Rubenstein EB, Boyer M, Elting L, Feld R, et al. The Multinational Association for Supportive Care in Cancer risk index: A multinational scoring system for identifying low-risk febrile neutropenic cancer patients. J Clin Oncol Off J Am Soc Clin Oncol 2000;18. https://doi.org/10.1200/JCO.2000.18.16.3038.

(4)   Korvick JA, Bryan CS, Farber B, Beam TR, Schenfeld L, Muder RR, et al. Prospective observational study of Klebsiella bacteremia in 230 patients: outcome for antibiotic combinations versus monotherapy. Antimicrob Agents Chemother 1992;36. https://doi.org/10.1128/AAC.36.12.2639.

(5)   Jarlier V, Högberg LD, Heuer OE, Campos J, Eckmanns T, Giske CG, et al. Strong correlation between the rates of intrinsically antibiotic-resistant species and the rates of acquired resistance in Gram-negative species causing bacteraemia, EU/EEA, 2016. Eurosurveillance 2019;24. https://doi.org/10.2807/1560-7917.ES.2019.24.33.1800538.

(6)   World Health Organization. Antimicrobial resistance: global report on surveillance. World Health Organization; 2014.

(7)   Cassini A, Högberg LD, Plachouras D, Quattrocchi A, Hoxha A, Simonsen GS, et al. Attributable deaths and disability-adjusted life-years caused by infections with antibiotic-resistant bacteria in the EU and the European Economic Area in 2015: a population-level modelling analysis. Lancet Infect Dis 2019;19. https://doi.org/10.1016/S1473-3099(18)30605-4.

(8)   Trecarichi EM, Tumbarello M. Antimicrobial-resistant Gram-negative bacteria in febrile neutropenic patients with cancer: current epidemiology and clinical impact. Curr Opin Infect Dis 2014;27. https://doi.org/10.1097/QCO.0000000000000038.

(9)   Rolston KVI. Infections in Cancer Patients with Solid Tumors: A Review. Infect Dis Ther 2017;6. https://doi.org/10.1007/s40121-017-0146-1.

(10)   Riley RD, Hayden JA, Steyerberg EW, Moons KGM, Abrams K, Kyzas PA, et al. Prognosis Research Strategy (PROGRESS) 2: Prognostic Factor Research. PLOS Med 2013;10. https://doi.org/10.1371/journal.pmed.1001380.

(11)   Steyerberg EW, Moons KGM, Windt DA van der, Hayden JA, Perel P, Schroter S, et al. Prognosis Research Strategy (PROGRESS) 3: Prognostic Model Research. PLOS Med 2013;10. https://doi.org/10.1371/journal.pmed.1001381.

(12)   Steyerberg EW. Clinical Prediction Models : A Practical Approach to Development, Validation, and Updating. 2nd ed. 2019. Cham: Springer International Publishing : Imprint: Springer; 2019.

(13)   Danielsen AS, Franconeri L, Kacelnik O, Tornes RA, Bjørnholt JV. Antimicrobial resistance and cancer patients: A systematic review of risk factor modelling. PROSPERO 2021. https://www.crd.york.ac.uk/prospero/display_record.php?RecordID=282769 (accessed April 11, 2022).

(14)   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021;372. https://doi.org/10.1136/bmj.n71.

(15)   Rayyan – Intelligent Systematic Review n.d. https://www.rayyan.ai/ (accessed April 11, 2022).

(16)   National Heart, Lung and Blood Institute. Study Quality Assessment Tools n.d. https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools (accessed April 11, 2022).

(17)   RStudio Team. RStudio: Integrated Development Environment for R. Boston, MA: RStudio, PBC.; 2020.

(18)   Bednarsch J, Czigany Z, Heij LR, Luedde T, van Dam R, Lang SA, et al. Bacterial bile duct colonization in perihilar cholangiocarcinoma and its clinical significance. Sci Rep 2021;11. https://doi.org/10.1038/s41598-021-82378-y.

(19)   Gianotti L, Tamini N, Gavazzi F, Mariani A, Sandini M, Ferla F, et al. Consequences of Increases in Antibiotic Resistance Pattern on Outcome of Pancreatic Resection for Cancer. J Gastrointest Surg Off J Soc Surg Aliment Tract 2017;21. https://doi.org/10.1007/s11605-017-3483-1.

(20)   Lee D-S, Ryu J-A, Chung CR, Yang J, Jeon K, Suh GY, et al. Risk factors for acquisition of multidrug-resistant bacteria in patients with anastomotic leakage after colorectal cancer surgery. Int J Colorectal Dis 2015;30. https://doi.org/10.1007/s00384-015-2161-6.

(21)   Szvalb AD, El Haddad H, Rolston KV, Sabir SH, Jiang Y, Raad II, et al. Risk factors for recurrent percutaneous nephrostomy catheter-related infections. Infection 2019;47. https://doi.org/10.1007/s15010-018-1245-y.

(22)   Zhou T, Yang W, Yang Q, Xuan B, Zhang L, Li X, et al. Distribution, diagnosis, and analysis of related risk factors of multidrug-resistant organism in patients with malignant neoplasms. Int J Clin Exp Pathol 2020;13.

(23)   Antonio M, Gudiol C, Royo-Cebrecos C, Grillo S, Ardanuy C, Carratalà J. Current etiology, clinical features and outcomes of bacteremia in older patients with solid tumors. J Geriatr Oncol 2019;10. https://doi.org/10.1016/j.jgo.2018.06.011.

(24)   Himmelsbach V, Knabe M, Ferstl PG, Peiffer K-H, Stratmann JA, Wichelhaus TA, et al. Colonization with multidrug-resistant organisms impairs survival in patients with hepatocellular carcinoma. J Cancer Res Clin Oncol 2022;148. https://doi.org/10.1007/s00432-021-03741-0.

(25)   Jiang A-M, Liu N, Ali Said R, Ren M-D, Gao H, Zheng X-Q, et al. Nosocomial Infections in Gastrointestinal Cancer Patients: Bacterial Profile, Antibiotic Resistance Pattern, and Prognostic Factors. Cancer Manag Res 2020;12. https://doi.org/10.2147/CMAR.S258774.

(26)   Marín M, Gudiol C, Castet F, Oliva M, Peiró I, Royo-Cebrecos C, et al. Bloodstream infection in patients with head and neck cancer: a major challenge in the cetuximab era. Clin Transl Oncol Off Publ Fed Span Oncol Soc Natl Cancer Inst Mex 2019;21. https://doi.org/10.1007/s12094-018-1905-5.

(27)   Matsutani N, Yoshiya K, Chida M, Sakaguchi H, Kikkawa T, Fukuda H, et al. Postoperative empyema following lung cancer surgery. Oncotarget 2018;9. https://doi.org/10.18632/oncotarget.25629.

(28)   Mei T, Yang X, Yu Y, Tian X, Deng Q, Xu Y, et al. Secondary Infections After Diagnosis of Severe Radiation Pneumonitis (SRP) Among Patients With Non-Small Cell Lung Cancer: Pathogen Distributions, Choice of Empirical Antibiotics, and the Value of Empirical Antifungal Treatment. Int J Radiat Oncol Biol Phys 2022;112. https://doi.org/10.1016/j.ijrobp.2021.08.022.

(29)   Royo-Cebrecos C, Gudiol C, García J, Tubau F, Laporte J, Ardanuy C, et al. Characteristics, aetiology, antimicrobial resistance and outcomes of bacteraemic cholangitis in patients with solid tumours: A prospective cohort study. J Infect 2017;74. https://doi.org/10.1016/j.jinf.2016.10.008.

(30)   Stratmann JA, Lacko R, Ballo O, Shaid S, Gleiber W, Vehreschild MJGT, et al. Colonization with multi-drug-resistant organisms negatively impacts survival in patients with non-small cell lung cancer. PloS One 2020;15. https://doi.org/10.1371/journal.pone.0242544.

(31) Ramanan P, Cummins NW, Wilhelm MP, Heimbach JK, Dierkhising R, Kremers WK, et al. Epidemiology, risk factors, and outcomes of infections in patients undergoing liver transplantation for hilar cholangiocarcinoma. Clin Transplant 2017;31. https://doi.org/10.1111/ctr.13023.

(32) Sugimachi K, Iguchi T, Mano Y, Morita M, Mori M, Toh Y. Significance of bile culture surveillance for postoperative management of pancreatoduodenectomy. World J Surg Oncol 2019;17. https://doi.org/10.1186/s12957-019-1773-7.

(33) Perdikouri EIA, Arvaniti K, Lathyris D, Apostolidou Kiouti F, Siskou E, Haidich AB, et al. Infections Due to Multidrug-Resistant Bacteria in Oncological Patients: Insights from a Five-Year Epidemiological and Clinical Analysis. Microorganisms 2019;7. https://doi.org/10.3390/microorganisms7090277.

(34) Golzarri MF, Silva-Sánchez J, Cornejo-Juárez P, Barrios-Camacho H, Chora-Hernández LD, Velázquez-Acosta C, et al. Colonization by fecal extended-spectrum β-lactamase-producing Enterobacteriaceae and surgical site infections in patients with cancer undergoing gastrointestinal and gynecologic surgery. Am J Infect Control 2019;47. https://doi.org/10.1016/j.ajic.2019.01.020.

(35) Liu NW, Shatagopam K, Monn MF, Kaimakliotis HZ, Cary C, Boris RS, et al. Risk for Clostridium difficile infection after radical cystectomy for bladder cancer: Analysis of a contemporary series. Urol Oncol 2015;33. https://doi.org/10.1016/j.urolonc.2015.07.007.

(36) Rodríguez Garzotto A, Mérida García A, Muñoz Unceta N, Galera Lopez MM, Orellana-Miguel MÁ, Díaz-García CV, et al. Risk factors associated with Clostridium difficile infection in adult oncology patients. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2015;23. https://doi.org/10.1007/s00520-014-2506-7.

(37) Zheng Y, Luo Y, Lv Y, Huang C, Sheng Q, Zhao P, et al. Clostridium difficile colonization in preoperative colorectal cancer patients. Oncotarget 2017;8. https://doi.org/10.18632/oncotarget.14424.

(38) Lin S, Melki S, Lisgaris MV, Ahadizadeh EN, Zender CA. Post-operative MRSA infections in head and neck surgery. Am J Otolaryngol 2017;38. https://doi.org/10.1016/j.amjoto.2017.03.013.

(39) Avcı N, Hartavi M, Kaçan T, Bayındır M, Avcı M, Özakın C, et al. Retrospective analysis of the microbiological spectrum of pneumonia in Turkish patients with lung cancer. Contemp Oncol Poznan Pol 2016;20. https://doi.org/10.5114/wo.2016.58502.

(40) Tarapan S, Matangkasombut O, Trachootham D, Sattabanasuk V, Talungchit S, Paemuang W, et al. Oral Candida colonization in xerostomic postradiotherapy head and neck cancer patients. Oral Dis 2019;25. https://doi.org/10.1111/odi.13151.

(41) Ballo O, Tarazzit I, Stratmann J, Reinheimer C, Hogardt M, Wichelhaus TA, et al. Colonization with multidrug resistant organisms determines the clinical course of patients with acute myeloid leukemia undergoing intensive induction chemotherapy. PloS One 2019;14. https://doi.org/10.1371/journal.pone.0210991.

(42) Cattaneo C, Zappasodi P, Mancini V, Annaloro C, Pavesi F, Skert C, et al. Emerging resistant bacteria strains in bloodstream infections of acute leukaemia patients: results of a prospective study by the Rete Ematologica Lombarda (Rel). Ann Hematol 2016;95. https://doi.org/10.1007/s00277-016-2815-7.

(43) Chen S, Lin K, Li Q, Luo X, Xiao M, Chen M, et al. A practical update on the epidemiology and risk factors for the emergence and mortality of bloodstream infections from real-world data of 3014 hematological malignancy patients receiving chemotherapy. J Cancer 2021;12. https://doi.org/10.7150/jca.50802.

(44) Di Domenico EG, Marchesi F, Cavallo I, Toma L, Sivori F, Papa E, et al. The Impact of Bacterial Biofilms on End-Organ Disease and Mortality in Patients with Hematologic Malignancies Developing a Bloodstream Infection. Microbiol Spectr 2021;9. https://doi.org/10.1128/Spectrum.00550-21.

(45) Facchin G, Candoni A, Lazzarotto D, Zannier ME, Peghin M, Sozio E, et al. Clinical characteristics and outcome of 125 polymicrobial bloodstream infections in hematological patients: an 11-year epidemiologic survey. Support Care Cancer Off J

Multinatl Assoc Support Care Cancer 2022;30.
https://doi.org/10.1007/s00520-021-06640-9.

(46) Garcia-Vidal C, Cardozo-Espinola C, Puerta-Alcalde P, Marco F, Tellez A, Agüero D, et al. Risk factors for mortality in patients with acute leukemia and bloodstream infections in the era of multiresistance. PloS One 2018;13. https://doi.org/10.1371/journal.pone.0199531.

(47) Scheich S, Reinheimer C, Brandt C, Wichelhaus TA, Hogardt M, Kempf VAJ, et al. Clinical Impact of Colonization with Multidrug-Resistant Organisms on Outcome after Autologous Stem Cell Transplantation: A Retrospective Single-Center Study. Biol Blood Marrow Transplant J Am Soc Blood Marrow Transplant 2017;23. https://doi.org/10.1016/j.bbmt.2017.05.016.

(48) Seong GM, Lee Y, Hong S-B, Lim C-M, Koh Y, Huh JW. Prognosis of Acute Respiratory Distress Syndrome in Patients With Hematological Malignancies. J Intensive Care Med 2020;35. https://doi.org/10.1177/0885066617753566.

(49) Waszczuk-Gajda A, Drozd-Sokołowska J, Basak GW, Piekarska A, Mensah-Glanowska P, Sadowska-Klasa A, et al. Infectious Complications in Patients With Multiple Myeloma After High-Dose Chemotherapy Followed by Autologous Stem Cell Transplant: Nationwide Study of the Infectious Complications Study Group of the Polish Adult Leukemia Group. Transplant Proc 2020;52. https://doi.org/10.1016/j.transproceed.2020.02.068.

(50) Kern WV, Roth JA, Bertz H, Götting T, Dettenkofer M, Widmer AF, et al. Contribution of specific pathogens to bloodstream infection mortality in neutropenic patients with hematologic malignancies: Results from a multicentric surveillance cohort study. Transpl Infect Dis Off J Transplant Soc 2019;21. https://doi.org/10.1111/tid.13186.

(51) Mendes FR, da Silva WF, da Costa Bandeira de Melo R, Silveira DRA, Velloso EDRP, Rocha V, et al. Predictive factors associated with induction-related death in acute myeloid leukemia in a resource-constrained setting. Ann Hematol 2022;101. https://doi.org/10.1007/s00277-021-04687-6.

(52) Pugliese N, Salvatore P, Iula DV, Catania MR, Chiurazzi F, Della Pepa R, et al. Ultrasonography-driven combination antibiotic therapy with tigecycline significantly increases survival among patients with neutropenic enterocolitis following cytarabine-containing chemotherapy for the remission induction of acute myeloid leukemia. Cancer Med 2017;6. https://doi.org/10.1002/cam4.1063.

(53) Calik S, Ari A, Bilgir O, Cetintepe T, Yis R, Sonmez U, et al. The relationship between mortality and microbiological parameters in febrile neutropenic patients with hematological malignancies. Saudi Med J 2018;39. https://doi.org/10.15537/smj.2018.9.22824.

(54) Andria N, Henig O, Kotler O, Domchenko A, Oren I, Zuckerman T, et al. Mortality burden related to infection with carbapenem-resistant Gram-negative bacteria among haematological cancer patients: a retrospective cohort study. J Antimicrob Chemother 2015;70. https://doi.org/10.1093/jac/dkv218.

(55) Tang Y, Wu X, Cheng Q, Li X. Inappropriate initial antimicrobial therapy for hematological malignancies patients with Gram-negative bloodstream infections. Infection 2020;48. https://doi.org/10.1007/s15010-019-01370-x.

(56) Tang Y, Xu C, Xiao H, Wang L, Cheng Q, Li X. Gram-Negative Bacteria Bloodstream Infections in Patients with Hematological Malignancies - The Impact of Pathogen Type and Patterns of Antibiotic Resistance: A Retrospective Cohort Study. Infect Drug Resist 2021;14. https://doi.org/10.2147/IDR.S322812.

(57) Shin DH, Shin D-Y, Kang CK, Park S, Park J, Jun KI, et al. Risk factors for and clinical outcomes of carbapenem non-susceptible gram negative bacilli bacteremia in patients with acute myelogenous leukemia. BMC Infect Dis 2020;20. https://doi.org/10.1186/s12879-020-05131-2.

(58) Scheich S, Weber S, Reinheimer C, Wichelhaus TA, Hogardt M, Kempf VAJ, et al. Bloodstream infections with gram-negative organisms and the impact of multidrug resistance in patients with hematological malignancies. Ann Hematol 2018;97.

https://doi.org/10.1007/s00277-018-3423-5.

(59) Criscuolo M, Marchesi F, Candoni A, Cattaneo C, Nosari A, Veggia B, et al. Fungaemia in haematological malignancies: SEIFEM-2015 survey. Eur J Clin Invest 2019;49. https://doi.org/10.1111/eci.13083.

(60) Satlin MJ, Cohen N, Ma KC, Gedrimaite Z, Soave R, Askin G, et al. Bacteremia due to carbapenem-resistant Enterobacteriaceae in neutropenic patients with hematologic malignancies. J Infect 2016;73. https://doi.org/10.1016/j.jinf.2016.07.002.

(61) Woerther P-L, Micol J-B, Angebault C, Pasquier F, Pilorge S, Bourhis J-H, et al. Monitoring antibiotic-resistant enterobacteria faecal levels is helpful in predicting antibiotic susceptibility of bacteraemia isolates in patients with haematological malignancies. J Med Microbiol 2015;64. https://doi.org/10.1099/jmm.0.000078.

(62) Liang T, Xu C, Cheng Q, Tang Y, Zeng H, Li X. Epidemiology, Risk Factors, and Clinical Outcomes of Bloodstream Infection due to Extended-Spectrum Beta-Lactamase-Producing Escherichia coli and Klebsiella pneumoniae in Hematologic Malignancy: A Retrospective Study from Central South China. Microb Drug Resist Larchmt N 2021;27. https://doi.org/10.1089/mdr.2020.0033.

(63) Alrstom A, Alsuliman T, Daher N, Abouharb R. The Impact of Modifying Empirical Antibiotic Therapy Based on Intestinal Colonization Status on Clinical Outcomes of Febrile Neutropenic Patients. Infect Chemother 2021;53. https://doi.org/10.3947/ic.2020.0111.

(64) Jaiswal SR, Gupta S, Kumar RS, Sherawat A, Rajoreya A, Dash SK, et al. Gut Colonization with Carbapenem-resistant Enterobacteriaceae Adversely Impacts the Outcome in Patients with Hematological Malignancies: Results of A Prospective Surveillance Study. Mediterr J Hematol Infect Dis 2018;10. https://doi.org/10.4084/MJHID.2018.025.

(65) Alonso CD, Braun DA, Patel I, Akbari M, Oh DJ, Jun T, et al. A multicenter, retrospective, case-cohort study of the epidemiology and risk factors for Clostridium difficile infection among cord blood transplant recipients. Transpl Infect Dis Off J Transplant Soc 2017;19. https://doi.org/10.1111/tid.12728.

(66) Ballo O, Kreisel E-M, Eladly F, Brunnberg U, Stratmann J, Hunyady P, et al. Use of carbapenems and glycopeptides increases risk for Clostridioides difficile infections in acute myeloid leukemia patients undergoing intensive induction chemotherapy. Ann Hematol 2020;99. https://doi.org/10.1007/s00277-020-04274-1.

(67) Ford CD, Lopansri BK, Webb BJ, Coombs J, Gouw L, Asch J, et al. Clostridioides difficile colonization and infection in patients with newly diagnosed acute leukemia: Incidence, risk factors, and patient outcomes. Am J Infect Control 2019;47. https://doi.org/10.1016/j.ajic.2018.09.027.

(68) Lavallée C, Labbé A-C, Talbot J-D, Alonso CD, Marr KA, Cohen S, et al. Risk factors for the development of Clostridium difficile infection in adult allogeneic hematopoietic stem cell transplant recipients: A single-center study in Québec, Canada. Transpl Infect Dis Off J Transplant Soc 2017;19. https://doi.org/10.1111/tid.12648.

(69) Morrisette T, Van Matre AG, Miller MA, Mueller SW, Bajrovic V, Abidi MZ, et al. Oral Vancomycin Prophylaxis as Secondary Prevention Against Clostridioides difficile Infection in the Hematopoietic Stem Cell Transplantation and Hematologic Malignancy Population. Biol Blood Marrow Transplant J Am Soc Blood Marrow Transplant 2019;25. https://doi.org/10.1016/j.bbmt.2019.06.021.

(70) Petteys MM, Kachur E, Pillinger KE, He J, Copelan EA, Shahid Z. Antimicrobial de-escalation in adult hematopoietic cell transplantation recipients with febrile neutropenia of unknown origin. J Oncol Pharm Pract Off Publ Int Soc Oncol Pharm Pract 2020;26. https://doi.org/10.1177/1078155219865303.

(71) Przybylski DJ, Reeves DJ. Moxifloxacin versus levofloxacin or ciprofloxacin prophylaxis in acute myeloid leukemia patients receiving chemotherapy. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2017;25. https://doi.org/10.1007/s00520-017-3797-2.

(72) Ran-Castillo D, Oluwole A, Abuaisha M, Banks Paulino AR, Alkhatatneh A, Jang J, et

al. Risk, Outcomes, and Trends of Clostridium Difficile Infection in Multiple Myeloma Patients from a Nationwide Analysis. Cureus 2019;11. https://doi.org/10.7759/cureus.4391.

(73)   Scappaticci GB, Perissinotti AJ, Nagel JL, Bixby DL, Marini BL. Risk factors and impact of Clostridium difficile recurrence on haematology patients. J Antimicrob Chemother 2017;72. https://doi.org/10.1093/jac/dkx005.

(74)   Amberge S, Kramer M, Schröttner P, Heidrich K, Schmelz R, Middeke JM, et al. Clostridium Difficile infections in patients with AML or MDS undergoing allogeneic hematopoietic stem cell transplantation identify high risk for adverse outcome. Bone Marrow Transplant 2020;55. https://doi.org/10.1038/s41409-019-0678-y.

(75)   Bhandari S, Pandey RK, Dahal S, Shahreyar M, Dhakal B, Jha P, et al. Risk, Outcomes, and Predictors of Clostridium difficile Infection in Lymphoma: A Nationwide Study. South Med J 2018;111. https://doi.org/10.14423/SMJ.0000000000000872.

(76)   Luo R, Greenberg A, Stone CD. Outcomes of Clostridium difficile infection in hospitalized leukemia patients: a nationwide analysis. Infect Control Hosp Epidemiol 2015;36. https://doi.org/10.1017/ice.2015.54.

(77)   Selvey LA, Slimings C, Joske DJL, Riley TV. Clostridium difficile Infections amongst Patients with Haematological Malignancies: A Data Linkage Study. PloS One 2016;11. https://doi.org/10.1371/journal.pone.0157839.

(78)   Shargian-Alon L, Gafter-Gvili A, Ben-Zvi H, Wolach O, Yeshurun M, Raanani P, et al. Risk factors for mortality due to Acinetobacter baumannii bacteremia in patients with hematological malignancies - a retrospective study. Leuk Lymphoma 2019;60. https://doi.org/10.1080/10428194.2019.1599113.

(79)   Tofas P, Samarkos M, Piperaki E-T, Kosmidis C, Triantafyllopoulou I-D, Kotsopoulou M, et al. Pseudomonas aeruginosa bacteraemia in patients with hematologic malignancies: risk factors, treatment and outcome. Diagn Microbiol Infect Dis 2017;88. https://doi.org/10.1016/j.diagmicrobio.2017.05.003.

(80)   Zhao Y, Lin Q, Liu L, Ma R, Chen J, Shen Y, et al. Risk Factors and Outcomes of Antibiotic-resistant Pseudomonas aeruginosa Bloodstream Infection in Adult Patients With Acute Leukemia. Clin Infect Dis Off Publ Infect Dis Soc Am 2020;71. https://doi.org/10.1093/cid/ciaa1522.

(81)   Aitken SL, Sahasrabhojane PV, Kontoyiannis DP, Savidge TC, Arias CA, Ajami NJ, et al. Alterations of the Oral Microbiome and Cumulative Carbapenem Exposure Are Associated With Stenotrophomonas maltophilia Infection in Patients With Acute Myeloid Leukemia Receiving Chemotherapy. Clin Infect Dis Off Publ Infect Dis Soc Am 2021;72. https://doi.org/10.1093/cid/ciaa778.

(82)   Kim S-H, Cho SY, Kang C-I, Seok H, Huh K, Ha YE, et al. Clinical predictors of Stenotrophomonas maltophilia bacteremia in adult patients with hematologic malignancy. Ann Hematol 2018;97. https://doi.org/10.1007/s00277-017-3178-4.

(83)   Cho S-Y, Lee D-G, Choi S-M, Park C, Chun H-S, Park Y-J, et al. Stenotrophomonas maltophilia bloodstream infection in patients with hematologic malignancies: a retrospective study and in vitro activities of antimicrobial combinations. BMC Infect Dis 2015;15. https://doi.org/10.1186/s12879-015-0801-7.

(84)   Kim S-H, Cha MK, Kang C-I, Ko J-H, Huh K, Cho SY, et al. Pathogenic significance of hemorrhagic pneumonia in hematologic malignancy patients with Stenotrophomonas maltophilia bacteremia: clinical and microbiological analysis. Eur J Clin Microbiol Infect Dis Off Publ Eur Soc Clin Microbiol 2019;38. https://doi.org/10.1007/s10096-018-3425-1.

(85)   Ford CD, Coombs J, Stofer MG, Lopansri BK, Webb BJ, Ostronoff F, et al. Decrease in vancomycin-resistant Enterococcus colonization associated with a reduction in carbapenem use as empiric therapy for febrile neutropenia in patients with acute leukemia. Infect Control Hosp Epidemiol 2019;40. https://doi.org/10.1017/ice.2019.93.

(86)   Ford CD, Lopansri BK, Haydoura S, Snow G, Dascomb KK, Asch J, et al. Frequency, risk factors, and outcomes of vancomycin-resistant Enterococcus colonization and infection in patients with newly diagnosed acute leukemia: different patterns in

patients with acute myelogenous and acute lymphoblastic leukemia. Infect Control Hosp Epidemiol 2015;36. https://doi.org/10.1017/ice.2014.3.

(87)   Herc ES, Kauffman CA, Marini BL, Perissinotti AJ, Miceli MH. Daptomycin nonsusceptible vancomycin resistant Enterococcus bloodstream infections in patients with hematological malignancies: risk factors and outcomes. Leuk Lymphoma 2017;58. https://doi.org/10.1080/10428194.2017.1312665.

(88)   Hefazi M, Damlaj M, Alkhateeb HB, Partain DK, Patel R, Razonable RR, et al. Vancomycin-resistant Enterococcus colonization and bloodstream infection: prevalence, risk factors, and the impact on early outcomes after allogeneic hematopoietic cell transplantation in patients with acute myeloid leukemia. Transpl Infect Dis Off J Transplant Soc 2016;18. https://doi.org/10.1111/tid.12612.

(89)   Heisel R, Sutton R, Mascara G, Winger D, Weber D, Lim S, et al. Vancomycin-resistant enterococci in acute myeloid leukemia and myelodysplastic syndrome patients undergoing induction chemotherapy with idarubicin and cytarabine. Leuk Lymphoma 2017;58. https://doi.org/10.1080/10428194.2017.1306645.

(90)   Klein E-M, Sauer S, Klein S, Tichy D, Benner A, Bertsch U, et al. Antibiotic Prophylaxis or Granulocyte-Colony Stimulating Factor Support in Multiple Myeloma Patients Undergoing Autologous Stem Cell Transplantation. Cancers 2021;13. https://doi.org/10.3390/cancers13143439.

(91)   Kamboj M, Cohen N, Huang Y-T, Kerpelev M, Jakubowski A, Sepkowitz KA, et al. Impact of Empiric Treatment for Vancomycin-Resistant Enterococcus in Colonized Patients Early after Allogeneic Hematopoietic Stem Cell Transplantation. Biol Blood Marrow Transplant J Am Soc Blood Marrow Transplant 2019;25. https://doi.org/10.1016/j.bbmt.2018.11.008.

(92)   Kirkizlar TA, Akalin H, Kirkizlar O, Ozkalemkas F, Ozkocaman V, Kazak E, et al. Vancomycin-resistant enterococci infection and predisposing factors for infection and mortality in patients with acute leukaemia and febrile neutropenia. Leuk Res 2020;99. https://doi.org/10.1016/j.leukres.2020.106463.

(93)   Ornstein MC, Mukherjee S, Keng M, Elson P, Tiu RV, Saunthararajah Y, et al. Impact of vancomycin-resistant enterococcal bacteremia on outcome during acute myeloid leukemia induction therapy. Leuk Lymphoma 2015;56. https://doi.org/10.3109/10428194.2014.1003557.

(94)   Papanicolaou GA, Ustun C, Young J-AH, Chen M, Kim S, Woo Ahn K, et al. Bloodstream Infection Due to Vancomycin-resistant Enterococcus Is Associated With Increased Mortality After Hematopoietic Cell Transplantation for Acute Leukemia and Myelodysplastic Syndrome: A Multicenter, Retrospective Cohort Study. Clin Infect Dis Off Publ Infect Dis Soc Am 2019;69. https://doi.org/10.1093/cid/ciz031.

(95)   Zhang P, Wang J, Hu H, Zhang S, Wei J, Yang Q, et al. Clinical Characteristics and Risk Factors for Bloodstream Infection Due to Carbapenem-Resistant Klebsiella pneumoniae in Patients with Hematologic Malignancies. Infect Drug Resist 2020;13. https://doi.org/10.2147/IDR.S272217.

(96)   Micozzi A, Gentile G, Santilli S, Minotti C, Capria S, Moleti ML, et al. Reduced mortality from KPC-K.pneumoniae bloodstream infection in high-risk patients with hematological malignancies colonized by KPC-K.pneumoniae. BMC Infect Dis 2021;21. https://doi.org/10.1186/s12879-021-06747-8.

(97)   Trecarichi EM, Pagano L, Martino B, Candoni A, Di Blasi R, Nadali G, et al. Bloodstream infections caused by Klebsiella pneumoniae in onco-hematological patients: clinical impact of carbapenem resistance in a multicentre prospective survey. Am J Hematol 2016;91. https://doi.org/10.1002/ajh.24489.

(98)   Micozzi A, Gentile G, Minotti C, Cartoni C, Capria S, Ballarò D, et al. Carbapenem-resistant Klebsiella pneumoniae in high-risk haematological patients: factors favouring spread, risk factors and outcome of carbapenem-resistant Klebsiella pneumoniae bacteremias. BMC Infect Dis 2017;17. https://doi.org/10.1186/s12879-017-2297-9.

(99)   Benanti GE, Brown ART, Shigle TL, Tarrand JJ, Bhatti MM, McDaneld PM, et al.

Carbapenem versus Cefepime or Piperacillin-Tazobactam for Empiric Treatment of Bacteremia Due to Extended-Spectrum-β-Lactamase-Producing Escherichia coli in Patients with Hematologic Malignancy. Antimicrob Agents Chemother 2019;63. https://doi.org/10.1128/AAC.01813-18.

(100) Trecarichi EM, Giuliano G, Cattaneo C, Ballanti S, Criscuolo M, Candoni A, et al. Bloodstream infections caused by Escherichia coli in onco-haematological patients: Risk factors and mortality in an Italian prospective survey. PloS One 2019;14. https://doi.org/10.1371/journal.pone.0224465.

(101) Heo ST, Tatara AM, Jiménez-Ortigosa C, Jiang Y, Lewis RE, Tarrand J, et al. Changes in In Vitro Susceptibility Patterns of Aspergillus to Triazoles and Correlation With Aspergillosis Outcome in a Tertiary Care Cancer Center, 1999-2015. Clin Infect Dis Off Publ Infect Dis Soc Am 2017;65. https://doi.org/10.1093/cid/cix297.

(102) Magira E, Jiang Y, Economides M, Tarrand J, Kontoyiannis D. Mixed mold pulmonary infections in haematological cancer patients in a tertiary care cancer centre. Mycoses 2018;61. https://doi.org/10.1111/myc.12830.

(103) Wang E, Farmakiotis D, Yang D, McCue DA, Kantarjian HM, Kontoyiannis DP, et al. The ever-evolving landscape of candidaemia in patients with acute leukaemia: non-susceptibility to caspofungin and multidrug resistance are associated with increased mortality. J Antimicrob Chemother 2015;70. https://doi.org/10.1093/jac/dkv087.

(104) Folan SA, Marx KR, Tverdek FP, Raad I, Mulanovich VE, Tarrand JJ, et al. Clinical Outcomes Associated With Linezolid Resistance in Leukemia Patients With Linezolid-Resistant Staphylococcus epidermidis Bacteremia. Open Forum Infect Dis 2018;5. https://doi.org/10.1093/ofid/ofy167.

(105) Valcarcel B, De-la-Cruz-Ku G, Malpica L, Enriquez-Vera D. Clinical features and outcome of Aeromonas sobria bacteremia in pediatric and adult patients with hematologic malignancies: A single-center retrospective study in Peru. PloS One 2021;16. https://doi.org/10.1371/journal.pone.0255910.

(106) Bastug A, Kayaaslan B, Kazancioglu S, But A, Aslaner H, Akinci E, et al. Emergence of multidrug resistant isolates and mortality predictors in patients with solid tumors or hematological malignancies. J Infect Dev Ctries 2015;9. https://doi.org/10.3855/jidc.6805.

(107) Chiang H-Y, Wu T-H, Hsu C-Y, Chao W-C. Association Between Positive Cultures During Admission and 1-Year Mortality in Patients With Cancer Receiving Perioperative Intensive Care. Cancer Control J Moffitt Cancer Cent 2018;25. https://doi.org/10.1177/1073274818794162.

(108) Freire MP, Pierrotti LC, Zerati AE, Benites L, da Motta-Leal Filho JM, Ibrahim KY, et al. Role of Lock Therapy for Long-Term Catheter-Related Infections by Multidrug-Resistant Bacteria. Antimicrob Agents Chemother 2018;62. https://doi.org/10.1128/AAC.00569-18.

(109) Islas-Muñoz B, Volkow-Fernández P, Ibanes-Gutiérrez C, Villamar-Ramírez A, Vilar-Compte D, Cornejo-Juárez P. Bloodstream infections in cancer patients. Risk factors associated with mortality. Int J Infect Dis IJID Off Publ Int Soc Infect Dis 2018;71. https://doi.org/10.1016/j.ijid.2018.03.022.

(110) Joncour A, Puyade M, Michaud A, Tourani J-M, Cazenave-Roblot F, Rammaert B. Is current initial empirical antibiotherapy appropriate to treat bloodstream infections in short-duration chemo-induced febrile neutropenia? Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2020;28. https://doi.org/10.1007/s00520-019-05113-4.

(111) Marín M, Gudiol C, Ardanuy C, Garcia-Vidal C, Jimenez L, Domingo-Domenech E, et al. Factors influencing mortality in neutropenic patients with haematologic malignancies or solid tumours with bloodstream infection. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis 2015;21. https://doi.org/10.1016/j.cmi.2015.01.029.

(112) Martinez-Nadal G, Puerta-Alcalde P, Gudiol C, Cardozo C, Albasanz-Puig A, Marco F,

et al. Inappropriate Empirical Antibiotic Treatment in High-risk Neutropenic Patients With Bacteremia in the Era of Multidrug Resistance. Clin Infect Dis Off Publ Infect Dis Soc Am 2020;70. https://doi.org/10.1093/cid/ciz319.

(113) Royo-Cebrecos C, Gudiol C, Ardanuy C, Pomares H, Calvo M, Carratalà J. A fresh look at polymicrobial bloodstream infection in cancer patients. PloS One 2017;12. https://doi.org/10.1371/journal.pone.0185768.

(114) Jiang A-M, Shi X, Liu N, Gao H, Ren M-D, Zheng X-Q, et al. Nosocomial infections due to multidrug-resistant bacteria in cancer patients: a six-year retrospective study of an oncology Center in Western China. BMC Infect Dis 2020;20. https://doi.org/10.1186/s12879-020-05181-6.

(115) Puerta-Alcalde P, Ambrosioni J, Chumbita M, Hernández-Meneses M, Garcia-Pouton N, Cardozo C, et al. Clinical Characteristics and Outcome of Bloodstream Infections in HIV-Infected Patients with Cancer and Febrile Neutropenia: A Case-Control Study. Infect Dis Ther 2021;10. https://doi.org/10.1007/s40121-021-00445-3.

(116) Costa P de O, Atta EH, Silva ARA da. Infection with multidrug-resistant gram-negative bacteria in a pediatric oncology intensive care unit: risk factors and outcomes. J Pediatr (Rio J) 2015;91. https://doi.org/10.1016/j.jped.2014.11.009.

(117) Marini BL, Hough SM, Gregg KS, Abu-Seir H, Nagel JL. Risk factors for piperacillin/tazobactam-resistant Gram-negative infection in hematology/oncology patients with febrile neutropenia. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2015;23. https://doi.org/10.1007/s00520-014-2582-8.

(118) Tossey JC, El Boghdadly Z, Reed EE, Dela-Pena J, Coe K, Williams SN, et al. Oral fluoroquinolones for definitive treatment of gram-negative bacteremia in cancer patients. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2021;29. https://doi.org/10.1007/s00520-021-06063-6.

(119) Kim Y, Jung S, Kang J, Ryoo S, Sohn S, Seo D, et al. Risk factors for extended-spectrum beta-lactamase-producing Enterobacteriaceae infection causing septic shock in cancer patients with chemotherapy-induced febrile neutropenia. Intern Emerg Med 2019;14. https://doi.org/10.1007/s11739-018-02015-x.

(120) Refay SM, Ahmed EH, Abd ELzaher AR, Morsy AM, Yasser MM, Mahmoud AM. Risk of Drug Resistance and Repeated Infection with Klebsiella pneumoniae and Escherichia coli in Intensive Care Unit Cancer Patients. Comb Chem High Throughput Screen 2022;25. https://doi.org/10.2174/1386207324666210121104724.

(121) Medboua-Benbalagh C, Touati A, Kermas R, Gharout-Sait A, Brasme L, Mezhoud H, et al. Fecal Carriage of Extended-Spectrum β-Lactamase-Producing Enterobacteriaceae Strains Is Associated with Worse Outcome in Patients Hospitalized in the Pediatric Oncology Unit of Beni-Messous Hospital in Algiers, Algeria. Microb Drug Resist Larchmt N 2017;23. https://doi.org/10.1089/mdr.2016.0153.

(122) Ben-Chetrit E, Eldaim MA, Bar-Meir M, Dodin M, Katz DE. Associated factors and clinical outcomes of bloodstream infection due to extended-spectrum β-lactamase-producing Escherichia coli and Klebsiella pneumoniae during febrile neutropenia. Int J Antimicrob Agents 2019;53. https://doi.org/10.1016/j.ijantimicag.2018.12.003.

(123) Ceken S, Iskender G, Gedik H, Duygu F, Mert D, Kaya AH, et al. Risk factors for bloodstream infections due to extended-spectrum β-lactamase producing Enterobacteriaceae in cancer patients. J Infect Dev Ctries 2018;12. https://doi.org/10.3855/jidc.9720.

(124) Martínez-Valencia A, Gómez Martínez B, Montañez Ayala A, García K, Sánchez Pedraza SP, Jiménez Cetina L, et al. Development and validation of a scoring system for predicting cancer patients at risk of extended-spectrum b-lactamase-producing Enterobacteriaceae infections. BMC Infect Dis 2020;20. https://doi.org/10.1186/s12879-020-05280-4.

(125) Abu-Sbeih H, Choi K, Tran CN, Wang X, Lum P, Shuttlesworth G, et al. Recurrent Clostridium difficile infection is associated with treatment failure and prolonged illness

in cancer patients. Eur J Gastroenterol Hepatol 2019;31.
https://doi.org/10.1097/MEG.0000000000001288.

(126) Chang GY, Dembry LM, Banach DB. Epidemiology of Clostridium difficile infection in hospitalized oncology patients. Am J Infect Control 2016;44.
https://doi.org/10.1016/j.ajic.2016.04.210.

(127) Daida A, Yoshihara H, Inai I, Hasegawa D, Ishida Y, Urayama KY, et al. Risk Factors for Hospital-acquired Clostridium difficile Infection Among Pediatric Patients With Cancer. J Pediatr Hematol Oncol 2017;39.
https://doi.org/10.1097/MPH.0000000000000742.

(128) Fuereder T, Koni D, Gleiss A, Kundi M, Makristathis A, Zielinski C, et al. Risk factors for Clostridium difficile infection in hemato-oncological patients: A case control study in 144 patients. Sci Rep 2016;6. https://doi.org/10.1038/srep31498.

(129) Hebbard AIT, Slavin MA, Reed C, Trubiano JA, Teh BW, Haeusler GM, et al. Risks factors and outcomes of Clostridium difficile infection in patients with cancer: a matched case-control study. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2017;25. https://doi.org/10.1007/s00520-017-3606-y.

(130) Shoaei P, Shojaei H, Khorvash F, M Hosseini S, Ataei B, Esfandiari Z, et al. Clostridium difficile infection in cancer patients with hospital acquired diarrhea at the teaching hospitals in Iran: Multilocus sequence typing analysis (MLST) and Antimicrobial resistance pattern. Ann Ig Med Prev E Comunita 2019;31.
https://doi.org/10.7416/ai.2019.2298.

(131) Willis DN, Huang FS, Elward AM, Wu N, Magnusen B, Dubberke ER, et al. Clostridioides difficile Infections in Inpatient Pediatric Oncology Patients: A Cohort Study Evaluating Risk Factors and Associated Outcomes. J Pediatr Infect Dis Soc 2021;10. https://doi.org/10.1093/jpids/piaa090.

(132) Willis ZI, Nicholson MR, Esbenshade AJ, Xu M, Slaughter JC, Friedman DL, et al. Intensity of Therapy for Malignancy and Risk for Recurrent and Complicated Clostridium difficile Infection in Children. J Pediatr Hematol Oncol 2019;41.
https://doi.org/10.1097/MPH.0000000000001411.

(133) Gupta A, Tariq R, Frank RD, Jean GW, Beg MS, Pardi DS, et al. Trends in the Incidence and Outcomes of Hospitalized Cancer Patients With Clostridium difficile Infection: A Nationwide Analysis. J Natl Compr Cancer Netw JNCCN 2017;15.
https://doi.org/10.6004/jnccn.2017.0046.

(134) Delgado A, Reveles IA, Cabello FT, Reveles KR. Poorer outcomes among cancer patients diagnosed with Clostridium difficile infections in United States community hospitals. BMC Infect Dis 2017;17. https://doi.org/10.1186/s12879-017-2553-z.

(135) Siddiqui NS, Khan Z, Khan MS, Khan Z, Haq KF, Solanki SD, et al. Trends in Incidence and Outcomes of Clostridium difficile Colitis in Hospitalized Patients of Febrile Neutropenia: A Nationwide Analysis. J Clin Gastroenterol 2019;53.
https://doi.org/10.1097/MCG.0000000000001171.

(136) Fan L, Wang Z, Wang Q, Xiong Z, Xu Y, Li D, et al. Increasing rates of Acinetobacter baumannii infection and resistance in an oncology department. J Cancer Res Ther 2018;14. https://doi.org/10.4103/jcrt.JCRT_737_17.

(137) Cornejo-Juárez P, Cevallos MA, Castro-Jaimes S, Castillo-Ramírez S, Velázquez-Acosta C, Martínez-Oliva D, et al. High mortality in an outbreak of multidrug resistant Acinetobacter baumannii infection introduced to an oncological hospital by a patient transferred from a general hospital. PloS One 2020;15.
https://doi.org/10.1371/journal.pone.0234684.

(138) Freire MP, de Oliveira Garcia D, Garcia CP, Campagnari Bueno MF, Camargo CH, Kono Magri ASG, et al. Bloodstream infection caused by extensively drug-resistant Acinetobacter baumannii in cancer patients: high mortality associated with delayed treatment rather than with the degree of neutropenia. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis 2016;22. https://doi.org/10.1016/j.cmi.2015.12.010.

(139) Ñamendys-Silva SA, Correa-García P, García-Guillén FJ, González-Herrera MO, Pérez-Alonso A, Texcocano-Becerra J, et al. Outcomes of critically ill cancer patients

with Acinetobacter baumannii infection. World J Crit Care Med 2015;4. https://doi.org/10.5492/wjccm.v4.i3.258.

(140) Gudiol C, Albasanz-Puig A, Laporte-Amargós J, Pallarès N, Mussetti A, Ruiz-Camps I, et al. Clinical Predictive Model of Multidrug Resistance in Neutropenic Cancer Patients with Bloodstream Infection Due to Pseudomonas aeruginosa. Antimicrob Agents Chemother 2020;64. https://doi.org/10.1128/AAC.02494-19.

(141) Smith ZR, Tajchman SK, Dee BM, Bruno JJ, Qiao W, Tverdek FP. Development of a combination antibiogram for Pseudomonas aeruginosa bacteremia in an oncology population. J Oncol Pharm Pract Off Publ Int Soc Oncol Pharm Pract 2016;22. https://doi.org/10.1177/1078155215586081.

(142) Velázquez-Acosta C, Zarco-Márquez S, Jiménez-Andrade MC, Volkow-Fernández P, Cornejo-Juárez P. Stenotrophomonas maltophilia bacteremia and pneumonia at a tertiary-care oncology center: a review of 16 years. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2018;26. https://doi.org/10.1007/s00520-017-4032-x.

(143) Aktürk H, Sütçü M, Somer A, Karaman S, Acar M, Ünüvar A, et al. Results of Four-Year Rectal Vancomycin-Resistant Enterococci Surveillance in a Pediatric Hematology-Oncology Ward: From Colonization to Infection. Turk J Haematol Off J Turk Soc Haematol 2016;33. https://doi.org/10.4274/tjh.2015.0368.

(144) Akhtar N, Sultan F, Nizamuddin S, Zafar W. Risk factors and clinical outcomes for vancomycin-resistant enterococcus bacteraemia in hospitalised cancer patients in Pakistan: A case-control study. JPMA J Pak Med Assoc 2016;66.

(145) Di Domenico EG, Cavallo I, Sivori F, Marchesi F, Prignano G, Pimpinelli F, et al. Biofilm Production by Carbapenem-Resistant Klebsiella pneumoniae Significantly Increases the Risk of Death in Oncological Patients. Front Cell Infect Microbiol 2020;10. https://doi.org/10.3389/fcimb.2020.561741.

(146) Freire MP, Pierrotti LC, Filho HHC, Ibrahim KY, Magri ASGK, Bonazzi PR, et al. Infection with Klebsiella pneumoniae carbapenemase (KPC)-producing Klebsiella pneumoniae in cancer patients. Eur J Clin Microbiol Infect Dis Off Publ Eur Soc Clin Microbiol 2015;34. https://doi.org/10.1007/s10096-014-2233-5.

(147) Nham E, Huh K, Cho SY, Chung DR, Peck KR, Lee NY, et al. Characteristics and Clinical Outcomes of Extended-Spectrum beta-lactamase-producing Klebsiella pneumoniae Bacteremia in Cancer Patients. Infect Chemother 2020;52. https://doi.org/10.3947/ic.2020.52.1.59.

(148) Bello-Chavolla OY, Bahena-Lopez JP, Garciadiego-Fosass P, Volkow P, Garcia-Horton A, Velazquez-Acosta C, et al. Bloodstream infection caused by S. aureus in patients with cancer: a 10-year longitudinal single-center study. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2018;26. https://doi.org/10.1007/s00520-018-4275-1.

(149) Khanzada MJ, Zianab I, Samreen A. A RESEARCH STUDY ON BACTERAEMIA PRODUCED THROUGH ESCHERICHIA COLI IN TUMOR PATIENTS AT THE SPECIFIC CENTER IN OUR COUNTRY. INDO Am J Pharm Sci 2019;o6. https://doi.org/10.5281/zenodo.2593737.

(150) Zhang Q, Zhang W, Li Z, Bai C, Li D, Zheng S, et al. Bacteraemia due to AmpC β-lactamase-producing Escherichia coli in hospitalized cancer patients: risk factors, antibiotic therapy, and outcomes. Diagn Microbiol Infect Dis 2017;88. https://doi.org/10.1016/j.diagmicrobio.2017.04.006.

(151) Parveen A, Sultan F, Raza A, Zafar W, Nizamuddin S, Mahboob A, et al. Bacteraemia caused by Escherichia coli in cancer patients at a specialist center in Pakistan. JPMA J Pak Med Assoc 2015;65.

(152) Zhang Q, Gao H-Y, Li D, Li Z, Qi S-S, Zheng S, et al. Clinical outcome of Escherichia coli bloodstream infection in cancer patients with/without biofilm formation: a single-center retrospective study. Infect Drug Resist 2019;12. https://doi.org/10.2147/IDR.S192072.

(153) Farmakiotis D, Kyvernitakis A, Tarrand JJ, Kontoyiannis DP. Early initiation of

appropriate treatment is associated with increased survival in cancer patients with Candida glabrata fungaemia: a potential benefit from infectious disease consultation. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis 2015;21. https://doi.org/10.1016/j.cmi.2014.07.006.

(154) Jung DS, Farmakiotis D, Jiang Y, Tarrand JJ, Kontoyiannis DP. Uncommon Candida Species Fungemia among Cancer Patients, Houston, Texas, USA. Emerg Infect Dis 2015;21. https://doi.org/10.3201/eid2111.150404.

(155) Le A, Farmakiotis D, Tarrand JJ, Kontoyiannis DP. Initial Treatment of Cancer Patients with Fluconazole-Susceptible Dose-Dependent Candida glabrata Fungemia: Better Outcome with an Echinocandin or Polyene Compared to an Azole? Antimicrob Agents Chemother 2017;61. https://doi.org/10.1128/AAC.00631-17.

(156) Paixao de Sousa da Silva AM, de Moraes-Pinto MI, Teofilo Pignati L, Barbosa Teixeira B, Cordeiro Lima AP, Costa Pimentel Germano P, et al. Candida spp bloodstream infections in a Latin American Pediatric Oncology Reference Center: Epidemiology and associated factors. Mycoses 2020;63. https://doi.org/10.1111/myc.13106.

(157) Puig-Asensio M, Ruiz-Camps I, Fernández-Ruiz M, Aguado JM, Muñoz P, Valerio M, et al. Epidemiology and outcome of candidaemia in patients with oncological and haematological malignancies: results from a population-based surveillance in Spain. Clin Microbiol Infect Off Publ Eur Soc Clin Microbiol Infect Dis 2015;21. https://doi.org/10.1016/j.cmi.2014.12.027.

(158) Raza A, Zafar W, Mahboob A, Nizammudin S, Rashid N, Sultan F. Clinical features and outcomes of Candidaemia in cancer patients: Results from Pakistan. JPMA J Pak Med Assoc 2016;66.

(159) Sun M, Chen C, Xiao W, Chang Y, Liu C, Xu Q. Increase in Candida Parapsilosis Candidemia in Cancer Patients. Mediterr J Hematol Infect Dis 2019;11. https://doi.org/10.4084/MJHID.2019.012.

(160) Ko J-H, Kang C-I, Lee WJ, Huh K, Yoo JR, Kim K, et al. Clinical features and risk factors for development of Bacillus bacteremia among adult patients with cancer: a case-control study. Support Care Cancer Off J Multinatl Assoc Support Care Cancer 2015;23. https://doi.org/10.1007/s00520-014-2382-1.

(161) Fontana NS, Ibrahim KY, Bonazzi PR, Rossi F, Almeida SCG, Tengan FM, et al. Fluoroquinolone treatment as a protective factor for 10-day mortality in Streptococcus pneumoniae bacteremia in cancer patients. Sci Rep 2021;11. https://doi.org/10.1038/s41598-021-81415-0.

(162) Altman DG. Systematic reviews of evaluations of prognostic variables. BMJ 2001;323. https://doi.org/10.1136/bmj.323.7306.224.

(163) Alevizakos M, Gaitanidis A, Andreatos N, Arunachalam K, Flokas ME, Mylonakis E. Bloodstream infections due to extended-spectrum β-lactamase-producing Enterobacteriaceae among patients with malignancy: a systematic review and meta-analysis. Int J Antimicrob Agents 2017;50. https://doi.org/10.1016/j.ijantimicag.2017.07.003.

(164) Alevizakos M, Gaitanidis A, Nasioudis D, Tori K, Flokas ME, Mylonakis E. Colonization With Vancomycin-Resistant Enterococci and Risk for Bloodstream Infection Among Patients With Malignancy: A Systematic Review and Meta-Analysis. Open Forum Infect Dis 2017;4. https://doi.org/10.1093/ofid/ofw246.

(165) Li Z, Zhuang H, Wang G, Wang H, Dong Y. Prevalence, predictors, and mortality of bloodstream infections due to methicillin-resistant Staphylococcus aureus in patients with malignancy: systemic review and meta-analysis. BMC Infect Dis 2021;21. https://doi.org/10.1186/s12879-021-05763-y.

(166) Montassier E, Batard E, Gastinne T, Potel G, de La Cochetière MF. Recent changes in bacteremia in patients with cancer: a systematic review of epidemiology and antibiotic resistance. Eur J Clin Microbiol Infect Dis Off Publ Eur Soc Clin Microbiol 2013;32. https://doi.org/10.1007/s10096-013-1819-7.

(167) Bender R, Lange S. Adjusting for multiple testing--when and how? J Clin Epidemiol 2001;54. https://doi.org/10.1016/s0895-4356(00)00314-0.

(168) Greenland S, Mansournia MA, Altman DG. Sparse data bias: a problem hiding in plain sight. BMJ 2016;352. https://doi.org/10.1136/bmj.i1981.

(169) Ioannidis JPA. Why Most Discovered True Associations Are Inflated. Epidemiology 2008;19. https://doi.org/10.1097/EDE.0b013e31818131e7.

(170) Hayden JA, Côté P, Bombardier C. Evaluation of the Quality of Prognosis Studies in Systematic Reviews. Ann Intern Med 2006;144. https://doi.org/10.7326/0003-4819-144-6-200603210-00010.

(171) Wolff RF, Moons KGM, Riley RD, Whiting PF, Westwood M, Collins GS, et al. PROBAST: A Tool to Assess the Risk of Bias and Applicability of Prediction Model Studies. Ann Intern Med 2019;170. https://doi.org/10.7326/M18-1376.

(172) Smith G. Step away from stepwise. J Big Data 2018;5. https://doi.org/10.1186/s40537-018-0143-6.

(173) Sun G-W, Shook TL, Kay GL. Inappropriate use of bivariable analysis to screen risk factors for use in multivariable analysis. J Clin Epidemiol 1996;49. https://doi.org/10.1016/0895-4356(96)00025-X.

(174) Hernán MA, Robins JM. Causal Inference: What If. Boca Raton: Chapman & Hall/CRC; 2020.

Supplementary material 1

First search

Database: Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations, Daily and Versions(R) <1946 to June 21, 2021>

Date: 22.06.21

Hits: 6878

1	exp Neoplasms/	3485612
2	(tumor? or tumour? or cancer? or neoplas* or paraneoplastic or precancer* or cyst? or malignanc* or carcinogenes#s or oncogenes#s or tumorigenes#s or cocarcinogenes#s or metastas#s or metastatic or incidentaloma?).tw,kf.	3441687
3	1 or 2	4533774
4	exp Drug Resistance, Microbial/ or exp Drug Resistance, Bacterial/	170727
5	(((antibacterial or bacterial or bacterium or antibiotic or antimicrobial or microbial or multidrug or multiple drug) adj2 (resistance or resistant)) or antibiotic non-susceptibility or antibiotic nonsusceptibility).tw,kf.	136901
6	4 or 5	254631
7	3 and 6	22036
8	limit 7 to (danish or english or interlingua or multilingual or norwegian or spanish or swedish)	20721
9	limit 8 to yr="2015 -Current"	6878

Database: Embase <1974 to 2021 June 22>

Date: 23.06.21

Hits: 7314

1	exp neoplasm/	4730064
2	(tumor? or tumour? or cancer? or neoplas* or paraneoplastic or precancer* or cyst? or malignanc* or carcinogenes#s or oncogenes#s or tumorigenes#s or cocarcinogenes#s or metastas#s or metastatic or incidentaloma?).tw,kw.	4525726
3	1 or 2	5821589

4	exp antibiotic resistance/	177653
5	(((antibacterial or bacterial or bacterium or antibiotic or antimicrobial or microbial or multidrug or multiple drug) adj2 (resistance or resistant)) or antibiotic non-susceptibility or antibiotic nonsusceptibility).tw,kw.	167050
6	4 or 5	267257
7	3 and 6	28672
8	limit 7 to (danish or english or norwegian or polyglot or spanish or swedish)	26746
9	limit 8 to yr="2015 -Current"	9903
10	limit 9 to embase	7314

Database: **Web of Science (Science Citation Index Expanded (SCI-EXPANDED) --1987-present, Social Sciences Citation Index (SSCI) --1987-present, Arts & Humanities Citation Index (A&HCI) --1987-present, Emerging Sources Citation Index (ESCI) --2015-present)**

Date: 24.06.21

Hits: 11,302

1	TS=("tumor$" or "tumour$" or "cancer$" or "neoplas*" or "paraneoplastic" or "precancer*" or "cyst$" or "malignanc*" or "carcinogenes?s" or "oncogenes?s" or "tumorigenes?s" or "cocarcinogenes?s" or "metastas?s" or "metastatic" or "incidentaloma$")	3,746,108
2	TS=((("antibacterial" or "bacterial" or "bacterium" or "antibiotic " or "antimicrobial" or "microbial" or "multidrug" or "multiple drug") NEAR/1 ("resistance" or "resistant")) or "antibiotic non-susceptibility" or "antibiotic nonsusceptibility")	172,855
3	(#2 AND #1) AND LANGUAGE: (English OR Danish OR Multiple Languages OR Norwegian OR Spanish OR Swedish)	11,302

Supplementary material 2

Second search

Database: Ovid MEDLINE(R) and Epub Ahead of Print, In-Process, In-Data-Review & Other Non-Indexed Citations, Daily and Versions(R) <1946 to November 18, 2021>

Date: 19.11.21

Hits: 12 478

1	exp Neoplasms/	3573100
2	(tumor? or tumour? or cancer* or neoplas* or paraneoplas* or preneoplas* or precancer* or premalign* or malign* or carcinogenes#s or oncogenes#s or tumorigenes#s or cocarcinogenes#s or metastas#s or metastatic or incidentaloma? or leukemia? or leucemia? or lymphoma? or sarcoma? or carcinoma* or melanoma? or glioma ? or blastoma?).tw,kf.	3958905
3	1 or 2	4744972
4	exp Drug Resistance, Microbial/ or exp Drug Resistance, Bacterial/	174771
5	((antibacterial or bacterial or bacterium or antibiotic or antifungal or antimicrobial or microbial or multidrug or multiple drug or multiple antibiotics or penicillin or meth#cillin? or met#cillin? or cephalosporin* or cephalosporanic acid? or cefepim? or cefuroxime or cefotaxim? or ceftazidime or ceftriaxon? or cefixime or ceftaroline or carbapenem? or aminoglycoside? or gentam#cin? or amikacin? or fluoroquinolone? or linezolid? or vancomycin? or echinocandin? or azole? or colistin? or colimycin? or polymyxin e or intrinsic* or penicillin or macrolide or tobramycin? or fluconazol? or voricanozole or posaconazole or caspofungin or anidulafungin or micafungin or ciproflox* or ciproxin or levofloxacin or clindamycin? or dalacin or clarithromycin or CLAR or erythromycin or azt?reonam or azactam) adj4 (resistance or resistant or nonsusceptib* or insusceptib* or non-susceptib* or susceptibility)).tw,kf.	235262
6	(piperacillin adj3 tazobactam adj4 resistant).tw,kf.	202
7	beta-Lactamases/ or Methicillin-Resistant Staphylococcus aureus/ or Vancomycin-Resistant Enterococci/ or Vancomycin-Resistant Staphylococcus aureus/	40508
8	(b*-lactamase? or "ec 3.5.2.6" or beta lactam?hydrolase or cefinase or ESBL or carbapenemase? or biofilm or non-fermenting).tw,kf.	88504
9	Pseudomonas aeruginosa/ or (pseudomonas aeruginosa or "P. aeruginosa").tw,kf.	74856

10	Acinetobacter/ or (acinetobacter* or "A. baumanii").tw,kf.	20486
11	Stenotrophomonas Maltophilia/ or ((Stenotrophomonas or Pseudomonas or Xanthomonas) adj malt#philia).tw,kf.	3477
12	Clostridioides difficile/ or ((clostridium or clostridioides) adj difficil*).tw,kf.	17332
13	Enterococcus faecium/ or ((enterococcus or streptococcus) adj faecium).tw,kf.	7433
14	(candida adj1 (non albicans or auris)).tw,kf.	1779
15	Aspergillus fumigatus/ or ((aspergillus or neosartorya) adj fumigat*).tw,kf.	12302
16	or/4-15	456787
17	3 and 16	35096
18	limit 17 to (danish or english or interlingua or multilingual or norwegian or spanish or swedish)	32942
19	limit 18 to yr="2015 -Current"	12478

Database: Embase <1974 to 2021 November 17>

Date: 19.11.21

Hits: 16471

1	exp Neoplasm/	4853484
2	(tumor? or tumour? or cancer* or neoplas* or paraneoplas* or preneoplas* or precancer* or premalign* or malign* or carcinogenes#s or oncogenes#s or tumorigenes#s or cocarcinogenes#s or metastas#s or metastatic or incidentaloma? or leukemia? or leucemia? or lymphoma? or sarcoma? or carcinoma* or melanoma? or glioma ? or blastoma?).tw,kf.	5177268
3	1 or 2	6060075
4	exp Antibiotic Resistance/ or Antifungal resistance/ or Antifungal susceptibility/ or Multidrug resistance/ or Extensive Drug Resistance/ or Antifungal susceptibility/	229586
5	((antibacterial or bacterial or bacterium or antibiotic or antifungal or antimicrobial or microbial or multidrug or multiple drug or multiple antibiotics or penicillin or meth#cillin? or met#cillin? or cephalosporin* or cephalosporanic acid? or cefepim? or cefuroxime or	287384

	cefotaxim? or ceftazidime or ceftriaxon? or cefixime or ceftaroline or carbapenem? or aminoglycoside? or gentam#cin? or amikacin? or fluoroquinolone? or linezolid? or vancomycin? or echinocandin? or azole? or colistin? or colimycin? or polymyxin e or intrinsic* or penicillin or macrolide or tobramycin? or fluconazol? or voricanozole or posaconazole or caspofungin or anidulafungin or micafungin or ciproflox* or ciproxin or levofloxacin or clindamycin? or dalacin or clarithromycin or CLAR or erythromycin or azt?reonam or azactam) adj4 (resistance or resistant or nonsusceptib* or insusceptib* or non-susceptib* or susceptibility)).tw,kf.	
6	(piperacillin adj3 tazobactam adj4 resistant).tw,kf.	334
7	beta-Lactamase/ or Extended Spectrum Beta Lactamase/ or Carbapenemase/ or Carbapenemase Producing Enterobacteriaceae/ or Carbapenem resistance/ or Methicillin Resistant Staphylococcus aureus/ or Vancomycin-Resistant Enterococcus/ or Vancomycin Resistant Staphylococcus aureus/	85966
8	(b*-lactamase? or "ec 3.5.2.6" or beta lactam?hydrolase or cefinase or ESBL or carbapenemase? or biofilm or non-fermenting).tw,kf.	105594
9	Pseudomonas aeruginosa/ or (pseudomonas aeruginosa or "P. aeruginosa").tw,kf.	122215
10	Acinetobacter/ or (acinetobacter* or "A. baumanii").tw,kf.	29911
11	Stenotrophomonas Maltophilia/ or ((Stenotrophomonas or Pseudomonas or Xanthomonas) adj malt#philia).tw,kf.	7160
12	Clostridioides difficile/ or ((clostridium or clostridioides) adj difficil*).tw,kf.	24295
13	Enterococcus faecium/ or ((enterococcus or streptococcus) adj faecium).tw,kf.	12422
14	Candida auris/ or (candida adj1 (non albicans or auris)).tw,kf.	2531
15	Aspergillus fumigatus/ or ((aspergillus or neosartorya) adj fumigat*).tw,kf.	20302
16	or/4-15	575608
17	3 and 16	57855
18	limit 17 to (danish or english or norwegian or polyglot or spanish or swedish)	54246
19	limit 18 to yr="2015 -Current"	23248
20	limit 19 to embase	16471

14	(#13) AND LA=(English OR Danish OR Multiple Languages OR Norwegian OR Spanish OR Swedish)	19,369
13	(#12) AND PY=(2015-2021)	19,528
12	#1 and #11	46,063
11	#2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10	454,205
10	TS=(("aspergillus" or "neosartorya") NEAR/0 "fumigat*")	13,687
9	TS=("candida" NEAR/0 ("non albicans" or "auris"))	1,906
8	TS=(("enterococcus" or "streptococcus") NEAR/0 "faecium")	8,677
7	TS=(("clostridium" or "clostridioides") NEAR/0 "difficil*")	20,940
6	TS=(("Stenotrophomonas" or "Pseudomonas" or "Xanthomonas") NEAR/0 "malt?philia")	4,062
5	TS=("acinetobacter*" or "A. baumanii")	23,475
4	TS=("pseudomonas aeruginosa" or "P. aeruginosa")	88,969
3	TS=("b*-lactamase$" or "ec 3.5.2.6" or "beta lactam hydrolase" or "beta lactamhydrolase" or "cefinase" or "ESBL" or "carbapenemase$" or "biofilm" or "non-fermenting")	118,459
2	TS=(("antibacterial" or "bacterial" or "bacterium" or "antibiotic$" or "antifungal" or "antimicrobial" or "microbial" or "multidrug" or "multiple drug" or "multiple antibiotics" or "penicillin" or "meth?cillin$" or "met?cillin$" or "cephalosporin*" or "cephalosporanic acid$" or "cefepim$" or "cefuroxime" or "cefotaxim$" or "ceftazidime" or "ceftriaxon$" or "cefixime" or "ceftaroline" or "carbapenem$" or "aminoglycoside$" or "gentam?cin$" or "amikacin$" or "fluoroquinolone$" or "linezolid$" or "vancomycin$" or "echinocandin$" or "azole$" or "colistin$" or "colimycin$" or "antibacterial" or "bacterial" or "bacterium" or "antibiotic" or "antifungal" or "antimicrobial" or "microbial" or "multidrug" or "multiple drug" or	274,926

| | "multiple antibiotics" or "penicillin" or "meth?cillin$" or "met?cillin$" or "cephalosporin*" or "cephalosporanic acid$" or "cefepim$" or "cefuroxime" or "cefotaxim$" or "ceftazidime" or "ceftriaxon$" or "cefixime" or "ceftaroline" or "carbapenem$" or "aminoglycoside$" or "gentam?cin$" or "amikacin$" or "fluoroquinolone$" or "linezolid$" or "vancomycin$" or "echinocandin$" or "azole$" or "colistin$" or "colimycin$" or "polymyxin e" or "intrinsic*" or "penicillin" or "macrolide" or "tobramycin$" or "fluconazol$" or "voricanozole" or "posaconazole" or "caspofungin" or "anidulafungin" or "micafungin" or "ciproflox*" or "ciproxin" or "levofloxacin" or "clindamycin$" or "dalacin" or "clarithromycin" or "CLAR" or "erythromycin" or "azt$reonam" or "azactam" or "polymyxin e" or "intrinsic*" or "penicillin" or "macrolide" or "tobramycin$" or "fluconazol$" or "voricanozole" or "posaconazole" or "caspofungin" or "anidulafungin" or "micafungin" or "ciproflox*" or "ciproxin" or "levofloxacin" or "clindamycin$" or "dalacin" or "clarithromycin" or "CLAR" or "erythromycin" or "azt$reonam" or "azactam") NEAR/3 ("resistance" or "resistant" or "nonsusceptib*" or "insusceptib*" or "non-susceptib*" or "susceptibility")) | |
| 1 | TS=("tumor$" or "tumour$" or "cancer$" or "neoplas*" or "paraneoplas*" or "preneoplas*" or "precancer*" or "premalign*" or "malign*" or "carcinogenes?s" or "oncogenes?s" or "tumorigenes?s" or "cocarcinogenes?s" or "metastas?s" or "metastatic" or "incidentaloma$" or "leukemia$" or "leucemia$" or "lymphoma$" or "sarcoma$" or "carcinoma*" or "melanoma$" or "glioma$" or "blastoma$") | 4,410,329 |

Supplementary material 3

Table S3 - All included articles in the systematic review with an infection outcome

Year	Title	Authors	Number of patients	Number of events in the final (and largest) model	Country/setting	Patient population	Factors included in the final model	Microbial etiology and resistance	Risk of bias grading (NIH tool)	Bivariable screening/ stepwise regression	Events per variable in the final (and largest) model	Number of variables included in the tested/screened/initial model	Number of variables included in the final (and largest) model	Comments
2019	Recurrent *Clostridium difficile* infection is associated with treatment failure and prolonged illness in cancer patients	Abu-Sbeih, Choi, Tran et al (125)	170	270	USA	Adult cancer patients who had both NAAT and EIA results positive for *C. difficile* were identified in our institutional laboratory database from January 2015 to May 2017	Antibiotic use, Comorbidities, NSAID use, PPI use, Immunosuppressant use, Chemotherapy, Organ transplantation, Gastrointestinal GVHD, Previous hospitalisation Haematological malignancy, Mean length of CDI treatment, CDI treatment failure, Severe/fulminant CDI disease	*Clostridioides difficile*	Low	+	38,6	13	7	Authors concluded that the risk factors for RCDI they found in cancer patients were the same as one can otherwise find in non-cancer patients (previous antibiotic use, NSAID, comorbidities, and immunosuppressant use), except for the risk use of chemotherapeutic therapy.
2020	Alterations of the oral microbiome and cumulative carbapenem exposure are associated with *Stenotrophomonas maltophilia* infection in patients with acute myeloid leukemia receiving	Aitken, Sahasrabhojane, Kontoyiannis et al (81)	90	8	USA	Patients with a new diagnosis of AML who were receiving remission-induction chemotherapy (RIC) between 9/2013 and 8/2015.	Ciprofloxacin, Levofloxacin, Meropenem, *S. maltophilia* oral abundance	*Stenotrophomonas maltophilia*	Low	-	2,0	8	4	Authors found that the abundance of *S. maltophilia* in the oral microbiome, in addition to fluoroquinolone use, is associated with *S. maltophilia* infection.

	chemotherapy													
2016	Results of Four-Year Rectal Vancomycin-Resistant Enterococci Surveillance in a Pediatric Hematology-Oncology Ward: From Colonization to Infection	Akturk, Sutcu, Somer et al (143)	72	7	Turkey	All patients admitted to the paediatric haematology/oncology ward with a documented Vancomycin-resistant enterococci colonisation 48-72 h after admission.	Severe neutropenia (<100/mm3), Previous bacteremia with another pathogen	Vancomycin-resistant enterococci	Medium	+	3,5	10	2	A brief report investigating the risk factors for progression from vancomycin-resistant enterococci colonisation to infection, which they find is associated with severe neutropenia and previous bacteraemia.
2017	A multicenter, retrospective, case-cohort study of the epidemiology and risk factors for *Clostridium difficile* infection among cord blood transplant recipients	Alonso, Braun, Patel et al (65)	226	226	USA	Adult Umbilical cord blood transplant (UCBT recipients at the three Boston centres from January 1, 2003 to December 31, 2012	Age, BMI, Conditioning regimen, Antipseudomonal penicillin, Bacterial infection post- UCBT	*Clostridioides difficile*	Low	+	45,2	21	5	Authors identified having had a bacterial infection post-transplant as a risk factor of CDI, without considering the treatment of the bacterial infection.
2016	Retrospective analysis of the microbiological spectrum of pneumonia in Turkish patients with lung cancer	Avcı, Hartavi, Kaçan et al (39)	119	22	Turkey	Patients with lung cancer and pneumonia between 2010 and 2011	Male sex, Age, Diabetes mellitus, COPD, Histology, Tumour stage, Tumour localisation, Surgical treatment, Pleural effusion, Hospital-acquired pneumonia, Chemotherapy dose modifications, Respiratory failure, Febrile neutropaenia, Development of pneumonia within the first 10 days	*Aspergillus fumigatus*	Medium	-	1,6	14	14	Authors found that febrile neutropenia and pneumonia within the first 10 days of admission may increase the risk of *A. fumigatus* pneumonia.
2020	Use of carbapenems and glycopeptides increases risk for *Clostridioides difficile* infections in acute myeloid leukemia patients undergoing intensive induction chemotherapy	Ballo, Kreisel, Eladly et al (66)	415	37	Germany	All patients with AML who underwent intensive induction chemotherapy between 2007 and 2019.	Cumulative glycopeptide exposure	*Clostridioides difficile*	Low	+	7,4	8	5	Authors found an association between cumulative exposure to some antibiotics and *C. difficile* infection

Year	Title	Authors			Country	Population	Risk factors	Organism	Quality	+/-				Findings
2021	Bacterial bile duct colonization in perihilar cholangiocarcinoma and its clinical significance	Bednarsch, Czigany, Heij et al (18)	95	52	Germany	All surgically treated patients with localised pCCA with available information on bile duct colonisation from intraoperative sampling	Portal vein embolization, Susceptibility to applied perioperative antibiotics	47 species identified, top 5 most common: *Enterococcus faecalis*, *Enterococcus faecium*, *Enterobacter cloacae*, *Escherichia coli*, *Klebsiella pneumoniae*	Medium	+	26,0	26	2	Authors found that portal vein embolization and susceptibility to perioperative antibiotics is associated with postoperative abdominal infections after excluding a number of variables
2016	Epidemiology of *Clostridium difficile* infection in hospitalized oncology patients	Chang, Dembry, Banach et al (126)	229	77	USA	Adult patients admitted to oncology units, those with *C. difficile* infection were cases.	Age, Prior history of *Clostridioides difficile*, Blood transfusion, β-lactam/β-lactamase inhibitor, Cephalosporin, Metronidazole	*Clostridioides difficile*	Medium	+	12,8	32	6	Authors found a very strong association between a history of *C. difficile infection* and current *C. difficile* infection, in addition to a selection of antibiotics.
2017	Risk Factors for Hospital-acquired *Clostridium difficile* Infection Among Pediatric Patients With Cancer	Daida, Yoshihara, Inai et al (127)	145	51	Japan	Paediatric patients with cancer admitted to the hospital between July 2003 and September 2012, CDI cases matched to controls	Age, Male (vs. female), Same ward, Duration of neutropenia	*Clostridioides difficile*	Low	+	10,2	9	5	Antibiotics use is omitted from the final model due to collinearity with duration of neutropenia, which is associated with an increased odds of CDI
2015	Infection with multidrug-resistant gram-negative bacteria in a pediatric oncology intensive care unit: risk factors and outcomes	de Oliveira Costa, Atta, da Silva (116)	76	47	Brazil	Infection episodes related to GNB that occurred between January 1, 2009 and December 31, 2012 in PICU patients, aged between 0 months and 18 years, who were hospitalised for more than 24 h in a tertiary oncology public hospital.	Healthcare-associated infection, Hematologic diseases, Length of neutropenia ≥3 days, Length of ICU stay >3 days, Previous antibiotic therapy	Multidrug-resistant Gram-negative bacteria	Medium	+	9,4	14	5	The authors found that there is a higher odds of a MDR-GNB infection compared to a non-MDR-GNB infection if the infection was healthcare-associated and the patient had a haematological cancer
2018	Increasing rates of *Acinetobacter*	Fan, Wang, Wang et al (136)	6154	60	China	Cancer patients with nosocomial	Gender, Age, Hematologic	*Acinetobacter baumannii*	Medium	-	8,6	7	7	Authors found that there is an association

	baumannii infection and resistance in an oncology department					infections in a general teaching hospital from January 2010 to December 2015	malignancy, Radiation, Hospitalisation time, Operation, Prior antibiotic use							between prior antibiotic use and *A. baumannii* infection
2018	Clinical Outcomes Associated With Linezolid Resistance in Leukemia Patients With Linezolid-Resistant *Staphylococcus epidermidis* Bacteremia	Folan, Marx, Tverdek et al (104)	82	33	USA	All adult leukaemia patients with *S. epidermidis* bacteremia treated with empiric linezolid between 2012 and 2015.	Linezolid-resistant *S. epidermidis*, Age, AML, Prior HSCT, Presence of CVC	*Staphylococcus epidermidis*, linezolid-resistance	Low	+	8,3	12	4	The infection outcome was persistent *S. epidermidis* bacteraemia, which the authors found was more common in patients that were infected by linezolid-resistant bacteria
2019	Decrease in vancomycin-resistant Enterococcus colonization associated with a reduction in carbapenem use as empiric therapy for febrile neutropenia in patients with acute leukemia	Ford, Coombs, Stofer et al (85)	342	79	USA	Patients admitted with newly diagnosed acute myelogenous leukaemia (AML), acute lymphoblastic leukaemia(ALL), biphenotypic leukaemia (BPL), or chronic myelogenous leukaemia in blast phase (CML-BP) between September 2011 and August 2017	Age, Days of empiric antibiotics, LOS, Cycling vs carbapenem period, Days of carbapenem, Days of cefepime or piperacillin/tazobactam	Vancomycin-resistant enterococci	Low	-	13,2	6	6	The infection outcome is VRE colonisation. The authors found that after the hospital changed policies to use less carbapenems, the VRE colonisation rates declined.
2015	Frequency, Risk Factors, and Outcomes of Vancomycin-Resistant Enterococcus Colonization and Infection in Patients with Newly Diagnosed Acute Leukemia: Different Patterns in Patients with Acute Myelogenous and Acute Lymphoblastic Leukemia	Ford, Lopansri, Haydoura et al (86)	214	15	USA	Patients with newly diagnosed acute leukaemia between 2006 and 2012	Severe neutropenia, Number of stools/day	Vancomycin-resistant enterococci	Low	+	7,5	7	2	Data extracted from the model with bloodstream infection as the infection outcome. Authors find that such infections are associated with prolonged neutropenia and diarrhoea.
2018	*Clostridioides difficile* colonization and	Ford, Lopansri, Webb et al (67)	509	31	USA	Patients with AL admitted	Age, Gender female, Diabetes,	*Clostridioides difficile*	Low	-	3,1	10	10	CDI during hospitalisation was

	infection in patients with newly diagnosed acute leukemia: Incidence, risk factors, and patient outcomes					between 2006 and 2017	Hospitalisation prior 2 mo, Antibiotics prior 3 mo, Karnofsky performance Status, Admit serum albumin, Severe neutropenia, Antibiotics (no.), Antibiotics (d)							found in 31 patients - more often in those who had recently been hospitalised or had recently been put on antibiotics
2016	Risk factors for *Clostridium difficile* infection in hemato-oncological patients: A case control study in 144 patients	Fuereder, Koni, Gleiss et al (128)	288	144	Austria	Patients that had a histologically or cytologically confirmed hemato-oncological disease diagnosed between 1st January 2004 and 31st December 2014. Cases were patients that had received chemotherapy during the course of their disease and suffered from CDI positive diarrhoea, controls were those with CDI negative diarrhoea.	Intervall chemotherapy to diarrhoea, Antibiotic therapy within 30 days	*Clostridioides difficile*	Medium	-	72,0	2	2	The authors found that chemotherapy is not associated with CDI positive diarrhoea, but that antimicrobial therapy is.
2017	Consequences of Increases in Antibiotic Resistance Pattern on Outcome of Pancreatic Resection for Cancer	Gianotti, Tamini, Gavazzi et al (19)	517	99	Italy	Adult patients who underwent elective pancreatic resections for cancer from January 1, 2013 to December 31, 2015, with an available sample of biological fluid.	Type of operation (pancreatoduodenectomy vs. others), Operation time, Estimated blood loss, Preoperative biliary stenting (yes vs. no)	Several bacteria, tested for resistance towards several antibiotics	Low	-	24,8	4	4	The authors found that there is an association between preoperative biliary stenting and multidrug or extensively drug resistant infectious complications
2019	Colonization by fecal extended-spectrum b-lactamase-producing *Enterobacteriaceae* and surgical site infections in patients	Golzarri, Silva-Sanchez, Cornejo-Juarez et al (34)	171	25	Mexico	Adult patients, with gynaecological or gastrointestinal malignancies,	ESBL-PE. Adjusted for the following confounders: age, sex, albumin, and type of malignancy.	Extended-spectrum beta-lactamase-producing *Enterobacteria*	Low	+ (with fixed set of confounders)	3,6	13	7	The risk ratio of several infectious outcomes in carriage ESBL-producing Enterobacteriaceae vs non-carriers are

Year	Title	Authors			Location	Population	Risk factors	Organism						Findings
	with cancer undergoing gastrointestinal and gynecologic surgery					who were admitted to the hospital for elective abdominal and pelvic surgical procedures from September 2014 to December 2015		ceae						modelled, after adjusting for confounders. The authors found that the risk of surgical site infection and blood stream infection is increased by such carriage, but not pneumonia and urinary tract infection.
2020	Clinical Predictive Model of Multidrug Resistance in Neutropenic Cancer Patients with Bloodstream Infection Due to *Pseudomonas aeruginosa*	Gudiol, Albasanz-Puig, Laporte-Amargos et al (140)	1217	309	34 centres in 12 countries	Adult neutropenic onco haematological patients diagnosed with at least one episode of *P. aeruginosa* BSI from 1 January 2006 to 31 May 2018.	Prior piperacillin-tazobactam therapy (within 1 mo), Prior fluoroquinolone prophylaxis (within 1 mo), Urinary catheter, Prior antipseudomonal carbapenem therapy (within 1 mo), Haematological disease	*Pseudomonas aeruginosa*, multidrug resistant	Low	-	51,5	32	6	The authors constructed a clinical prediction model to predict resistance in P. aeruginosa bloodstream infections in onco haematological patients, achieving an area under the curve of 0.72 (0.63, 0.80) in internal validation. A risk score calculator of this model is available on https://ubidi.shinyapps.io/ironic/.
2017	Risks factors and outcomes of *Clostridium difficile* infection in patients with cancer: a matched case-control study	Hebbard, Slavin, Reed et al (129)	200	50	Australia	Patients with CDI at a cancer centre from May 2013 to May 2015	Administration of chemotherapy, Gastro-intestinal/abdominal surgery, Proton pump inhibitors, Antibiotic days of therapy	*Clostridioides difficile*	Low	+	12,5	27	4	The authors did not find that the administration of any antibiotics is associated with a *C. difficile* infection, but that chemotherapy is.
2016	Vancomycin- resistant *Enterococcus* colonization and bloodstream infection: prevalence, risk factors, and the impact on early outcomes after allogeneic hematopoietic cell transplantation in patients with acute myeloid leukemia	Hefazi, Damlaj, Alkhateeb et al (88)	203	12	USA	All consecutive patients who underwent their first allogeneic HCT for AML at ta clinic between April 2004 and December 2014	Age (≥60 years vs <60), HCI-CI (≥3 vs 0- 2), KPS (>80 vs ≤80), Conditioning (RIC vs MAC), VRE colonisation (within 30 days of HCT)	Vancomycin-resistant enterococci	Low	+	2,4	10	5	The authors found that there is an association between VRE colonisation and VRE bloodstream infection
2017	Vancomycin-resistant	Heisel, Sutton,	229	134	USA	Adult patients	Gender (male),	Vancomycin-re	Mediu	+	44,7	14	3	The authors found an

Year	Title	Authors			Country	Description	Risk factors	Organism/Resistance						Conclusions
	enterococci in acute myeloid leukemia and myelodysplastic syndrome patients undergoing induction chemotherapy with idarubicin and cytarabine	Mascara et al (89)				who received intensive induction chemotherapy with the standard 7+3 regimen of cytarabine and idarubicin for newly diagnosed AML or MDS from January 2012 to December 2015	Vancomycin IV, Cephalosporin	sistant enterococci	m					association between both cephalosporins and intravenous vancomycin, and a VRE infection
2017	Changes in In Vitro Susceptibility Patterns of Aspergillus to Triazoles and Correlation With Aspergillosis Outcome in a Tertiary Care Cancer Center, 1999–2015	Heo, Tatara, Jimenez-Ortigos a et al (101)	107	19	USA	A case-control study of patients with haematological malignancy comparing azole-resistant strains is nested within a cohort of all Aspergillus species clinical isolates recovered from respiratory sources in patients treated at a cancer centre between January 1999 and December 2015	Asian race, Culture specimen–BAL fluid, Previous azole exposure history	Four *Aspergillus* species (fumigatus, flavus, terreus, niger), azole-resistance	Low	+	6,3	58	3	The authors concluded that there are three independent risk factors of invasive pulmonary aspergillosis caused by Aspergilli with a non-wild type azole MIC - asian race, bronchoalveolar lavage fluid culture and a history of azole exposure. There is also a univariate analysis of mortality in the article.
2017	Daptomycin nonsusceptible vancomycin resistant Enterococcus bloodstream infections in patients with hematological malignancies: risk factors and outcomes	Herc, Kauffman, Marini et al (87)	60	20	USA	Adult patients treated on an inpatient haematology service from January 2011 to December 2015 with a haematological malignancy and a blood culture positive for an enterococcus species.	Congestive heart failure, Typhilitis, Gastrointestinal bleeding, Daptomycin, Cefepime received within 90 days, Cumulative Daptomycin days within 90 days, Cumulative Fluoroquinolone days within 90 days, Prior enterococcal infection, Hospital LOS prior to positive culture,, Stage of malignancy, Number	*Enterococcus*, vancomycin and daptomycin resistance	Mediu m	+	1,4	46	14	The authors found that the only risk factor associated with a daptomycin-resistant VRE infection is daptomycin exposure within 90 days

							induction chemotherapy regimens, Clofarabine within 30 days							
2019	A RESEARCH STUDY ON BACTERAEMIA PRODUCED THROUGH ESCHERICHIA COLI IN TUMOR PATIENTS AT THE SPECIFIC CENTER IN OUR COUNTRY	Khanzada, Zianab, Samreen (149)	229	99	Pakistan	Cancer patients with *E. coli* bacteremia from October 2014 to September 2015	Man sex, Age fewer than 19 years, Haematological distortion, Hospitalization inside 1 month before contagion, ICU admission, Carlson score, Dominant intravenous tube use, Previous chemotherapy inside 1 month, Previous surgery inside 1 month, Preceding fallout inside 1 month, ANC less than 110 cells/mm3, Usage of quinolones inside 95 days of catalogue sample, Usage of third group cephalosporins inside 95 days of index sample, Usage of Piperacilin/Tazobactam within 95 days of index sampling, Usage of carbapenems inside 95 days of index sample	*Escherichia coli*, multidrug-resistant	High	-	7,1	14	14	The authors find that multidrug-resistant *E. coli* bacteraemia was associated with young age, central venous catheter, and piperacillin/tazobactam exposure within 95 days of infection
2018	Clinical predictors of *Stenotrophomonas maltophilia* bacteremia in adult patients with hematologic malignancy	Kim, Cho, Kang et al (82)	236	118	South Korea	Cases were *S. maltophilia* bacteremia in adult patients with hematologic malignancy. Controls were other Gram-negative (*Escherichia coli, Klebsiella* species, *Acinetobacter* species,	Polymicrobial infection, Previous *S. maltophilia* isolation, Breakthrough infection during carbapenem therapy, No. of previous antibiotic use ≥3, Previous TMP/SMX use	*Stenotrophomonas maltophilia*	Medium	+	23,6	18	5	The authors concluded that all independent variables included in the final multivariable model were associated with *S. maltophilia* bacteraemia.

					Pseudomonas species, and Enterobacter species) bacteraemias.								
2019	Risk factors for extended-spectrum beta-lactamase-producing Enterobacteriaceae infection causing septic shock in cancer patients with chemotherapy-induced febrile neutropenia	Kim, Jung, Kang et al (119)	179	32	South Korea	Adult (≥ 18 years) patients with chemotherapy-induced febrile neutropenia admitted to an emergency department with septic shock	Profound neutropenia	Enterobacteriaceae, extended-spectrum beta-lactamase	Low	-	6,4	5	5	After adjustment, the authors found that only profound neutropenia was associated with ESBL-PE infection
2021	Antibiotic Prophylaxis or Granulocyte-Colony Stimulating Factor Support in Multiple Myeloma Patients Undergoing Autologous Stem Cell Transplantation	Klein, Sauer, Klein et al (90)	298	25	Germany	Multiple myeloma inpatients who received high-dose therapy with melphalan followed by autologous stem cell transplantation at a hospital between March 2016 and July 201	Antibiotic prophylaxis (vs. G-CSF support), No prophylaxis (vs. G-CSF support), Age (per ten years), ASCT at relapse (vs. first-line treatment), ≥VGPR before ASCT (vs. ≤PR), Stem cell amount ≥2.5 * (vs. <2.5)	Enterococcus faecium, vancomycin-resistant	Medium	-	4,2	6	6	The article includes analyses of several microbes, but the assessed model investigates the association between the listed risk factors and VRE detection. Authors concluded that Granulocyte-Colony Stimulating Factor instead of antibiotic prophylaxis is associated with a fewer detected VRE cases.
2015	Clinical features and risk factors for development of Bacillus bacteremia among adult patients with cancer: a case-control study	Ko, Kang, Lee et al (160)	258	86	South Korea	Cases were adult patients with cancer who were diagnosed with Bacillus bacteremia during the period of January 1995 through December 2012. Controls were patients with bacteraemia due to other pathogens.	Hospital stay of longer than 14 days, History of HSCT, Presence of a CVC, Hematologic malignancy, Diarrhoea at presentation, Prior use of extended-spectrum cephalosporin, Prior use of glycopeptides	*Bacillus spp.*	Low	-	9,6	9	9	The authors found that in addition to a central venous catheter, the use of cephalosporins within a month of admission was associated with Bacillus bacteraemia
2016	Risk factors for the development of	Lavallee, Labbe, Talbot et al (68)	188	65	Canada	Adults allo−HSCT	Mucositis, CMV reactivation,	*Clostridioides difficile*	Low	+	13,0	30	5	The authors concluded that

Year	Title	Authors			Country	Population	Variables	Pathogen	Quality	+/-				Conclusion
	Clostridium difficile infection in adult allogeneic hematopoietic stem cell transplant recipients: A single-center study in Québec, Canada					recipients. Cases were CDI positive, controls were CDI negative.	Herpesviridae reactivation other than CMV, Antibiotic use within 30 days before transplantation (No antibiotic, Any antibiotic except TMP-SMX, TMP-SMX)							reactivation of CMV and other Herpesviridae was associated with CDI. There is also a multivariable model stratified on the time of CDI positivity, which shows that high-risk antibiotics were associated with CDI late in the treatment.
2015	Risk factors for acquisition of multidrug-resistant bacteria in patients with anastomotic leakage after colorectal cancer surgery	Lee, Ryu, Chung et al (20)	143	46	South Korea	Patients with colorectal cancer that underwent surgery and had anastomotic leakage and a subsequent positive blood or intra abdominal fluid culture from January 2009 to April 2013	Diabetes mellitus, The total length of antibiotic administration for more than 5 days before diagnosis of anastomosis site leakage. Adjustment variables indeterminable.	Several bacteria and fungi, tested for resistance towards several antimicrobials	Medium	+	Indeterminable	Indeterminable	Indeterminable	The authors found that compared to non-MDR pathogens, diabetes mellitus and the total length of antibiotic administration for more than 5 days before diagnosis of anastomosis site leakage were associated with acquisition of a MDR pathogen.
2017	Post-operative MRSA infections in head and neck surgery	Lin, Melki, Lisgaris et al (38)	113	24	USA	Patients who were >18 years of age and underwent major head and/or neck surgery between January 1, 2008 and January 1, 2014, and were found to have a documented surgical site infection during their postoperative hospitalisation	Indeterminable	*Staphylococcus aureus*, methicillin-resistance	Medium	-	Indeterminable	Indeterminable	Indeterminable	The authors describe a multivariable model in the methods chapter, but we were unable to locate a regression table. Nonetheless, the authors conclude that hospital exposure prior to the procedure, a history of chemotherapy, or immunosuppression were risk factors for developing a MRSA SSI.
2015	Risk for *Clostridium difficile* infection after radical cystectomy for bladder cancer: Analysis of a contemporary series	Liu, Shatagopam, Monn et al (35)	552	49	USA	Patients undergoing radical cystectomy between January 2010 and December 2013 for bladder cancer	Diversion type (Ileal conduit, Indiana pouch, Orthotopic neobladder), Preoperative haemoglobin, Preoperative antacid use, Days of antibiotics received (1 day or less, 2–7	*Clostridioides difficile*	Low	+	8,2	22	6	The authors found that chronic antacid therapy and long duration of antibiotic exposure in the perioperative setting were associated with a *C. difficile* infection in these patients

							days, Greater than 7 days)							
2018	Mixed mold pulmonary infections in haematological cancer patients in a tertiary care cancer centre	Magira, Jiang, Economides et al (102)	54	27	USA	Cases were adult (>18 years old) patients with haematological malignancy who had proven or probable pulmonary infection with >1 mould grown concurrently in sputum or bronchoalveolar lavage (BAL) cultures, controls were patients with haematological cancer (from the same time period) with IPA caused by Aspergillus fumigatus.	Significant use of corticosteroids, culture from sputum specimens	Mixed moulds, *Aspergillus fumigatus*	Medium	+	Indeterminable	29	Indeterminable	We were unable to locate a printed regression table, but the authors concluded that there is an association between use of corticosteroids and sputum samples, and mixed moulds instead of *A. fumigatus* infection
2015	Risk factors for piperacillin/tazobactam-resistant Gram-negative infection in hematology/oncology patients with febrile neutropenia	Marini, Hough, Gregg et al (117)	171	25	USA	Adult haematology/oncology patients with a diagnosis of FN and any culture positive for a Gram-negative rod from November 2007 to November 2013	Demographic model: Unit ICU, Source of positive culture (Respiratory). Resistance model: History of PTZ-R, Antibiotic use previous 90 days (Antibiotic therapy >14 days), Specific antibiotics previous 90 days (PTZ, Vancomycin), LOS >14 days prior to culture, Healthcare exposures previous 90 days (ICU admission), Transfer from outside hospital, Urinary catheter at culture. Haematology/oncology model: Stage of disease (Hematologic malignancy–newly	Gram-negative bacilli, Piperacillin/tazobactam resistance	Medium	+ (collinear variables removed)	Indeterminable	69	Indeterminable	There are three models - one with demographic variables, one with resistance variables, and one with haematology/oncology-specific variables. A final model with variables from all these models are described and the area under the curve of this model is reported, but a regression table is not printed. The authors concluded that prolonged antibiotic use and critical illness increase the risk of piperacillin/tazobactam resistance.

Year	Title	Author			Country	Population		Organism							Comments
						diagnosed, Hematologic malignancy–continuation/consolidation), Oncology treatment previous 90 days (Rituximab, Clofarabine).									
2020	Development and validation of a scoring system for predicting cancer patients at risk of extended-spectrum b-lactamase-producing Enterobacteriaceae infections	Martínez-Valencia, Martínez, Ayala et al (124)	710	265	Colombia	Cancer patients with documented microbiological isolation during hospitalisation at a national cancer institute	Prolonged hospitalisation (≥7 days), Hospitalisation during previous year, Immunosuppressive therapy, Neutropenia, Beta-lactams during the previous month, Invasive devices at the time of culture, Neoplasia in remission, No chemotherapy	*Enterobacteriaceae*, ESBL	Low	+	33,1	39	8	The authors did develop a clinical prediction model, which is validated in a separate cohort and assessed with an area under the curve. The variable selection in the prediction model is based on the statistical significance of the candidate predictors in a stepwise regression. The authors translated the model into a scoring system, with prolonged hospitalisation and previous hospitalisation the last year as the highest scores.
2019	Oral Vancomycin Prophylaxis as Secondary Prevention Against *Clostridium difficile* Infection in the Hematopoietic Stem CellTransplantation and Hematologic Malignancy Population	Morrisette, Van Matre, Miller et al (69)	50	11	USA	Patients aged >18 years with a history of autologous or allogeneic HSCT and patients with HM who did not undergo HSCT, who have been treated for the initial episode of CDI first with planned oral vancomycin monotherapy and must have been receiving a broad-spectrum antibiotic at the time of CDI diagnosis and/or	Oral vancomycin prophylaxis, Outpatient CDI diagnosis.	*Clostridioides difficile*	Low	+	5,5	Indeterminable	2	The infection outcome here is recurrent CDI infection, which the authors found was associated with no oral vancomycin prophylaxis and an inpatient CDI diagnosis.

Year	Title	Authors			Country	Population	Covariates	Pathogen							Conclusion
						during the course of CDI treatment/prophylaxis.									
2019	Antimicrobial de-escalation in adult hematopoietic cell transplantation recipients with febrile neutropenia of unknown origin	Petteys, Kachur, Pillinger et al (70)	107	3	USA	Adult HCT recipients with FN at an academic teaching hospital between March 2014 and April 2018.	Early de-escalation vs hematopoietic recovery de-escalation, Gender, Race, Age	Clostridioides difficile	Medium	-	0,8	4	4		The authors did not find that the early de-escalation of broad-spectrum antibiotics was associated with a Clostridioides difficile infection
2017	Moxifloxacin versus levofloxacin or ciprofloxacin prophylaxisin acute myeloid leukemia patients receiving chemotherapy	Przybylski & Reeves (71)	141	141	USA	Adult patients admitted to the hospital with a diagnosis of AML who received induction or reinduction chemotherapy from January 2008 to June 2015	Prophylactic antibiotherapy received, treatment antibiotic duration, mucositis, proton pump inhibitor, neutropenic days	Clostridioides difficile	Medium	-	28,2	Indeterminable	5		Authors found that treatment antibiotic duration was the only statistically significant association with with C. difficile infection
2019	Risk, Outcomes, and Trends of Clostridium Difficile Infection in Multiple Myeloma Patients from a Nationwide Analysis	Ran-Castillo, Oluwole, Abuaisha et al (72)	114249	3549	USA	Patients with a diagnosis of MM between 2010 and September 2015	Year (2010, 2011, 2012, 2013, 2014, 2015), Age categories (18-49, 50-59, 60-69, 70-79, ≥ 80), Race/Ethnicity (White, Black, Hispanic, Others), Female, Obesity, Peripheral vascular disease, Chronic renal disease, Atrial fibrillation, Stem cell transplantation, Neutropenia, Chemotherapy, Inflammatory bowel disease, Congestive heart failure, Hypertension, Chronic liver failure, Elixhauser score (0, 1-3, ≥ 4), Hospital bed size (Small, Medium, Large), Hospital teaching status (Rural, Urban non-teaching, Urban	Clostridioides difficile	Medium	-	177,5	20	20		The authors balanced multiple myeloma patients with non-multiple myeloma using propensity scores, then modelled the risk of CDI and of mortality given CDI. The unadjusted OR for mortality is also reported. They found several risk factors for CDI, e.g. neutropenia.

						teaching), Expected primary payer (Medicare, Medicaid, Private, Others), Median household income in quartile (1st, 2nd, 3rd, 4th).								
2021	Risk of Drug Resistance and Repeated Infection with *Klebsiella pneumoniae* and *Escherichia coli* in Intensive Care Unit Cancer Patients	Refay, Ahmed, ELzaher et al (120)	107	38	Egypt	Cancer patients who were treated at either the Paediatric Oncology or the Medical Oncology ICU with an initial infection or repeated infection with either *K. pneumoniae* or *E. coli* from February 2017 to May 2018	Age at diagnosis, Type of Malignancy, Type of Organism, Total prior Hospital Stay, Total Prior Admission in ICU, Prior Mechanical Ventilation, Prior Urinary Catheterization, Peripheral or Central CVP Line, Infection	*Klebsiella pneumoniae* and *Escherichia coli*, tested for resistance towards several antibiotics	Low	-	4,8	9	8	The infection outcome is drug resistance in the previous infection. A ROC curve is printed to show predictive accuracy. The authors conclude that restricting unnecessary antibiotic prescriptions is important to prevent the emergence of multidrug-resistance in these infections
2015	Risk factors associated with *Clostridium difficile* infection in adult oncology patients	Rodriguez Garzotto, Merida Garcia, Munoz Unceta et al (36)	225	39	Spain	Hospitalised cancer patient with diarrhoea between January 2009 through April 2013	Gastrointestinal cancer, Breast cancer, Lung cancer, Age (>65), Antibiotic exposure, Corticosteroid use, PP inhibitor use	*Clostridioides difficile*	Medium	-	9,8	4	4	The authors stratified three tables based on cancer types and include 4 previously identified risk factors. They concluded that antibiotic exposure was associated with CDI infection
2016	Bacteremia due to carbapenem-resistant Enterobacteriaceae in neutropenic patients with hematologic malignancies	Satlin, Cohen, Ma et al (60)	301	43	USA	Cases were adult neutropenic patients with haematological malignancies with carbapenem-resistant Enterobacteriaceae bacteraemias. Primary control group was patients with bacteraemias caused by pathogens other than carbapenem-resistant	b-lactam/b-lactamase inhibitor within previous 30 days, Carbapenem within previous 30 days, Receiving TMP-SMX at BSI onset, Receiving glucocorticoids at BSI onset, Prior CRE at any site within the previous 90 days, ICU stay within previous 30 days, Receiving cephalosporin at BSI onset, Receiving fluoroquinolone at BSI onset, Renal	*Enterobacteriaceae*, carbapenem-resistance	Medium	+	4,8	81	9	The infection outcome here was bacteraemia. The authors find that previous CRE infection and antibiotic exposure was associated with CRE bacteraemia

Year	Title	Authors	N	N2	Country	Population	Model/Variables	Bacteria	Quality	+/-				Findings
						Gram-negative bacteria, and the secondary control group was patients with carbapenem-susceptible Gram-negative bacteria.	disease							
2017	Risk factors and impact of *Clostridium difficile* recurrence on haematology patients	Scappaticci, Perissinotti, Nagel et al (73)	100	41	USA	Adult (age >18 years) patients with an active haematological malignancy and CDI from June 2010 to December 2014	Demographic and traditional model: Severe CDI, salvage lymphoma. Longitudinal model: Hospitalised (days), Antibiotic exposure (ceftriaxone, fluoroquinolones, piperacillin/tazobactam, other (daptomycin), other (vancomycin IV), number of antibiotic exposures), Haematological risk factors (neutropenia (days), chemotherapy exposure, decitabine, CHOP, salvage lymphoma)	*Clostridioides difficile*	Medium	+	20,5	40	2	The infection outcome is recurring CDI, and there is one model containing demographical and traditional risk factors, and one containing longitudinal risk factors. Data are extracted from the "traditional" model. The authors found that exposure to a number of antibiotics were associated with recurrent CDI.
2019	*Clostridium difficile* infection in cancer patients with hospital acquired diarrhea at the teaching hospitals in Iran: Multilocus sequence typing analysis (MLST) and Antimicrobial resistance pattern	Shoaei, Shojaei, Khorvash et al (130)	67	19	Iran	Adult (>18 years old) cancer patients with presence of hospital-acquired diarrhea at major teaching hospitals between April 2015 and May 2017	Chronic renal diseases, Gastroenteritis diseases, Previous surgery	*Clostridioides difficile*	Low	+	6,3	14	3	The authors found that patients with gastrointestinal diseases had an association with lower CDI incidence
2015	Development of a combination antibiogram for *Pseudomonas aeruginosa* bacteremia in an oncology population	Smith, Tajchman, Dee et al (141)	123	36	USA	*Pseudomonas aeruginosa* isolates from an oncology population at least 18 years of age who were hospitalised between 1	Prophylactic antibiotic (None, Fluoroquinolone, Cefpodoxime, Amoxicillin-clavulanate), LOS prior to culture (<5 days, >5 days), IV antibiotics in previous 90 day,	*Pseudomonas aeruginosa*, multidrug resistant	Low	+	9,0	5	4	The authors did develop an antibiogram for multidrug-resistant P. aeruginosa. In multivariable analysis they found that such bacteraemias were associated with

					January 2012 and 31 December 2012	Hemodialysis in previous 30 day								receiving IV antibiotics 90 days prior
2019	Significance of bile culture surveillance for postoperative management of pancreatoduodenectomy	Sugimachi, Iguchi, Mano et al (32)	69	43	Japan	Patients who underwent pancreatoduodenectomy at a single institute between the years 2014 and 2017	Smoking history (+ vs -), Pancreatic texture (soft vs hard)	*Enterobacter spp., Staphylococcus aureus, Pseudomonas spp.,* multidrug-resistance	Medium	+	21,5	8	2	The authors do not find that having a multidrug-resistant bacteria in the preoperative bile culture was associated with surgical site infection following a pancreatoduodenectomy, and as such did not include it in the multivariable model
2019	Risk factors for recurrent percutaneous nephrostomy catheter-related infections	Szvalb, El Haddad, Rolston et al (21)	81	10	USA	Patients who had undergone an initial percutaneous nephrostomy catheter placement at the hospital between July 2014 and February 2017	Concurrent antibiotics use for PCN infection, PCN exchange within 4 days of infection	Several bacteria	Medium	+	5,0	10	2	The infection outcome here is recurrent catheter-related infection. The authors did not find any association between intrinsically resistant pathogens and recurrent infections
2019	Oral Candida colonization in xerostomic postradiotherapy head and neck cancer patients	Tarapan, Matangkasombut, Trachootham et al (40)	72	72	Thailand	Xerostomic post radiotherapy head and neck cancer patients, aged from 30 to 70 years old, who finished radiotherapy for at least 1 month and/or chemotherapy for at least 2 weeks, from 3 Thai institutes	Sex, denture, antifungals, objective dry mouth scores, NACS (?)	*C. albicans, Candida spp.*	Medium	+	14,4	8	5	Authors found that increased signs of dry mouth, female sex and dental prostheses may promote NACS colonization.
2021	*Clostridioides difficile* Infections in Inpatient Pediatric Oncology Patients: A Cohort Study Evaluating Risk Factors and Associated Outcomes	Willis, Huang, Elward et al (131)	952	109	USA	Paediatric patient with malignancies admitted at St Louis hospital between July 2009 and February 2018	Model 1 Risk for CDI: Length of stay before CDI, Leukaemia, Lymphoma, Autologous transplant, Allogeneic transplant, CDI exposure, Antibiotic	*Clostridioides difficile*	Low	+	15,6	16	7	Leukaemia, Lymphoma, autologous transplant, antibiotic exposure, CDI exposure were associated with CDI. The authors also showed that LOS and chemotherapy delay were outcomes linked

						exposure prior to CDI ; Model 2 outcomes linked to CDI : LOS, Chemotherapy delay OR and mean difference.						to CDI.
2019	Intensity of Therapy for Malignancy and Risk for Recurrent and Complicated *Clostridium difficile* Infection in Children	Willis, Nicholson, Esbenshade et al (132)	122	122	USA	Children >12 months and <18 years of age undergoing treatment for cancer and diagnosed with CDI between August 1st, 2008 and May 1st, 2015	Level 2 of patient's Intensity of Treatment Rating Scale, level 3 with HCST, level 3 without HCST and the HCST status	*Clostridioides difficile*	Low	-	30,5	Indeterminable	4	The authors concluded that they did not manage to find which paediatric oncology patients were at increased risk of CDI recurrence.
2015	Monitoring antibiotic-resistant enterobacteria faecal levels is helpful in predicting antibiotic susceptibility of bacteraemia isolates in patients with haematological malignancies	Woerther, Micol, Angebault et al (61)	104	16	France	Patient with neutropenia, haematological malignancies and EB-BSI from a probable intestinal origin	Death during neutropenia, non beta lactam antibiotic previous exposure, Carbapenem previous exposure, empirical treatment of EB-BSI with carbapenems, empirical treatment with antibiotics combination, presence of BSBL-Resistant EnteroBacteria, ESBL-EB, and non E. coli EB in blood cultures	Enterobacteria, BSBL or ESBL	Low	+	2,0	20	8	The authors showed that to have a BSBL-RE carriage increased risk to have BSBL-RE in blood cultures.
2020	Clinical Characteristics and Risk Factors for Bloodstream Infection Due to Carbapenem-Resistant *Klebsiella pneumoniae* (CRKP) in Patients with Hematologic Malignancies	Zhang, Wang, Hu et al (95)	734	22	China	Patients with haematological malignancies, hospitalised for more than 7 days between January 2017 and January 2018	CRKP rectal colonisation, Severe neutropenia,IMV.	*Klebsiella pneumonia*, carbapenem-resistance	Medium	+ (+ clinically relevant variables)	7,3	42	3	Variables with p<0.2 in the bivariable screening were included in the multivariable model, but also variables that were "clinically significant", which were not disclosed. The authors concluded that rectal colonisation with carbapenem-resistant *K. pneumoniae* increases the risk of

														such bloodstream infections.
2017	*Clostridium difficile* colonization in preoperative colorectal cancer patients	Zheng, Luo, Lv et al (37)	205	33	China	Consecutive preoperative CRC patients who were scheduled to accept radical cancer resection in a hospital in China, October, 2014 and August, 2015.	Age (<60 vs >60), Site (Colon vs Rectum), Morphology (Ulcerative vs Exophytic), Differentiation (Poor vs Well), T stage (Non-T4 vs T4), N stage (LNM neg. Vs LN pos.), Albumin (<35 vs >35), FBG (>7 vs <7), BMI, Triglyceride, Total cholesterol, HDL, Haemoglobin, OB	*Clostridioides difficile*	Medium	-	2,4	14	14	The authors concluded that colorectal cancer patients in China with large tumours or metastases were associated with a higher risk of *C. difficile* infections.
2020	Distribution, diagnosis, and analysis of related risk factors of multidrug-resistant organism in patients with malignant neoplasms	Zhou, Yang, Yang et al (22)	278	128	China	Patients aged >18 years old and <75 years old with malignant tumour, positive bacterial culture during hospitalisation at Zhuji Hospital from January 2017 to December 2019	Length of hospitalisation, Fever days, Invasive operation, Excessive bed rest, Hypoproteinemia, Procalcitonin, C Reactive Protein, Serum Amyloid	MDRO, non MDRO	Medium	+	16,0	13	8	Variables with p<0.001 in the bivariable screening were included in the multivariable model, except for excessive bedstay (p=0.002). The authors found that invasive operation, excessive bedstay, hypoproteinemia, PCT and SA were associated with MDRO infection in cancer patients. Measurements of such variables should help to prevent MDRO infection.

Supplementary material 4

Table S4 - All included articles in the systematic review with a mortality outcome

Year	Title	Authors	Number of patients	Number of events in the final (and largest) model	Country/setting	Patient population	Factors included in the final model	Microbial etiology and resistance	Risk of bias grading (NIH tool)	Bivariable screening/ stepwise regression	Events per variable in the final (and largest) model	Number of variables included in the tested/screened/initial model	Number of variables included in the final (and largest) model	Comments
2016	Risk factors and clinical outcomes for vancomycin-resistant enterococcus bacteraemia in hospitalised cancer patients in Pakistan: A case-control study	Akhtar, Sultan, Nizamuddin et al. (144)	111	57	Pakistan	Cancer patients whose blood culture grew either vancomycin-sensitive or resistant enterococcus (VSE or VRE) from January 2012 to December 2014	Paediatric (Compared to adults), sex, type of malignancies, location in hospital at onset of bacteraemia, inpatient, others(OPD/ ER), APACHE-II score at onset of bacteraemia, length of hospital stay in days before onset of bacteraemia, shock at onset of bacteraemia), disseminated Intravascular Coagulation (DIC) score, received vancomycin within 4 weeks prior to bacteraemia	*Enterococcus spp.*, vancomycin-resistance	Medium	No	5.2	11	11	The only risk factor that seems to be of importance for the mortality of patients with VRE vs VSE bacteraemia is shock at onset of bacteraemia.
2019	*Clostridium difficile* Infections in patients with AML or MDSundergoing allogeneic hematopoietic stem cell transplantationidentify high risk for adverse outcome	Amberge, Kramer, Schröttner et al (74)	727	Indeterminable	Germany	All patients with acute myeloid leukaemia or myelodysplastic syndrome who underwent allo hæmotological cell transplant at the institution between January 1st 2004 and March 31st 2015.	*Clostridium difficile* status (no infection/asymptomatic/symptomatic), donor type (related/unrelated), human Leukocyte Antigen compatibility (match/mismatch), disease risk (favourable/intermediate/high), age, meropenem exposition (yes/no)	*Clostridioides difficile*	Low	No	#VALUE!	6	6	The study indicated symptomatic *C. difficile* infection increases the risk of death

2015	Mortality burden related to infection with carbapenem-resistant Gram-negative bacteria among haematological cancer patients: a retrospective cohort study	Andria, Henig, Kotler et al (54)	330	95	Israel	Adult patients with Gram-negative aerobic bacteraemia hospitalised in the haemato-oncological/bone marrow transplantation departments actively treated for their malignancy between 2008-2014	Dependent functional capacity at baseline, pulmonary disease, liver disease, hæmotologiv stem cell transplant comorbidities index, renal failure, admission-days before bacteraemia, type of the last chemotherapy treatment, functional capacity at bacteraemia onset, urinary catheter, nasogastric tube, type of bacteria, polymicrobial bacteraemia, appropriate empirical antibiotic treatment, CRGNB, systolic blood pressure , diastolic blood pressure, septic shock	Carbapenem-resistant Gram-negative bacteria	Medium	Yes	5.6	28	9	The authors found that a bacteraemia with a Carbapenem-resistant Gram-negative bacteria increases the risk of death, but the propensity score for resistance did not.
2019	Current etiology, clinical features and outcomes of bacteremia in olderpatients with solid tumors	Antonio, Gudiol, Royo-Cebrecos et al (23)	217	77	Spain	All consecutive patients with solid tumours hospitalised with one or more episodes of bacteremia from January 2006 to November 2015.	Age years, lung cancer, prostate cancer, advance neoplasm, comorbidities, current corticosteroid therapy, bacteremia origin (healthcare/nosocomial), clinical manifestations (Low grade fever (<38 °C/Shock at presentation), source of bacteraemia (Urinary tract), inadequate empirical antibiotic therapy	Multidrug-resistant organisms	Low	Yes	6.4	23	8	There is both a model for early case-fatality (at 7 days) and 30-day mortality. The risk factors in this table is from the final 30-day mortality model, in which the author did not include resistant pathogens due to automatic variable selection.
2019	Colonization with multidrug resistant organisms determines the clinical course of patients with acute myeloid leukemia undergoing intensive induction chemotherapy	Ballo, Tarazzit, Stratmann et al (41)	220	104	Germany	All patients with acute myeloid leukaemia who underwent intensive induction chemotherapy between 2007 and 2015.	Age >60, adverse genetic group acute myeloid leukaemia, day 15 bone marrow blast clearance, stem cell transplantation as consolidation therapy, colonisation with carbapenem resistant enterococci	Multidrug-resistant organisms	Medium	Yes	20.8	9	6	Finds an increased risk of dying for AML patients colonised with carbapenem-resistant *enterobacteriaceae*
2015	Emergence of multidrug resistant isolates and mortality predictors in patients with solid tumors or hematological	Bastug, Kayaaslan, Kazancioglu et al (106)	205	48	Turkey	Patients admitted to hemato-oncology wards of the 1,140-bed tertiary care	Age > 65 years, prior hæmatologic stem cell transplant, prolonged neutropenia prior to infection, inadequate initial antibiotic treatment,	Multidrug-resistant organisms	Medium	Yes	4.8	10	10	Finds an increased risk of dying among patients with predominantly haematological

Year	Title	Authors			Country	Population	Variables		Quality					Conclusions
	malignancies					hospital between 2008 and 2013.	gram-negative bacterial infection, infection with coagulase negative Staphylococci, infection with *Acinetobacter baumannii,* infection with *Klebsiella spp.*, infection with extended spectrum beta lactamase-producing bacteria, infection with Gram-negative multi drug resistant bacteria							malignancies when infected with either *A. baumannii* or *Klebsiella spp.*
2018	The relationship between mortality and microbiological parameters in febrile neutropenic patients with hematological malignancies	Çalık, Arı Bilgir et al (53)	106	58	Turkey	Patients diagnosed with hematologic malignancies	Absolute neutrophil count, duration of neutropenia (days), catheterization focus of infection, central catheterization, isolation of Gram + bacteria in culture, isolation of Gram - bacteria in culture, extended spectrum beta lactamase *Enterobacteriaceae*, carbapenem resistance, imipenem empirical antibiotic regimen, bacterial growth during antibiotic treatment, administration of appropriate antibiotic treatment for minimum 72 hours, septic shock	Several bacterias, tested for sensitivity to all major classes of antibiotics.	Medium	Yes	4.8	18	12	Does not find an association between the resistant bacteria and mortality.
2016	Emerging resistant bacteria strains in bloodstream infections of acute leukaemia patients: results of a prospective study by the Rete Ematologica Lombarda (Rel)	Cattaneo, Zappasodi, Mancini et al (42)	433	37	Italy	All febrile/infectious episodes were recorded in patients with acute leukaemia from December 2012 to December 2014	Complete remission, relapsed/refractory disease, central venous catheter-related bloodstream infection (BSI), multi-resistant Gram-negative BSI, pneumonia	Several bacteria and fungi, tested for various antibiotics	Medium	Yes	7.4	20	4	Authors found that multi-resistant Gram-negative bloodstream infections are associated with higher mortality
2021	A practical update on the epidemiology and risk factors for the emergence and mortality of bloodstream infections from real-world data of 3014 hematological	Chen, Lin, Li et al (43)	725	171	China	Patients with haematological malignancies, between 2013 and 2016	Age>45, Hospital length of stay, Duration of neutropenia before blood culture, Hemograms of bloodstream infections (white blood cell, Haemoglobin, platelet), Disease status, Use of antibiotics (≥3 agents),	Several bacteria and fungi, some intrinsically resistant	Medium	Yes	6.3	27	27	The authors found that the intrinsically resistant bacteria *S. maltophilia* and *A. baumannii* is associated with an increased hazard for

Year	Title	Authors	N	n	Country	Population	Variables	Organism						Conclusion
	malignancy patients receiving chemotherapy						Co-infections (respiratory), Numbers of co-infected locations, Pathogens for bloodstream infections (among them 11 bacterias, fungi, polymicrobial and other), gram negative organisms, gram positive organisms, polyorganisms, and fungi							mortality
2018	Association Between Positive Cultures During Admission and 1-Year Mortality in Patients With Cancer Receiving Perioperative Intensive Care	Chiang, Wu, Hsu et al (107)	638	134	Taiwan	All adult patients who were registered in the hospital cancer registry and had at least one intensive care unit admission to an from 2011 through 2016 following their cancer diagnosis	Age, male gender, type II diabetes mellitus, cerebrovascular disease, congestive heart failure, metastatic cancer, emergent surgery, APACHE II, mechanical ventilation, positive culture of multi drug resistant organism, positive culture in blood/respiratory tract/urinary tract/skin and soft tissue	Several bacteria and fungi are studied, including MDROs (methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant Enterococci, and carbapenem-resistant Gram-negative bacilli)	Medium	No	12.2	11	11	After adjusting for other factors, the authors did not find that MDROs were associated with a higher hazard of death.
2015	*Stenotrophomonas maltophilia* bloodstream infection in patients with hematologic malignancies: a retrospective study and in vitro activities of antimicrobial combinations	Cho, Lee, Choi et al (83)	31	20	South Korea	All consecutive episodes of *S. maltophilia* bloodstream infections in adult patients with hematologic malignancies from June 2009 to May 2014.	Neutropenia at the onset of bloodstream infection, severe neutropenia at the onset of bloodstream infection (<100/mm3), duration of neutropenia >21 days, hospital stay >30 days, SAPS II score >40, Source of infection (Pneumonia), polymicrobial bloodtream infection, shock	*Stenotrophomonas maltophilia*	Medium	Yes	6.7	9	3	The authors found that shock, neutropenia and combined *S. maltophilia* pneumonia are associated with mortality in these patients.
2020	High mortality in an outbreak of multidrug resistant *Acinetobacter baumannii* infection introduced to an oncological hospital by a patient transferred from a general hospital	Cornejo-Juarez, Cevallos, Castro-Jaimes et al (137)	106	53	Mexico	Patients with multi drug resistant *A. baumannii* in a tertiary care oncology hospital from January 2011 to December 2015	Age <60, colonised/Infected, solid tumour/Hematologic malignancy, recent diagnosis (vs relapse), no recent chemotherapy (vs recent), appropriate treatment/Non-appropriate treatment, SOFA score <10/>10	Multidrug-resistant *Acinetobacter baumannii*	Low	Yes	7.6	7	7	In this outbreak analysis, the authors found that haematological cancer patients had a higher risk of death than solid tumour patients

2019	Fungaemia in haematological malignancies: SEIFEM-2015 survey	Criscuolo, Marchesi, Candoni et al (59)	215	82	Italy	Hospitalised patients with haematological malignancies in which a documented fungaemia was diagnosed between January 2011 and December 2015	Performance status >2, chronic kidney disease, steroid administration, gastrointestinal symptoms, respiratory symptoms, septic shock, multiorgan failure, concurrent bacterial sepsis, central venous catheter removal, neutrophil recovery, albicans vs non albicans	Several fungi, including intrinsically resistant	Low	Yes	9.1	9	7	In a univariate analysis, authors founnd that non-albicans candidaemias had a higher odds of death, but the factor is not retained in the final model
2020	Biofilm Production by Carbapenem-Resistant *Klebsiella pneumoniae* Significantly Increases the Risk of Death in Oncological Patients	Di Domenico, Cavallo, Sivori et al (145)	53	19	Italy	Oncological patients colonised or infected with carbapenem resistant *K. pneumoniae*	Biofilm (strong vs weak), colistin resistance, fungal infection, phenotype, site (respiratory vs other)	Carbapenem-resistant *Klebsiella pneumoniae*	Medium	No	3.8	5	5	The authors did find that the production of biofilm in carbapenem-resistant *K. pneumoniae* is associated with a higher hazard of death
2021	The Impact of Bacterial Biofilms on End-Organ Disease and Mortality in Patients with Hematologic Malignancies Developing a Bloodstream Infection	Di Domenico, Marchesi, Cavallo et al (44)	96	8	Italy	All consecutive adult patients (aged >18 years) affected by haematological malignancies and experiencing a bloodstream infection as detected from April 2016 through April 2019	Antimicrobial prophylaxis, multi drug resistant (MDR) vs non-MDR bloodstream infection (BSI), initial antimicrobial failure, P. aeruginosa vs other BSI, strong vs weak biofilm-producer, septic shock, end organ disease	Several bacteria, including intrinsically resistant tested for resistance towards several antibiotics	Medium	Yes	1.1	16	6	The authors found that a multidrug-resistant bloodstream infection is associated with a higher hazard of death in these patients
2021	Clinical characteristics and outcome of 125 polymicrobial bloodstream infections in hematological patients: an 11-year epidemiologic survey	Facchin, Candoni, Lazzarotto et al (45)	125	19	Italy	Patients affected by haematological neoplasms with a documented polymicrobial bloodstream infection	Age, infection related mortality, septic shock rate, multi drug resistant-polymicrobial bloodstream infections mortality	Several bacteria tested for resistance towards several antibiotics	Medium	Yes	4.8	13	4	The authors found that in these patients, having a multidrug-resistant bacteria as part of a polymicrobial bloodstream infection is associated with a higher odds of death

2015	Early initiation of appropriate treatment is associated with increased survival in cancer patients with *Candida glabrata* fungaemia: a potential benefit from infectious disease consultation	Farmakiotis, Kyvernitakis, Tarr et al (153)	146	58	USA	Patients with at least one blood culture(s) positive for C. Glabrata at a cancer centre between March 2005 and September 2013	Intensive care unit stay, monocytopenia, APACHE II score, mechanical ventilation, septic shock, acute kidney infection or acute renal failure, fever, corticosteroids, catheter-related candidemia, mixed bloodstream infection, caspofungin resistance, echinocandin pre-exposure, central venous catheter removal within 48 h after blood culture collection, appropriate treatment within 48 h after blood culture collection	*Candida glabrata*	Medium	Yes	4.1	18	7	The authors concluded that patients with a blood culture positive for *C. glabrata* should consult an ID specialist within 48 hours
2021	Fluoroquinolone treatment as a protective factor for 10-day mortality in *Streptococcus pneumoniae* bacteremia in cancer patients	Fontana, Ibrahim, Bonazzi et al (161)	161	34	Brazil	Patients attended between January 2009 and July 2015 at a cancer institute with haematological malignancies and / or solid tumours, over 18 years of age, and who had at least one episode of *S. pneumoniae* bacteremia during the referred period	Ethnicity, diabetes Mellitus, neutropenia past month, current neutropenia, severe neutropenia past month, current severe neutropenia, febrile neutropenia, SOFA score, polymicrobial bacteremia, vaccine serotype VPV23, 3rd and 4th generation cephalosporins, fluoroquinolones.	*Streptococcus pneumoniae*, tested for resistance against several antibiotics	Low	Yes	2.8	53	21	While the authors test for association between antimicrobial susceptibility and mortality, they conclude that there is no association and do not include it in the final model
2016	Bloodstream infection caused by extensively drug-resistant *Acinetobacter baumannii* in cancer patients: high mortality associated with delayed treatment rather than with the degree of neutropenia	Freire, de Oliveira Garcia, Garcia et al. (138)	92	77	Brazil	All patients diagnosed with extensively drug resistant *A. baumannii* bacteraemia, acquired during hospitalisation in the intensive care unit at a cancer institute	Serum albumin, septic shock at diagnosis of XDR-ABC bacteraemia (extended drug resistant), SOFA score at diagnosis, received appropriate therapy for XDR-ABC bacteraemia, combination therapy, days from 1rst positive culture to initial treatment, removal of invasive devices within the first 48 h	*Acinetobacter baumannii*, extensively drug resistance	Low	Yes	11.0	20	6	The authors found that 68% of all healthcare-associated bacteraemias are caused by extensively drug resistant *A. baumannii*, which is associated with a 30-day mortality of more than 80 %. The authors

														concluded that the only risk factor of importance was appropriate antibiotic treatment.
2018	Role of Lock Therapy for Long-Term Catheter-Related Infections by Multidrug-Resistant Bacteria	Freire, Pierotti, Zerati et al (108)	212	212	Brazil	All LTCVC-associated infections diagnosed at the a cancer institute between January 2009 and December 2016	Haematological malignancies, Palliative care, SOFA score, MDR bacteria, Adherence to protocol, Antibiotic lock therapy	Several bacteria and fungi, tested for resistance towards several antibiotics	Low	+	15,1	28	14	The authors found that there is an increased hazard of death when the patient is infected by multidrug-resistant bacteria, but that antibiotic lock therapy is associated with a decrease in death.
2015	Infection with *Klebsiella pneumoniae* carbapenemase (KPC)-producing *Klebsiella pneumoniae* in cancer patients	Freire, Pierrotti, Filho et al (146)	83	49	Brazil	All patients diagnosed with an healthcare associated infection due to Klebsiella producing carbapenemase-*Klebsiella pneumoniae* acquired during hospitalisation at a cancer institute between January 2009 and July 2013	Serum albumin, concomitant infection, bloodstream infection, urinary tract infection, SOFA score at healthcare associated infection (HAI) diagnosis, removal of invasive device or control deep infection, received effective therapy for Klebsiella producing carbapenemase-Kp infection, intensive care unit admission for healthcare associated infection (HAI), mechanical ventilation, acute kidney injury within 30 days after HAI diagnosis	*Klebsiella pneumoniae*, carbapenemase-producing	Low	Yes	4.5	27	11	The authors concluded that there is high mortality in patients with KPC-Kp infection, and that mortality is associated with the severity of the infection rather than the underlying disease
2018	Risk factors for mortality in patients with acute leukemia and bloodstream infections in the era of multiresistance	Garcia-Vidal, Cardozo-Espinola, Puerta-Alcalde et al (46)	589	87	Spain	All consecutive episodes of bloodstream infection occurring in patients with acute leukaemia from July 2004 to February 2016.	Older age (>65 years old), diabetes mellitus, chronic lung disease, chronic liver disease, prolonged neutropenia, mucositis, fatal prognosis according to McCabe index, community site of acquisition, catheter as a source of bacteremia, pulmonary source of infection, intra abdominal	Several bacteria and fungi, tested for resistance towards several antibiotics. A special focus on multidrug-resistant	Low	Yes	5.1	34	17	The authors found an association between multidrug-resistant *P. aeruginosa* and mortality, after adjusting for some other confounders and risk factors.

							source of infection, cONs, *E.coli, P.aeruginosa*, shock at onset, inappropriate antibiotic therapy for multi drug resistant strains, inappropriate antibiotic therapy for multi drug resistant *P. aeruginosa*	*Pseudomonas aeruginosa*.						
2017	Trends in the Incidence and Outcomes of Hospitalized Cancer Patients With Clostridium difficile Infection: A Nationwide Analysis	Gupta, Tariq, Frank et al (133)	20,100,000	Indetermi nable	USA	Cancer hospitalizations in a national cancer database from 2001 to 2010.	*C. difficile* infection, age, sex, admission status, and comorbidities.	*Clostridioide s difficile*	Mediu m	No	Indeterm inable	Indeterminabl e	Indeterm inable	The authors found that CDI is associated with a higher mortality among cancer patients. It was not clear how many factors variables were included in this model but we assumed 12 confounders and 2 factors (CDI and dismissal to care facility)
2021	Colonization with multidrug-resistant organisms impairs survival in patients with hepatocellular carcinoma	Himmelsbach, Knabe, Ferstl et al (24)	954	476	Germany	Patients with confirmed hepatocellular carcinoma presenting between January 2008 and December 2017	Age, sex, underlying cirhhosis, Child–Pugh class C vs. A or B, BCLC stage C or D vs. A or B, ALBI grade 3 vs. grade 1 or 2, Alpha-fetoprotein > 400 ng/ml, sorafenib treatment, resection as hepatocellular carcinoma treatment, multi drug resistant organism-colonization	Several bacteria, tested for resistance towards several antibiotics	Mediu m	Yes	95.2	10	5	In this study, the authors concluded that there is an association between colonisation with a multidrug-resistan t organism and mortality in patients with hepatocellular carcinoma.
2018	Bloodstream infections in cancer patients. Risk factors associated with mortality	Islas-Muñoz, Volkow-Fernánd ez, Ibanes-Gutiérre z et al (109)	496	109	Mexico	From August 2016 to July 2017, all positive blood cultures detected at the Microbiology Laboratory at a cancer centre were recorded, and all episodes	Age <60 years vs. Age >60 years, Solid tumour vs Haematological, Recent diagnosis or remission vs Progression/relapse, Gram-positive vs Gram-negative, Monomicrobial vs Polymicrobial, Appropriate treatment vs Inappropriate treatment, Non-multidrug resistant (MDR) vs MDR	Several bacteria, tested for resistance towards several antibiotics	Low	No	13.6	8	8	Authors included every variable of the univariate in the multivariate despite it being written that inclusion depended on p<0.5. The authors concluded that there was an

Year	Title	Authors	N		Country	Population	Variables/Factors	Resistant pathogens						Comments
						confirmed as bloodstream infection were included in the study.	and/or extended spectrum beta lactamase, Neutrophils >500 vs Neutrophils 500							association in the 30-day mortality among the cancer patients and bloodstream infection with multidrug-resistant Gram-negative bacteria, particularly if it was inappropriately treated.
2020	Nosocomial Infections in Gastrointestinal Cancer Patients: Bacterial Profile, Antibiotic Resistance Pattern, and Prognostic Factors	Jiang, Liu, Said et al (25)	428	Indeterminable	China	Hospitalised gastrointestinal cancer patients with nosocomial infections at a cancer treatment centre from August 2013 to June 2019	Gram negative bacteria, *E.coli, K. pneumoniae, Enterobacter* spp, Fungi, *C. albicans. Candida* spp, enterococcus	Several bacteria and fungi, tested for resistance towards several antimicrobials	Medium	Yes	Indeterminable	24	8	There are three full multivariable models in this article that all estimate the OR for mortality associated with different factors; a demographic/clinical model, an infection-related model, and a causative pathogens model. Only the causative pathogens model is assessed as the other models don't explicitly include antimicrobial resistance. One is about resistant pathogens. The authors concluded that Gram-negative bacteria were most common.
2020	Nosocomial infections due to multidrug-resistant bacteria in cancer patients: a six-year retrospective study of an oncology Center in Western China	Jiang, Shi, Liu et al (114)	257	28	China	Cancer patients with nosocomial infections due to multi drug resistant bacteria who received medical care during	Clinical model: Demographic data (age), Smoking history (Never smoker/Former smoker/Current smoker), ECOG performance status (0,1/2,3,4), Existence of distant metastasis, Comorbidities (Liver	Methicillin resistant *S. aureus* (MRSA), Extended spectrum beta lactamase producing	Low	Yes	2.5	40	11	There are two full multivariable models in this article that both are assessed, as both have antimicrobial resistance as factors. Numbers

Year	Question	Author			Country	Population	Factors	Bacteria	Quality					Notes
						hospitalisation from August 2013 to May 2019	disease), Charlson comorbidity index CCI (1–2), CCI≥3, Surgery, Chemotherapy, Intrapleural/abdominal infusion (within 30 days), Presence of indwelling urinary catheters, presence of Drains postoperative; Infection-related model: Sample type (Urine/Blood culture), Primary sites of infection (Respiratory tract/Urinary tract/BSI), Length of hospitalisation (≥21.0), Septic shock, Laboratory examination results (Haemoglobin g/L <110.0, Platelet count ×109/L <100.0, Lymphocytes count ×109/L <1.0, PCT ng/mL ≥1.0, Albumin g/L <30.0)	*Enterobacteriaceae* (ESBL-PE), multi drug resistant (MDR) *Pseudomonas aeruginosa, Acinetobacter baumannii, Stenotrophomonas maltophilia*, Carbapenem-resistant *Enterobacteriaceae*						are extracted from the nested model with the less variables tested, hence the highest number of events by variable in theory. Authors explained they included variables form univariate in MVA when p<0.1 but they included other factors as well. The most common bacteria were ESBL-PE, and the authors conclude that former smokers, intrapleural/abdominal infusion history within 30 days, presence of indwelling urinary catheters, and anaemia were independent risk factors for in-hospital mortality of nosocomial infections caused by MDR bacteria.
2020	Is current initial empirical antibiotherapy appropriate to treat bloodstream infections in short-duration chemo-induced febrile neutropenia?	Joncour, Puyade, Michaud et al (110)	27	13	France	Febrile neutropenic patients at a hospital undergoing chemotherapy for solid tumours or malignant haemopathy with expected duration of neutropenia ≤7 days from 1 January 2015 to 31 December 2016.	Signs of severity, documented clinical source, qSOFA score ≥2	Several bacteria, tested for resistance towards several antibiotics	Medium	Yes	4.3	18	3	The authors found that the only risk factor associated with mortality was the qSOFA score, and found no association between mortality and a history of infection/colonisation with MDR *P. aeruginosa*. Numbers are extracted from the model table in Supplementary

														material
2015	Uncommon Candida Species Fungemia among Cancer Patients, Houston, Texas, USA	Jung, Farmakiotis, Jiang et al (154)	66	40	USA	Patients ≥18 years of age hospitalised in the cancer centre during January 1998–September 2013.	Underlying leukaemia, steroid exposure, intensive care unit admission, intubation, persistent neutropenia, APACHE IIscore≥19, hypoalbuminemia, breakthrough fungemia	*Candida* spp.	Medium	Yes	5.0	9	8	The authors showed that ICU, neutropenia and high APACHE score, were the strongest risk factors for mortality.
2019	Impact of empiric treatment for vancomycin-resistant Enterococcus (VRE) in colonized patients early after allogeneic hematopoietic stem cell transplantation	Kamboj, Cohen, Huang et al (91)	95	5	USA	Adults 18 years and older who underwent allo-hæmotologic stem cell transplantation and developed vancomycin resistant enterococcus bacteremia between January 1, 2005 and December 31, 2014, at the cancer centre	T cell depletion, prior vancomycin resistant enterococcus bacteremia (VREB), hypotension (<90/50), persistent VREB >48 hours	*Enterococcus* spp., vancomycin-resistance	Medium	Yes	1.3	13	4	The authors found there is no association with higher mortality at day 30 and any of tested variables in VREB cancer patients
2019	Contribution of specific pathogens to bloodstream infection mortality in neutropenic patients with hematologic malignancies: Results from a multicentric surveillance cohort study	Kern, Roth, Bertz et al (50)	1424	36	Germany, Austria, Switzerland	Patients with bloodstream infection during neutropenia subsequent to high-dose chemotherapy for acute leukaemia (HDC) or autologous or allogeneic hematopoietic stem cell transplantation (auto-HSCT or allo-HSCT) from the beginning of 2002 through April 2015	Mortality 30 days model : Auto hematologic stem cell transplant treatment, CoNS (coagulase negative staphylococci). Mortality 7 days model: coNS	Several bacteria, tested for resistance towards several antibiotics	Low	Yes	18.0	5	2	There are 2 mortality models in Annex, and one model about ICU admission which we did not assess. Numbers are extracted from the mortality 30 days model. In this relatively large cohort, the authors found that there is an association between the infecting pathogen and 7 day mortality after BSI - particularly *Klebsiella, Enterobacter, Serratia*, and *Pseudomonas*.

2019	Pathogenic significance of hemorrhagic pneumonia in hematologic malignancy patients with *Stenotrophomonas maltophilia* bacteremia: clinical and microbiological analysis	Kim, Cha, Kang et al (84)	118	72	South Korea	Patients ≥18 years of age with hematologic malignancy who were diagnosed with *S. maltophilia* bacteremia	Age, hospital stay, polymicrobial infection, previous isolation of *S. maltophilia*, Focus of infection catheter related infection (CRI), focus of infection hemorrhagic, CRI non hemorrhagic, leukaemia, lymphoma, refractory or recurrent disease, Charlson comorbidity index I>3, neutropenia, intensive care unit stay, mechanical ventilation, renal Replacement Therapy, initial SOFA score, platelet count, previous antibiotics <3, early empirical TMP/SMX antibiotic use within 72 h	*Stenotrophomonas maltophilia*	Low	Yes	3.8	27	19	The authors also included clinically relevant variables in the MVA. The authors concluded that the mortality of haematological malignancy patients with *S. maltophilia* bacteraemia is high, and that hemorrhagic pneumonia is associated with an increased mortality
2020	Vancomycin-resistant enterococci infection and predisposing factors for infection and mortality in patients with acute leukaemia and febrile neutropenia	Kirkizlar, Akalin, Kirkizlar et al (92)	179	17	Turkey	Adult acute leukaemia patients with vancomycin resistant enterococcus colonised who had febrile neutropenia	Age, sex, primary disease acute myeloid leukemia, remission, previous hospitalisation, previous colonization, exposure to empirical antibiotics, mucositis, central venous catheter, invasive procedures, < 0,5x109/L neutrophil count while vancomycine resistant enterococcus (VRE) +, coinfection, acute kidney injury, >15 days VRE	*Enterococcus* spp. , vancomycin-resistance	Low	No	1.2	14	14	Apart from age and gender, the authors found an association between in-hospital mortality in VRE-colonised patients and low neutrophil count and coinfection
2017	Initial Treatment of Cancer Patients with Fluconazole-Susceptible Dose-Dependent Candida glabrata Fungemia: Better Outcome with an Echinocandin or Polyene Compared to an Azole?	Le, Farmakiotis, Tarrand et al (155)	68	26	USA	Patients with *C. glabrata* fungemia caused by strains with dose-dependent in vitro susceptibility to fluconazole at a cancer centre between March 2005 and September 2013.	APACHE II score, absolute monocyte count<100, intensive care unit, azole monotherapy	*Candida glabrata*	Medium	Yes	2.9	17	9	The authors found that azole monotherapy was associated with higher mortality than compared to the addition of an echinocandin or polyene in these patients
2021	Epidemiology, Risk Factors, and Clinical	Liang, Xu, Cheng et al (62)	449	Indeterminable	China	Patients with haematological	Myelodysplastic syndrome, MASCC score	*Escherichia coli* and	Medium	Yes	Indeterminable	Indeterminable	Indeterminable	The authors did not find a

Year	Title	Author			Country	Population	Factors	Bacteria						Comments
	Outcomes of Bloodstream Infection due to Extended-Spectrum Beta-Lactamase-Producing *Escherichia coli* and *Klebsiella pneumoniae* in Hematologic Malignancy: A Retrospective Study from Central South China					malignancies who were diagnosed with extended spectrum beta lactamase (ESBL)-producing and non-ESBL-producing *E. coli* and *K. pneumoniae* bacteremia between January 2010 and July 2018	<21, Charlson Comorbidity index>3	*Klebsiella pneumoniae*, ESBL						difference in overall mortality between the ESBL and non-ESBL groups, but they do find that the Charlson Comorbidity Index and the MASCC score were associated with an increased mortality in the ESBL-group. Number of deaths extracted from supplementary material.
2015	Factors influencing mortality in neutropenic patients with haematologic malignancies or solid tumours with bloodstream infection	Marin, Gudiol, Ardanuy et al (111)	510	61	Spain	Hospitalised adult neutropenic cancer patients with at least one episode of bloodstream infection	Haematological malignancies: age, sex, advanced neoplasm, MASCC score <21, corticosteroid therapy, coagulase-negative staphylococci, multi drug resistant gram negative bacteria, Empirical antibiotic combination therapy, intensive care unit admission. Solid tumours: age, sex, advanced neoplasm, corticosteroid therapy, septic shock.	Several bacteria, tested for resistance towards several antibiotics	Low	Yes	6.8	40	9	The authors used stepwise regression but also included some clinically relevant variables. The authors stratified the population and made one model for haematological malignancies and one model for solid tumours. The numbers were extracted from the hematologic model, with a higher number of events. They only found an association between antimicrobial resistance and mortality among haematological cancer patients
2019	Bloodstream infection in patients with head and neck cancer: a	Marin, Gudiol, Castet et al (26)	51	7	Spain	Consecutive episodes of bloodstream	Early case fatality: age, sex, Charlson scoring system, median albumin	Several bacteria and fungi, tested	Low	Yes	1.2	39	6	The authors used stepwise regression but

	major challenge in the cetuximab era					infection occurring in patients with head and neck cancer and bloodstream infection from January 2006 to April 2017	level, corticosteroid therapy. Overall case fatality: age, sex, Charlson scoring system, comorbidities, median Albumin level, corticosteroid therapy.	for resistance towards several antimicrobials					also included some clinically relevant variables. Both models investigating fatality include antimicrobial resistance as a factor, but it is removed after the bivariable screening. Numbers were extracted from the early fatality model. There is also a matched case-control study investigating the risk factors of all BSIs but it does not include any resistance.
2020	Inappropriate Empirical Antibiotic Treatment in High-risk Neutropenic Patients With Bacteremia in the Era of Multidrug Resistance	Martinez-Nadal, Puerta-Alcalde, Gudiol et al (112)	251	87	Spain	All consecutive episodes of bacteremia in patients with high-risk febrile neutropenia from January 2006 to January 2017	Inappropriate empirical antibiotic therapy, septic shock at onset, pneumonia	Several bacteria tested for resistance towards several antibiotics. The model is on *Pseudomonas aeruginosa*.	Low	Yes	29.0	Indeterminable	3	The authors found that inappropriate empirical antibiotic therapy, septic shock and pneumonia is associated with mortality among patients with *P. aeruginosa* bloodstream infection. The area under the curve of this model is printed.
2018	Postoperative empyema following lung cancer surgery	Matsutani, Yoshiya, Chida et al (27)	43	5	Japan	Patients who underwent thoracic surgery to treat primary lung cancer and had empyema between January, 2008 and December,	Exacerbation of interstitial pneumonia, bronchial stump fistula, administration of steroid, interstitial pneumonia, histology with or without squamous cells, microbiology with or without non-fermenting Gram(-) bacilli	Several bacteria, tested for resistance towards several antibiotics	Medium	Yes	0.8	35	6	The authors used stepwise regression but it is unclear whether they used p-value<0.05 to include factors of the univariate in the final model. they exclude one

Year	Title	Authors			Country	Population	Variables	Pathogen	Quality					Comments
						2012, from 9 institutions								significant factor with no observation in one group. No association with poor prognosis in empyema patients after lung cancer surgery were found through the multivariable model.
2017	Fecal Carriage of Extended-Spectrum b-Lactamase-Producing Enterobacteriaceae Strains Is Associated with Worse Outcome in Patients Hospitalized in the Pediatric Oncology Unit of Beni-Messous Hospital in Algiers, Algeria	Medboua-Benbalagh, Touati, Kermas et al (121)	171	28	Algeria	Paediatric oncology patients from February 2012 to May 2013	Age, gender, previous hospital admission, haematological malignancies, antibiotic treatment for the last 3 months before admission, haematological malignancies, Nephroblastoma, neuroblastoma, sarcomas, extended spectrum beta lactamase enterobacteria (ESBL-E)	*Enterobacteriaceae*, ESBL	Medium	Yes	Indeterminable	Indeterminable	Indeterminable	No regression table is printed, but the variables that the authors report to be associated with the two outcomes are reported here. During screening, the authors find that more than half of the patients are carriers of ESBL-producing *Enterobacteriaceae*.
2021	Secondary Infections After Diagnosis of SevereRadiation Pneumonitis (SRP) Among Patients With Non-Small Cell Lung Cancer: Pathogen Distributions, Choice of Empirical Antibiotics, and the Value of Empirical Antifungal Treatment	Mei, Yang, Yu et al (28)	777	174	China	Patients with severe radiation pneumonitis treated at a hospital between January 2009 and December 2020 with non-small cell lung cancer, who were infected after at least 48 hours of hospitalisation and received at least 1 antibiotic before susceptibility testing.	Performance status score, chronic obstructive pulmonary disease, diabetes, hypertension, radiotherapy dose in Gy, multiple bacterial infections, bacteria/fungal coinfections, appropriate antibiotic, empirical antifungal	Several bacteria and fungi, tested for resistance towards several antimicrobials	Medium	Yes	19.3	21	9	The authors used stepwise regression but it is unclear if the authors used a p-value<0.05 to include the factors tested in the univariate in the final model. The authors did not find that there is any association between intrinsically resistant bacteria and infection-related mortality

2021	Predictive factors associated with induction-related death in acute myeloid leukemia in a resource-constrained setting	Mendes, da Silva, da Costa Bandeira de Melo et al (51)	206	53	Brazil	Patients aged >=16 years with newly diagnosed acute myeloid leukaemia who started any regimen of intensive treatment	ECOG score binary, sex, age, intensive care unit entrance, Charlson comorbidities index (CCI), colonisation by *Klebsiella* producing carbapenemase, colonisation by *Acinetobacter*, colonisation by *Pseudomonas*, DM, tumor lysis, pre-chemo infection, Albumin to Globulin Ratio, monocytic acute myeloid leukaemia, monocytes, C reactive protein, prothrombine activity, D-Dimer, creatinin, albumin, alkaline phosphatase, total bilirubine, aciclovir, antibiotiv prophylaxis, anthracycline	Several bacteria and fungi, tested for resistance towards several antimicrobials	Low	Yes	2.1	56	25	The authors used stepwise regression but excluded ECOG in the stepwise approach due to the high missing rate found during the chart reviewing. The final multivariable model showed that age>60 years, Gram-negative colonisation, monocytic AML, CRP>15 mg/dL, and an adverse risk in the genetic stratification were independently associated with early mortality in AML patients
2021	Reduced mortality from KPC-*K.pneumoniae* bloodstream infection in high-risk patients with hematological malignancies colonized by KPC-*K.pneumoniae*	Micozzi, Gentile, Santilli et al (96)	34	10	Italy	Patients with haematological malignancies identified as Klebsiella producing carbapenemase -*K. pneumoniae* carriers attending a haematology department between March 2012-December 2013 (Period 1) and January 2017-October 2018 (Period 2).	Model 1: *Klebsiella* producing carbapenemase-*K. pneumoniae* (KPC-KP) bloodstream infection (BSI) developing during inactive antibiotic treatment, acute myeloid leukaemia, shock, intensive chemotherapy. Model 2: Initial active treatment, KPC-KP BSI developing during inactive antibiotic treatment, acute myeloid leukaemia, shock, intensive chemotherapy.	*Klebsiella pneumoniae*, carbapenemase-producing	Medium	Yes	2.5	22	4	The numbers were extracted from the Model 1. It is unclear if authors used stepwise with p-value<0.05 to include factors of the univariate in the final model, and it is also unclear which factors are included or kept in the model. The authors found that 3 of 10 patients that died were put on initially active treatment
2015	Outcomes of critically ill cancer patients with Acinetobacter baumannii infection	Ñamendys-Silva, Correa-García, García-Guillén et al (139)	23	6	Mexico	Cancer patients who acquired *Acinetobacter baumannii* infections during their stay	Length of hospital stay, duration of vasopressors, blood lactate level (mmol/L), number of organ dysfunction, creatinine,	*Acinetobacter baumannii*	Medium	Yes	1.2	14	5	The authors found that the blood lactate level was associated with mortality in these

Year	Title	Authors			Country	Population	Factors	Organism						Comments
						at the intensive care unit of a cancer institute								patients
2020	Characteristics and Clinical Outcomes of Extended-Spectrum beta-lactamase-producing *Klebsiella pneumoniae* Bacteremia in Cancer Patients	Nham, Huh, Cho et al (147)	267	46	South Korea	Patients above the age of eighteen with cancer who had an episode of *K. pneumoniae* bacteremia between 2010 and 2012	30-day mortality model: Neutropenia, foley catheter, nasogastric tube, mechanical ventilation, hemodialysis, catheter related infection, 14-day mortality model: solid tumour, extended spectrum beta lactamase, chronic liver disease, mechanical ventilation, hemodialysis, catheter related infection, PITT bacteremia score, carbapenem empirical therapy, vancomycin empirical therapy	*Klebsiella pneumoniae*, ESBL	Medium	Yes	7.7	51	6	There are two models, one on 14 days mortality and one about 30 days mortality. Numbers were extracted from the 30 day mortality model. The authors found that in these patients and with this statistical approach, there was an association between ESBL-production and 14-day mortality, but not with 30-day mortality
2015	Impact of vancomycin-resistant enterococcal bacteremia on outcome during acute myeloid leukemia induction therapy	Ornstein, Mukherjee, Keng et al (93)	350	84	USA	Patients with de novo and secondary acute myeloid leukaemia who received cytarabine-based induction chemotherapy at Cleveland Clinic between 1 January 2000 and 1 April 2008	Age (>60 vs. <60), cytology Poor vs. Favourable, cytology Poor vs. Intermediate, cytology Poor vs. Unknown, VRE bacteremia, other bloodstream infections, aetiology (secondary vs. de novo), white blood cell at diagnosis, year of diagnosis	*Enterococcus* spp., vancomycin-resistance	Medium	Yes	9.3	14	9	The authors included factors which were significant in the univariate analysis plus blood cell count, year of diagnosis and non VRE BSI. conclude that the survival is poorer in AML patients who had a VRE bacteraemia.
2020	*Candida* spp bloodstream infections in a Latin American Pediatric Oncology Reference Center: Epidemiology and associated factors	Paixao de Sousa da Silva, de Moraes-Pinto, Teofilo Pignati et al (156)	90	90	Brazil	Paediatric patients admitted to a paediatric oncology institute who presented with *Candida* spp. BSI from January 2004	Age, Hospitalisation, ICU, Corticosteroid, Prior antifungal use, Skin lesion, Dissemination, *Candida* spp Species (*C parapsilosis, C tropicalis,* Others).	*Candida* spp.	Low	+	9,0	28	10	The authors find that a *Candida parapsilosis* infection was associated with a lower mortality, while there was no association between *Candida tropicalis* and

Year	Title	Authors			Location	Population	Variables	Bacteria						Conclusion
						to December 2016.								mortality.
2019	Bloodstream Infection Due to Vancomycin-resistant *Enterococcus* Is Associated With Increased Mortality After Hematopoietic Cell Transplantation for Acute Leukemia and Myelodysplastic Syndrome: A Multicenter, Retrospective Cohort Study	Papanicolaou, Ustun, Young et al (94)	7128	2774	450 transplant centres worldwide	Patients who received their first allogeneic hematologic cell transplantation for acute myelogenous leukaemia (AML), acute lymphoblastic leukaemia (ALL), or myelodysplastic syndrome (MDS) between January 2008 and December 2012	Vancomycin resistant enterococcus bloodstream infection, age (21-40, 41-50, 51-60, >60), AML/ALL intermediate, AML/ALL/MDS advanced, MDS advanced, Cord blood allograft, Human leukocyte antigen 7/8, Karnofsky performance scale <90, cytomegalovirus donor or receptor positive, hematologic cell transplant during 2010-2012	*Enterococcus* spp. , vancomycin-resistance	Low	Indeterminable	277.4	Indeterminable	10	The overall survival model was examined here. We were unable to determine the method used for variable selection, other than that several variables were "examined" in a multivariable model. The authors conclude that VRE BSI was associated with mortality.
2019	Infections Due to Multidrug-Resistant Bacteria in Oncological Patients: Insights from a Five-Year Epidemiological and Clinical Analysis	Perdikouri, Arvaniti, Lathyris et al (33)	73	22	Greece	All cancer patients aged 18 years and older who presented an infection due to multi drug resistant bacteria from 1 January 2013 to 31 December 2017.	Model 1: neutropenia, C reactive protein (CRP). Model 2: creatinine, CRP. Model 3: guided antibiotics, duration of guided antibiotics. Model 4: empirical antibiotics, guided antibiotics. Model 5: CRP, guided antibiotics. Model 6: duration of empirical antibiotics, days of hospitalisation. Model 7: guided antibiotics, effective empirical antibiotics (there are 34 models overall, more models are in the Supplementary table)	*E. coli, Pseudomonas aeruginosa, Klebsiella pneumoniae, Acinetobacter baumannii, Enterococcus faecium, Staphylococcus aureus*, multidrug-, vancomycin- or oxacillin-resistant	Low	Yes	11.0	37	2	34 models with each time 2 different variables included were tested. The authors selected some variables thanks to those significant in the univariate but also based on previous literature. In univariate analyses they found several associations with death, but none of these held up in multivariable analysis.
2021	Clinical Characteristics and Outcome of Bloodstream Infections in HIV-Infected Patients with Cancer and Febrile Neutropenia: A Case–Control	Puerta-Alcalde, Ambrosioni, Chumbita et al (115)	180	43	Spain	All episodes of febrile neutropenia following chemotherapy occurring in patients with cancer and HIV from January	Hodgkin's lymphoma, Myelodysplastic syndrome, Solid neoplasia, Pulmonary source, Abdominal source, Shock, Candidemia, HIV-infection. Adjusted for: chronic liver disease, diabetes mellitus, chronic	Several bacteria and fungi, tested for resistance towards several antimicrobials	Medium	Yes	5.4	Indeterminable	8	The authors did not find any association between resistant microbes and mortality, other than candidemia, but include CoNS and S.

Year	Title	Author			Country									
	Study					1997 to March 2018.	renal failure, HIV, corticosteroid use, catheter-related source, inappropriate empirical antibiotic treatment, coagulase-negative staphylococci bacteremia, *S. pneumoniae* bacteremia, intensive care unit requirement.							pneumoniae as adjustment factors.
2017	Ultrasonography-driven combination antibiotic therapy with tigecycline significantly increases survival among patients with neutropenic enterocolitis following cytarabine-containing chemotherapy for the remission induction of acute myeloid leukemia	Pugliese, Salvatore, Iula et al (52)	100	23	Italy	Adult patients with newly diagnosed with acute myeloid leukaemia hospitalised at a haematological department from 1 January 2002 to 31 December 2012 in order to receive cytotoxic agent induction courses for haematological remission	High-dose cytarabine-containing chemotherapy, standard dose cytarabine chemotherapy, ultrasonography-driven necrotising enterocolitis NEC therapy with antibiotic regimens including tigecycline.	*Escherichia coli, Klebsiella pneumoniae, Enterococcus* spp., *Pseudomonas aeruginosa, Candida albicans*	Medium	Yes	7.7	24	3	It was unclear which variables were included in the final model. The authors did not find that mortality was associated with the different microbes
2015	Epidemiology and outcome of candidaemia in patients with oncological and haematological malignancies: results from a population-based surveillance in Spain	Puig-Asensio, Ruiz-Camps, Fernandez-Ruiz et al (157)	193	22	Spain	*Candida* bloodstream infection episodes in adult patients (>16 years) with underlying solid organ tumours or haematological malignancies	Charlson index, primary source of infection, catheter related source of infection, septic shock, *C. tropicalis*, central venous catheter removal within 48 hours, adequate combined treatment within 48 hours	*Candida* spp.	Low	Yes	3.1	27	7	The authors found that more than 1/4 of all *Candida* spp. isolates were non-susceptible to fluconazole, and that removal of the catheter and adequate antifungal treatment were associated with lower mortality.
2016	Clinical features and outcomes of Candidaemia in cancer patients: Results from Pakistan	Raza, Zafar, Mahboob et al (158)	311	166	Pakistan	Cancer patients with a positive blood culture for *Candida* species between January 1995 and December	Time period (1996-2005, 2006-2009, 2010-2013), Age group (18-59 year, Younger than 18 years, 60 years or older), sex, haematological malignancy (Compared to solid tumours), *Candida*	*Candida* spp.	Low	No	10.4	16	16	The authors wrote that the model was developed through forward selection, but do not report a stop criterion. We

Year	Title	Author	N		Country	Inclusion	Variables	Microorganisms						Comments
						2013	*albicans* (Compared to non-*C.albicans* or mixed), bacteraemia, shock, receiving chemotherapy, receiving total parenteral nutrition, receiving steroids, absolute neutrophile count 500 or more, duration of stay in hospital, receiving Amphotericin B (Compared to those not receiving), empirically, after positive culture							were unable to find the number of deaths. The authors did not find an association between death and different *Candida* species.
2017	A fresh look at polymicrobial bloodstream infection in cancer patients	Royo-Cebrecos, Gudiol, Ardanuy et al (113)	194	62	Spain	Consecutive episodes of primary bloodstream infections occurring in patients with cancer, including haematopoietic stem cell transplant recipients, from January 2006 to December 2015	Sex, age, solid tumor, MASCC risk score< 21, corticosteroid therapy, persistent bacteremia, respiratory source, septic Shock, malignancy-related complications	Several bacteria and fungi, tested for resistance towards several antimicrobials	Low	Yes	6.9	11	9	The authors used stepwise regression but also included clinically relevant variables in the final model regardless if they were significant or not in the univariate model. There is another model in this article with more patients analysing risk factors for polymicrobial bloodstream infections, which we did not consider to be intrinsically resistant. In the mortality model, the authors did not find that multidrug-resistant bacteria factor met the criterion in the univariable analysis to be included in the multivariable model.
2017	Characteristics, aetiology, antimicrobial	Royo-Cebrecos, Gudiol, Garcia et al (29)	170	45	Spain	All consecutive episodes of bacteraemic	Sex, age, pancreatic tumour, biliary tumour, corticosteroid therapy,	Several bacteria, tested for	Low	Yes	6.4	10	7	The authors used stepwise regression but

	resistance and outcomes of bacteraemic cholangitis in patients with solid tumours: A prospective cohort study					cholangitis occurring in patients with solid tumours from January 2006 to September 2015	fever >38C, malignancy-related complications	resistance towards several antibiotics						also included clinically relevant variables in the final model regardless if they were significant or not in the univariate model. The authors did not find an association between multidrug-resistance and mortality, and do not include it in the final, multivariable model
2017	Clinical Impact of Colonization with Multidrug-Resistant Organisms on Outcome after Autologous Stem Cell Transplantation: A Retrospective Single-Center Study	Scheich, Reinheimer, Brandt et al (47)	184	45	Germany	Patients admitted to a haematological and oncological department for auto-hæmatopoietic stem cell transplantation between January 2012 and October 2015, screened for multi drug resistant organisms	multi drug resistant organism colonisation (MDRO), age >55 yr, prior therapies >2, ECOG	MDRO	Medium	No	11.3	Indeterminable	4.0	The authors find an association between higher mortality and colonisation by multidrug-resistant organisms
2018	Bloodstream infections with gram-negative organisms and the impact of multidrug resistance in patients with hematological malignancies	Scheich, Weber, Reinheimer et al (58)	109	25	Germany	Patients with haematological malignancies and bloodstream infection with Gram-negative rods between January 2008 and December 2016	Sex, age > 65 years, multidrug resistant gram negative bloodstream infections (BSI), nonfermenter BSI, ANC < 500/µl duration until BSI (days), antibiotic prophylaxis, ICU admission, Charlson index > 4 points	Gram-negative bacteria, multidrug-resistance	Low	Yes	3.1	18	8	The authors used stepwise regression but they also included sex factor in the final model. The authors found that both being infected by a multidrug-resistant Gram-negative bacterium or a non-fermenter is associated with higher mortality
2018	Prognosis of Acute	Seong, Lee,	185	106	South Korea	All adult	Initial low tidal volume	Several	Mediu	Yes	8.8	33	12	The authors used

Year	Title	Authors			Country	Population	Variables	Organism						Conclusions
	Respiratory Distress Syndrome in Patients With Hematological Malignancies	Hong et al (48)				patients admitted to the Intensive care unit with haematological malignancy and acute respiratory distress syndrome between January 1, 2008, and December 31, 2015	ventilation, severe acute respiratory distress syndrome, carbapenem resistant gram negative bacteria (CR - GNB), age, underlying malignancies, status of disease, neutrophil count, platelet count, SOFA score, prone positioning treatment, vasopressors treatment, renal replacement therapy, gram negative bacteria, blood transfusion, pulmonary aspergillosis	bacteria and fungi, tested for resistance towards several antimicrobials	m					stepwise regression but also included clinically relevant variables in the final model regardless if they were significant or not in the univariate. The authors found that the mortality of patients with haematological malignancies that are moved to the ICU with ARDS was associated with carbapenem-resistant Gram-negative bacteria
2019	Risk factors for mortality due to *Acinetobacter baumannii* bacteremia in patients with hematological malignancies – a retrospective study	Shargian-Alon, Gafter-Gvili, Ben-Zvi et al (78)	46	33	Israel	Hospitalised adult patients with either active haematological malignancy or past treatment for haematological malignancy in the preceding five years and Acinetobacter baumannii bacteremia between January 2010 and August 2018	Appropriate empirical antibiotics within 48 h, SOFA score, ventilated at infection onset, urinary catheter at infection onset, duration of hospitalisation prior to bacteremia, absolute neutrophil count, creatinine level	*Acinetobacter baumannii*	Low	Yes	4.7	25	7	The authors found that there was a much lower mortality associated with being put on appropriate empirical antibiotic therapy within 48 hours
2020	Colonization with multi-drug-resistant organisms negatively impacts survival in patients with non-small cell lung cancer	Stratmann, Lacko, Ballo et al (30)	295	31	Germany	Patients diagnosed with stage 1-4 non-small cell lung cancer between 2012 and 2016 and screened for multi drug resistant	Multi drug resistant organism colonisation (MDRO), male gender, age at diagnosis, extensive diagnosis, histology: squamous cell carcinoma, histology: others, ECOG 1, ECOG 2, ECOG 3, ECOG 4, diabetes mellitus	MDRO	Low	No	2.8	Indeterminable	11	The authors concluded that the patients who were colonised by MDROs had a poorer survival than patients without such an infection

						organisms								
2020	Inappropriate initial antimicrobial therapy for hematological malignancies patients with Gram-negative bloodstream infections	Tang, Wu, Cheng et al (55)	361	45	China	Patients admitted to three university-affiliated tertiary care hospitals from January 2010 to April 2015	Age > 60 years, acute respiratory failure, disease relapsed or uncontrolled, platelet < 10 × 103 mm−3, Pitt score ≥ 4, 72-h IIAT	Gram-negative bacteria, tested for resistance towards several antibiotics	Medium	Indeterminable	7.5	Indeterminable	6	The authors did not disclose how they selected the variables to be included in the multivariable model of mortality. The authors did not find an association between intrinsically resistant pathogenic species and mortality, but found that inappropriate initial antimicrobial therapy was associated with higher mortality.
2021	Gram-Negative Bacteria Bloodstream Infections in Patients with Hematological Malignancies – The Impact of Pathogen Type and Patterns of Antibiotic Resistance: A Retrospective Cohort Study	Tang, Xu, Xiao et al (56)	835	313	China	Patients aged >16 years old with haematological malignancies and Gram negative bacteria-bloodstream infection, in 3 hospitals of Hunan province, from January 2010 to May 2018.	Age >60, sex, relapsed or uncontrolled malignancy, MASCC score<21, urine tube, use of vasopressors, acute respiratory failure, renal insufficiency, prior antimicrobial exposure, carbapenem resistant gram negative bacteria , non fermentative bacteria, inadequate antibiotic treatment, hemoglobin <70g/Dl, platelet <10×103mm−3, albumin <30g/L, AST >120U/L, total bilirubin >34.2µmol/L, prothrombin time >14s	Gram-negative bacteria, carbapenem resistance	Medium	Yes	17.4	18	18	The authors used stepwise regression but also included clinically relevant variables in the final model regardless if they were significant or not in the univariate. Authors highlighted that BSI caused by non-fermenting GNB, disease state, presence of acute respiratory failure, use of vasopressors, and inadequate antibiotic treatment were risk factors for a poor prognosis at 7 days.

2016	Bloodstream infections caused by *Klebsiella pneumoniae* in onco-hematological patients: clinical impact of carbapenem resistance in a multicentre prospective survey	Trecarichi, Pagano, Martino et al (97)	278	101	Italy	All episodes of bloodstream infections caused by *K. pneumoniae* that occurred in hospitalised haematological malignancy patients	Age>55 years old, ANC<100/mmc, ANC<500/mmc, acute myeloid leukaemia, non Hodgkin's lymphoma, Hodgkin's lymphoma, haematological disease newly diagnosed / relapsed after one remission, hematologic stem cell transplant HSCT, autologous HSCT, septic shock, altered state of consciousness, acute renal failure, acute respiratory failure, acute hepatic failure, inadequate initial antimicrobial therapy, carbapenem resistant by Klebsiella pneumoniae	*Klebsiella pneumonia*, carbapenem -resistance	Medium	Yes	6.3	31	16	Contains one model for carbapenem-susceptible and one model for carbapenem-resistant *K. pneumoniae* infections. Finds higher mortality among patients with a resistant infection than a susceptible infection.
2021	Clinical features and outcome of *Aeromonas sobria* bacteremia in pediatric and adult patients with hematologic malignancies: A single-center retrospective study in Peru	Valcarcel, De-la-Cruz-Ku, Malpica et al (105)	37	10	Peru	Adult and paediatric patients with hematologic diseases and *A. sobria* bacteremia hospitalised in the National institute of Neoplastic diseases from January 2000 to December 2017	Age group, comorbidities, qSOFA score, type of bacteremia	*A. sobria*	Low	No	2.5	4	4	The authors showed that organ failure and community acquired bacteremia increased the risk of mortality
2018	*Stenotrophomonas maltophilia* bacteremia and pneumonia at a tertiary-care oncology center: a review of 16 years	Velázquez-Acosta, Zarco-Márquez, Jiménez-Andrade et al (142)	76	38	Mexico	Patients aged >18 years old with cancer hospitalised in Instituto Nacional de Cancerologia with *S. maltophilia* bloodstream infection or pneumonia, from January 2000 to 31 December 2016	Model 1 in BSI: central venous catheter removal; Model 2 in pneumonia: age <65 years, received antibiotherapy with Sulfamethoxazole / trimethoprim, received appropriate treatment for *S. maltophilia*	*S. maltophilia*	Medium	Yes	12.7	5	3	This study contains 2 mortality models: one for BSI 30 days mortality and one for pneumonia 30 days mortality. Numbers were extracted from the pneumonia mortality model. The authors showed that for patients with pneumonia by S. maltophilia,

Year	Title	Authors			Country	Population	Variables	Species						Comments
														age>65 and not having received an appropriate treatment were risk factors for higher mortality within 30 days.
2015	The ever-evolving landscape of candidaemia in patients with acute leukemia: non-susceptibility to caspofungin and multidrug resistance are associated with increased mortality	Wang, Farmakiotis, Yang et al (103)	67	28	USA	Patients ≥18 years of age with Acute myeloid leukaemia, Acute lymphoid leukaemia or myelodysplastic syndrome, with Candidemia from 1 January 2008 to 31 October 2012	14 days mortality models 1 and 2 : age >65 years old, intensive care unit stay, APACHE II score >20, mechanical ventilation, vasopressors, caspofungin non-susceptibility, multidrug resistance ; 30 days mortality models 1 and 2 : sex, age >65 years old, ICU stay, APACHE II score >20, mechanical ventilation, vasopressors, caspofungin non-susceptibility, multidrug resistance	*Candida* spp.	Medium	Yes	4.0	7	7	There are four models displayed in the study. Numbers were extracted from the 14 days mortality model 1: The authors showed, after multivariable analysis, association between 14 or 30 days mortality and *Candida* spp. BSI that were non-susceptible to caspofungin. However, no statistically significant association was found between mortality and removal of CVC.
2020	Infectious Complications in Patients With Multiple Myeloma After High-Dose Chemotherapy Followed by Autologous Stem CellTransplant: Nationwide Study of the Infectious Complications StudyGroup of the Polish Adult Leukemia Group	Waszczuk-Gajda, Drozd-Sokołowska, Basak et al (49)	336	16	Poland	Patients with multiple myeloma having received autologous hematopoietic stem cell transplant during the 3 years period from 2012 to 2014, within one of 13 Polish transplant centres.	Bacterial multi drug resistant infection, coexistence of fungal infection, clinically documented infection	Several fungi and bacterial species, multidrug resistance	Medium	Yes	5.3	9	3	The authors highlighted 3 fatal outcome risk factors: MDR infection, fungal coinfection, Clinically documented infection, to adapt infection control in transplants recipients, with small sample size (16 infection related deaths).
2017	Bacteraemia due to AmpC β-lactamase-producin	Zhang, Zhang, Li et al (150)	248	62	China	Patients aged >18 years old with solid or	Abdominal infection, Urinary tract infection, unknown origin of	*E. coli*, AMPc Beta-Lacta	Low	Yes	5.6	29	11	The death model includes all bacteremias that

					Patient population	Factors	Microbial						Comments
g *Escherichia coli* in hospitalized cancer patients: risk factors, antibiotic therapy, and outcomes					haematological tumour, with *E. coli* bacteremia between September 2012 and December 2015	infection, intensive care unit stay, invasive procedures, previous blood transfusion, cancer with metastasis, septic shock, organ failure, length of stay, timing of appropriate antibiotherapy	mase positive or negative						were AMPc positive and negative but the study design excludes ESBL *E. coli* bacteremia.

Supplementary material 5

Table S5 - All included articles in the systematic review with both an infection outcome and a mortality outcome

Year	Title	Authors	Number of patients	Number of events in the final (and largest) model	Country/setting	Patient population	Factors included in the final model	Microbial aetiology and resistance	Risk of bias grading (NIH tool)	Bivariable screening/ stepwise regression	Events per variable in the final (and largest) model	Number of variables included in the tested/screened/initial model	Number of variables included in the final (and largest) model	Comments
2021	The Impact of Modifying Empirical Antibiotic Therapy Based on Intestinal Colonization Status on Clinical Outcomes of Febrile Neutropenic Patients	Alrstom, Alsuliman, Daher et al (63)	201	68	Syria	Adults older than 18 years of age with haematological malignancies who received chemotherapy and developed febrile neutropenia	Infection model: age, gender, prior quinolone use, prior hospitalization during previous month ; Mortality model: age, sex, extended spectrum beta lactamase carriage, carbapenem resistant *Enterobacteriaceae* carriage	Extended-spectrum β-lactamase-producing *Eschericia coli* and Carbapenem-Resistant *Enterobacteriaceae*	Medium	Yes	17.0	17	4	The stepwise regression also includes the age and sex variables. The authors did not find that carriage of the two resistant microbes increases the risk of dying. Numbers are taken from the infection model.
2018	Bloodstream infection caused by *S.aureus* in patients with cancer:a 10-year longitudinal single-center study	Bello-Chavolla, Bahena-Lopez, Garciadiego-Foss ass et al (148)	450	92	Mexico	All positive, non-duplicate, blood cultures for S.aureus	Infection model: hematologic neoplasm, solid neoplasia, healthcare associated	Methicillin-resistant *Staphylococcus aureus*	Medium	Yes	5.8	28	16	Authors created a risk stratification based on the regression coefficients from

Year	Title	Authors			Country	Population	Model variables	Organism						Comments
							bloodstream infections (BSI), hospital acquired BSI, foci as healthcare associated pneumonia, central venous catheter (CVC), neutropenia, gram negative bacteria coinfection, gram positive bacteria coinfection, inpatient management, specific anti SA treatment, CVC removal, days from infection to CVC removal, intensive care admission, septic shock, relapse after *S. aureus* bacteremia. Mortality model: abdominal source, hematologic malignancy, Methicillin-resistant *Staphylococcus aureus* (MRSA), glucose > 140 mg/dL, catheter removal, anti-staph treatment < 48 h, infectious endocarditis							the mortality model. The final model table can be found in supplementary material. The final model is also adjusted for CCI and glycemia >140 mg/dL. Numbers extracted from the infection model of MRSA infection in hematologic malignancies.
2019	Carbapenem versus Cefepime or Piperacillin-Tazobactam for Empiric Treatment of Bacteremia Due to Extended-Spectrum-β-Lactamase-Producing *Escherichia coli* in Patients with Hematologic Malignancy	Benanti, Brown, Shigle et al (99)	103	8	USA	Patients with hematologic malignancy and a first episode of mono-microbial extended spectrum beta lactamase-producing *E. coli* bacteremia	Infection model: cefepime treatment, piperacillin-tazobactam treatment (all vs Carbapenem treatment), age. Mortality model: cefepime treatment, Pitt bacteremia score, intensive care unit residence	Extended-Spectrum-β-Lactamase-Producing *Escherichia coli*	Medium	Yes	4.0	11	2	The stepwise regression included the cefepime treatment variable. The infection outcome is persistent bacteraemia. Authors found that only Pitt bacteraemia score is associated with an increased mortality. Numbers were

Year	Title	Authors			Country	Population	Model/Variables	Pathogen						Notes
														extracted from 14 days mortality model
2019	Associated factors and clinical outcomes of bloodstream infection due to extended-spectrum β-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* during febrile neutropenia	Ben-Chetrit, Eldaim, Bar-Meir et al (122)	80	80	Israel	All patients with hematologic or oncologic (solid tumours) diseases who were diagnosed with extended spectrum beta lactamase (ESBL) -positive and ESBL-negative *Escherichia coli* or *Klebsiella pneumoniae* bacteremia during febrile neutropenia between January 2010 and October 2017	Infection model: pathogene type, fungal infection within 30 days, presence of central venous catheter long term during index culture, admission days prior index culture, index culture >48h, previous antimicrobial therapy (6 months). Mortality model: extended spectrum beta lactamase-positive bacteremia, appropriate empirical antimicrobial therapy, charlson comorbidity index (CCI) >4, Pitt bacteremia score ≥4, underlying disease.	Extended-spectrum β-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*	Low	Yes	16.0	25	5	Authors found that a central venous catheter and previous antibiotic use is associated with ESBL-positive Gram-negative bacteraemia, but that the ESBL-positivity is not associated with an increase in mortality. Both models are the same size hence we chose the infection model to extract the data : extended-spectrumβ-lactamase(ESBL)-positive Gram-negative bacteremia.
2018	Risk, Outcomes, and Predictors of *Clostridium difficile* Infection in Lymphoma: A Nationwide Study	Bhandari, Pandey, Dahal et al (75)	236312	19358	USA	All adult patients aged 18 years or older with the primary diagnosis of lymphoma between 2007 and 2011	Age, sex, race, Hodgkin lymphoma, hematologic stem cell transplant, chemotherapy, radiotherapy, gastro intestinal surgery, infection, GVHD, IBD, CCI, year of admission. Mortality: *C. difficile* infection.	*Clostridioides difficile*	High	No	19358.0	Indeterminable	1	Authors found that HSCT, chemotherapy and surgery in the gut is associated with CDI, which increases the risk of death. Numbers were extracted from the mortality model. Most of the dataset is simulated using weights to create national estimates.
2018	Risk factors for bloodstream infections due to extended-spectrum β-lactamase producing	Ceken, Iskender, Gedik et al (123)	122	31	Turkey	Haematology/oncology patients with culture positive bloodstream infections with *E.*	Infection model: Quinolone prophylaxis, piperacillin-tazobactam previous 90 days, carbapenem	Extended-spectrum β-lactamase producing *Eschericia coli* and	Medium	Yes	10.3	4	3	It is not clear whether the authors used p<0.05 to include the variables in the multivariable

Year	Title	Authors			Country	Inclusion criteria	Factors	Bacteria						Comments
	Enterobacteriaceae in cancer patients					*coli* or *Klebsiella* spp from January 2013 to December 2014	previous 90 days, GNB infection previous 3 months and total parenteral nutrition. Mortality model : neutropenia, duration of neutropenia before infection, inappropriate treatment, delay for appropriate treatment, complications (hypoxia, septic shock etc.).	*Klebsiella* spp.						models. Numbers were extracted from the 30 days mortality model. For the infection model, authors found out that quinolone prophylaxis, total parenteral nutrition, piperacillin or carbapenem for the previous 90 days, or previous infection with GNB previous 90 days were risk factors to develop ESBL-P BSIs. Authors found that two factors were associated with an increased mortality in BSI cancer patients - prolonged neutropenia and complications.
2017	Poorer outcomes among cancer patients diagnosed with *Clostridium difficile* infections in United States community hospitals	Delgado, Reveles, Cabello et al (134)	30244426	Indeterminable	USA	Patients ≥18 years old with a principal or secondary ICD-9-CM code (discharge diagnosis of cancer) obtained from the U.S. National Hospital Discharge Surveys from 2001 to 2010	Indeterminable	*Clostridioides difficile*	High	Indeterminable	Indeterminable	Indeterminable	Indeterminable	Authors found that CDI significantly increased the risk for mortality and prolonged hospital stays among cancer patients. But overall the authors do not give any details about the model which was computed here.
2018	Role of Lock Therapy for Long-Term Catheter-Related Infections by Multidrug-Resistant Bacteria	Freire, Pierotti, Zerati et al (108)	212	75	Brazil	All long-term central venous catheter (LTCVC)-associated infections diagnosed at the a cancer institute	Haematological malignancies, type of CT last 3 months, palliative care, ICU admission, hemodynamic instability, SOFA	Several bacteria and fungi, tested for resistance towards several	Low	Yes	6.3	28	12	Model of relapse and mortality model. Numbers were extracted from the treatment failure or relapse model.

Year	Title	Authors			Country	Population	Variables	Organism						Conclusions
						between January 2009 and December 2016	score, CVC duration, *A.baumanii*, polymicrobial infection, Nadir WBC count, adherence to protocol, antibiotic lock therapy	antibiotics						The authors found that ALT is a protective factor for relapse risk and that there is an increased hazard of death when the patient is infected by multidrug-resistant bacteria, but that antibiotic lock therapy is associated with a decrease in death.
2018	Gut Colonization with Carbapenem-resistant Enterobacteriaceae Adversely Impacts the Outcome in Patients with Hematological Malignancies: Results of A Prospective Surveillance Study	Jaiswal, Gupta, Kumar et al (64)	225	48	India	Patients with newly diagnosed haematological malignancy admitted from October 2013 to January 2016	Age, gender, acute leukaemia, acute myeloid leukaemia, Charlson comorbidity index, performance status	*Enterobacteriaceae*, carbapenem-resistance	Low	No	8.0	6	6	The authors concluded that there is an association between having acute leukaemia and being at risk of both contracting and dying to a CRE infection. Numbers were extracted from the model for risk to be colonised to CRE before hospital admission.
2015	Outcomes of Clostridium difficile Infection in Hospitalized Leukemia Patients: A Nationwide Analysis	Luo, Greenberg, Stone (76)	1243107	95355	USA	Adults, aged ≥18 years, with a primary diagnosis code for leukaemia	Infection model: age, gender, race (Caucasian, Black, Hispanic, Asian/Pacific Islander, Native American), length of stay (days: 2, 3–5, 6–10, 11–15, ≥16), hospital teaching status, leukaemia type, complications (several), Charlson index (0, 1, 2, 3, 4, >5). Mortality model: age <65, gender, race, Charlson index	*Clostridioides difficile*	High	No	13622.1	7	7	The authors found a number of demographic or clinical associations with CDI, and that CDI was also associated with mortality. It is not reported how much of the dataset is individual data or projections based on weights. Numbers were extracted from

						score, teaching status sof hospital, the presence of complications or comorbidities (eg, hypertension, diabetes,pneumonia, etc.), and the performance of bone marrow vs stem cell transplant during hospitalisation.								the mortality model because no details are given concerning the other infection model.
2017	Carbapenem-resistant *Klebsiella pneumoniae* in high-risk haematological patients: factors favouring spread, risk factors and outcome of carbapenem-resistant *Klebsiella pneumoniae* bacteremias	Micozzi, Gentile, Minotti et al (98)	19	10	Italy	All patients with haematological malignancies infected or colonised with carbapenem resistant *K. pneumoniae* (CRKP), hospitalised between 24 February 2012 and 31 May 2013	Infection model: underlying disease, age, gender, intensive chemotherapy, >= 7 days colonised by carbapenem resistant *K. pneumoniae*. Mortality model 1: *Klebsiella* producing carbapenemase-*K.p neumoniae* bloodstream infection developing during inactive antibiotic treatment. Mortality model 2 and 3: initial adequate therapy, breakthrough bacteremia occurrence within 48 h of ongoing antibiotics, intensive Chemotherapy.	*Klebsiella pneumoniae*, carbapenem-resistance	Low	Yes	3.3	19	3	The infection outcome is bacteraemia. The authors found an association between AML and CRKP bacteraemia, and further that there was an association between survival and initial adequate therapy. Data was extracted from the final mortality model.
2015	Bacteraemia caused by *Escherichia coli* in cancer patients at a specialist center in Pakistan	Parveen, Sultan, Raza et al (151)	227	80	Pakistan	Cancer patients with *E.coli* bacteraemia during the 12 months from December 2012 to November 2013.	Infection model: sex, age less than 18 years, haematological malignancy, hospitalization within 30 days prior to infection, intensive care unit admission, Charlson score, central venous catheter use, previous chemotherapy within	*Escherichia coli*, multidrug-resistant	Low	No	6.7	12	12	The infection outcome is MDR in *E. coli* bacteraemia. The authors concluded that MDR *E. coli* is not associated with an increase in mortality. Numbers were extracted from the mortality model.

Year	Title	Authors			Country	Population	Model	Organism						Notes
							30 days, previous surgery within 30 days, previous radiation within 30 days, absolute neutrophil count (ANC)less than 100 cells/mm3, use of quinolones within 90 days of index sampling, use of third generation cephalosporins within 90 days of index sampling, use of Piperacilin/Tazobactam within 90 days of index sampling, use of carbapenems within 90 days of index sampling. Mortality model: sex, age less than 18 years, haematological malignancy, hospitalisation within 30 days prior to infection, admission to intensive care unit, previous chemotherapy within 30 days, previous surgery within 30 days, previous radiation within 30 days, ANC less than 100 cells/mm³, Charlson score, multi drug resistant *E.coli*, inappropriate empiric antibiotic therapy.							
2020	*Candida* spp bloodstream infections in a Latin American Pediatric Oncology Reference Center: Epidemiology and associated factors	Paixao de Sousa da Silva, de Moraes-Pinto, Teofilo Pignati et al (156)	90	90	Brazil	Paediatric patients admitted to a paediatric oncology institute who presented with *Candida* spp. BSI from January 2004 to December 2016.	Model of treatment failure or infection model: age, hospitalisation, corticosteroid, prior antifungal use, skin lesion, dissemination, *Candida* spp	*Candida* spp.	Low	Yes	10.0	31	9	There are 2 models in this study, one for antifungal treatment failure (then an infection model), the other one for 30 days mortality.

						Species, *C. tropicalis*, treatment. 30 days Mortality model: Age, Hospitalisation, intensive care unit, corticosteroid, prior antifungal use, skin lesion, dissemination, *Candida* spp Species, *C. parapsilosis*								Numbers were extracted from the infection model. The authors found that older age and thrombocytopenia were good independent predictors of therapeutic failure and that *Candida parapsilosis* infection was associated with a lower mortality, while there was no association between *Candida tropicalis* and mortality.
2017	Epidemiology, risk factors, and outcomes of infections in patients undergoing liver transplantation for hilar cholangiocarcinoma	Ramanan, Cummins, Wilhelm et al (31)	124	255	USA	Adult patients who underwent liver transplantation for biopsy-proven hilar cholangiocarcinoma at a clinic between January 2004 and March 2013	Infection model: vancomycin resistant enterococcus (VRE) colonisation pre-transplant, living donor transplant, cold ischemia time, cytomegalovirus donor seropositivity, hepatic artery thrombosis post-transplant, biliary stricture post-transplant, intra-abdominal fluid collection post-transplant, number of re-operations in 1st month post-transplant. Mortality model: abdominal infection, recurrent Hilar cholangiocarcinoma, Post-transplant *C. difficile* infection, Bloodstream infection	Several bacteria and fungi, tested for resistance towards several antimicrobials	Medium	No	15.9	16	16	There are two more models with an infection outcome in the article, of abdominal infection and bloodstream infection, but the model of any post-transplant infection is included here. The authors modelled subsets of 10 variables for infections and 2 variables for death and selected the best model based on score. They found that infections are associated with increased mortality, and that VRE colonisation pre-transplant is associated with

Year	Title	Authors			Country	Population	Details	Organism						Notes
														infection post-transplant. Numbers were extracted from the infection model.
2016	*Clostridium difficile* Infections amongst Patients with Haematological Malignancies: A Data Linkage Study	Selvey, Slimings, Joske et al (77)	2085	65	Australia	Patients admitted at least once to any hospital in the region for treatment or management of a haematological malignancy in the period 1 July 2011 to 30 June 2012	Infection model: acute lymphocytic leukaemia not in remission, neutropenia at any time, any bacterial pneumonia, other bacterial infection, any bacterial pneumonia and neutropenia at any time. Mortality model: acute myeloid leukaemia not in remission, *C. difficile* infection, age 65 and over, cerebrovascular disease, dementia, neutropenia at any time.	*Clostridioides difficile*	Low	Yes	4.1	30	16	There are two infection models, one with all patients and one with patients from October 2011. There are also two mortality models, one with death within 60 days and one with death within 90 days. The authors found that a *C. difficile* infection is associated with an increased mortality. Univariate analysis included both table 1 and 2. It is unclear where the dementia variable was taken from, but it is included in the mortality model. Numbers were extracted from the CDI model in all patients
2020	Risk factors for and clinical outcomes of carbapenem non-susceptible gram negative bacilli bacteremia in patients with acute myelogenous leukemia	Shin, Shin, Kang et al (57)	489	63	South Korea	Adult patients (aged ≥18 years) with acute myeloid leukaemia that contracted gram negative bacteremia during induction or consolidation chemotherapy, from January 2000 to	Infection model: age, chemotherapy, isolation of VRE in the prior 1 year, isolation of ESBL in the prior 1 year, isolation of carbapenem resistant *A. baumanii* (CRAB) in the prior 1 year, presence of preceding bacteremia during	Gram-negative bacilli, carbapenem-resistant	Medium	+	4.5	22	14	The authors found that carbapenem treatment at the onset of Gram-negative bacteraemia and infection with carbapenem-resistant *Acinetobacter baumannii* was associated with a

					December 2015	the hospitalisation, central line associated infection, *E.coli, Klebsiella* spp., *Pseudomonas* spp, *S. maltophilia, Acinetobacter* spp, antibiotics at the onset of gram negative bacteria (GNB) bacteremia, inappropriate empiric antibiotics, days from chemotherapy to GNB bacteremia, hospital days to GNB bacteremia. Mortality model: age, mean (± SD), induction or re-induction chemotherapy, isolation of vancomycin resistant enterobacteriaceae (VRE)in the prior 1 year, isolation of extended spectrum beta lactamase (ESBL) in the prior 1 year, isolation of CRAB in the prior 1 year, history of GNB bacteremia in the prior 1 year, presence of preceding bacteremia during the hospitalisation, pneumonia, Pitt score, septic shock, carba-non sensitive, inappropriate empiric antibiotics, days from chemotherapy to GNB bacteremia, hospital days to GNB bacteremia.								carbapenem-resistant Gram-negative bacteraemia, and also that infection with VRE the previous year and carbapenem-resistance was associated with an increased mortality in patients with Gram-negative bacteraemia. Numbers were extracted from the mortality model.
2019	Trends in Incidence and Outcomes of *Clostridium difficile* Colitis in Hospitalized Patients of Febrile	Siddiqui, Khan, Khan et al (135)	394987	Indeterminable	USA	All patients with primary and secondary diagnosis of febrile	Age, race, gender, Charleston Comorbidity Index, malignancy type, admission days,	*Clostridioides difficile*	High	No	Indeterminable	23	23	Authors showed that patients older than 65 years old,Hispanic

	Neutropenia A Nationwide Analysis					neutropenia, with and without *C. difficile* infection from years 2008 to 2014, according to the Nationwide Inpatient Sample (NIS)	hospital region, hospital location, sepsis							race, and presence of sepsis were associated with increased risk of mortality in cancer patients with FN. In addition, Hispanic race, haematological malignancies, and sepsis were significant risk factors for development of CDI in these patients. Both models are the same size.
2019	Increase in *Candida Parapsilosis* Candidemia in Cancer Patients	Sun, Chen, Xiao et al (159)	323	47	China	All patients with positive blood culture for *Candida* species at a cancer hospital from 1 March 2012 to 28 February 2018	Infection by *C. parapsilosis* model: age, hematologic malignancy, in intensive care unit (ICU) at diagnosis, parenteral nutrition, neutropenia, previous surgery last 3 months, abdominal surgery, chemotherapy, previous antifungal use, duration of antifungal; Infection by *C. albican*s model: age, hematologic malignancy, ICU at diagnosis, neutropenia, abdominal surgery, central venous catheter (CVC) related candidemia, chemotherapy, previous antifungal use, duration of antifungal; Mortality model: age, ICU at diagnosis, mechanical ventilation, CVC	*Candida* spp.	Medium	Yes	5.2	23	9	The authors concluded that there are several risk factors associated with a *Candida* non-albicans infection, but did not model the role of these infections in the mortality of patients. To compute the number of events per variable we chose the infection model by *C. parapsilosis*.

							related candidemia, removal of CVC.						
2017	*Pseudomonas aeruginosa* bacteraemia in patients with hematologic malignancies: risk factors, treatment and outcome	Tofas, Samarkos, Piperaki et al (79)	64	23	Greece	Cases: Consecutive patients with hematologic malignancy who developed *P. aeruginosa* (PA) bloodstream infection (BSI) between January 2012 and December 2014. Controls: Patients with hematologic malignancies without PA infection.	Infection with *P. aeruginosa* model: severity of underlying disease (Ultimately fatal, Rapidly fatal), hospitalisation in preceding 3 months, surgery in preceding 3 months, neutropenia, use of steroids, Prior use of any antimicrobial, prior use of anti-pseudomonal lactams including carbapenems, prior use of Aminoglycoside, prior use of Fluoroquinolones. Infection with carbapenem-resistant PA model: hospitalization in preceding 3 months, anti-pseudomonal lactams, carbapenems, aminoglycoside, fluoroquinolones. Mortality model : severity of sepsis (Sepsis, Severe sepsis or septic shock), empirical treatment (No active drug, At least one active drug), definitive treatment (One active drug, More than one active drugs).	*Pseudomonas aeruginosa*, carbapenem resistant	Low	Yes	7.7	19	3	There are three models here, one for PA BSI, one for carbapenem-resistant PA BSI and one for mortality in patients with PA BSI. All numbers are extracted from the 28 days mortality model. The authors found that prior use of antibiotics were associated with PA BSI, and that prior hospitalisation was associated with carbapenem-resistance.
2021	Oral fluoroquinolones for definitive treatment of gram-negative bacteremia in cancer patients	Tossey, El Boghdadly, Reed et al (118)	211	39	USA	Patients at a cancer centre with positive blood cultures for gram-negative organisms during the study period	Intravenous group, hematologic malignancy, neutropenia, *P. aeruginosa* infection, history of hematologic stem	Gram-negative bacteria	Medium	Indeterminable	5.6	Indeterminable	7	The authors modelled both infection recurrence and mortality as a composite outcome

Year	Title	Authors	Col4	Col5	Country	Population	Variables	Outcome						Comments
						from November 1, 2011, to September 30, 2017	cell transplantation, Pitt bacteremia score, intensive care unit admission							"treatment failure". No method for variable selection is disclosed. They found that having a *P. aeruginosa* infection is associated with this composite outcome.
2019	Bloodstream infections caused by Escherichia coli in onco-haematological patients: Risk factors and mortality in an Italian prospective survey	Trecarichi, Giuliano, Cattaneo et al (100)	342	24	Italy	All episodes of bloodstream infection caused by E. coli (EC BSI) occurring in hospitalised haematologic malignancies adult patients from January 2016 to December 2017 in 15 haematology wards	Infection model: recent endoscopic procedures, parenteral nutrition, polymorphonuclear leukocytes < 500/mmc for at least 10 days, previous antibiotic therapy, multiple myeloma, allogeneic matched unrelated transplant, multi drug resistant bacteria culture-positive surveillance rectal swabs, antibiotic prophylaxis with fluoroquinolones. Mortality model: sex, complete remission, autologous transplant, septic shock, altered consciousness, acute renal failure, avute respiratory failure, acute hepatic failure, 3rd generation cephalosporins resistance by *E. coli* isolate.	*Escherichia coli*, 3rd generation cephalosporin-resistant	Medium	Yes	2.7	34	9	The infection outcome here is resistance towards 3rd generation cephalosporins in *E. coli* bacteraemia. The authors found that such infections are associated with an increased mortality in these patients. All the numbers were extracted from the mortality model
2019	Clinical outcome of Escherichia coli bloodstream infection in cancer patients with/without biofilm formation: a	Zhang, Gao, Li et al (152)	324	71	China	Adult patients (aged ≥18 years) with cancer and first episode of E. coli bloodstream infection	Infection models 1 (extended spectrum beta lactamase yes no) and 2 (biofilm formation yes no) : sex, age, hypertension,	ESBL-producing *Escherichia coli* or biofilm-producer *Escherichia*	Low	Yes	8.9	18	8	The authors concluded biofilm is a risk factor for mortality , and also to be ESBL or not (and vice versa). No clear

	single-center retrospective study					hospitalised at Tianjin Cancer Institute from January 2013 to September 2017	chronic heart disease, diabetes mellitus, solid tumor, hematological cancer, abdominal infection, urinary tract infection, biliary infection, pulmonary infection, catheter-related infection, unknown origin infection, others, C3G (or cephalosporin (model1) exposure previous month, betalactam with or without inhibitor of betalactamase exposure previous month, carbapanems exposure previous month, combined exposure previous month, aminoglycoside exposure previous month, invasive procedure, surgery previous month, chemotherapy, ESBL, LOS (model 1). Mortality model: sex, age, hypertension, chronic heart disease, diabetes mellitus, solid tumor, hematological cancer, LOS, ICU, invasive procedure, mechanical ventilation, metastasis, chemotherapy, surgery, previous blood transfusion, ESBL, BF, organ failure and sepsis shock	*coli*						association between infection with biofilm-producer and antibiotics used. Numbers were extracted from the mortality model.
2020	Risk Factors and Outcomes of	Zhao, Lin, Liu et al (80)	293	293	China	Adult patients hospitalised with	Infection model 1 : Previous use of	*P. aeruginosa*,	Mediu m	+	48,8	14	6	Outcomes for infection are 1)

	Antibiotic-resistant Pseudomonas aeruginosa Bloodstream Infection in Adult Patients With Acute Leukemia					Acute Leukaemia and P. aeruginosa bloodstream infection from January 2014 to December 2019	fluoroquinolones, previous use of piperacillin-tazobactam, previous use of carbapenems, HSCT. Infection model 2 : Previous use of fluoroquinolones, previous use of piperacillin-tazobactam. Death model : Age>55, Perianal infection, Pulmonary infection, MDR-PA	carbapenem resistant or MDR						Carbapenem-resistant *P. aeruginosa* infection or 2) MDR *P. aeruginosa* infection. Variables with p<0.1 in the univariate analysis were included in the multivariable model. The authors found that MDR is associated with bad prognosis in BSI *P. aeruginosa* patients, and they have tried to show as well that exposure to certain classes of antibiotics could lead to developing MDR *P. aeruginosa* BSI in these patients. They aimed to enhance the role of an appropriate antibioprophylaxis in acute leukaemia patients. Numbers were taken from infection model MDR PA BSI.

Supplementary material 6

PRISMA 2020 Main Checklist

Topic	No.	Item	Location where item is reported
TITLE			
Title	1	Identify the report as a systematic review.	pp. 1
ABSTRACT			
Abstract	2	See the PRISMA 2020 for Abstracts checklist	
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of existing knowledge.	pp. 2
Objectives	4	Provide an explicit statement of the objective(s) or question(s) the review addresses.	pp. 3
METHODS			
Eligibility criteria	5	Specify the inclusion and exclusion criteria for the review and how studies were grouped for the syntheses.	pp. 4
Information sources	6	Specify all databases, registers, websites, organisations, reference lists and other sources searched or consulted to identify studies. Specify the date when each source was last searched or consulted.	pp. 3
Search strategy	7	Present the full search strategies for all databases, registers and websites, including any filters and limits used.	S1, S2
Selection process	8	Specify the methods used to decide whether a study met the inclusion criteria of the review, including how many reviewers screened each record and each report retrieved, whether they worked independently, and if applicable, details of automation tools used in the process.	pp. 4

Data collection process	9	Specify the methods used to collect data from reports, including how many reviewers collected data from each report, whether they worked independently, any processes for obtaining or confirming data from study investigators, and if applicable, details of automation tools used in the process.	pp. 4-5
Data items	10 a	List and define all outcomes for which data were sought. Specify whether all results that were compatible with each outcome domain in each study were sought (e.g. for all measures, time points, analyses), and if not, the methods used to decide which results to collect.	pp. 4-5
	10 b	List and define all other variables for which data were sought (e.g. participant and intervention characteristics, funding sources). Describe any assumptions made about any missing or unclear information.	pp. 4-5
Study risk of bias assessment	11	Specify the methods used to assess risk of bias in the included studies, including details of the tool(s) used, how many reviewers assessed each study and whether they worked independently, and if applicable, details of automation tools used in the process.	pp. 4
Effect measures	12	Specify for each outcome the effect measure(s) (e.g. risk ratio, mean difference) used in the synthesis or presentation of results.	NA
Synthesis methods	13 a	Describe the processes used to decide which studies were eligible for each synthesis (e.g. tabulating the study intervention characteristics and comparing against the planned groups for each synthesis (item 5)).	NA
	13 b	Describe any methods required to prepare the data for presentation or synthesis, such as handling of missing summary statistics, or data conversions.	NA
	13 c	Describe any methods used to tabulate or visually display results of individual studies and syntheses.	pp. 4-5
	13 d	Describe any methods used to synthesize results and provide a rationale for the choice(s). If meta-analysis was performed, describe the model(s), method(s) to identify the presence and extent of statistical heterogeneity, and software package(s) used.	pp. 4-5
	13 e	Describe any methods used to explore possible causes of heterogeneity among study results (e.g. subgroup analysis, meta-regression).	NA
	13f	Describe any sensitivity analyses conducted to assess robustness of the synthesized results.	NA

Reporting bias assessment	14	Describe any methods used to assess risk of bias due to missing results in a synthesis (arising from reporting biases).	NA
Certainty assessment	15	Describe any methods used to assess certainty (or confidence) in the body of evidence for an outcome.	NA

RESULTS

Study selection	16 a	Describe the results of the search and selection process, from the number of records identified in the search to the number of studies included in the review, ideally using a flow diagram.	pp. 5-6
	16 b	Cite studies that might appear to meet the inclusion criteria, but which were excluded, and explain why they were excluded.	pp. 12-13
Study characteristics	17	Cite each included study and present its characteristics.	S4, S5, S6
Risk of bias in studies	18	Present assessments of risk of bias for each included study.	S4, S5, S6
Results of individual studies	19	For all outcomes, present, for each study: (a) summary statistics for each group (where appropriate) and (b) an effect estimate and its precision (e.g. confidence/credible interval), ideally using structured tables or plots.	pp. 7-8
Results of syntheses	20 a	For each synthesis, briefly summarise the characteristics and risk of bias among contributing studies.	NA
	20 b	Present results of all statistical syntheses conducted. If meta-analysis was done, present for each the summary estimate and its precision (e.g. confidence/credible interval) and measures of statistical heterogeneity. If comparing groups, describe the direction of the effect.	pp. 7-8
	20 c	Present results of all investigations of possible causes of heterogeneity among study results.	pp. 8-12
	20 d	Present results of all sensitivity analyses conducted to assess the robustness of the synthesized results.	NA
Reporting biases	21	Present assessments of risk of bias due to missing results (arising from reporting biases) for each synthesis assessed.	NA
Certainty of evidence	22	Present assessments of certainty (or confidence) in the body of evidence for each outcome assessed.	NA

DISCUSSION

Discussion	23 a	Provide a general interpretation of the results in the context of other evidence.	pp. 12
	23 b	Discuss any limitations of the evidence included in the review.	pp. 14
	23 c	Discuss any limitations of the review processes used.	pp. 14
	23 d	Discuss implications of the results for practice, policy, and future research.	pp. 15
OTHER INFORMATION			
Registration and protocol	24 a	Provide registration information for the review, including register name and registration number, or state that the review was not registered.	pp. 3
	24 b	Indicate where the review protocol can be accessed, or state that a protocol was not prepared.	pp. 3
	24 c	Describe and explain any amendments to information provided at registration or in the protocol.	NA
Support	25	Describe sources of financial or non-financial support for the review, and the role of the funders or sponsors in the review.	pp. 15
Competing interests	26	Declare any competing interests of review authors.	pp. 15
Availability of data, code and other materials	27	Report which of the following are publicly available and where they can be found: template data collection forms; data extracted from included studies; data used for all analyses; analytic code; any other materials used in the review.	pp. 15

PRISMA Abstract Checklist

Topic	No.	Item	Reported?
TITLE			

Title	1	Identify the report as a systematic review.	Yes
BACKGROUND			
Objectives	2	Provide an explicit statement of the main objective(s) or question(s) the review addresses.	Yes
METHODS			
Eligibility criteria	3	Specify the inclusion and exclusion criteria for the review.	Yes
Information sources	4	Specify the information sources (e.g. databases, registers) used to identify studies and the date when each was last searched.	Yes
Risk of bias	5	Specify the methods used to assess risk of bias in the included studies.	Yes
Synthesis of results	6	Specify the methods used to present and synthesize results.	Yes
RESULTS			
Included studies	7	Give the total number of included studies and participants and summarise relevant characteristics of studies.	Yes
Synthesis of results	8	Present results for main outcomes, preferably indicating the number of included studies and participants for each. If meta-analysis was done, report the summary estimate and confidence/credible interval. If comparing groups, indicate the direction of the effect (i.e. which group is favoured).	Yes
DISCUSSION			
Limitations of evidence	9	Provide a brief summary of the limitations of the evidence included in the review (e.g. study risk of bias, inconsistency and imprecision).	No
Interpretation	10	Provide a general interpretation of the results and important implications.	Yes
OTHER			
Funding	11	Specify the primary source of funding for the review.	Yes
Registration	12	Provide the register name and registration number.	Yes