Toward unraveling the correlates of fatigue in glioma

Jantine G. Röttgering[1,2]*, Vera Belgers[2,3], Philip C. De Witt Hamer[2,4], Hans Knoop[1,5,6], Linda Douw[2,7], Martin Klein[1,2]

*Corresponding author, j.rottgering@amsterdamumc.nl, +31625005952, address: Amsterdam UMC location Vrije Universiteit Amsterdam, Medical Psychology, Boelelaan 1117, Amsterdam, The Netherlands

[1] Amsterdam UMC location Vrije Universiteit Amsterdam, Medical Psychology, Boelelaan 1117, Amsterdam, The Netherlands (JR, HK, MK)

[2] Cancer Center Amsterdam, Brain Tumor Center, Amsterdam, The Netherlands (JR, VB, PWH, LD, MK)

[3] Amsterdam UMC location Vrije Universiteit Amsterdam, Neurology, Boelelaan 1117, Amsterdam, The Netherlands (VB)

[4] Amsterdam UMC location Vrije Universiteit Amsterdam, Neurosurgery, Boelelaan 1117, Amsterdam, The Netherlands (PWH)

[5] Amsterdam Public Health Research Institute, Amsterdam, The Netherlands (HK)

[6] Amsterdam UMC location University of Amsterdam, Medical Psychology, Meibergdreef 9, Amsterdam, The Netherlands (HK)

[7] Amsterdam UMC location Vrije Universiteit Amsterdam, Anatomy and Neurosciences, Boelelaan 1117, Amsterdam, The Netherlands (LD)

Funding: Funding for this work was provided by the Stichting Anita Veldman Foundation (CCA-2019-2-21).

Conflict of interests: The authors declare that they have no competing interests.

Manuscript word-count: 5828

Authorship*:* *Conceptualization and design*: JR, MK, LD. *Data collection and curation*: JR, VB. *Data analysis and interpretation*: JR, VB, LD, MK. *Visualization*: JR. *Writing of the manuscript*: JR. *Revising of the manuscript*: JR, VB, PWH, HK, LD, MK. *Final approval of manuscript*: JR, VB, PWH, HK, LD, MK. *Accountable for all aspects of the work*: JR, VB, PWH, HK, LD, MK.

Abstract

Background: Even though fatigue is one of the most prevalent and burdensome symptoms in patients with glioma, its etiology and determinants are still poorly understood. We aimed to identify which demographic, tumor- and treatment-related characteristics and patient-reported outcome measures (PROMs) are associated with or are predictors of fatigue in glioma.

Methods: In this retrospective observational study, we included glioma patients with preoperative and postoperative assessments including PROMs on fatigue, depression, cognitive functioning, and health-related quality of life (HRQoL). Linear mixed models were used to identify which clinical factors and PROMs were associated with fatigue and linear multiple regression was used to detect predictors of postoperative fatigue.

Results: In this study, 222 patients were included (78% grade II-III glioma, 22% grade IV). These patients had performed 333 assessments (193 preoperative and 116 one year postoperatively). Of all assessments, 39% was indicative of severe fatigue. Several HRQoL domains, depression, and right-sided tumors were significantly associated with fatigue (marginal R^2 = 0.63). Contrary to common expectations, tumor type, treatment-related factors and timing of the assessment, were not associated with fatigue. In a subgroup of seventy patients with follow-up assessments, preoperative fatigue and physical functioning were predictors of postoperative fatigue (adjusted R^2 = 0.31).

Conclusion: Fatigue is a complex symptom, which should not solely be attributed to the tumor or its treatment, but is instead related to different aspects of mood and HRQoL. These insights are of importance in understanding fatigue and could guide symptom management, especially in patients with lower grade tumors.

Keywords

Fatigue; Patient Reported Outcome Measures; Brain Neoplasms; Quality of Life; Depression

Key points

Up to 40% of patients are severely fatigued, regardless of tumor type or timing of assessment

HRQoL, depression, and right-sided tumor explain 63% of variance in fatigue severity

Demographics, tumor type and treatment are not associated with fatigue severity

Importance of the Study

Despite fatigue being one of the most frequent and burdensome symptoms in patients with glioma, its etiology remains poorly understood. We aimed to unravel the impact of demographic, tumor- and treatment-related factors, symptoms, and HRQoL to grasp the complexity of fatigue in 222 glioma patients. Our findings demonstrate that fatigue is associated with several domains of HRQoL and depression, as well as having a tumor in the right hemisphere. What stands out is that fatigue is not associated with tumor type, treatment-related characteristics or phase of the disease. These findings could be important in symptom management since tumor- and treatment-related factors are non-modifiable, whilst depressive symptoms and social and physical functioning could be more suitable treatment targets. These results underscore the need for timely screening and symptom management focusing not only on fatigue but also on mood and diminished HRQoL.

Background

Patients with diffuse glioma experience a multitude of symptoms during the course of the disease including fatigue, cognitive impairment, and neurological deficits.[1] Fatigue is one of the most frequently reported and burdensome symptoms and is associated with a significant loss of health-related quality of life (HRQoL).[2] The prevalence of fatigue fluctuates over time and reported numbers vary between studies.[3] Of the lower-grade glioma patients 39-77% report fatigue,[4] while 48% of glioblastoma patients report fatigue after surgery, and this number increases to up to 90% in patients with tumor recurrence.[5,6] Because gliomas are still incurable and many patients suffer from fatigue and its sequelae, it is important to understand the interaction between fatigue and other symptoms and HRQoL to guide rational symptom management.

The etiology of fatigue in brain tumor patients is complex and poorly understood. Demographic, biomedical, neuropsychological, and psychosocial factors are thought to be related to fatigue.[7] It is unclear whether those factors are causally linked to the emergence and perpetuation of fatigue, or that symptom burden and diminished HRQoL are a result of severe fatigue. Several studies have reported conflicting variables to be associated with fatigue in glioma patients, often in small samples including different types of primary brain tumor patients at different time points during the disease. A relatively large study identified poor performance status, female sex, and active disease (compared to stable disease) to be associated with fatigue in two-hundred primary brain tumor patients.[8] Contrarily, another study in 65 postoperative glioblastoma patients found that performance status was not associated with fatigue, but that depression played an important role.[5] Furthermore, two smaller studies in lower-grade glioma patients found that fatigued patients experienced more cognitive impairment, were older and used more anti-epileptic drugs.[9,10] It is important to note, that all these studies were cross-sectional and did not take longitudinal changes of fatigue into account, and thus did not investigate any potential predictors of fatigue.

Fatigue is also a prominent symptom in patients with cancer, cancer survivors, and patients with neurological diseases, including multiple sclerosis and Parkinson's disease.[11,12] A meta-analysis combined fifteen studies on fatigue in common chronic disorders and found that over half of all patients suffered from severe fatigue and that type of disease could only explain a small proportion of variance

in fatigue severity.[13] Interestingly, when sex, age, motivational and concentration problems, pain, sleep disturbances, physical functioning, reduced activity and lower self-efficacy were added to the model, up to half of the variance in fatigue severity could be explained. The common denominator in those studies is that disease-specific factors were not associated with fatigue, whilst psycho-social symptoms, physical functioning and HRQoL were. It is not clear whether this would be the same in patients with glioma and whether tumor- and treatment-related factors play a role in glioma-related fatigue.

The current study aims to evaluate fatigue in a large sample of glioma patients in relation to demographic, tumor- and treatment-related variables, as well as symptoms and HRQoL measured with patient-reported outcome measures (PROMs). Furthermore, in a subset of patients with follow-up data, we aim to find predictors of postoperative fatigue and demonstrate how fatigue severity changes over time within the patient.

Methods

Participants and procedures

This is a retrospective observational study conducted at the Amsterdam UMC location Vrije Universiteit Amsterdam, a tertiary referral hospital in the Netherlands, consisting of a convenience sample of patients diagnosed with diffuse glioma. Ethical approval for the use of clinical data for research purposes was granted by the Medical Ethics Review Committee (METc VUmc 2010.126) and written informed consent was obtained from the patients.

Between June 2010 and April 2021, patients who were scheduled for elective tumor surgery were referred for assessment to the Department of Medical Psychology by their treating physician as part of routine clinical care. Both before and one year after surgery, patients conducted an assessment consisting of questionnaires on fatigue, depression, self-perceived cognitive functioning, brain tumor-specific symptoms, and HRQoL. If a patient was scheduled for re-resection, again assessments were conducted before and one year after re-resection. For the current study, patients who had conducted at least one assessment were included in the so-called **Glioma cohort**. Patients in the Glioma cohort conducted a varying number of assessments (between one and six) as the result of re-resection or loss to follow-up. Assessments with a missing fatigue questionnaire were excluded. Additionally, to assess fatigue longitudinally, a selection of patients in the *Glioma cohort* were included in the **Longitudinal subgroup**. These patients had performed both an assessment 0 to 6 months before surgery and an assessment 6 to 24 months after that same surgery.

Patient-reported outcome measures

Fatigue was assessed with the Checklist Individual Strength (CIS), a 20-item questionnaire answered on a seven-point Likert scale focusing on symptoms in the past two weeks.[14] The questionnaire consists of four subscales: fatigue severity (CIS-fatigue), concentration problems, reduced motivation, and reduced activity level. A cut-off of ≥35 on the CIS-fatigue subscale was used to indicate severe fatigue. The questionnaire and cut-off score have been validated in cancer survivors.[14] A change of ≥7 points on the CIS-fatigue subscale was defined as clinically relevant improvement or deterioration.[15]

Depression was assessed with the Center for Epidemiologic Studies Depression questionnaire (CES-D).[16] The Medical Outcomes Study Cognitive Functioning Scale (MOS-cog) was used to measure *self-perceived cognitive functioning.*[17] *Health-related quality of life* was assessed with the Medical Outcomes Study Short-Form Health Survey (SF-36).[18] The SF-36 is organized into nine scales assessing physical functioning, bodily pain, role limitations due to physical health problems, role limitations due to personal or emotional problems, emotional well-being, social functioning, energy/vitality, general health perceptions, and changes in health. In this study, we did not include the energy/vitality subscale. *Brain tumor-specific symptoms* were assessed with the European Organization for Research and Treatment of Cancer brain tumor module (BN20) questionnaire.[19] We only included the items future uncertainty, motor dysfunction, and seizures in the analyses, because the other items were already assessed with the other questionnaires. All questionnaires were recoded and scored according to the relevant scoring manuals. A higher score of the CIS, CES-D, and BN-20 reflects more symptoms or problems, and a lower score of the MOS-cog and the SF-36 reflects more disability and worse functioning.

Statistical analysis

The scripts for the statistical analyses are available in an online repository on Github (https://github.com/multinetlab-amsterdam/projects/tree/master/fatigue_glioma_manuscript_2022). These scripts can be run with the provided mock data in the script. Analyses were conducted using Rstudio (version 4.0.3).[20] Two-sided *p*-values < 0.05 were considered significant. First, we presented the percentages of the assessments and patients who were classified as severely fatigued based on the CIS-fatigue cut-off score. To clarify, in the *Glioma cohort* the presented percentages reflect the number of assessments indicative of severe fatigue, and not the number of patients since multiple assessments per patient were included. In the *Longitudinal subgroup,* the percentages do reflect the number of fatigued patients preoperatively and postoperatively.

Our primary aim was to identify which demographic, tumor- and, treatment-related characteristics and PROMs were associated with fatigue severity in the *Glioma cohort* with a linear mixed model. Patients were included as random intercepts, so we could control for the inclusion of multiple assessments per patient. We took three distinct steps: first, with univariate linear mixed models we evaluated associations between the CIS-fatigue score as a continuous variable and age, sex, education, KPS, location of the

tumor, histology (oligodendroglioma, astrocytoma, oligoastrocytoma, glioblastoma), isocitrate dehydrogenase (IDH) status (IDH mutation versus wildtype), phase of the disease (defined as pre-surgery, during chemoradiation and/or chemotherapy, stable disease, or progressive disease), the timing of the assessment (preoperative or postoperative), dexamethasone use (yes or no), radiotherapy before the assessment, chemotherapy before the assessment, and all the PROMs. Secondly, all above mentioned variables that showed a statistically significant association with fatigue were included in a linear mixed model as independent variables. As a third and final step, we performed backward selection based on the so-called Akaike Information Criterion to find the best-fitting model. After this, the model consisted of only variables that are significantly associated with fatigue severity. Together these variables can explain a proportion of the variance in the level of fatigue, demonstrated with an R^2 value.

Our secondary aim was to find predictors of postoperative fatigue. The within-subject data of the *Longitudinal subgroup* was used to compute a multiple linear regression model with the postoperative CIS-fatigue score as the dependent variable and the clinical characteristics and preoperative PROMs as potential predictors. We used the same three-step approach as described above. First, we evaluated univariate associations between the postoperative CIS-fatigue score and the individual variables. Subsequently, all variables that were associated with the postoperative CIS-fatigue score were used to compute a multiple linear regression model. After backward selection, the final model included only clinical characteristics and preoperative PROMs that were statistically significant predictors of postoperative fatigue severity. Additionally, to assess within-subject similarity between preoperative and postoperative CIS-fatigue scores in the *Longitudinal subgroup*, we computed a two-way random effects single measure intraclass correlation coefficient with absolute agreement (ICC 2,1).[21] The 95% confidence interval of the ICC was classified as poor (<0.5), moderate (0.5 – 0.75) or good (>0.75).[22]

Results

In this study, 222 patients with diffuse glioma were included in the *Glioma cohort* (Figure 1). They had performed 343 assessments, of which ten were excluded because the CIS questionnaire was missing. Of the 333 assessments, 193 were conducted preoperatively and 116 postoperatively. The mean age of the 222 patients in the *Glioma cohort* was 45 years and 57% were male. The majority of tumors were located in the left hemisphere (62%) and the frontal lobe (60%) and were classified as a WHO grade II or III (Table 1). Patients in the *Glioma cohort* performed between 1 and 6 assessments. Of the *Glioma cohort,* seventy patients were included in the *Longitudinal subgroup* (Figure 1, Table 1). Almost all of these patients had a WHO grade II or III tumor. See Table S1 for an overview of the PROMs in the *Glioma cohort* and the *Longitudinal subgroup.*

Prevalence of fatigue in the Glioma cohort and the Longitudinal subgroup

Of all assessments in the *Glioma cohort*, 39% (130/333) were indicative of severe fatigue (Figure 2). Of the preoperative assessments 38% were classified as severely fatigued, and 41% of the postoperative assessments. During treatment, 37% of the assessments (30/81) were indicative of severe fatigue, and 43% (33/76) of the assessments in patients with stable disease and 38% (9/24) of the assessments in patients with progressive disease. In the *Longitudinal subgroup* preoperatively 34% of the patients and postoperatively 40% of the patients were severely fatigued.

Explaining variance in fatigue severity in the Glioma cohort

In the *Glioma cohort*, depressive symptoms, four HRQoL domains, and having a tumor in the right hemisphere were significantly associated with fatigue severity (CIS-fatigue score), and explained a large proportion of variance in fatigue severity between patients. First, with univariate comparisons, we determined that age, dexamethasone use, chemotherapy, radiotherapy, tumor grade, histology, involved lobes, IDH status, phase of the disease, and timing of the assessment were not associated with fatigue severity (Table 2). On the other hand, depressive symptoms, self-perceived cognitive functioning, brain-tumor related symptoms, all eight HRQoL domains, sex, education, tumor hemisphere, and KPS were independently associated with fatigue severity. These significantly associated variables were included in a linear mixed model, and after backward selection, the final

model included depressive symptoms, four HRQoL domains and having a tumor in the right hemisphere (Table 3). These six variables were significantly associated with fatigue severity and together explained 63% of the variance in fatigue severity (marginal $R^2 = 0.63$).

Within-subject changes and predictors of postoperative fatigue

In the *Longitudinal subgroup*, fatigue improved postoperatively in 24% of the patients (17/70) and worsened in 30% (21/70, Figure 3). Within-patient similarity between the preoperative and postoperative fatigue severity was poor to moderate (ICC = 0.522, 95% CI 0.329 - 0.673), indicating that the level of fatigue indeed often fluctuated within the patient.

Moreover, when patients experience high levels of fatigue preoperatively or lowered physical functioning (SF36-PF) preoperatively, these patients are more likely to experience high levels of fatigue one year postoperatively. First, with univariate comparisons, we found that demographic and tumor- and treatment-related variables were not predictive of postoperative fatigue severity (Table S2), but that the preoperative fatigue subscales, depressive symptoms, self-perceived cognitive functioning, brain-tumor related symptoms, and HRQoL domains were significant predictors of postoperative fatigue severity. After combining these significantly associated variables in one multiple linear regression model and applying backward selection, the final model identified preoperative fatigue severity (estimated effect 0.32, $p = 0.02$) and preoperative physical functioning (estimated effect -0.26, $p = 0.02$) as significant predictors of postoperative fatigue (adjusted $R^2 = 0.31$).

Discussion

The current study aimed to identify which demographic, tumor- and treatment-related characteristics and PROMs were associated with fatigue severity in a large sample of glioma patients and which factors were predictive of postoperative fatigue in a subgroup of patients with both preoperative and postoperative assessments. We showed that a large proportion of the variance in fatigue severity could be explained by patient-reported limitations in HRQoL, depressive symptoms, and a right-sided tumor. Preoperative fatigue and physical functioning were independent predictors of postoperative fatigue.

Interestingly, there were no direct associations between fatigue severity and age, tumor type, treatment characteristics, phase of the disease, or timing of the assessment. These results corroborate studies in brain tumor patients that found no link between fatigue and disease duration, type of neurosurgical intervention, and chemo- and radiotherapy.[8-10] Also in cancer survivors and patients with chronic disease, the persistence of fatigue is not determined by tumor- and treatment-related characteristics.[11,23-25] More importantly, sleep disturbances, reduced physical functioning and activity, and lower self-efficacy concerning fatigue, do seem to explain the differences between fatigued and non-fatigued patients.[11,13,26] In line with these studies, we found limitations in HRQoL and depressive symptoms to be associated with fatigue, rather than tumor- and treatment-related factors. Functional and physical impairments have been identified as important determinants of fatigue and HRQoL before.[8,27,28] Similarly, depression is also known to be associated with fatigue in patients with cancer, meningioma, and neurological disease, with generally high correlations between the two.[29-31] However, it does remain to be elucidated whether fatigue is a direct result of depression and diminished HRQoL, or whether the association between the two is bidirectional or even reversed. Future studies should focus on why fatigue persists or emerges in some patients, while it decreases in others, for example through the incorporation of patients' daily life reports in research using ecological momentary assessments.[32]

We found tumors in the right hemisphere to be associated with elevated levels of fatigue. Tumor laterality has inconsistently been shown to be associated with HRQoL and symptoms like fatigue, depression and anxiety.[33-35] Also in stroke patients, there is no conclusive evidence on the association between fatigue and depression, and hemisphere and lesion location.[36-38] A recent study showed right-

hemisphere gliomas to be associated with worse HRQoL, compared to left-hemisphere tumors.[39] The authors hypothesized that tumor infiltration in areas involved with controlling empathy could result in higher self-focus with diminished HRQoL as an effect. Several models explain the possible hemispheric lateralization of emotions, for example how primary emotions are controlled by the right hemisphere and social emotion by the left hemisphere.[40] Perhaps these proposed mechanisms could also explain the association between fatigue and tumor laterality.

In the *Longitudinal subgroup*, about 40% of the patients reported severe fatigue preoperatively, and half of the patients showed clinical improvement or deterioration of the CIS-fatigue score. We identified preoperative fatigue and physical functioning as independent predictors of postoperative fatigue. Fatigue at baseline has been identified as an important predictor of fatigue at follow-up in cancer patients before.[41] In line with our study, research in women with breast cancer found that preoperative fatigue and lower role functioning before surgery, and not cancer characteristics, were associated with heightened postoperative fatigue levels.[42] Clinically, these results underline that preoperative identification of glioma patients suffering from fatigue or impaired physical functioning is important. Potentially, preoperative fatigue treatment and symptom management could also result in less symptom burden later on. However, this remains to be investigated with intervention studies with adequate follow-up.

Throughout this study, we have made a distinction between PROMs and demographic, and tumor- and treatment-related factors. This distinction is of importance in symptom management and when developing interventions targeting fatigue. Even though tumor- and treatment-related factors are important in disease management, these factors are usually non-modifiable and therefore not suitable as targets in ameliorating fatigue.[43] On the other hand, factors such as depression, social and physical functioning and health perception are modifiable to some extent and therefore could be relevant targets. Moreover, these results call for the timely and integrative assessment and treatment of fatigue, psychological symptoms and impaired HRQoL in glioma patients. Additionally, it could be of interest to understand whether mastery, or sense of control, coping style and catastrophizing are associated with fatigue in glioma patients. It is already known, that among patients with cancer a high sense of personal mastery is associated with less fatigue.[44] Also, decreased catastrophizing is a mediator in trials

investigating the effectiveness of cognitive-behavioral therapy targeting fatigue. Currently, there are no evidence-based interventions targeting fatigue in glioma patients.[7] However, there are several effective interventions targeting fatigue in cancer patients, including cognitive behavioral therapy, mindfulness, exercise, activity enhancement, problem-solving therapy, and acceptance and commitment therapy.[24,45] Those interventions might also be promising in treating fatigue in patients with glioma.[45-47] Unfortunately, we did not assess whether antiepileptic drugs (AEDs) were associated with fatigue, since some studies do suggest that AEDs could result in fatigue.[5,10,48] Therefore it might be worthwhile to investigate whether switching between AEDs or lowering the dosage would be a useful approach to ameliorate fatigue.[48]

An important limitation of this study is the potential selection bias introduced by the procedure of active referral. Patients included in this study were referred to the Department of Medical Psychology by their treating physician, and therefore this study consists of a convenience sample. It is likely that physicians more often referred younger patients, patients with tumors in eloquent areas, or patients scheduled for awake craniotomy. Furthermore, caution is warranted since not all patients participated in follow-up assessments. Almost all of the patients included in the *Longitudinal subgroup* had a WHO grade II or III tumor, hence these longitudinal results are mainly applicable to patients with a lower grade tumor. Additionally, we acknowledge that a wide range of questionnaires and PROMs are available to address symptoms, all with varying psychometric properties. With regards to the CIS questionnaire, it is unknown how changes in the CIS score during a follow-up period reflect actual changes in everyday life in the individual patient. The CIS questionnaire does have high test-retest reliability (r=0.74-0.86) but unfortunately has only been validated in cancer survivors, and not in glioma patients.[14] In the Dutch population,15% of the patients without chronic disease report severe fatigue based on the CIS-fatigue score,[11] compared to around 40% in the current study, indicating the prevalence of fatigue is indeed much higher in glioma patients. However, it would be of value to replicate the current results using other questionnaires and PROMs.

In conclusion, fatigue is a highly prevalent symptom throughout the disease in patients with glioma. Tumor type, treatment-related factors, the timing of the assessment, and phase of the disease were not associated with fatigue severity, whilst physical functioning, physical role limitations, general health perceptions, social functioning, depressive symptoms, and a right-sided tumor were. In lower grade

glioma patients, levels of fatigue vary substantially over time, and preoperative fatigue and physical functioning are important predictors of fatigue one year after surgery. The current study demonstrates that fatigue is a complex symptom, that should not solely be contributed to the tumor itself or its treatment but is related to a broad range of symptoms. Since the current study included patients with predominantly lower grade tumors, we recommend corroborating the current findings in a sample with more glioblastoma patients with an extensive longitudinal approach. In clinical care, the interplay between fatigue, depression, and HRQoL factors, should be taken into account when providing care for fatigued glioma patients and when designing interventions to target fatigue.

Abbreviations

Antiepileptic drugs	AEDs
The European Organization for Research and Treatment of Cancer brain tumor module	BN20
Center for Epidemiologic Studies Depression	CES-D
Checklist Individual Strength	CIS
Checklist Individual Strength subscale fatigue severity	CIS-fatigue
Health-related quality of life	HRQoL
Intraclass correlation coefficient	ICC
Isocitrate dehydrogenase	IDH
Medical Outcomes Study Cognitive Functioning Scale	MOS-cog
Patient-reported outcome measure	PROM
Medical Outcomes Study Short-Form Health Survey	SF-36

References

1.   Taphoorn MJ, Sizoo EM, Bottomley A. Review on quality of life issues in patients with primary brain tumors. *Oncologist.* 2010; 15(6):618-626.

2.   Gustafsson M, Edvardsson T, Ahlström G. The relationship between function, quality of life and coping in patients with low-grade gliomas. *Support Care Cancer.* 2006; 14(12):1205-1212.

3.   M IJ-K, Snijders TJ, de Graeff A, Teunissen S, de Vos FYF. Prevalence of symptoms in glioma patients throughout the disease trajectory: a systematic review. *Journal of neuro-oncology.* 2018; 140(3):485-496.

4.   van Coevorden-van Loon EMP, Coomans MB, Heijenbrok-Kal MH, Ribbers GM, van den Bent MJ. Fatigue in patients with low grade glioma: systematic evaluation of assessment and prevalence. *Journal of neuro-oncology.* 2017; 133(2):237-246.

5.   Valko PO, Siddique A, Linsenmeier C, Zaugg K, Held U, Hofer S. Prevalence and predictors of fatigue in glioblastoma: a prospective study. *Neuro-Oncology.* 2014; 17(2):274-281.

6.   Osoba D, Brada M, Prados MD, Yung WK. Effect of disease burden on health-related quality of life in patients with malignant gliomas. *Neuro Oncol.* 2000; 2(4):221-228.

7.   Asher A, Fu JB, Bailey C, Hughes JK. Fatigue among patients with brain tumors. *CNS Oncol.* 2016; 5(2):91-100.

8.   Armstrong TS, Cron SG, Bolanos EV, Gilbert MR, Kang DH. Risk factors for fatigue severity in primary brain tumor patients. *Cancer.* 2010; 116(11):2707-2715.

9.   van Coevorden-van Loon EMP, Heijenbrok-Kal MH, Horemans HLD, et al. The relationship between mental fatigue, cognitive functioning, and employment status in patients with low-grade glioma: a cross-sectional single-center study. *Disability and Rehabilitation.* 2021:1-7.

10.   Struik K, Klein M, Heimans JJ, et al. Fatigue in low-grade glioma. *Journal of neuro-oncology.* 2009; 92(1):73-78.

11.   Goërtz YMJ, Braamse AMJ, Spruit MA, et al. Fatigue in patients with chronic disease: results from the population-based Lifelines Cohort Study. *Scientific Reports.* 2021; 11(1):20977.

12.   Kluger BM, Krupp LB, Enoka RM. Fatigue and fatigability in neurologic illnesses: proposal for a unified taxonomy. *Neurology.* 2013; 80(4):409-416.

13.   Menting J, Tack CJ, Bleijenberg G, et al. Is fatigue a disease-specific or generic symptom in chronic medical conditions? *Health Psychol.* 2018; 37(6):530-543.

14. Worm-Smeitink M, Gielissen M, Bloot L, et al. The assessment of fatigue: Psychometric qualities and norms for the Checklist individual strength. *Journal of psychosomatic research.* 2017; 98:40-46.

15. Sobral Rebelo PF, Oliveira A, Paixão C, Marques A. Minimal clinically important difference for measures of fatigue, cough and sputum. *European Respiratory Journal.* 2019; 54(suppl 63):PA1216.

16. Radloff LS. The CES-D Scale:A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement.* 1977; 1(3):385-401.

17. Stewart A WJ, Jr, Sherbourne C, Wells K. Durham, NC. *Psychological distress/well-being and cognitive functioning measures.*: Duke University Press; 1992.

18. Aaronson NK, Muller M, Cohen PD, et al. Translation, validation, and norming of the Dutch language version of the SF-36 Health Survey in community and chronic disease populations. *Journal of clinical epidemiology.* 1998; 51(11):1055-1068.

19. Taphoorn MJ, Claassens L, Aaronson NK, et al. An international validation study of the EORTC brain cancer module (EORTC QLQ-BN20) for assessing health-related quality of life and symptoms in brain cancer patients. *European journal of cancer (Oxford, England : 1990).* 2010; 46(6):1033-1040.

20. (2020). RT. RStudio, PBC, Boston, MA. RStudio: Integrated Development Environment for R. 2020.

21. Matthias Gamer, Jim Lemon, Singh IFP. irr: Various Coefficients of Interrater Reliability and Agreement. 2019.

22. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *Journal of Chiropractic Medicine.* 2016; 15(2):155-163.

23. Schmidt ME, Chang-Claude J, Seibold P, et al. Determinants of long-term fatigue in breast cancer survivors: results of a prospective patient cohort study. *Psychooncology.* 2015; 24(1):40-46.

24. Bower JE. Cancer-related fatigue—mechanisms, risk factors, and treatments. *Nature Reviews Clinical Oncology.* 2014; 11(10):597-609.

25. Maass SWMC, Brandenbarg D, Boerman LM, Verhaak PFM, de Bock GH, Berendsen AJ. Fatigue among Long-Term Breast Cancer Survivors: A Controlled Cross-Sectional Study. *Cancers.* 2021; 13(6):1301.

26. Nap-van der Vlist MM, Dalmeijer GW, Grootenhuis MA, et al. Fatigue among children with a chronic disease: a cross-sectional study. *BMJ Paediatrics Open.* 2021; 5(1):e000958.

27. Taveira FM, Braz NFT, Comini-Frota ER, Teixeira AL, Domingues RB. Disability as a determinant of fatigue in MS patients. *Arq Neuropsiquiatr.* 2019; 77(4):248-253.

28. Sturgeon JA, Darnall BD, Kao M-CJ, Mackey SC. Physical and Psychological Correlates of Fatigue and Physical Function: A Collaborative Health Outcomes Information Registry (CHOIR) Study. *The Journal of Pain.* 2015; 16(3):291-298.e291.

29. Brown LF, Kroenke K. Cancer-related fatigue and its associations with depression and anxiety: a systematic review. *Psychosomatics.* 2009; 50(5):440-447.

30. Skorvanek M, Gdovinova Z, Rosenberger J, et al. The associations between fatigue, apathy, and depression in Parkinson's disease. *Acta Neurol Scand.* 2015; 131(2):80-87.

31. van der Linden SD, Gehring K, Rutten GM, Kop WJ, Sitskoorn MM. Prevalence and correlates of fatigue in patients with meningioma before and after surgery. *Neuro-oncology practice.* 2020; 7(1):77-85.

32. Kampshoff CS, Verdonck-de Leeuw IM, van Oijen MG, Sprangers MA, Buffart LM. Ecological momentary assessments among patients with cancer: A scoping review. *Eur J Cancer Care (Engl).* 2019; 28(3):e13095.

33. Liu R, Page M, Solheim K, Fox S, Chang SM. Quality of life in adults with brain tumors: Current knowledge and future directions. *Neuro-Oncology.* 2009; 11(3):330-339.

34. Palese A, Lamanna F, Di Monte C, Calligaris S, Doretto M, Criveller M. Quality of life in patients with right- or left-sided brain tumours: literature review. *J Clin Nurs.* 2008; 17(11):1403-1410.

35. Mainio A, Hakko H, Niemelä A, Tuurinkoski T, Koivukangas J, Räsänen P. The effect of brain tumour laterality on anxiety levels among neurosurgical patients. *J Neurol Neurosurg Psychiatry.* 2003; 74(9):1278-1282.

36. Kutlubaev MA, Duncan FH, Mead GE. Biological correlates of post-stroke fatigue: a systematic review. *Acta Neurologica Scandinavica.* 2012; 125(4):219-227.

37. Ulrichsen KM, Kolskår KK, Richard G, et al. Structural brain disconnectivity mapping of post-stroke fatigue. *Neuroimage Clin.* 2021; 30:102635-102635.

38. Bhogal SK, Teasell R, Foley N, Speechley M. Lesion Location and Poststroke Depression. *Stroke.* 2004; 35(3):794-802.

39. Fortin D, Iorio-Morin C, Tellier A, Goffaux P, Descoteaux M, Whittingstall K. High-Grade Gliomas Located in the Right Hemisphere Are Associated With Worse Quality of Life. *World Neurosurgery.* 2021; 149:e721-e728.

40. Ross ED. Differential Hemispheric Lateralization of Emotions and Related Display Behaviors: Emotion-Type Hypothesis. *Brain Sci.* 2021; 11(8).

41. Susanne K, Michael F, Thomas S, Peter E, Andreas H. Predictors of fatigue in cancer patients: a longitudinal study. *Supportive Care in Cancer.* 2019; 27(9):3463-3471.

42. Rotonda C, Guillemin F, Bonnetain F, Velten M, Conroy T. Factors associated with fatigue after surgery in women with early-stage invasive breast cancer. *Oncologist.* 2013; 18(4):467-475.

43. Drijver AJ, Oort Q, Otten R, Reijneveld JC, Klein M. Is poor sleep quality associated with poor neurocognitive outcome in cancer survivors? A systematic review. *J Cancer Surviv.* 2022.

44. Jacobsen PB, Andrykowski MA, Thors CL. Relationship of catastrophizing to fatigue among women receiving treatment for breast cancer. *J Consult Clin Psychol.* 2004; 72(2):355-361.

45. Boele FW, Klein M, Reijneveld JC, Verdonck-de Leeuw IM, Heimans JJ. Symptom management and quality of life in glioma patients. *CNS Oncol.* 2014; 3(1):37-47.

46. Röttgering JG, Douw L, de Witt Hamer PC, et al. Reducing severe fatigue in patients with diffuse glioma: a study protocol for an RCT on the effect of blended cognitive behavioural therapy. *Trials.* 2022; 23(1):568.

47. Daun JT, Capozzi LC, Roldan Urgoiti G, et al. ACE-Neuro: A tailored exercise oncology program for neuro-oncology patients - Study protocol. *Contemp Clin Trials Commun.* 2022; 28:100925.

48. St Louis EK. Minimizing AED adverse effects: improving quality of life in the interictal state in epilepsy care. *Curr Neuropharmacol.* 2009; 7(2):106-114.

49. Louis DN, Ohgaki H, Wiestler OD, et al. The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol.* 2007; 114(2):97-109.

50. Allen M, Poggiali D, Whitaker K, Marshall TR, Kievit RA. Raincloud plots: a multi-platform tool for robust data visualization. *Wellcome Open Res.* 2019; 4:63.

*The longitudinal subgroup is
a subset of all 222 patients*

Glioma cohort: 222 patients	Longitudinal subgroup: 70 patients

Consists of the following patients and assessments:

1 assessment: 137 patients

- 1 preop: 103 patients
- 1 postop: 34 patients

2 assessments: 68 patients

- 1 preop, 1 postop: 57 patients
- 2 preop, 0 postop: 3 patients
- 0 preop, 2 postop: 8 patients

3 assessments: 10 patients

- 2 preop, 1 postop: 5 patients
- 1 preop, 2 postop: 4 patients
- 0 preop, 3 postop: 1 patient

4 assessments: 6 patients

- 2 preop, 2 postop: 4 patient
- 3 preop, 1 postop: 1 patient
- 1 preop, 3 postop: 1 patient

6 assessments: 1 patient

- 2 preop, 4 postop: 1 patient

Consists of the following patients and assessments:

2 assessments: 70 patients

- 1 preop, 1 postop after a single surgery: 70 patients

Figure 1. Included patients and assessments in *Glioma cohort* **and the** *Longitudinal subgroup*

Abbreviations: Preop, preoperative assessment; Postop, postoperative assessment

	Glioma cohort (N=222)	Longitudinal subgroup (n=70)
Age, mean (SD)	45.6 (13.3)	41.0 (11.6)
Sex, n (%)		
Male	126 (56.8%)	42 (60.0%)
Female	96 (43.2%)	28 (40.0%)
Level of education (Verhage), n (%)		
Low (1-4)	41 (18.5%)	11 (15.7%)
Middle (5)	75 (33.8%)	26 (37.1%)
High (6-7)	106 (47.7%)	32 (45.7%)
Missing		1 (1.4%)
Tumor hemisphere, n (%)		
Left	137 (61.7%)	46 (65.7%)
Right	81 (36.5%)	23 (32.9%)
Both	4 (1.8%)	1 (1.4%)
Tumor location, n (%)		
Frontal	132 (59.5%)	41 (58.6%)
Non-frontal	90 (40.5%)	29 (41.4%)
WHO grade, n (%)		
II	114 (51.4%)	47 (67.1%)
III	60 (27.0%)	20 (28.6%)
IV	48 (21.6%)	3 (4.3%)
Tumor type, n (%)		
Oligodendroglioma	74 (33.3%)	33 (47.1%)

	Glioma cohort (N=222)	Longitudinal subgroup (n=70)
Astrocytoma, IDH mutated	67 (30.2%)	24 (34.3%)
Astrocytoma, IDH wildtype	10 (4.5%)	3 (4.3%)
Astrocytoma, IDH status unknown	14 (6.3%)	3 (4.3%)
Oligoastrocytoma*	9 (4.1%)	4 (5.7%)
Glioblastoma, IDH mutated	5 (2.3%)	1 (1.4%)
Glioblastoma, IDH wildtype	21 (9.5%)	2 (2.9%)
Glioblastoma, IDH status unknown	22 (9.9%)	0 (0%)
Using anti-epileptic drugs, n (%)		
No	5 (2.3%)	1 (1.4%)
Yes	163 (73.4%)	52 (74.3%)
Missing	54 (24.3%)	17 (24.3%)
Number of included assessments per patient, n (%)		
1	137 (61.7%)	
2	68 (30.3%)	70 (100%)
3	10 (4.5%)	
4	6 (2.7%)	
6	1 (0.5%)	

Table 1. Sociodemographic and clinical characteristics *Glioma cohort* and *Longitudinal subgroup*

* Histological diagnosis of these patients was based on the 2007 WHO classification of the central nervous system tumors, and could not be re-classified because molecular markers were not available[49]

Abbreviations: SD, standard deviation; n, number.

Figure 2. Raincloud plots preoperative and postoperative Checklist Individual Strength

Each panel shows a raincloud plot, which is a combination of a scatterplot, a boxplot with median and interquartile range, and half a violin plot [50]. The grey lines between the dots of the scatterplots are combinations of scores of a preoperative and postoperative assessment within one patient. The grey dotted horizontal lines indicate the cut-off scores for the scales with available validated cut-off values. All panels include all assessments of the Glioma cohort. Panel A reflects the Checklist Individual Strength (CIS) total score. Panels B-E reflect the four CIS subscales. Abbreviations: CIS, checklist individual strength.

Variable	N^	Estimated Effect	Standard Error	*T* value	*P* value	*Included in first LMM*	*Included in final LMM*
SF36 subscales							
Physical functioning	324	-0.40	0.030	-13.3	<0.001	**Yes**	**Yes**
Role limitation - Physical	325	-0.19	0.013	-15.0	<0.001	**Yes**	**Yes**
General health perceptions	326	-0.34	0.029	-11.5	<0.001	**Yes**	**Yes**
Social functioning	331	-0.28	0.022	-12.9	<0.001	**Yes**	**Yes**
Role limitation - Emotional	322	-0.16	0.014	-10.9	<0.001	**Yes**	No
Mental health	328	-0.34	0.035	-9.5	<0.001	**Yes**	No
Bodily pain	327	-0.20	0.028	-7.1	<0.001	**Yes**	No
Change in health	332	-0.16	0.022	-7.2	<0.001	**Yes**	No
CES-D score depression	320	0.76	0.066	11.5	<0.001	**Yes**	**Yes**
Hemisphere, right	327	5.073	1.719	2.952	0.0035	**Yes**	**Yes**
KPS, 70-80 vs. 90-100	320	5.918	2.416	2.45	0.015	**Yes**	No
BN20 items/subscales							
Future uncertainty	322	0.22	0.025	8.99	<0.001	**Yes**	No
Motor dysfunction	328	0.35	0.039	9.02	<0.001	**Yes**	No
Seizures	331	0.12	0.023	5.25	<0.001	**Yes**	No
MOS-cog	332	-1.06	0.11	-9.43	<0.001	**Yes**	No
Female sex, versus male sex	333	5.95	1.63	3.65	<0.001	**Yes**	No
Verhage education, score 1-4 (low)	331						
Score 5 (middle)		-3.52	2.22	-1.58	0.12	No	
Score 6-7 (high)		-5.64	2.17	-2.59	0.01	**Yes**	No
Age	333	0.064	0.062	1.02	0.31	No	
Dexamethasone use, yes vs. no	322	0.36	2.68	0.13	0.89	No	
IDH status, wildtype vs. mutated	328	-2.42	2.60	-0.93	0.35	No	
Preoperative or postoperative assessment, yes vs. no	333	1.25	1.21	1.04	0.30	No	
Had chemotherapy before assessment, yes vs. no	333	1.34	1.66	0.81	0.42	No	

	N^					
Had radiotherapy before assessment, yes vs. no	333	2.04	1.44	1.41	0.16	No
Tumor location, frontal	333					
Temporal		-0.10	2.056	-0.05	0.96	No
Parietal		-1.44	2.424	-0.60	0.55	No
Parieto-temporal		1.28	4.996	0.26	0.80	No
Multi-focal and other		-6.29	5.114	-1.23	0.22	No
Histology, astrocytoma versus	333					
Oligodendroglioma		0.78	1.91	0.41	0.68	No
Oligoastrocytoma*		8.30	4.29	1.94	0.05	No
Glioblastoma		-1.40	2.23	-0.63	0.53	No
Disease status, baseline	333					
Treatment		-0.39	1.663	-0.23	0.82	No
Stable		1.32	1.562	0.85	0.40	No
Progression		-2.38	2.744	-0.87	0.39	No

Table 2. Associations between CIS-fatigue subscale and clinical factors and PROMs

For each variable, a linear mixed model was computed with CIS-fatigue as the dependent factor and patient as a random effect. For example, the SF-36 physical functioning subscale was significantly associated with the CIS-fatigue score. This variable was then included in the next step of the analysis and after backward selection remained included in the final linear mixed model.

^ N is the number of assessments included in the analysis

* Histological diagnosis of these patients was based on the 2007 WHO classification of the central nervous system tumors, and could not be re-classified because molecular markers were not available [49]

Abbreviations: CES-D, Center for Epidemiologic Studies Depression; CIS-fatigue, Checklist Individual Strength subscale fatigue severity; IDH, Isocitrate dehydrogenase; MOS-cog, Medical Outcomes Study Cognitive Functioning Scale; SF36, Medical Outcomes Study Short-Form Health Survey; BN20, The European Organization for Research and Treatment of Cancer brain tumor module.

Fixed effects	Estimated effect	Standard error	*T* value	*P* value
Intercept: CIS-fatigue	53.568	3.181	*16.8*	*<0.001*
SF-36 role limitation - physical	-0.086	0.015	*-5.8*	*<0.001*
SF-36 physical functioning	-0.169	0.032	*-5.2*	*<0.001*
SF-36 general health perceptions	-0.121	0.028	-4.3	*<0.001*
SF-36 social functioning	-0.054	0.026	-2.1	*0.039*
CES-D	0.234	0.069	*3.4*	*<0.001*
Right hemisphere	3.353	1.096	*3.1*	*0.003*

Table 3. HRQoL, depression, lesion in the right hemisphere associated with fatigue severity

This table shows the best-fit linear mixed model with six variables explaining a large proportion of variance in the CIS-fatigue score. Patient was included as a random effect to control for multiple measurements.

Abbreviations: CES-D, Center for Epidemiologic Studies Depression; CIS-fatigue, Checklist Individual Strength subscale fatigue severity; SF-36, Medical Outcomes Study Short-Form Health Survey.

Figure 3. Improvement and deterioration of CIS-fatigue scores in the *Longitudinal subgroup*

The horizontal and vertical grey dashed lines reflect the cut-off score of the CIS-fatigue of 35 points. Deteriorating is defined as an increase in CIS-fatigue of ≥ 7, clinically significant improvement as a decrease in CIS-fatigue of ≥ 7.[15]