Host-pathogen sympatry and differential transmissibility of *Mycobacterium tuberculosis complex*

Authors: Matthias I Gröschel[1,14]*, Francy J. Pérez-Llanos[2,14], Roland Diel[3,4], Roger Vargas Jr[1], Vincent Escuyer[5], Kimberlee Musser[5], Lisa Trieu[6], Jeanne Sullivan Meissner[6], Jillian Knorr[6], Don Klinkenberg[7], Peter Kouw[8], Susanne Homolka[2], Wojciech Samek[9,10], Barun Mathema[11], Dick van Soolingen[7], Stefan Niemann[2,15], Shama Ahuja[6], Maha R Farhat[1,13,15]*.

Affiliations:

[1]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, USA

[2]Molecular and Experimental Mycobacteriology, Research Center Borstel, Borstel, Germany

[3]Institute for Epidemiology, University Medical Hospital Schleswig-Holstein, Kiel, Germany

[4]Lungenclinic Grosshansdorf, Airway Research Center North (ARCN), Member of the German Center for Lung Research (DZL), Grosshansdorf, Germany

[5]Wadsworth Center, New York State Department of Health, Albany, New York, USA

[6]New York City Department of Health and Mental Hygiene, Bureau of Tuberculosis Control, Queens, New York

[7]National Tuberculosis Reference Laboratory, Center for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

[8]Department of Infectious Diseases, Public Health Service, Amsterdam, The Netherlands

[9]Department of Electrical Engineering and Computer Science, Technical University Berlin

[10]Department of Artificial Intelligence, Fraunhofer Heinrich Hertz Institute, Berlin, Germany

[11]Mailman School of Public Health, Columbia University, New York City, NY, USA

[12]German Center for Infection Research, Partner Site Hamburg-Lübeck-Borstel-Riems, Borstel, Germany

[13]Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital, Boston, MA, USA

[14,15]These authors contributed equally

Keywords

Tuberculosis, Transmission, Sympatry, Contact Tracing, Whole Genome Sequencing, Host-pathogen interaction

Correspondence: matthias_groeschel@hms.harvard.edu and maha_farhat@hms.harvard.edu

SUMMARY

The obligate human pathogen *Mycobacterium* tuberculosis complex (*Mtbc*) separates

genetically into nine lineages several of which demonstrate sympatry with their human host

i.e. have distinct and restricted patterns of geographical distribution globally.[1–3]

Geographically restricted *Mtbc* lineages have been hypothesized to be adapted to infect

and/or transmit among sympatric human hosts, *i.e.* to be niche specialists, but this is yet to

be confirmed while controlling for exposure, social networks and risk of disease after

exposure.[1,4] Here we show that strains of geographically restricted (*Mtbc* lineages

L1,L2$_{restricted}$,L3,L4$_{restricted}$,L5,L6 are intrinsically less transmissible than widespread *Mtbc*

lineages (L2$_{widespread}$,L4$_{widespread}$) across Western European and North American cosmopolitan

populations. Comparing transmissibility between sympatric and allopatric contact-pathogen

pairs, we find the first controlled evidence for a biological impact of sympatry between *Mtbc*

strains and their human hosts; allopatric host-pathogen exposures has a 38% decrease in

the odds of infection among contacts compared with sympatric exposures. We measure 10-

fold lower phagocytosis and growth rates of L6 geographically restricted strains compared to

L4$_{widespread}$ in *in vitro* allopatric macrophage infections. Long-term co-existence of *Mtbc*

strains and humans has resulted in differential transmissibility between allopatric and

sympatric hosts for strains of geographically restricted lineages. Understanding the specific

genetic and immunological underpinnings of sympatry in TB may inform rational vaccine

design and TB control.

INTRODUCTION

Humans and their ancestors have continuously inhabited an earth replete with a wide diversity of microbial species. While many of these species are harmless or even beneficial to humans, some have the ability to infect human tissues and inflict harm, including pathogenic species that evolved the concurrent ability to transmit between humans (e.g., via contact, droplets or airborne transmission). Until recently, the study of host-pathogen relationships at long-time scales was methodologically limited to model organisms in model environments, as direct investigation of naturally occurring pathogen adaptation and long-term generational effects was technically and logistically difficult.[5]

High-throughput whole genome sequencing (WGS) has since enabled the direct study of hundreds of thousands of pathogens sampled directly from their human host, including pathogens of the *Mycobacterium tuberculosis* complex (*Mtbc*). Strains of the genetic lineages of the *Mtbc* exhibit distinct geographical distributions across the world regions:[6] Tuberculosis (TB) disease in Europe and North-America is mainly caused by Lineage (L) 4 (Euro-American) strains and specifically the three sublineages 4.1.2/Haarlem, 4.10/PGG3 and L4.3/LAM.[2] *Mtbc* sublineage 4.1.1/X has mostly been isolated from the Americas and sublineages 4.1.3/Ghana, 4.6.1/Uganda and 4.6.2/Cameroon are largely restricted to sub-Saharan Africa.[1–3] TB disease in Asia is usually caused by the Beijing lineage (L2, more than 70% in China). [3,7]. In West-Africa, TB disease can be caused by *M. africanum* West African 1 and 2 (*Mtbc* Lin5 and 6), lineages phylogenetically ancestral to *Mtb* sensu stricto (Lineages 1-4 & L7). *Mtbc* L5 and L6 have to-date not been isolated outside of West-Africa.[8,9] Similarly, *Mtbc* L7 and 8 have only been isolated in East Africa.[10,11]

It has been suggested that as an obligate human pathogen, Mtbc, co-diverged with its human host and the resulting geographical spread of its genetic lineages is a result of co-evolution.[1,12] However, this would place bacterial evolution on a similar timescale as that of humans. This possibility has been questioned in other works including those that applied

formal tests for co-divergence to small contemporaneous samples of Mtb and representative

human Y chromosome data.[13–17] With or without co-divergence, studies describing the phylo-

geographical distribution of *Mtbc*, strains within geographically-restricted lineages of *Mtbc*

are hypothesized to preferentially infect local human hosts with a restricted range of

genotypes, while strains of other lineages are isolated worldwide, capable of infecting hosts

with a wide range of genotypes.[1–3,6,18–20] The geographic overlap of two distinct species is

also termed sympatry, but it has not been studied whether the long term co-existence of

Mtbc lineages and local hosts has resulted in any biological effect that could explain the

geographic restriction and global spread of the *Mtbc* lineages.

Intriguingly, stable pathogen-host ancestry associations among secondary TB cases have

also been observed in two cosmopolitan cities where mixing of host and pathogen

populations occurs, suggesting different transmission dynamics attributable to both pathogen

and host ancestry .[4,21] However, as *Mtbc* is an airborne pathogen, observations suggesting a

biological effect resulting from long term co-existence of host and pathogen have been

alternatively attributed to differential exposure to *Mtbc* lineages as a result of social mixing

patterns, genetic drift, or founder effects rather than an effect of host-pathogen sympatry.[1]

These studies often measured transmission using data derived only from individuals with TB

after the development of active disease and did not control for social mixing or other aspects

of exposure.

Here, we set out to study if the observed sympatry (geographic co-localization) between

Mtbc and local hosts has a measurable impact on the TB epidemic. To test the hypothesis

whether geographically restricted lineages can infect hosts with a restricted range of

genotypes based on geographic co-localization, while widespread lineages can infect hosts

with a wider range of genotypes, we leverage detailed public health contact tracing data and

whole genome sequencing from three low TB incidence cosmopolitan cities. We first study

transmissibility between geographically restricted and globally spread *Mtbc* strains while

controlling for founder effects, host factors, and extent of exposure. We then quantify the

effect of sympatry on transmissibility, and follow up the epidemiological analysis with *in vitro*

macrophage infection experiments.

RESULTS

A multi-site cosmopolitan cohort of tuberculosis cases and close contacts

A major barrier to investigating a biological role of sympatry in *Mtbc* is the limitation in size

and depth of contact tracing data directly measuring transmissibility in cosmopolitan

societies with mixing of human and pathogen populations. To address this, we built a large

cohort of TB cases (n = 5,256) and their contacts (n=28,889) from New York City (NYC),

Amsterdam, and Hamburg. The three settings routinely and systematically surveyed for TB

cases and contacts as part of TB control activities and performed whole genome sequencing

(WGS) for strains from nearly all active culture-positive TB cases in this cohort (>99%,

5,219/5,256). WGS allowed *Mtbc* lineage classification and informed transmission

assessment. In the study of transmissibility, we were able to control for founder effects by

considering close contacts of each active TB index case individually. We were able to control

for exposure by modeling transmissibility using variables of TB index case infectiousness

and contact exposure time.

After excluding extrapulmonary TB as potential source cases, TB index cases without linked

contacts, and removing contacts without significant exposure (see **Methods** for exposure

times considered significant by site), the transmission analysis cohort comprised 2,279

patients with pulmonary TB linked to 12,749 close contacts (**Methods, Table 1, Figure 1a**).

As extrapulmonary TB cases without clinical or bacterial evidence of pulmonary TB are not

thought to be contagious they were excluded as potentially transmitting index cases. The

median number of close contacts investigated per index TB case was respectively 4

contacts (IQR 2 – 6), 4 (IQR 2 – 8), and 3 (IQR 1 – 7) in NYC, Amsterdam, and Hamburg

(**Figure 1b**).

Patient characteristics differed across study sites

The three source cities differed in key drivers of TB case burden and transmission, including

birth in a country with high TB incidence, proportion of TB in the adolescent to young adult

age group, sputum *Mtbc* smear positivity, alcohol or substance use disorder, and

homelessness (**Table 1**).[22] On average across study sites, 69% of TB cases and 48% of

close contacts were born in a country with a high TB incidence country, defined as

>30/100,000 TB incidence based WHO's 2016 data;[23] TB patients and close contacts from

NYC were more likely to be born in a high-incidence country (Chi squared, p<0.001 for both

comparisons corrected for the false discovery rate). TB patients in NYC were older by 9 and

14 years on median compared with patients in Hamburg or Amsterdam, respectively

(Wilcoxon rank sum, p<0.001). TB patients from Hamburg were less often sputum smear

positive (Chi squared, p=0.002 for Amsterdam and p<0.001 NYC), but more often used

alcohol (p < 0.001 for both comparisons), or were homeless (p = 0.036 Amsterdam, p <

0.001 NYC). NYC TB cases more often used illicit substances (p = 0.009 Hamburg, p =

0.018 Amsterdam). Factors recognized to increase the contact risk of progression to active

TB after infection, including HIV and other immunosuppression, did not differ by city.

Diabetes was more common among TB cases in NYC (p < 0.001 for both comparisons). TB

patients in NYC were more often isoniazid mono-resistant than in Hamburg (p = 0.005);

there was no difference in multidrug resistance (MDR, resistance to isoniazid and rifampicin)

across the cities.

Geographically restricted lineages cause a measurable proportion of TB disease

burden

We next used the WGS data to characterize the circulating *Mtbc* lineages in the three cities.

Among the 5,087 isolate sequences from patients infected with *Mtbc sensu stricto* or *M.*

africanum (including extrapulmonary TB cases and those without linked contact information),

we identified 297,061 single nucleotide substitutions (SNSs) that allowed the calculation of a

maximum likelihood phylogeny (**Figure 1c**). Using canonical signature SNSs, the *Mtbc*

strains analyzed were classified into six of the nine known human-adapted *Mtbc* complex

lineages:[18] the most common being strains of the Euro-American Lineage (L4, n = 2,976,

59%), followed in order by strains of the East Asia Lineage (L2, n = 802, 16%), East Africa

Central Asia (L3, n = 709, 14%), and East African, The Philippines, rim of the Indian Ocean

lineages (L1, n = 551, 11%). A minority of strains belonged to *M. africanum* 1 (L5, n = 17) or

2 (L6, n = 32). We categorized *Mtbc* strains as geographically restricted either if their

Simpson's diversity index was ≤0.3 (ancestral branches of L2: 2.1 'Proto-Beijing', 2.1.2.2.1

'Asia Ancestral 2', and L4 sub-lineages: 4.11, 4.2.1.1, 4.3.i2, 4.5, 4.6.2.2 'Cameroon',

4.6.1.1.1 'Uganda')[18] or if they belonged to lineages found to be phylo-geographically

restricted in previous literature[1,12,18–20] (L1, L3, L5, L6 as well as the ancestral L2 branch

2.2.2 'Asia Ancestral 1') (**Suppl. Table S1**) and compared them to the remaining widespread

sublineages of L2 and L4.[1,2,18]

Of 2,279 pulmonary TB patients, 28% (n = 643) were infected with strains of a

geographically restricted lineage. The proportion of restricted strain infections remained

stable over the study period, and there was no association with patient age or sex (Chi

squared, p>0.17, **Figure 1d-e**). Strains of geographically restricted lineages were equally

likely to harbor isoniazid mono-resistance (p = 0.8) and slightly less likely to harbor MDR

compared with widespread strains (OR 0.98, CI_{95} 0.97 to 1, Wald p = 0.021), but the overall

rate of MDR was low (n = 60, 2.6%, **Figure 1c**).

Strains of geographically restricted *Mtbc* lineages differ in phylogenetic metrics of

transmissibility

We hypothesized that strains of geographically restricted lineages were less transmissible

than widespread strains in the cosmopolitan source communities. To investigate this, we

compared genetic diversity metrics between strains of geographically restricted and

widespread lineages using the total sample phylogeny. Terminal branch lengths, which

represent an upper bound of evolutionary time since transmission for each *Mtbc* case, were

significantly longer for strains of geographically restricted lineages (median 66 SNSs,

Interquartile Range (IQR) 22 to 111, vs 41 SNSs, IQR 10 to 76) for widespread lineages, p <

2.2×10^{-16} using Wilcoxon rank sum test, **Figure 2a**). This difference was driven by a

combination of the long terminal branch lengths of strains of L1 (median 80 SNSs, IQR 52 to

129), L2$_{restricted}$ (91 SNSs, IQR 50 to 133) and L4$_{restricted}$ (68 SNSs, IQR 22 to 109) compared

to shorter branch lengths for L2$_{widespread}$, L3, and L4$_{widespread}$ (medians 52 (IQR 26 to 78), 38

(IQR 6 to 91), 36 (IQR 7 to 74) SNSs, respectively) (**Suppl. Figure S1a**). Strains of L2$_{restricted}$

and L4$_{restricted}$ had longer terminal branch lengths than strains of L2$_{widespread}$ and L4$_{widespread}$.

Respectively ($p < 3.9 \times 10^{-8}$ Wilcoxon rank sum test for both comparisons). Possible-

transmission clusters were defined as strains within 12-SNSs distance of each other based

on an updated filtering scheme that allows to call genetic variation across ˜7% more

positions in the *Mtbc* genome compared to previous filtering scheme (**Methods**).[24] Clustering

was more common for strains of geographically widespread lineages than restricted lineages

(822/3,778 of widespread lineage strains clustered in 239/292 total clusters vs 240/1,309 of

restricted lineage strains clustered in 53/292 clusters, p = 0.035, Chi Squared, **Figure 2b**).

The geographically restricted lineages L5/L6 comprised one L6 12-SNS cluster with two

strains. The average size of clusters did not significantly differ between geographically

widespread lineages compared with restricted lineages (3.4 patients vs 4.5 patients per

cluster, p = 0.07, two-sample t-test, and in a sensitivity analysis using a 5-SNSs distance 3.5

patients vs 4.6 patients per cluster, p = 0.08).

Strains of geographically restricted *Mtbc* lineages demonstrate intrinsically lower

transmissibility

We isolated the effect of the infecting *Mtbc* lineage on the rate of infection among close

contacts from other exposure characteristics and contact susceptibility as follows: we

focused on TB infection as measured by a positive QuantiFERON®-TB Gold In-Tube test

(IGRA ≥0.35 IU/ml) among close contacts (**Methods**), and (1) modeled the count of new TB

infections in each index cases' group of close contacts as a quasi-Poisson distributed

random variable with the mean estimated as a linear function of *Mtbc* lineage and index case

characteristics and a subset of contact susceptibility characteristics that was available

across the three source sites, and (2) modelled the probability of TB infection of each

individual close contact as a function of index case infectiousness and a subset of contact

susceptibility variables using generalized estimating equation (GEE) with a logistic link

function (**Methods**). The latter approach allowed for more fine-grain control of contact

characteristics but is expected to have lower power than the quasi-Poisson model. Exposure

to a geographically restricted lineage strain was associated with a lower rate of TB infections

among close contacts (adjusted incidence rate ratio (IRR_{adj}) 0.86, CI_{95} 0.78 to 0.95, Wald P-

value = 0.004) (**Table 2**, **Figure 2c**). We adjusted for a higher proportion of close contacts

born in a high TB-incidence country as a confounder and found its effect on IGRA positivity

to be significant (IRR_{adj} 1.98, CI_{95} 1.75 to 2.25, Wald P-value < 0.001). Host predictors of TB

infection among close contacts that were significantly associated in the model included

younger age and sputum smear positivity of the index case (**Table 2, Figure 2c**). In the

logistic GEE, exposure to a geographically restricted lineage was associated with a lower

odds of TB infection among close contacts (OR_{adj} 0.82, CI_{95} 0.72 to 0.94, Wald P-value =

0.004), in agreement with the models above (**Suppl. Table S2**).

We similarly modeled the interplay of *Mtbc* lineage and host characteristics on the rate of

secondary active TB among close contacts using a quasi-Poisson model. Secondary

pulmonary and extrapulmonary cases were linked to the source case based on a Bayesian

posterior probability of transmission ≥0.5 estimated using Phybreak that uses genetic

distance, sampling interval, and priors on incubation period and within-host pathogen

dynamics (**Methods**).[25] Phybreak estimated that 364 of the 2,279 pulmonary TB index cases

had one close contact progress to active disease, 53 had two close contacts progress, and

10 had three progress (**Figure 2d**). Most of these transmitting TB index cases (77%,

322/427) were infected with a geographically widespread *Mtbc* lineage strain. Exposure to a

geographically restricted lineage was associated with significantly lower risk of secondary

active TB cases among close contacts controlling for index characteristics (IRR_{adj} 0.55, CI_{95}

0.32 to 0.90, Wald P-value = 0.023) (**Table 2, Figure 2c**). In the logistic GEE, the exposure

to a geographically restricted lineage had a similar trend towards lower risk of secondary

active TB but was not statistically significant (OR_{adj} 0.77, CI_{95} 0.57 to 1.04, Wald P-value =

0.09) (**Suppl. Table S2**). We used the effect estimates of all variables of the quasi-Poisson

model of disease progression to calculate a host infectiousness score.[22] The infectiousness

score did not differ between index cases infected with strains of geographically restricted or

widespread *Mtbc* lineages (Wilcoxon signed-rank test, p = 0.072), supporting the conclusion

that host factors alone do not explain the observed differences in transmissibility (**Figure 2f**).

The above models only partially account for the effect of contact susceptibility on TB

transmission and disease progression, as data on contact characteristics was limited for

contacts in Hamburg and Amsterdam. We hence performed a secondary logistic GEE

analysis using only NYC data to confirm the association between geographically restricted

Mtbc lineages and intrinsic transmissibility controlling for the additional characteristics of

diabetes and HIV (n = 8,481 close contacts). The model confirmed young age, male sex,

and diabetes as independent predictors of secondary active TB among close contacts

(**Suppl. Table S3**). HIV status among contacts was not associated with secondary active TB

possibly because the majority of contacts had well controlled HIV (78% of all estimated

people with HIV in New York City were virally suppressed based on data from 2020).[26] Only

four contacts reported using illicit drugs and hence this variable was not included in the

model. Exposure to a geographically restricted *Mtbc* lineage was associated with lower odds

of secondary active TB than estimated using the logistic GEE on the larger cohort with

control for a subset of contact characteristics (OR_{adj} 0.54, CI_{95} 0.37 to 0.78, Wald P-value =

0.001 *vs.* OR_{adj} 0.77, CI_{95} 0.57 to 1.04, Wald P-value = 0.09). The effect estimate did not

reach statistical significance for the model predicting TB infection (**Suppl. Table S3**).

We next evaluated the specific contributions of *Mtbc* sub-lineages on the observed

differential transmissibility between geographically restricted and widespread strains across

the three cosmopolitan cities using the quasi-Poisson model. L1 strains were associated with

a lower rate of TB infections and secondary active cases among close contacts compared to

L4$_{\text{widespread}}$ strains (IRR$_{\text{adj}}$ 0.71, CI$_{95}$ 0.60 to 0.85, P-value < 0.001, and 0.23, CI$_{95}$ 0.05 to 0.69,

P-value = 0.027 respective) (**Figure 2e, Suppl. Table S4**). Although L3 strains

demonstrated high proportions of clustering based on genetic distance in the Netherlands

(**Suppl. Figure S1b**), the adjusted incidence rate ratio of L3 strains was 0.41 (CI$_{95}$ 0.16 to

0.88) indicating a lower rate of secondary active cases among close contacts compared with

L4$_{\text{widespread}}$ strains (P-value < 0.032, **Figure 2e, Suppl. Table S4**). We did not include close

contacts exposed to L5/L6 strains in models of secondary active TB due to the small

numbers with the outcome (one of the 136 close contacts developed secondary active TB).

Allopatric host-pathogen relationships decrease susceptibility of TB infection

The observation of lower transmissibility of strains of geographically restricted lineages in

Western cosmopolitan cities may relate to their poor ability to transmit among allopatric

human populations, i.e., where there has not been long-term host-pathogen co-localization.

To investigate this, we first mapped countries linked to geographically restricted *Mtbc*

lineages and assessed previously reported stable host ancestry-*Mtbc* lineage combinations

among the TB index cases of the present analysis (**Figure 3a-b**).[4,21] Seventy-five percent of

index cases were infected with a sympatric *Mtbc* geographically restricted strain, significantly

more than expected by chance (**Figure 3c-d**). To add validity to our definition of sympatry,

we evaluated the geographic origin and lineage of 25,243 *Mtbc* strains with publicly available

WGS (**Suppl. Figure S2a).** To test this co-localization hypothesis, we measured the

transmissibility of geographically restricted *Mtbc* lineages among exposed sympatric *vs.*

allopatric close contacts using contact place of birth as a proxy for host ancestry (**Methods**).

We used a logistic GEE to study the effect of contact sympatry on infection with a

geographically restricted *Mtbc* strain controlling for index and other contact characteristics

(**Figure 3e**). We excluded 108/12,749 close contacts with mixed ancestry, and 667/12,749

close contacts without information on country of birth, and excluded contacts exposed to a

geographically widespread lineage (L2$_{\text{widespread}}$ and L4$_{\text{widespread}}$). Exposure location (e.g.

household vs work) did not differ between allopatric and sympatric exposures (**Suppl. Table**

S5). Contact allopatry was associated with a lower odds of *Mtbc* strain infection among close

contacts of index cases infected with a geographically restricted lineage strain (OR_{adj} for

pooled L1,L3, L5,L6: 0.62, CI_{95} 0.43 – 0.91, n = 2556, p = 0.013, N by lineage: L1=1,080,

L3=1,353, L5=35, L6=88, **Figure 3f, Suppl. Figure S2b, Suppl. Table S6**) with similar

additional host factors associated with *Mtbc* strain infection as measured above in models

comparing geographically restricted strains across all contacts. The effect of contact

allopatry was stable when adding contacts exposed to $L2_{restricted}$ or $L4_{restricted}$, but extinguished

when assuming the modern widespread L2 sub lineages to be sympatric to East Asian

ancestry, in line with evidence of modern L2's global spread (Suppl. Table S6).[27] We found

the association of allopatry to *Mtbc* strain infection to be robust to ancestry misclassification

due to e.g. second-generation immigration in a sensitivity analysis (OR_{adj} 0.8, CI_{95} 0.65 –

0.99) (**Methods, Figure 3g**) including among the subset of data from NYC with the

additional contact susceptibility variables diabetes, and HIV (**Suppl. Table S7**).

Human macrophages differ in susceptibility to infection with *Mtbc* L4 and *M.*

***africanum* strains by self-reported ancestry**

We looked for evidence of a biological effect of sympatry using *in vitro* macrophage infection

models using strains of different *Mtbc* lineages. Human blood monocyte derived

macrophages (MDMs) were obtained from donors of self-identified German, Nigerian,

Cameroonian or Ghanaian ancestry (see **Figure 4a** for the approach). We compared the

Mtbc phagocytosis/uptake of two clinical isolates of the geographically widespread L4 LAM

lineage with that of two clinical strains of the geographically restricted *Mtbc* L6 lineage while

including the ATCC reference strains of each lineage in each group (**Suppl. Figure S3**). We

ensured equal infection doses by comparing colony forming units (CFUs) of the inoculum.

Macrophages of donors of self-reported ancestry co-localizing with L6 phagocytosed strains

of L6 on average at 30.4% compared to 2.9% among macrophages of donors whose self-

reported ancestry was Germany, not co-localizing with L6 (p < 0.001, unpaired T-test)

(**Figure 4b**). There was no significant difference in growth of L4 strains in macrophages of

donors of ancestry that do or do not co-localize with L6 (33.9% vs 29.5%, p = 0.8) (**Figure**

4c). This difference in uptake of L6 strains between L6 co-localizing and non-co-localizing

donors was driven by all three L6 lineages (all comparisons p < 0.001) (**Figure 4d**) while we

found both intra host and inter host variability in uptake efficiency of the three L4 and L6

strains (**Figure 4e, Suppl. Figure S4 and S5**).

We further investigated whether L4 or L6 strains differed in their intracellular growth kinetics

after uptake into macrophages of different ancestries over time (4h, 24h, 96h, and 168h post

infection). In macrophages from donors with self-reported ancestry of Nigeria, Cameroon, or

Ghana there was no significant difference in growth between L4 and L6 at two of the three

time points (**Figure 4f-h, Suppl. Fig S6**). However, growth in macrophages from these

donors was higher for L6 compared with L4 strains at 24hpi (comparing average growth

across the three strains for each lineage for the Ghanaian donor, p < 0.001, Cameroonian

donor p < 0.05, no difference in macrophages of the Nigerian donor) (**Figure 4 e-f**).

Conversely, in the macrophages from donors of self-reported German ancestry, L4 strains

grew at higher rates than L6 at all the time points tested (p < 0.05 at 24hpi, 96hpi and

168hpi, unpaired T-test and ANOVA with Bonferroni correction) (**Suppl. Figure S6**).

Overall, these results provide evidence that differential transmissibility between allopatric

and sympatric pairs may relate to the observed weaker uptake and growth of geographically

restricted lineages in allopatric macrophages than increased growth and uptake of these

lineages in sympatric macrophages. This suggests that sympatry predominantly manifests in

diminished infectivity of geographically restricted *Mtbc* lineages in allopatric hosts.

DISCUSSION

It is well established that pathogens of genetic lineages of the *Mycobacterium tuberculosis*

complex co-localize with human hosts, but it remains unclear if co-localization has any

biological effect for infection with *M. tuberculosis.* In this study, we provide the first evidence

for a measurable impact of sympatry on the TB epidemic in a controlled study. Our results

demonstrate intrinsically reduced transmissibility of strains of geographically restricted *Mtbc*

lineages compared with strains of widespread *Mtbc* lineages in three cosmopolitan cities.

While strains of geographically restricted lineages are able to infect sympatric hosts, they

display lower transmissibility in allopatric hosts. We validate our epidemiological

observations using a macrophage infection model, finding evidence that the reduced

transmissibility of geographically restricted strains may relate to lower uptake and growth of

Mtbc in allopatric macrophages after initial exposure. Our central discovery is that the

reduced infectivity of strains of geographically restricted *Mtbc* lineages in allopatric

populations is likely responsible for their limited success in driving the global spread of TB

compared to strains of modern widespread L2/L4 lineages.

Our results support the theory that geographic co-localization, or sympatry, between

pathogen and host can have biological implications and suggests that strains of

geographically restricted lineages face a barrier in their ability to migrate to other

geographical locations. Patterns of human migration alone are not sufficient in accounting for

this geographical spread seen today as evidenced by the little disease burden caused by

strains of the geographically restricted *Mtbc* lineages in the Americas. The ecological

observations of differing global strain diversity thus do not seem to be only the effect of

founder effects and human migration, but a differing human host range of *M. tuberculosis.*

Since bacterial uptake and transmissibility are preserved in sympatric vs allopatric pairs in

the analyses presented here suggests a specific role for the host-pathogen pair beyond *Mtbc*

lineage alone. Further, TB disease in India, East Africa, and many parts of South East Asia

is largely caused by geographically restricted lineages (L1 and L3),[1] indicating effective

transmission of strains of these lineages among local populations. A biological effect of

sympatry was also shown for *H. pylori*, a major risk factor of human gastric cancers. A recent

study revealed that while colonization with allopatric *H. pylori* strains was correlated with

severe gastric cancer, sympatric *H. pylori* infections carry the best prognosis.[28]

Previous attempts at quantifying the biological effect of sympatry in tuberculosis were limited

by measuring transmission at the index case level and often did not control for host level

variables resulting in inadequate control for social network structure, extent of exposure, and

susceptibility to disease. We used a Poisson framework to quantify the number of

transmissions among close contacts of each index case. Recent exposure is a substantial

risk for all contacts independent of their prior IGRA status (i.e., previous exposure) and TB

transmission generally requires significant contact time between a contact and an active

case.[29–31] Our modeling approach and data allowed us to control for contact exposure

between lineages and, separately, between sympatric and allopatric contacts because all

contacts had a known and similar high-risk exposure to *Mtbc*.

We validated our epidemiological findings in an in vitro macrophage infection assay and

found low efficiency in phagocytic uptake in macrophages from donors who did not co-

localize with the infecting strain. We cannot test the hypothesis that host ancestry related

genes are responsible for phenotypic difference in uptake as we did not obtain human

sequence data, yet recent work suggested that up to 9.3% of macrophage expressed genes

show ancestry-associated differences in the response to infection.[32] While more work is

warranted to compare more strain-host combinations, including interrogating the host

genetics, we had chosen L6 and L4 strains based on their phylogenetic distance and

availability of sympatric host macrophage donors. Despite causing a considerable proportion

of TB disease in West Africa, L6 strains are associated with reduced virulence in animal

models[33], HIV[34] and diabetes[35], and as such could represent a lineage of generally reduced

virulence.

Elucidating molecular mechanisms of resistance to *Mtbc* infection will take time, but our

findings can be used to improve existing models of TB susceptibility used in public health

practice. Although several patient risk factors (e.g., age, sex, HIV, and diabetes) are

currently recognized to influence individual risk of TB infection and disease after exposure,

they are rarely used quantitatively to support decisions in contact tracing or the initiation of

chemoprophylaxis after exposure. Here, we demonstrate that allopatric infections constitute

about 52% of total contact exposures to strains of geographically restricted *Mtbc* lineages in

low prevalence settings and influence the odds of infection to a similar extent as recognized

risk factors such as index case smear grade. The incorporation of *Mtbc* sympatry for

geographically restricted *Mtbc* lineages as well as host factors into the baseline assessment

of risk using statistical models may help refine the use of chemoprevention and guide

contact tracing activities. Moreover, identifying the components of Mtbc genetic variation

responsible for host specificity can inform vaccine design to elucidate pathogen targets of

host susceptibility.

The investigation of the effects of sympatry between humans and *Mtbc* strains requires a

diverse cohort that ensures both infections with sympatric and allopatric strains are

adequately represented in the sample. This study was enabled by the availability of robust

TB case data, pathogen WGS, and contact tracing data and illustrates that these data can

successfully be repurposed and pooled across several public health systems. Yet, this study

was not without limitations. The TB index case and contact information relied on social

contact tracing that may be subject to recall bias. Information on contacts of TB cases is not

routinely collected in Hamburg and Amsterdam and as a result some key risk factors of TB

susceptibility (contact HIV, contact diabetes) were not available. The analysis of secondary

active TB relied on modeling rather than epidemiological linkage which may include false

positive transmission links, and some contacts of TB index cases may not be visible to the

public health departments due to contact relocation between cities or states. We used

country of birth, or in case of minors, the parent's countries of birth, as a surrogate for

ancestry. As self-reported ancestry is a poor proxy for human genetics our approach did not

allow us to directly measure host-genetic adaptation in this study. We performed several

sensitivity analyses to address these limitations. We randomly assigned the sympatry label

for 14% of our sample (the proportion of Dutch residents who are foreign-born, Statistics

Netherlands) to account for ancestry misclassification. Estimated transmission links were

only accepted based on sample size and support value thresholds that were shown to

provide robust effect estimates in our study and other work.[36] We built models of

transmission at contact level in a subset of data where HIV and diabetes information was

available, as these are important effect modifiers of TB susceptibility. Finally, the differing

uptake efficiencies by macrophage donor – pathogen strain combinations may impact the

growth kinetics of *Mtbc* strains. To enable comparability, all macrophages were infected with

the same amount of *Mtbc* bacilli, and the infection dose was quantified by plating the

inoculum on agar plates for CFU counting.

In summary, we demonstrate an epidemiological impact of sympatry that manifests in lower

infectivity of geographically restricted *Mtbc* lineages in allopatric host-pathogen

combinations. This suggests that strains of widespread lineages are likely to increase in

prevalence relative to strains of geographically restricted lineages in a progressively

globalized world. Future work should aim to understand the molecular underpinnings of

allopatric versus sympatric infections.

Data availability

Bioprojects and accessions for each fastq file have been requested and will be available in
Supplementary table S8.

Acknowledgements

We thank Pascal Lapierre, Han de Neeling, and Thomas Kohl for transferring sequence
data, Henrieke Schimmel and Erika Slump for help with data linkage of the Dutch dataset,
the Dutch TB Registry (NTR commissie) as well as all study participants. We thank the
Wadsworth Center Applied Genomics Technology Cluster for whole genome sequencing.

Author contributions

MRF, and MIG conceived the idea for the epidemiological analysis. SN, SH, FJPL conceived
the idea for the *in vitro* experiments. MRF supervised the project. MIG performed data
curation and data analysis. MIG and MRF wrote the first draft. FJPL performed data curation
and data analysis. RVJr and DK carried out data analysis. LT, PK, RD carried out data
acquisition. VE, KM, JSM, SH, DvS, SA, SN supervised data acquisition and curation. WS
and BM critically reviewed the drafts. All authors reviewed the draft and assisted in the
manuscript preparation.

Funding

This work was funded by National Institutes of Health / National Institute of Allergy and
Infectious Diseases R21 AI154089 to MRF. MIG is supported by the German Research
Foundation (GR5643/1-1). FJPL, SN, and SH were funded by the Leibniz Science Campus
EvoLUNG (Evolutionary Medicine of the Lung) http://evolung.fz-471borstel.de/, grant number
W47/2019. SN received additional funding by the German Research Foundation under
Germanys Excellence Strategy – EXC 2167 Precision Medicine in Inflammation, and the
German Ministry of Education and Research (BMBF) for the German Center of Infection

REFERENCES

1.   Brites, D. & Gagneux, S. Co-evolution of Mycobacterium tuberculosis and Homo sapiens. *Immunol. Rev.* **264**, 6–24

(2015).

2.   Stucki, D. *et al.* Mycobacterium tuberculosis lineage 4 comprises globally distributed and geographically restricted

sublineages. *Nat. Genet.* **48**, 1535–1543 (2016).

3.   Merker, M. *et al.* Evolutionary history and global spread of the Mycobacterium tuberculosis Beijing lineage. *Nat. Genet.*

47, 242–249 (2015).

4.   Gagneux, S. *et al.* Variable host–pathogen compatibility in Mycobacterium tuberculosis. *Proc. Natl. Acad. Sci. U. S. A.*

103, 2869–2873 (2006).

5.   Woolhouse, M. E. J., Webster, J. P., Domingo, E., Charlesworth, B. & Levin, B. R. Biological and biomedical implications

of the co-evolution of pathogens and their hosts. *Nat. Genet.* **32**, 569–577 (2002).

6.   Hirsh, A. E., Tsolaki, A. G., DeRiemer, K., Feldman, M. W. & Small, P. M. Stable association between strains of

Mycobacterium tuberculosis and their human host populations. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 4871–4876 (2004).

7.   Roetzer, A. *et al.* Whole genome sequencing versus traditional genotyping for investigation of a Mycobacterium

tuberculosis outbreak: a longitudinal molecular epidemiological study. *PLoS Med.* **10**, e1001387 (2013).

8.   De Jong, B. C. *et al.* Use of spoligotyping and large sequence polymorphisms to study the population structure of the

Mycobacterium tuberculosis complex in a cohort study of consecutive smear-positive tuberculosis cases in The Gambia.

J. Clin. Microbiol. **47**, 994–1001 (2009).

9.   Asante-Poku, A. *et al.* Molecular epidemiology of Mycobacterium africanum in Ghana. *BMC Infect. Dis.* **16**, 385 (2016).

10. Ngabonziza, J. C. S. *et al.* A sister lineage of the Mycobacterium tuberculosis complex discovered in the African Great

Lakes region. *Nat. Commun.* **11**, 2917 (2020).

11. Firdessa, R. *et al.* Mycobacterial lineages causing pulmonary and extrapulmonary tuberculosis, Ethiopia. *Emerg. Infect.*

Dis. **19**, 460–463 (2013).

12. Comas, I. *et al.* Out-of-Africa migration and Neolithic coexpansion of Mycobacterium tuberculosis with modern humans.

Nat. Genet. **45**, 1176–1182 (2013).

13. Bos, K. I. *et al.* Pre-Columbian mycobacterial genomes reveal seals as a source of New World human tuberculosis.

Nature **514**, 494–497 (2014).

14. Kay, G. L. *et al.* Eighteenth-century genomes show that mixed infections were common at time of peak tuberculosis in

Europe. *Nat. Commun.* **6**, 6717 (2015).

15. Sabin, S. *et al.* A seventeenth-century Mycobacterium tuberculosis genome supports a Neolithic emergence of the

Mycobacterium tuberculosis complex. *Genome Biol.* **21**, 201 (2020).

16. Vågene, Å. J. *et al.* Geographically dispersed zoonotic tuberculosis in pre-contact South American human populations.

Nat. Commun. **13**, (2022).

17. Pepperell, C. S. *et al.* The role of selection in shaping diversity of natural M. tuberculosis populations. *PLoS Pathog.* **9**,

e1003543 (2013).

18. Freschi, L. *et al.* Population structure, biogeography and transmissibility of Mycobacterium tuberculosis. *Nat. Commun.*

12, (2021).

19.   Baker, L., Brown, T., Maiden, M. C. & Drobniewski, F. Silent nucleotide polymorphisms and a phylogeny for

Mycobacterium tuberculosis. *Emerg. Infect. Dis.* **10**, 1568–1577 (2004).

20.   Asante-Poku, A. *et al.* Mycobacterium africanum is associated with patient ethnicity in Ghana. *PLoS Negl. Trop. Dis.* **9**,

e3370 (2015).

21.   Reed, M. B. *et al.* Major Mycobacterium tuberculosis lineages associate with patient country of origin. *J. Clin. Microbiol.*

47, 1119–1128 (2009).

22.   Nebenzahl-Guimaraes, H., Borgdorff, M. W., Murray, M. B. & van Soolingen, D. A novel approach - the propensity to

propagate (PTP) method for controlling for host factors in studying the transmission of Mycobacterium tuberculosis. *PLoS*

One **9**, e97816 (2014).

23.   World Health Organization. *Global Tuberculosis Report 2017*. (2017).

24.   Marin, M. *et al.* Benchmarking the empirical accuracy of short-read sequencing across the M. tuberculosis genome.

Bioinformatics **38**, 1781–1787 (2022).

25.   Klinkenberg, D., Backer, J. A., Didelot, X., Colijn, C. & Wallinga, J. Simultaneous inference of phylogenetic and

transmission trees in infectious disease outbreaks. *PLoS Comput. Biol.* **13**, e1005495 (2017).

26.   *HIV surveillance annual report 2020*. chrome-

extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www1.nyc.gov/assets/doh/downloads/pdf/dires/hiv-surveillance-

annualreport-2020.pdf (2020).

27.   Holt, K. E. *et al.* Frequent transmission of the Mycobacterium tuberculosis Beijing lineage and positive selection for the

EsxW Beijing variant in Vietnam. *Nat. Genet.* **50**, 849–856 (2018).

28.   Kodaman, N. *et al.* Human and Helicobacter pylori coevolution shapes the risk of gastric disease. *Proc. Natl. Acad. Sci.*

U. S. A. **111**, 1455–1460 (2014).

29.   Gounder, P. P. *et al.* Risk for Tuberculosis Disease Among Contacts with Prior Positive Tuberculin Skin Test: A

retrospective Cohort Study, New York City. *J. Gen. Intern. Med.* **30**, 742–748 (2015).

30.   Getahun, H. *et al.* Management of latent Mycobacterium tuberculosis infection: WHO guidelines for low tuberculosis

burden countries. *Eur. Respir. J.* **46**, 1563–1576 (2015).

31.   Mathema, B. *et al.* Drivers of Tuberculosis Transmission. *J. Infect. Dis.* **216**, S644–S653 (2017).

32.   Nédélec, Y. *et al.* Genetic ancestry and natural selection drive population differences in immune responses to pathogens.

Cell **167**, 657-669.e21 (2016).

33.   Yeboah-Manu, D., de Jong, B. C. & Gehre, F. The biology and epidemiology of Mycobacterium africanum. *Adv. Exp.*

Med. Biol. **1019**, 117–133 (2017).

34.   de Jong, B. C. *et al.* Mycobacterium africanum: a new opportunistic pathogen in HIV infection? *AIDS* **19**, 1714–1715

(2005).

35.   Asante-Poku, A. *et al.* TB-diabetes co-morbidity in Ghana: The importance of Mycobacterium africanum infection. *PLoS*

One **14**, e0211822 (2019).

36.   Gygli, S. M. *et al.* Prisons as ecological drivers of fitness-compensated multidrug-resistant Mycobacterium tuberculosis.

Nat. Med. **27**, 1171–1177 (2021).

Table 1: Pulmonary tuberculosis patients characteristics.

	N	Overall N = 2,279[1]	Hamburg N = 498[1]	Netherlands N = 266[1]	New York City N = 1,515[1]
Sex	2,279				
Female		849 (37%)	191 (38%)	98 (37%)	560 (37%)
Male		1,430 (63%)	307 (62%)	168 (63%)	955 (63%)
Age	2,279	48 (31, 66)	43 (29, 61)	38 (26, 55)	52 (33, 68)
Country of birth	2,279				
High TB incidence		1,574 (69%)	224 (45%)	158 (59%)	1,192 (79%)
Medium/low TB incidence		705 (31%)	274 (55%)	108 (41%)	323 (21%)
Sputum microscopy	2,279				
Negative		809 (35%)	230 (46%)	90 (34%)	489 (32%)
Positive		1,470 (65%)	268 (54%)	176 (66%)	1,026 (68%)
Symptomatic	2,254	1,985 (88%)	448 (92%)	203 (81%)	1,334 (88%)
Homeless	2,275	72 (3.2%)	32 (6.5%)	7 (2.6%)	33 (2.2%)
Drug use	2,279	108 (4.7%)	13 (2.6%)	6 (2.3%)	89 (5.9%)
Alcohol use	2,272	137 (6.0%)	77 (16%)	5 (1.9%)	55 (3.6%)
HIV	1,975	77 (3.9%)	21 (4.7%)	8 (3.6%)	48 (3.7%)
Diabetes	2,279	418 (18%)	24 (4.8%)	22 (8.3%)	372 (25%)
Immunosuppression	1,829	44 (2.4%)	6 (3.4%)	6 (4.3%)	32 (2.1%)
Isoniazid monoresistance	2,278	242 (11%)	35 (7.0%)	25 (9.4%)	182 (12%)
Multidrug resistance	2,278	60 (2.6%)	13 (2.6%)	3 (1.1%)	44 (2.9%)
***M. tuberculosis* complex lineage**	2,279				
Geographically widespread		1,636 (72%)	396 (80%)	183 (69%)	1,057 (70%)
Geographically restricted		643 (28%)	102 (20%)	83 (31%)	458 (30%)

[1]n (%); Median (IQR); High TB incidence defined as >30/100.000 incidence based on 2016 WHO data

538 **Table 2: Associations of participant characteristics and bacterial factors on**

539 **transmission** (quasi-Poisson model with overdispersion φ = 1.2 and φ = 1.4, respectively, N

540 = 2,279)

Characteristic	Tuberculosis Infection			Secondary Active Tuberculosis		
	IRR[1]	95% CI[1]	p-value	IRR[1]	95% CI[1]	p-value
Age (index)	0.96	0.94, 0.99	**<0.001**	0.90	0.82, 1.00	**0.042**
Sex (index)	1.04	0.95, 1.14	0.4	1.15	0.77, 1.75	0.5
Smear positive (index)	1.34	1.21, 1.48	**<0.001**	1.42	0.92, 2.28	0.13
Alcohol use (index)	1.10	0.87, 1.36	0.4	2.25	1.16, 4.01	**0.010**
Drug use (index)	0.93	0.74, 1.15	0.5	1.40	0.66, 2.65	0.3
High incidence contact group (%)	1.98	1.75, 2.25	**<0.001**	0.81	0.47, 1.39	0.5
One child contact (index)	1.03	0.94, 1.13	0.5	0.71	0.47, 1.05	0.091
Mtbc geographically restricted lineage (index)	0.86	0.78, 0.95	**0.004**	0.55	0.32, 0.90	**0.023**

[1]IRR = Incidence Rate Ratio, CI = Confidence Interval, *Mtbc* = M. tuberculosis complex, P-values from Wald test, High TB incidence defined as >30/100.000 incidence based on 2016 WHO data

Legend Figure 1: Properties of circulating *M. tuberculosis* complex geographically restricted and widespread strains. a) Flowchart of included tuberculosis index cases and linked contacts; b) Density plot of contact group sizes per city; c) Maximum likelihood

phylogenetic tree of included strains, clades are coloured by genetic lineage, the rings

denote (from inner to outer) the city, clustering based on a 12 Single Nucleotide Substitution

cut-off, phenotypic drug resistance to Rifampicin and Isoniazid; d) Proportion of isolated

strains by geographically restricted or widespread *Mtbc* lineages over the study period; e)

Bar plot showing the proportions of isolated restricted or widespread *Mtbc* lineages by age

groups. TB = tuberculosis, IGRA = Interferon Gamma Release Assay, SNS = Single

Nucleotide Substitution, Mtbc = *M. tuberculosis* complex.

Legend Figure 2: *M. tuberculosis* complex strain and tuberculosis index case

characteristics associated with transmissibility and clustering. a) Violin plot of the terminal

branch lengths of *Mtbc* strains, median and interquartile ranges are represented by overlaying

box plots; b) Numbers of genetic clusters and number of clustered strains by geographically

restricted or widespread *Mtbc* lineage (legend as in a); c) Forest plot of a multivariable Poisson

rate model quantifying the effect of several predictor variables (y-axis) on the outcome 'Count of

tuberculosis infected contacts' (in orange) or 'Count of secondary active tuberculosis cases

among contacts' (in yellow); d) Bar plot of secondary active tuberculosis among close contacts

inferred by Phybreak stratified by geographically restricted or widespread *Mtbc* lineage (legend

564   as in a)); e) Forest plot of a multivariable Poisson rate model quantifying the effect of Mtbc

565   lineage on tuberculosis infection (orange) or secondary active tuberculosis (yellow) among close

566   contacts (same legend as in c); f) Host Propensity to Propagate by geographically restricted or

567   widespread Mtbc lineage; f) Host propensity to propagate stratified by *M. tuberculosis* complex

568   lineage. Mtbc = *M. tuberculosis* complex

Legend for Figure 3: *M. tuberculosis* **complex sympatry and its effect on**

transmissibility. a - b) Global maps displaying the countries that were considered endemic

(or sympatric) to the geographically restricted *Mtbc* lineages detailed in the figure legends.

For maps of the distribution of restricted Lineage 4 sublineages see Reference [18]. c) Bar plot

showing the proportions of *Mtbc* lineages isolated from the 2,279 pulmonary tuberculosis

index cases included in this study shown for index cases with counts > 10; d) Permutation

analysis (n = 1000) of the expected proportion of sympatric host ancestry – *Mtbc* lineage

proportions. Vertical dashed line represents the proportion (0.75) among the index cases

included in the study; e) Modelling approach to quantify the probability of infection for

sympatric and allopatric contact exposures while including index case characteristics of

infectiousness, contact characteristics of susceptibility and the correlation introduced by

contact tracing. f) Forest plot of the effects estimated using a Generalized Estimating

Equation model to quantify the effect of host country of birth - pathogen lineage co-

localization (sympatry or allopatry). The outcome is tuberculosis infection per contact, and

sympatry was defined using the maps in panel a) and b). Four models assuming four

different co-localized lineage-host country of birth combinations are presented by different

colors. High-incidence country defined as >30/100.000 tuberculosis incidence based on

WHO 2016 data. g) Permutation test assuming 14% mixed ancestry in the sample (flipping

the sympatry variable from 1 to 0 and vice versa in 14% of the sample) while rerunning the

model in panel f). Mtbc = Mycobacterium tuberculosis complex.

591
Figure 4. **Comparative analysis of phagocytosis and intracellular growth of *M.***
***tuberculosis* complex strains in human macrophages.** a. General workflow: 1) Upon
obtaining whole blood from healthy donors, CD14+ monocytes were differentiated to
macrophages. 2) Macrophages were infected with *M. tuberculosis* complex L4 and L6 strain.
3) Incubation of macrophages with *M. tuberculosis* complex strains for 4h, 24 h, 96h, and
168h. 4) Infected macrophages were lysed, uptake and intracellular growth was assessed by
colony forming unit (CFU) assays. (**b-c**) *Mtbc* uptake averaged across three strains of the
Euro American Lineage 4 LAM (**b**) and the West African II Lineage 6 **(c)** in blood derived
monocyte macrophages (MDMs) from donors of self-reported ancestry that co-localizes with
the endemicity of L6**.** *Mtbc* uptake by MDMs was determined immediately at 4hpi (CFU
count at 4hpi x 100%/ CFU count of infection dose) **(d-e)** Phagocytic uptake of two clinical

strains and the reference strains of L6 **(d)** and L4 **(e)** lineages in six donors of differing self-
reported ancestry. **(f)** *M. tuberculosis* complex intracellular growth of two clinical and the
reference strain of L4 and L6 lineages after 24 hours post infection (hpi) in MDMs of donors
of different self-reported ancestries (see **Suppl. Figure S6** for 96hpi and 168hpi) **(g)**
Intracellular growth at 24 hpi averaged across three L4 and L6 strains by donor. Bars show
the mean and standard deviation of the average of CFU values from three infection wells per
strain (**Suppl. Figure S3**) and the mean of nine infection wells per lineage respectively.
Three CFU measurements from each infection well were performed. MDMs were processed
and infected at MOI ~1:1 (0.5 x10^6 cells: 0.5 x10^6 *Mtbc* bacilli) under the same conditions
and in the same laboratory. Statistical results are based on unpaired T-test (b, c, g) or One-
way ANOVA with Bonferroni post hoc test correction among donors (d and e), or between
strains and H37Rv (f). Mean, standard deviation, and non-significant (ns, $p > 0.05$) are
depicted.

METHODS

Study design, participants, and associated metadata

We harmonized TB surveillance data from public health departments tracing TB in three

large cities Amsterdam, Hamburg, and New York City, where TB is reportable to the

respective public health agencies. The data consists of clinical characteristics and pathogen

whole genome sequence data of TB index cases, as well as information on their close

contacts including exposure time (close contact versus non-close contact), demographics,

relevant comorbidities, and Interferon Gamma Release Assay (IGRA) results. We detail

each TB surveillance sample by site below:

Amsterdam and the Netherlands: The National Institute for Public Health and the

Environment (RIVM) in Bilthoven, The Netherlands, contains the Mycobacterial reference

laboratory for the diagnosis of all TB cases in the Netherlands. Since retrospective data in

this study was de-identified by name, WGS data from the RIVM and clinical and

demographic data for patients were linked based on sex, date of birth, year of diagnosis and

postal code. All notified *M. tuberculosis* culture-positive cases from the city of Amsterdam

between December 2015 and December 2019 and their contacts were included. In addition,

we obtained information on all active cases diagnosed elsewhere in The Netherlands in this

period. In the case of patients with multiple isolates, only the isolate with the earliest date of

diagnosis was included. Close contact was defined as daily contact in a room of in total ≥48

hours or ≥6 hours in a car per week.

Hamburg: We obtained data from the Public Health Office Hamburg, Germany for TB index

cases from 2008 through 2017. Sequencing of Hamburg cases was performed at the

National Reference Center for Mycobacteria at the Research Center Borstel. Close contact

was defined as a contact time of in total ≥8 hours.

New York City: The New York City Department of Health and Mental Hygiene notifies, and

traces contacts of all infectious TB index cases. We obtained clinical and demographic data

of TB index cases diagnosed between 2016 and 2020 and their contacts. Pathogen WGS is

performed at the New York State Wadsworth Center in Albany, NY, for culture-positive

cases. In the case of patients with multiple isolates, only the isolate with the earliest date of

diagnosis was included. Close contact was defined as a contact time of ≥8 hours per week.

We compared characteristics of index cases, and their contact group networks to explore

comparability of contact tracing activity across the three cities' public health departments.

Across the three cities, index case age positively correlated with mean contact group age

(pooled Pearson's correlation coefficient 0.4 [CI_{95} 0.39 - 42], and the count of TB infected

close contacts correlated moderately with the contact group size (Pearson's 0.5 [CI_{95} 0.53 –

0.56]) (**Figure S7a-b**). We excluded extrapulmonary TB cases from the source case analysis

due to their recognized significantly lower infectiousness (n = 903/5,256,17%). We excluded

strains not typed as *Mtb* sensu stricto or *M. africanum* (i.e., *M. canettii* or *M. bovis*, n =

132/5,219, 0.03%). Among contacts, we excluded those without blood test result for *Mtbc*

infection (Interferon Gamma Release Assay, or IGRA, 12%, 3,483/28,889), excluded

contacts not considered close (42%, 12,228/28,889) and contacts not linkable to an index

case (17%, 4,891/28,889). Our rationale for excluding contacts without significant exposure

time (<8 hours) independent of exposure location (household versus work or leisure) is

based on previous evidence that households account for <20% of TB transmission[1], and

logistic regression models indicating a significant effect of >8h exposure on IGRA positivity

(Odds Ratio 2.24, 95% Confidence Interval 2.06 to 2.82, Wald p-value <0.001) (Suppl. Table

S8).

Index case and contact data linkage

Demographics, comorbidity information, and data on social risk factors for TB were extracted

from the three public health databases. The investigated contacts were considered to have

TB infection if their IGRA values were above 0.35 IU/ml based on existing test specifications.

Contacts were linked to TB index cases by epidemiological investigation. This linkage of

contacts and active cases is coded as a unique identifier that was subsequently used to

compute aggregated contact group information (count and percent of high-incidence country

birthplaces) per index case. Conversely, index case information was linked to the contact

data (age, sex, sputum microscopy, co-morbidities) as these were important covariates to

control for in the modeling.

Ethical approval

The study is approved by the Institutional Review Boards at Harvard Medical School (IRB19-

1910), the University of Lübeck (AZ-19-071), the New York City Department of Hygiene and

Mental Health (20-078), and the Registration Committee of the Netherlands Tuberculosis

Register NTR (11-2019).

Phenotypic drug resistance

At RIVM (The Netherlands), the Research Center Borstel (Germany) and the New York

State Wadsworth Center phenotypic drug susceptibility testing was performed with the MGIT

960 method using the World Health Organization recommended critical concentrations of

1mg/L for rifampicin and 0.1 mg/L for isoniazid.

Whole Genome Sequencing and analysis

At the time of writing, whole genome sequencing of all culture positive TB cases is routinely

performed in the Netherlands and New York State since 2016.

Isolates from the Netherlands were sequenced at the RIVM Bilthoven as follows: genomic

DNA was extracted from positive MGIT culture tubes (Becton Dickinson, NJ, USA) using the

QIAamp DNA mini kit (QIAGEN GmbH, Hilden, Germany) and sequencing was performed

on Illumina HiSeq2500 with 2 x 125bp reads following Nextera XT library preparation.

At the Wadsworth Center of the New York State Department of Health, genomic DNA was

extracted from MGIT culture tubes (Becton Dickinson, NJ, USA) using the

InstaGene/FastPrep method as previously described.[2] Paired-end 250-bp DNA sequencing

was carried out using the Illumina MiSeq platform (Illumina, San Diego, CA, USA) after

Nextera XT library preparation.

Patient isolates from Hamburg were sequenced for a previous research project at the

Research Center Borstel as follows:[3] Genomic DNA was extracted from solid Löwenstein

Jensen culture tubes using the Cetyltrimethylammonium Bromide (CTAB) method.

Sequencing libraries were constructed from genomic DNA using a modified Illumina Nextera

protocol for 2 x 151 bp reads on the Illumina NextSeq 500 (Illumina, San Diego, CA, USA).

Raw sequencing reads across the three sites were pooled and analyzed on Harvard Medical

Schools High Performance Cluster O2. Analysis and quality control was performed using the

same bioinformatics pipeline. Reads were first trimmed to remove low quality bases and

adapter sequences using *fastp* v0.20.1.[4] The trimmed reads were aligned to the reference

genome H37Rv (accession NC_000962.3) using *BWA mem* v0.7.17 (parameters: default) [5].

Duplicate reads were removed with *Picard* v2.25.1 (http://broadinstitute.github.io/picard/).

We called single nucleotide substitutions (SNSs) and indels with *Pilon* v1.24 (parameters:

default) requiring confidence base calls (*pass* filter).[6]

Variable position alignment and phylogenetic analysis

To analyze the genetic variation in our sample we created a genotypes matrix based on the

variant call format files. We detected SNS sites at 347,471 H37Rv reference positions (of

which 9,641 SNSs were not biallelic) among our global sample of 5,219 strains. We

constructed a 347,471x5,219 genotypes matrix (coded as 0:A, 1:C, 2:G, 3:T, 9:Missing) and

filled in the matrix for the allele supported at each SNS site (row) for each isolate. We

excluded 7,406 SNS sites that had an Empirical Base-level Recall (EBR) score of <0.9,[7]

3,555 SNS sites in mobile genetic elements (e.g., transposases, integrases, phages, or

insertion sequences), and 933 SNS sites located in overlapping coding sequences. We next

dropped 10,169 SNS sites that are missing in >10% of our strains. An additional 133 SNS

sites were dropped because the minor allele is present in <1 isolate. If a base call at a

specific reference position for an isolate did not meet the filter criteria that allele was coded

as missing. Our filtered genotypes matrix had dimensions of 325,275x5,219 SNSs

representing 325,275 SNS sites across 5,219 strains. The phylogenetic lineage was

determined using a refined lineage-calling scheme based on 96-SNSs with *fast-lineage-*

caller.[8]

To infer the population structure of our strains we merged the variant call format files using

bcftools v1.12[9] and removed regions with repetitive sequence content with low EBR scores,[7]

mobile genetic elements, coding for genes involved in antibiotic resistance as well as all

invariant sites. A multiple sequence alignment of 297,061 characters found in 5,087 *Mtbc* L1-

L6 strains was created using *vcf2phylip* (v1.5, https://doi.org/10.5281/zenodo.1257057) and

fed to *iqtree* v2.1.2 for phylogenetic inference.[10] The final maximum likelihood tree was

inferred using a general time reversible model of nucleotide substitution, a Gamma model of

rate heterogeneity, empirical base frequencies (GTR+F+R2) and 1000 ultrafast bootstrap

replicates. The tree was annotated using iTol.[11]

Definition of clusters based on genetic distance

Mtbc strains from patients linked by transmission using epidemiological data rarely differ by

more than 5 SNSs.[3,12] To define clustered strains in our dataset, we first excluded the small

number of non-biallelic SNSs to simplify distance calculation. Of the 325,275 SNS sites,

316,274 were biallelic, i.e., had only a single alternate allele among our strains. We

calculated the pairwise distance as the Hamming distance of SNSs across the total 5,219

strains and subset this matrix for to obtain one pairwise distance matrix for each *Mtbc*

lineage and city.

We next performed graph-based clustering using the *gengraph* function implemented in the

adegenet R package v2.1.5[13] for each *Mtbc* lineage and city specific pairwise distance

matrix as we were only interested in transmission within each city. We used a 12 SNS cut-off

as possible transmission with single linkage clustering, i.e., strains in a cluster differed at

most 12 SNSs from at least one other isolate in the cluster.

Index case inference and infector/infectee relationships

We used the R package phybreak v0.5.2 to identify the infector case for each active TB case

in a cluster.[14] Cases that were not in clusters were assumed to not have secondary active

TB cases. (If contacts of TB cases moved to a different city and developed secondary active

TB this would not have been detected using this approach.) Phybreak identifies infector-

infectee relationships by combining models for transmission, case observation, within-host

pathogen dynamics, and mutation assuming that all cases of a transmission cluster have

been sequenced. The most likely transmission tree is sampled from the posterior distribution

through Bayesian inference and Monte Carlo Markov Chain sampling. The priors for the

mean and shape of the sampling interval (time between infection and sampling of a case)

and generation interval (i.e., time between infection of a primary case and a secondary case

by this primary case) were assumed to follow a gamma distribution as in previous work [15,16].

We ran five independent MCM chains with 10,000 cycles of burn-in and 50,000 sampling

cycles for each cluster. We dropped each estimated infector-infectee relationship where the

effective sample size was <200 which can occur if sampling dates occur within a short

period of time. We excluded active cases that did not have an infector with a posterior

probability (PP) of at least 0.5 from further analysis after a sensitivity analysis evaluating

different PP values in the transmission models (**Suppl. Fig S8**).

Statistical analysis and models of transmission

The python libraries *pandas* v1.3.5[17] and *numpy* v1.21.5[18] were used to manipulate

epidemiological data. All statistical analyses and visualizations were performed in R v4.1.2[19]

using libraries included in the *tidyverse* v.1.3.1.[20] Effect estimates of regression analyses

were visualized using the *sJPlot* package 2.8.10. Missing values for the covariates HIV

coinfection and alcohol were imputed using a bootstrapped logistic regression model

implemented in the package *mice* v3.14.0[21]. We coded age into brackets (0 to 14, 15 to 24,

25 to 34, 35 to 44, 45 to 54, 55 to 64, 65 to 74, and 75 and above).

We used a multivariate Quasi-Poisson rate regression model (Eq1) to identify predictors of

the number of secondary active TB cases or *Mtbc* infections among close contacts. We

ensured there was no collinearity between predictor variables (Suppl. Figure S8). The group

size of close contacts per index case was used as the offset to denote the exposed group in

the Poisson regression. Adjusted incidence rate ratios are presented with 95% confidence

intervals.

Eq1:

$$\boldsymbol{log}(Count\ TB\ infections_{index})$$
$$= \beta_0 + \beta_1 age_{index} + \beta_2 sex_{index} + \beta_3 sputum\ smear_{index} + \beta_4\ alcohol_{index}$$
$$+\ \beta_5 drug\ use_{index} + \beta_6 high\ incidence\ country_{contact}$$
$$+\ \beta_7 one\ child\ contact_{index}\ +\ \beta_8 Mtb\ lineage_{index}$$
$$+\ log(offset(contact\ group\ size))$$

Generalized estimating equations (GEE) were used to validate the Quasi-Poisson models

and to quantify the effect of host-pathogen sympatry on TB infection for each close contact

individually (Eq2). The *gee* package v4.13-20 was used for these analyses with the option

'exchangeable' to imply that any two observations per contact group are equally correlated

without correlation between observations from different contact groups [22]. This allowed to

adjust the effect estimates for correlations in the data. Adjusted odds ratios and 95% CI are

presented.

Eq2:

$$log\left[\frac{P(ContactTBinfection)}{1 - P(ContactTBinfection)}\right]$$

$$= \beta_0 + \beta_1 age_{contact} + \beta_2 sex_{contact} + \beta_3 diabetes_{contact} + \beta_4 HIV_{contact}$$

$$+ \beta_5 high\ incidence\ country_{contact} + \beta_6 sputum\ smear_{index} + \beta_7 alcohol_{index}$$

$$+ \beta_8 drug\ use_{index} + \beta_9 sympatric\ exposure_{contact}$$

Frequencies or proportions of binary variables were compared using Chi Squared, binary to

continuous variables were compared using the Wilcoxon Rank Sum test, and continuous

data was compared using ANOVA or t-tests. All statistical tests were two-sided and a *P*

value cut-off of 0.05 corrected for multiple testing with the false discovery rate.

Host infectiousness scores

The adjusted estimates from the multivariate Quasi-Poisson regression model were each

multiplied as weights to calculate each index case's infectiousness score, similar to previous

work on the propensity to propagate.[23]

Human macrophage donors

Blood donation was approved by the ethics committee of the Research Center Borstel (RCB)

(Borstel, Germany) and the University of Luebeck (Luebeck, Germany) (AZ-19-071). A total

of ~ 300 ml of whole blood (heparin- anti-coagulated) was obtained via venepuncture by the

blood service unit at the RCB. Human healthy donors self-reported ancestries from

Germany, Nigeria, Ghana, and Cameroon with a medical check-up no longer than two years

ago.

Mtbc **strains and culture conditions**

A panel of six *Mtbc* strains representing the modern and ancient human-adapted clades 1

and 2 (L4 and L6) was selected from a reference collection at the RBC. Strains were

recovered from clinical samples or provided by the American Type Culture Collection

(ATCC). Strains were characterized by drug susceptibility testing and sequencing as

previously described.[24,25] *Mtbc* strains were grown on Loewenstein-Jensen (LJ) medium and

further passaged in 10 ml of mycobacteria liquid media (MLM): Middlebrook 7H9 broth

(Becton Dickinson) supplemented with 10% oleic acid albumin dextrose catalase (OADC),

0.2% glycerol, 0.05% Tween 80. *Mtbc* bacilli were grown until mid-log phase at an Optical

Density $(OD)_{600nm}$ of 0.5 at 37 °C, harvested, and stored as stocks in MLM at -80 °C.

Aliquots were finally titrated by Colony-Forming Units (CFU) using Middlebrook 7H10 agar

(Becton Dickinson) plates supplemented with 10% heat-inactivated bovine serum and

cycloheximide (10 mg/ml).

In vitro generation of macrophages

Blood-derived human monocytes were obtained by density gradient centrifugation and

separation with anti-human CD14 antibodies (Milteny Biotec, Germany) following the

manufacturers instructions. Isolated monocytes had a purity and viability greater than 85-

90% which was assessed by flow cytometry and electronic current exclusion technology with

CASY. Monocytes were seeded in 24 well plates at $0.5x10^6$ cells in 500 µl of Macrophage

Culture Media (MCM): RPMI 1640 (Biochrom), penicillin/streptomycin 1% (Biochrom), L-

glutamine 1% (Biochrom) with AB + heat-inactivated human serum 1% and further

differentiated into monocyte-derived macrophages (MDM) after seven days of culture in the

presence of 10 ng/ml of human recombinant macrophage colony stimulating factor (M-CSF).

Cells were incubated at 37°C /5% CO_2, and MCM was changed every three days until cells

achieved ~ 80% confluence.

Cell infection assays

CFU-titrated *Mtbc* stocks were used to prepare the bacterial suspension for infection

experiments in macrophage infection media consisting of 1% L-Glutamine and 10% FSC in

RPMI medium (all Biochrom). For growth experiments, each experimental setup comprised

four plates for different time points (4 h, 24 h, 96 h, and 168 h post-infection (hpi)). **0.5×10^6**

Mtbc bacilli per 500 µl Macrophage infection media were used to infect **0.5×10^6** MDM per

well. Macrophage infection media without bacteria was used as infection control. Upon

infection, plates were incubated at 37 °C in 5% CO_2 for 4h. We quantified CFUs of the

inoculate to be able to compare the infection dose. The percentage of *Mtbc* uptake was

calculated as colonies count at 4hpi*100%/colonies count of infectious dose. Each time point

included three replicates per *Mtbc* strain and three non-infected controls. In all *in vitro* cell

infection experiments, the number of intracellular *Mtbc* bacilli was determined by CFU

assays four hpi and at these different time points: 24 hpi, 96 hpi, and 168 hpi. Cells were first

lysed with sodium dodecyl sulfate (SDS) 0.05% and lysate was then plated in triplicates onto

Middlebrook 7H10 agar plates for CFU counting.

Three independent infection experiments of donors with self-reported ancestry of Germany

were performed representing a total of nine technical replicates of infection of which each

was plated three times for CFU quantification. Due to insufficient cell yield two experiments

compromised two donors each. Of these experiments, in one donor (Germany 2), L4 and L6

strains were assayed in macrophages of one single donor while the other two experiments

(Germany 1 and 3) comprised two donors. Of these, the first donor was infected with L4 and

H37Rv, the second donor with L6 and H37Rv. These data are represented together in one

figure as we detected no significant differences in H37Rv uptake, growth, and inter-intra host

variation among donors (**Suppl. Figure S9 a-h and Figure S10 a-f).** Three independent

infection experiments were conducted including donors from Nigeria, Cameroon, and

Ghana. Each compromised one single donor infected with L4 and L6 (including nine

technical replicates of infection, and 27 CFU technical replicates per strain).

878
879
880   METHODS REFERENCES
881

1.   Mathema, B. *et al.* Drivers of Tuberculosis Transmission. *J. Infect. Dis.* **216**, S644–S653 (2017).

2. Shea, J. *et al.* Comprehensive whole-genome sequencing and reporting of drug resistance profiles on clinical cases of

Mycobacterium tuberculosis in New York State. *J. Clin. Microbiol.* **55**, 1871–1882 (2017).

3. Diel, R. *et al.* Accuracy of whole genome sequencing to determine recent tuberculosis transmission: an 11-year

population-based study in Hamburg, Germany. *Eur. Respir. J.* (2019) doi:10.1183/13993003.01154-2019.

4. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890

(2018).

5. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]* (2013).

6. Walker, B. J. *et al.* Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly

improvement. *PLoS One* **9**, e112963 (2014).

7. Marin, M. *et al.* Genomic sequence characteristics and the empiric accuracy of short-read sequencing. *bioRxiv* (2021)

doi:10.1101/2021.04.08.438862.

8. Freschi, L. *et al.* Population structure, biogeography and transmissibility of Mycobacterium tuberculosis. *Nat. Commun.*

12, (2021).

9. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

10. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective stochastic algorithm for

estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).

11. Letunic, I. & Bork, P. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and

other trees. *Nucleic Acids Res.* **44**, W242-5 (2016).

12. Walker, T. M. *et al.* Whole-genome sequencing to delineate Mycobacterium tuberculosis outbreaks: a retrospective

observational study. *Lancet Infect. Dis.* **13**, 137–146 (2013).

13. Jombart, T. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**, 1403–1405

(2008).

14. Klinkenberg, D., Backer, J. A., Didelot, X., Colijn, C. & Wallinga, J. Simultaneous inference of phylogenetic and

transmission trees in infectious disease outbreaks. *PLoS Comput. Biol.* **13**, e1005495 (2017).

15. Gygli, S. M. *et al.* Prisons as ecological drivers of fitness-compensated multidrug-resistant Mycobacterium tuberculosis.

Nat. Med. **27**, 1171–1177 (2021).

16. Behr, M. A., Edelstein, P. H. & Ramakrishnan, L. Revisiting the timetable of tuberculosis. *BMJ* **362**, k2738 (2018).

17. Reback, J. *et al. pandas-dev/pandas: Pandas 1.3.5.* (Zenodo, 2021). doi:10.5281/ZENODO.3509134.

18. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).

19. R Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical Computing* (2013).

20. Wickham, H. *et al.* Welcome to the Tidyverse. *JOSS* **4**, 1686 (2019).

21. Finch, W. H., Finch, M. E. H. & Singh, M. Data imputation algorithms for mixed variable types in large scale educational

assessment: a comparison of random forest, multivariate imputation using chained equations, and MICE with recursive

partitioning. *Int. J. Quant. Res. Educ.* **3**, 129 (2016).

22. Yan, J. & Fine, J. Estimating equations for association structures. *Stat. Med.* **23**, 859–74; discussion 875-7,879-80

(2004).

23.   Nebenzahl-Guimaraes, H., Borgdorff, M. W., Murray, M. B. & van Soolingen, D. A novel approach - the propensity to

920   propagate (PTP) method for controlling for host factors in studying the transmission of Mycobacterium tuberculosis. *PLoS*

921   *One* **9**, e97816 (2014).

922   24.   Homolka, S., Niemann, S., Russell, D. G. & Rohde, K. H. Functional genetic diversity among Mycobacterium tuberculosis

923   complex clinical isolates: delineation of conserved core and lineage-specific transcriptomes during intracellular survival.

924   *PLoS Pathog.* **6**, e1000988 (2010).

925   25.   Reiling, N. *et al.* Clade-specific virulence patterns of Mycobacterium tuberculosis complex strains in human primary

926   macrophages and aerogenically infected mice. *MBio* **4**, (2013).

928   Supplementary Information is available for this paper.

930   Correspondence and requests for materials should be addressed to Matthias I Gröschel or

931   Maha R Farhat.