

Assessing Predictors of Mortality Among Children admitted with Sepsis at a Referral Tertiary Health Center, Northwestern Nigeria

Authors: Fatimah Hassan-Hanga^{1,2}, Baffa Sule Ibrahim³, Halima Kabir^{1,2}, Hafsat Ibrahim U^{1,2}, Kabiru Abdulsalam^{1,2}, Zainab Datti Ahmed^{1,2}, Halima Salisu Kabara¹, Sule Abdullahi Gaya^{1,2}, Dalha Gwarzo Haliru^{1,2}, Nasiru Magaji Sadiq¹, Salisu Inuwa¹, Mohammad Aminu Mohammad^{1,2}

1. Aminu Kano Teaching Hospital, Kano Nigeria
2. Bayero University Kano, Nigeria
3. Malaria Consortium, Abuja Nigeria

Corresponding author: Baffa Sule Ibrahim, baffasule@gmail.com, +234 809 611 1961.

Abstract

Background: Sepsis is a life-threatening infection that can lead to organ failure and death. We aim to assess predictors of mortality among children admitted with Septicemia at a referral health facility in Northwestern Nigeria.

Methods: We conducted a prospective cross-sectional study of children aged 0-14 years admitted to various units of the pediatrics department of the health facilities. Children were recruited between September 2018 and November 2019. All recruited children were followed up on clinical progress until either discharge, abscondment, or death. We assessed the children clinically daily and collected whole-blood samples for laboratory tests. We conducted a univariate and multivariable analysis using STATA-16 to assess identified predictive factors with our outcome variable.

Results: A total of 326 children were recruited, median age: 2-years. About 54.0% of the children were boys, and 53.1% were within 1-5 years age-group. Predominant organisms cultured from the blood of the children were *Salmonella typhi* (5.7%), *Klebsiella pneumoniae* (2.3%), and *Staphylococcus aureus* (2.0%). A total of 35 deaths were recorded with a case fatality rate (CFR) of 10.7%. CFR is highest in children <1years (13.6%).

Child's vaccination status, mother's education level as well as blood lactate levels, GCS, qSOFA score and positive blood culture were significantly associated with child's mortality. Factors associated with increase mortality include; children with incomplete vaccination history [OR=1.72, 95%CI: 2.74–15.53] versus those with full vaccination; children whose mothers had

no formal education [OR=14.39, 95%CI: 3.24–63.99] when compared to those children whose mothers have tertiary level of education. Furthermore, children with whole blood lactate level between 4-8mmol/l [OR=3.23, 95%CI: 1.15–9.07], or greater than 8mmol/l [OR=10.54, 95%CI: 3.68–30.14] versus children with whole blood lactate level less than 4mmol/l; children with qSOFA score of 3 [OR=15.62, 95%CI: 3.31–73.60] versus children with qSOFA score of 1; and children who had a positive blood culture [OR=6.90, 95%CI: 3.04–15.64].

Conclusion: We found a high prevalence of severe sepsis at pediatrics department of AKTH. Serum lactate levels, GCS, and qSOFA scores were predictive of mortality. Routine measurement and monitoring of these parameters will improve case management and reduce sepsis related mortality in the hospital.

Keywords: Mortality, Nigeria, Pediatrics, Sepsis, Serum lactate.

Introduction

Sepsis is one of the main causes of death and morbidity worldwide, with a particularly devastating impact in low-income nations. Tissue damage, organ failure, and death are all possible outcomes of the body's excessive and life-threatening response to infection.^{[1][2][3]}

Pediatric sepsis is a term that refers to a group of illnesses caused by bacteria, viruses, fungi, or parasites, as well as the harmful products of these pathogens. Infants and children benefit greatly from early detection and intervention^{[4][5]}. Sepsis-causing infections are a significant cause of death and morbidity in children worldwide^[6]. It affects people of all ages and is caused by a variety of infectious diseases, including pneumonia, malaria, meningitis, tuberculosis (TB), HIV, and COVID-19^[2,7].

Sepsis affects all age groups, accounting for 20% of global deaths^[8]. In 2017, children accounted for half of all sepsis cases worldwide, with an estimated 20 million cases and 2.9 million fatalities in under 5yrs children. An estimated 17 million of these cases and 3.5 million of the deaths occurring in Africa^{[9][10]}. Sepsis also accounts for 15% of neonatal deaths, and is the most common cause of death in infants globally. According to the World Health Organization, sepsis due to severe pneumonia, severe diarrhea, severe malaria, and severe measles, are responsible for the highest deaths in children^[11].

Significant regional disparities in Sepsis incidence and mortality exist between high income (HIC) and low- and middle-income countries (LMIC)^[7,8,12], with 85% of sepsis cases and related deaths occurring in Africa^[13]. Studies have shown that, children in LMICs are 18 times more likely to die under the age of five than children in higher income countries (HICs)^[7,8,12]. The observed discrepant survival may be caused by the interplay between socio-economic, factors and poor medical care and resource limitations^[6]. Nigeria is the second highest contributor to under-five mortality in the world, majority of these deaths were due to sepsis^[14].

Several factors are reported to be associated with severe sepsis in children. In the United States, infants are at highest risk of sepsis, with a rate ten times higher than that of older children^[15]. About 25% of severe sepsis occurs in Low- and very low-birth-weight (VLBW) children. Also, boys less than 10yrs of age had significantly higher rates of severe sepsis than girls, particularly among infants^[15].

Blood culture remains the gold standard in the microbiological diagnosis of sepsis, however, other prognostic scoring systems, laboratory investigations and biomarkers play a key role^[16]. Early identification and understanding the signs and symptoms of sepsis, along with the detection of important prognostic biomarkers (such as blood lactate and procalcitonin), are crucial elements for early diagnosis of sepsis and the timely establishment of its appropriate clinical management. Illness scoring systems (such as the Glasgow Coma Score, and the Quick Sequential Organ Assessment Score) are widely used in critically ill patients^[17]. However, their use in children with sepsis has largely been limited and shrouded in ambiguity^[18].

Biomarkers can be classified based on their clinical application. Prognostic biomarkers provide information on the likely outcome in an untreated child and help in establishing the intensity of treatment. Monitoring biomarkers serve to evaluate the disease process and are especially informative of the effectiveness of response to the therapeutic intervention, allowing the clinician to titrate or change the management protocol^[16,19].

Early intervention of sepsis especially in children is critical, and delays in commencing treatment worsen prognosis. Even HICs, with the most advanced healthcare systems struggle to deliver recommended care in sepsis timely^[20]. With Nigeria facing similar challenges to other African countries with poor infrastructure and delivery system, it is important for clinicians to have an established protocol for early detection of prognostic parameters in pediatric sepsis. This study aimed to review and assess prognostic factors for mortality in admitted children at a referral tertiary health center in North western Nigeria, and to evidently advocate for a review of the current sepsis management protocol.

Methods

Study Design: We conducted a prospective cross-sectional study on children (ages 0-14 years) who were admitted with sepsis to pediatrics unit of Aminu Kano Teaching Hospital (AKTH), from the first of September 2018 to 30th November 2019. Enrolled children were subsequently evaluated clinically until they were either discharge, absconded or dead.

Sampling: All admitted children were eligible. Enrolment of consenting patients for the study was done for all eligible children. Enrollment commenced from the 1st of September and continued until four weeks prior to end of the study.

Study Population: Children ages 0-14 years admitted with sepsis to pediatrics department of AKTH Kano.

Data Collection: A pre-tested interviewer administered questionnaire was used to collect patient information from the parents or guardians and entered into Open Data Kit (ODK) using smart phones. The following socio-demographic information were collected; hospital ID, name, age, sex, date of admission, ward of admission, parents' occupation and educational levels, child's vaccination history, and ethnicity. A team of experts (Pediatricians) conducted clinical examination on the enrolled children and documented signs and symptoms of sepsis. Clinical parameters recorded include Glasgow Coma Score (GCS), and the Quick Sequential Organ Failure Assessment (qSOFA).

Anthropometric parameters (weight, height, mid upper-arm circumference, head circumference) and vital signs (body temp, respiratory rate, heart rate) were measured and recorded at enrollment. At least 4ml of whole blood was collected from peripheral vein of each child after disinfection using iodine solution, then left to dry and wiped off with (70%) alcohol. Blood culture was done using automated BACTEC[®] 9050 blood-culture system (Becton Dickinson, Temse, Belgium) for a maximum of 5 days. About 2mls of the blood samples were inoculated into a commercially produced vials containing Fastidious Antimicrobial Neutralization Plus media (FAN[®] Plus) for aerobic and anaerobic media, and incubated in the BACTEC[®] incubator. The FAN[®] Plus media contains resins (Adsorbent Polymeric Beads) which absorb out prior antibiotics used, thus allowing the growth of organisms and enhancing the recovery of isolates. Positive vials were then aliquoted and sub-cultured onto Blood, Chocolate and MacConkey agar

plates. Isolated pathogens were gram-stained and results recorded. The remaining 2mls were sent (in tubes containing EDTA) to both Microbiology and Chemical Pathology departments for analysis. Experienced laboratory scientists tested for malaria using RDT kit, thick and thin film microscopy. Complete blood count was done using automated 3-parts hematology analyzer. Estimation for whole blood lactate were done using Roche Accutrend plus[®] point of care testing (POCT) analyzer. Urea, Electrolytes and Creatinine were determined using the Abbott Architect c4000[®] chemistry analyzer.

Data Management and Storage: Patients' information were collected using pre-tested questionnaires via open data kit software pre-installed into secured smartphones. Data were backed up daily to a secured laptop stored (in csv format) at a different location. Data entry was monitored daily and corrected for errors, and a final data cleaning was done at the end of the study.

Data Analysis: Statistical analysis was done using STATA-16. Descriptive statistics were used to summarize frequencies and percentages for categorical data and median and interquartile range for continuous data. In bivariate analysis, comparison of proportions was performed using Chi-square and exact Fisher's test, P-value of less than 0.05 was considered as statistically significant. Logistic regression was used to analyze sociodemographic and sepsis risk factors associated with clinical outcome. We conducted a manual backward stepwise regression to identify significant factors for the regression models. Two regression models were done. A reduced model (only included factors identified as significant using the backwards stepwise regression) and the full model which combined all socio-demographic, clinical and laboratory factors, adjusting for potential confounders and eliminating multicollinearity.

Ethical Clearance:

The study protocol and all data collection tools were approved by Research Ethical Review Subcommittee of the Kano State Ministry of Health, with approval number: MOH/Off/797/T.I/732, written and dated 1st June 2018.

Results

Socio-demographic characteristics

We recruited a total of 326 eligible children between September 2018 and November 2019. The median age was 2 years, with age range of 0 – 14 years. There were 176 (54.0%) boys and 150 (46.0%) girls. About 53.1% of the children were within the 1 – 5 years age bracket, while 27.0% were under one year. Majority, 296 (90.8%) of the patients were enrolled from the Emergency Pediatrics Unit (EPU) while 3 (0.9%) and 9 (2.8%) were enrolled from Oncology and Pediatrics Medical Wards (PMW) respectively. Also, most of the patients 284 (89.3%) were Hausa, while Yoruba and Igbo tribes constituted 7 (2.8%) and 8 (3.2%) respectively. Among recruited children, 165 (53.4%) were fully vaccinated, 90 (29.1%) had incomplete vaccination, while 54 (17.5%) had no any history of vaccination. Thirty-five (35) deaths were recorded (**Table 1**).

Clinical and laboratory findings

Out of the 326 enrolled children, 103 (48.4%) had whole blood lactate levels of <4.0mmol/l, while 72 (33.8%) and 38 (17.8%) had whole blood lactate levels between 4-8mmol/l and >8mmol/l respectively. Also, Glasgow Coma Score (GCS) at enrollment was mild in 59.6%, moderate in 19.6%, and severe in 20.8% of the children. Quick Sequential Organ Failure Assessment (qSOFA) revealed a score-1 in 45.4%, score-2 in 39.2% and score-3 in 15.4% of the children that had the assessment. (**Table 1**).

Severe malaria (103), pneumonia (54) and meningitis (48) constituted the most prevalent causes of sepsis at the center (**Figure 1**). *Salmonella typhi* (5.7%), *Klebsiella pneumoniae* (2.3%), and *Staphylococcus aureus* (2.0%), were the predominant organisms cultured from the blood of the children over the study period (**Table 1**).

Of the 326 enrolled children, 35 deaths were recorded which gives a case fatality rate (CFR) for sepsis of 10.7%. Disease specific fatality rate (DSFR) was highest in children diagnosed with neonatal sepsis (35.0%), followed by those diagnosed with Cerebrospinal meningitis (18.8%) and PEM (18.8%). There were no recorded deaths in children diagnosed with sepsis due to gastroenteritis or osteomyelitis (**Table 2**).

Bivariate and Multivariate

In bivariate analysis, the following factors were significantly associated with high chance of mortality among the children: child's vaccination status ($p<0.001$), mother's education ($p<0.001$), and father's education ($p<0.001$) (**Table 3**). Likewise, whole blood lactate levels ($p<0.001$), GCS ($p<0.001$), qSOFA score ($p<0.001$) and blood culture ($p<0.001$) were found to be significantly associated with increase mortality among the enrolled children (**Table 4**).

Table 5 and **Table 6** contain results of the full and reduced regression models for the socio-demographic and clinical outcomes respectively. In the reduced model for socio-demographic, several factors were associated with increase mortality among the patients: children with incomplete vaccination history [OR=1.72, 95% CI: 2.74–15.53] versus those with full vaccination; children whose mothers have no formal education [OR=14.39, 95% CI: 3.24–63.99] when compared to those children whose mothers have tertiary level of education; children whose fathers have no formal education [OR=7.89, 95% CI: 2.70–23.56] when compared to those children whose fathers have tertiary level of education.

In the reduced model for the clinical and laboratory outcomes, several factors were also associated with increase mortality among the patients: children with whole blood lactate level between 4–8mmol/l [OR=3.23, 95% CI: 1.15–9.07], or greater than 8mmol/l [OR=10.54, 95% CI: 3.68–30.14] versus children with whole blood lactate level less than 4mmol/l; children with qSOFA score of 3 [OR=15.62, 95% CI: 3.31–73.60] versus children with qSOFA score of 1; children who spend between 2–6days on admission [OR=4.59, 95% CI: 1.01–20.94], or 7–4days on admission [OR=7.64, 95% CI: 1.33–43.75], compared to children who spend less than 24hrs on admission; children who have a positive blood culture [OR=6.90, 95% CI: 3.04–15.64]. A GCS score between 14–15 (mild) at enrollment is associated with improved survival [OR=0.19, 95% CI: 0.08–0.45] when compared to children with GCS of 3–8 (severe) at enrolment.

Table 1: Distribution of Socio-demographic characteristics, clinical and laboratory outcomes of children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Socio-demographic Characteristics	Freq.	Percent	Clinical and Laboratory Outcomes	Freq.	Percent
Gender			Outcomes		
Boys	176	54.0	alive	291	89.3
Girls	150	46.0	dead	35	10.7
Age-group			Serum Lactate Levels		
<1	88	27.0	<4mmol/l	103	48.4
1-5	173	53.1	4-8mmol/l	72	33.8
6-10	46	14.1	>8mmol/l	38	17.8
11-14	19	5.8	Glasgow Coma Score (GCS)		
Ward			Severe	66	20.8
A/E	2	0.6	Moderate	62	19.6
EPU	296	90.8	Mild	189	59.6
FSSW	2	0.6	qSOFA Scores		
Oncology	3	0.9	Score-1	44	45.4
PMW	9	2.8	Score-2	38	39.2
SCBU	14	4.3	Score-3	15	15.4
Ethnicity			Vaccination Status		
Hausa	284	89.3	Not vaccinated	54	17.5
Yoruba	9	2.8	Partially vaccinated	90	29.1
Igbo	10	3.1	Fully vaccinated	165	53.4
Others	15	4.7	Blood Isolates		
Father's Occupation			<i>Citrobacter freundii</i>	1	0.3
Self-employed	120	37.5	<i>Escherichia coli</i>	2	0.7
Private sector	85	26.6	<i>Klebsiella pneumoniae</i>	7	2.3
Civil servant	104	32.5	<i>Salmonella typhi</i>	17	5.7
Others	11	3.4	<i>Staphylococcus aureus</i>	6	2.0
Mother's Education			<i>Streptococcus pneumoniae</i>	1	0.3
No Formal Education	75	23.8	Negative	266	88.7
Primary	50	15.9	Patient's Outcomes		
Secondary	114	36.2	Alive	291	89.3
Tertiary	76	24.1	Died	35	10.7
Father's Education					
No Formal Education	54	17.3			
Primary	44	14.1			
Secondary	95	30.5			
Tertiary	119	38.1			




Figure 1: Clinical diagnosis disaggregated by outcome of children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Table 2: Distribution of disease specific fatality rate (DSFR) and age specific fatality rate (ASFR) among children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326)

Clinical diagnosis	Patient's outcomes			DSFR
	alive	dead	Total	
Neonatal Sepsis	13	7	20	35.0%
Meningitis	39	9	48	18.8%
PEM	13	3	16	18.8%
Severe Malaria	93	10	103	9.7%
Sickle Cell Disease	12	1	13	7.7%
Bacterial Pneumonia	51	3	54	5.6%
Typhoid sepsis	17	1	18	5.6%
Others	32	1	33	3.0%
Gastroenteritis	16	0	16	0.0%
Osteomyelitis	4	0	4	0.0%

Age-groups	Patient's outcomes			ASFR
	alive	died	Total	
<1yr	76	12	88	13.6%
1-5yrs	154	19	173	11.0%
6-10yrs	43	3	46	6.5%
11-14yrs	18	1	19	5.3%

Table 3: Bivariate relationships between socio-demographic characteristics and patient's outcome among children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Socio-demographic Characteristics	Patient's Outcome			Chi-square (p-value)
	Alive (%)	Died (%)	Total	
Gender				
Male	152 (86.4)	24 (13.6)	176	3.3571 (0.067)
Female	139 (92.7)	11 (7.3)	150	
Age-group				
<1	76 (86.4)	12 (13.6)	88	2.2297 (0.526)
1-5	154 (89.0)	19 (11.0)	173	
6-10	43 (93.5)	3 (6.5)	46	
11-14	18 (94.7)	1 (5.3)	19	
Vaccination Status				
Not vaccinated	38 (70.4)	16 (29.6)	54	22.7243 (<0.001)**
Incomplete vaccination	81 (90.0)	9 (10.0)	90	
Completed vaccination	155 (93.9)	10 (6.1)	165	
Father's Occupation				
Business	121 (89.6)	14 (10.4)	135	1.7076 (0.426)
Private sector	70 (86.4)	11 (13.6)	81	
Civil servant	96 (92.3)	8 (7.7)	104	
Mother's Education				
No formal education	54 (72.0)	21 (28.0)	75	32.8358 (<0.001)**
Primary	43 (86.0)	7 (14.0)	50	
Secondary	109 (95.6)	5 (4.4)	114	
Tertiary	74 (97.4)	2 (2.6)	76	
Father's Education				
No formal education	40 (74.1)	14 (25.9)	54	19.4666 (<0.001)**
Primary	38 (86.4)	6 (13.6)	44	
Secondary	87 (91.6)	8 (8.4)	95	
Tertiary	114 (95.8)	5 (4.2)	119	

Table 4: Bivariate relationships between clinical and laboratory findings and patient's outcome among children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Clinical and Laboratory Outcomes	Patient's Outcome			Chi-square (p-value)
	Alive (%)	Died (%)	Total	
Serum Lactate Levels				
<4mmol/l	97 (94.2)	6 (5.8)	103	24.1196 (<0.001)**
4-8mmol/l	60 (83.3)	12 (16.7)	72	
>8mmol/l	23 (60.5)	15 (39.5)	38	
Glasgow Coma Scale (GCS)				
Severe	51 (77.3)	15 (22.7)	66	16.6885 (<0.001)**
Moderate	53 (85.5)	9 (14.5)	62	
Mild	179 (94.7)	10 (5.3)	189	
qSOFA Scores				
Score-1	41 (93.2)	3 (6.8)	44	27.0768 (<0.001)**
Score-2	33 (86.8)	5 (13.2)	38	
Score-3	7 (46.7)	8 (53.3)	15	
Duration on admission				
<=24hrs	28 (93.3)	2 (6.7)	30	6.7076 (0.082)
2-6days	64 (75.3)	21 (24.7)	85	
7-14days	11 (64.7)	6 (35.3)	17	
>14days	6 (66.7)	3 (33.3)	9	
Blood Culture Result				
Positive for blood pathogen	23 (63.9)	13 (36.1)	36	26.3493 (<0.001)**
Negative for blood pathogen	244 (92.4)	20 (7.6)	264	

Table 5: Multivariate logistic regression models with odds ratios, P-value and confidence intervals for socio-demographic factors, for children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Variables	Reduced Model			Full Model		
	OR	p-value	95%CI	OR	p-value	95%CI
Gender						
Male	1.00			1.00		
Female	0.50	0.071	0.23 – 1.06	0.47	0.072	0.20 – 1.07
Age-group						
<1	1.00			1.00		
1-5	0.78	0.532	0.36 – 1.70	0.96	0.921	0.39 – 2.32
6-10	0.44	0.225	0.12 – 1.65	0.46	0.304	0.11 – 2.02
11-14	0.35	0.330	0.04 – 2.88	0.57	0.626	0.06 – 5.43
Vaccination Status						
Completed vaccination	1.00			1.00		
Incomplete vaccination	1.72	<0.001	2.74 – 15.52	1.37	0.544	0.49 – 3.81
Not vaccinated	6.53	0.257	0.67 – 4.41	4.09	0.005	1.52 – 11.00
Mother's Education						
Tertiary	1.00			1.00		
Secondary	1.70	0.534	0.32 – 8.98	1.40	0.712	0.23 – 8.48
Primary	6.02	0.029	1.20 – 30.31	5.84	0.075	0.83 – 40.82
No formal education	14.39	<0.001	3.24 – 63.99	5.83	0.099	0.72 – 47.25
Father's Education						
Tertiary	1.00			1.00		
Secondary	2.10	0.208	0.66 – 6.63	0.79	0.749	0.19 – 3.27
Primary	3.60	0.043	1.04 – 12.47	0.82	0.812	0.16 – 4.30
No formal education	7.89	<0.001	2.70 – 23.56	1.58	0.575	0.32 – 7.81
Reduced Model: Log-likelihood= -71.635619 Pseudo R ² = 0.0514				Full Model: Log-likelihood = -85.592181 Pseudo R ² = 0.1785		
OR: Odds Ratio						
CI: Confidence Interval						

Table 6: Multivariate logistic regression models with odds ratios, P-value and confidence intervals for clinical and laboratory outcomes, for children (aged 0 – 14years) admitted with sepsis at pediatrics units of AKTH, Kano (n=326).

Variables	Reduced Model			Full Model		
	OR	p-value	95%CI	OR	p-value	95%CI
Serum Lactate Levels						
<4mmol/l	1.00			1.00		
4-8mmol/l	3.23	0.026	1.15 – 9.07	3.93	0.013	1.34 – 11.56
>8mmol/l	10.54	<0.001	3.68 – 30.14	10.24	<0.001	3.36 – 31.17
Glasgow Coma Scale (GCS)						
Severe (3-8)	1.00			1.00		
Moderate (9-13)	0.58	0.238	0.023 – 1.44	0.39	0.083	0.14 – 1.13
Mild (14-15)	0.19	<0.001	0.08 – 0.45	0.19	0.001	0.07 – 0.50
qSOFA Scores						
Score-1	1.00			1.00		
Score-2	2.07	0.343	0.46 – 9.31	1.57	0.588	0.31 – 7.92
Score-3	15.62	0.001	3.31 – 73.60	9.86	0.006	1.95 – 49.87
Duration on admission						
<=24hrs	1.00			1.00		
2-6days	4.59	0.049	1.01 – 20.94	0.63	0.720	0.05 – 7.97
7-14days	7.64	0.022	1.33 – 43.75	0.53	0.753	0.01 – 26.79
>14days	7.00	0.056	0.95 – 51.45	2.80	0.512	0.13 – 61.02
Blood Culture Result						
Negative for blood pathogen	1.00			1.00		
Positive for blood pathogen	6.90	<0.001	3.04 – 15.64	4.60	0.031	1.15 – 18.48
Reduced Model: Log-likelihood= -22.112108 Pseudo R ² = 0.3605				Full Model: Log-likelihood = -30.54064 Pseudo R ² = 0.2251		
OR: Odds Ratio						
CI: Confidence Interval						

Discussion

This study found case fatality rate for sepsis to be 10.7%. This was comparable to similar studies in Nigeria^[21,22] and studies by El-Mashad et al in Egypt^[23] where the mortality rate was 8.8% but lower than 22% reported by Novosad et al^[24] and 23.8% reported in Saudi Arabia^[25]. This difference could be due to difference in the studied population where their patients were those admitted into the pediatric intensive care unit only, indicating the level of severity of the disease. Majority of our patients were males (59.9%) and within 1-5 years of age (53.3%). A quarter (25.9%) of the patients were less than one year of age. This is similar to many Nigerian studies where most of their patients were less than five years^[21,22,26]. It is also similar to the study by Novosad et al where 39% of their patients were less than one year^[24]. This was expected as sepsis has a biphasic peak, first in early childhood and the second peak in the elderly.

Only 58.1% of our patients were fully immunized, which is less than the 67% reported in the 2018 Nigerian National Nutrition and Health Survey^[27]. This is worrisome because many of the sepsis cases are from infectious diseases. In this study, malaria (30.7%), Pneumonia (16.7%) and Meningitis (14.8%) were the most prevalent causes of sepsis. This was similar to a study conducted in Saudi Arabia^[25] where the respiratory system was the most common infection site but different from Novosad et al^[24] where most of their patients had congenital heart disease and cerebral palsy. This also reflects the difference in epidemiology between patients in high income countries where chronic diseases are more prevalent and those from low- and middle-income countries (LMIC) where infectious diseases are still more prevalent. Vaccination has been proven to be a highly effective prevention strategy against many infectious diseases in children like Pneumonia, meningitis, Tuberculosis etc.

Neonatal sepsis had the highest disease fatality rate in our study, this was similar to findings in Abakaliki (23.9%)^[28] and Jos (27.3%)^[29] but lower than that obtained in Ile-Ife (33.3%)^[30]. This finding also strengthens the fact that sepsis is one of the commonest causes of hospitalization among neonates. Meningitis and severe acute malnutrition followed next with disease fatality rates of 11.8% and 11.1% respectively. This was similar to findings by Ling et al where they identified malnutrition as one of the risk factors for sepsis^[31].

There was significant association between mortality and not being immunized, having high serum lactate, poor Glasgow Coma Score (GCS) and a high qSOFA score. Blood lactate levels

has been used to measure disease severity in patients with sepsis/septic shock. This finding was similar to a study by Hatherill et al that showed that hyperlactemia and its persistence after 24 hours of treatment can predict death^[32]. Though high Serum lactate levels have been associated with increased mortality especially in adult patients presenting with septic shock, some studies (23,14) in children have not been so conclusive. Gorgis et al did not find any association between an elevated early lactate and mortality in children with sepsis but it however correlated strongly with PRISM-III, a measure of mortality risk in pediatrics^[33]. In this study, poor GCS was significantly associated with mortality, this was similar to findings by Crippa et al who carried out a post hoc analysis of intensive care over Nations database investigating the effect of brain dysfunction, measured by the GCS on hospital mortality in critically ill patients^[34]. They concluded that hospital mortality was significantly higher in patients with brain dysfunction measured with GCS and particularly in the presence of sepsis^[34].

Higher qSOFA scores were found to be associated with increased mortality in this study, similar to the study by EL-Mashad et al where they found that the score was higher in non-survivors and the mortality increased progressively across their patient subgroups from lower to higher scores^[23]. However, some studies have reported that the qSOFA score has a poor sensitivity, 37%-70% in predicting in- hospital mortality^[35,36].

In our study, *Salmonella typhi*, *Staphylococcus aureus* and *Klebsiella pneumonia* were the predominant organisms cultured, this was similar to findings by Obaro et al where *Salmonella* species account for 24%-59.8% of bacteremia in Central and Northwest Nigeria. This however, differs from findings by Ogunkunle et al where *Staphylococcus aureus* was the most predominant organism isolated from their patients^[21]. This further signifies that there is a significant variation in the bacterial organisms causing sepsis even within the same region and country.

Conclusion

Pediatric sepsis is common in our hospital with a case fatality rate of 10.7%. Mortality from sepsis was significantly associated with not being fully immunized, poor GCS and high serum lactate levels. Neonates were more affected while in older children, infectious diseases like malaria, Pneumonia and meningitis were the predominant causes. *Salmonella typhi*, *Staphylococcus aureus* and *Klebsiella spp* were the predominant organisms cultured.

Data Availability:

Dataset for this study is available on request to the corresponding author (address above) or the PI of the study: Dr. Fatima Hasan-Hanga: fhassanhanga@gmail.com, +234 803 714 1639.

Conflict of Interest:

The authors do not report any conflicts of interest in relation to this study.

Funding:

Financial support for this study was by individual contributions from members of the SIDOK team which consist of all authors listed on this manuscript.

Acknowledgements:

The authors will like to acknowledge and appreciate the Kano State Ministry of Health, AKTH management, all clinical and nursing departments, and our team of data entrants for their unfiltered support and dedication to the completion of this study.

Authors Contributions

Dr. Fatima Hassan-Hanga contributed to the conception and design of the study and acquisition of data. She also contributed to revising the manuscript for important intellectual content and final approval of the version to be published.

Dr. Baffa Sule Ibrahim contributed to conception and design of the study, acquisition of data, data analysis and interpretation. He substantially contributed to drafting the article and final approval of the version of the manuscript to be published.

Dr. Halima Kabir, Dr. Hafsat Ibrahim, Dr. Kabiru Abdulsalam, Dr. Zainab Datti Ahmed, Halima Salisu Kabara, Sule Abdullahi Gaya, Dalha Gwarzo Halliru, Nasiru Magaji Sadiq, Salisu Inuwa, and Prof. Mohammad Aminu Mohammad, all contributed to the conception and design of the study, acquisition of data, revising the manuscript for important intellectual content, and approval of the version to be published.

Reference

1. World Health Organization. (2020). Global report on the epidemiology and burden of sepsis: current evidence, identifying gaps and future directions [Internet]. [cited 2022 Feb 7]; Available from: <https://apps.who.int/iris/handle/10665/334216>
2. Reinhart K, Daniels R, Kisoorn N, Machado FR, Schachter RD, Finfer S. Recognizing Sepsis as a Global Health Priority - A WHO Resolution. *N Engl J Med* 2017;377(5):414–7.
3. Vincent JL, Martinez EO, Silva E. Evolving concepts in sepsis definitions. *Crit Care Clin* 2009;25(4):665–75.
4. De Souza DC, Barreira ER, Faria LS. The epidemiology of sepsis in childhood. *Shock* 2017;47:2–5.
5. Da Costa São Pedro T, Morcillo AM, Baracat ECE. Etiology and prognostic factors of sepsis among children and adolescents admitted to the intensive care unit. *Revista Brasileira de Terapia Intensiva* 2015;27(3):240.
6. Nwankwor OC, McKelvie B, Frizzola M, Hunter K, Kabara HS, Oduwole A, et al. A national survey of resources to address sepsis in children in tertiary care centers in Nigeria. *Frontiers in Pediatrics* 2019;7(JUN):234.
7. Kisoorn N, Reinhart K, Daniels R, Machado MFR, Schachter RD, Finfer S. Sepsis in Children: Global Implications of the World Health Assembly Resolution on Sepsis. *Pediatric Critical Care Medicine* 2017;18(12):e625–7.
8. Fleischmann-Struzek C, Goldfarb DM, Schlattmann P, Schlapbach LJ, Reinhart K, Kisoorn N. The global burden of paediatric and neonatal sepsis: a systematic review. *Lancet Respir Med* 2018;6(3):223–30.
9. Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and national sepsis incidence and mortality, 1990-2017: analysis for the Global Burden of Disease Study. *Lancet* 2020;395(10219):200–11.
10. Keeley AJ, Nsutebu E. Improving sepsis care in Africa: an opportunity for change? *PAMJ* 2021; 40:204 2021;40(204).

11. Weiss SL, Fitzgerald JC, Pappachan J, Wheeler D, Jaramillo-Bustamante JC, Salloo A, et al. Global epidemiology of pediatric severe sepsis: the sepsis prevalence, outcomes, and therapies study. *Am J Respir Crit Care Med* 2015;191(10):1147–57.
12. Dagher GA, Saadeldine M, Bachir R, Zebian D, Chebl RB. Descriptive analysis of sepsis in a developing country. *Int J Emerg Med* 2015;8(1).
13. World Health Organization. Sepsis [Internet]. 2022 [cited 2022 Feb 9]; Available from: <https://www.who.int/news-room/fact-sheets/detail/sepsis>
14. Wen LS, Oshiomogho JI, Eluwa GI, Steptoe AP, Sullivan AF, Camargo CA. Characteristics and capabilities of emergency departments in Abuja, Nigeria. *Emerg Med J* 2012;29(10):798–801.
15. Watson RS, Carcillo JA. Scope and epidemiology of pediatric sepsis. *Pediatric Critical Care Medicine* 2005;6(3 SUPPL.).
16. Marshall JC, Reinhart K. Biomarkers of sepsis. *Crit Care Med* 2009;37(7):2290–8.
17. Adonis Muganza N, Olivier M, André Kabamba M, Charles Wembonyama M, Oscar Numbi L, Stanis Okitotsho W. Predictors of mortality in neonatal sepsis in a resource-limited setting. *Journal of Advanced Pediatrics and Child Health* 2021;4(1):057–61.
18. Yeshaneh A, Tadele B, Dessalew B, Alemayehu M, Wolde A, Adane A, et al. Incidence and predictors of mortality among neonates referred to comprehensive and specialized hospitals in Amhara regional state, North Ethiopia: a prospective follow-up study. *Italian Journal of Pediatrics* 2021;47(1).
19. Reinhart K, Bauer M, Riedemann NC, Hartog CS. New approaches to sepsis: Molecular diagnostics and biomarkers. *Clinical Microbiology Reviews* 2012;25(4):609–34.
20. Bruijns SR, Wallis LA, Burch VC. A prospective evaluation of the Cape triage score in the emergency department of an urban public hospital in South Africa. *Emerg Med J* 2008;25(7):398–402.
21. Ogunkunle T, Abdulkadir M, Katibi O, Bello S, Raheem R, Olaosebikan R. Pediatric blood culture isolates and antibiotic sensitivity pattern in a Nigerian tertiary hospital.

- Nigerian Journal of Medicine [Internet] 2020 [cited 2022 Jun 4];29(2):261. Available from: <http://www.njmonline.org/article.asp?issn=1115-2613;year=2020;volume=29;issue=2;spage=261;epage=264;aulast=Ogunkunle>
22. Ogundare E, Akintayo A, Aladekomo T, Adeyemi L, Ogunlesi T, Oyelami O. Presentation and outcomes of early and late onset neonatal sepsis in a Nigerian Hospital. African Health Sciences [Internet] 2019 [cited 2022 Jun 4];19(3):2390–9. Available from: <https://www.ajol.info/index.php/ahs/article/view/190753>
 23. Mohamed El-Mashad G, Said El-Mekrawy M, Helmy Zayan M. [Paediatric sequential organ failure assessment (pSOFA) score: A new mortality prediction score in the paediatric intensive care unit]. Anales de pediatria [Internet] 2020 [cited 2022 Jun 4];92(5):277–85. Available from: <https://pubmed.ncbi.nlm.nih.gov/31784324/>
 24. Novosad SA, Sapiano MRP, Grigg C, Lake J, Robyn M, Dumyati G, et al. Vital Signs: Epidemiology of Sepsis: Prevalence of Health Care Factors and Opportunities for Prevention. MMWR Morb Mortal Wkly Rep [Internet] 2016 [cited 2022 Jun 4];65(33):864–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/27559759/>
 25. Abdullah SO bin, Sørensen RH, Nielsen FE. Prognostic Accuracy of SOFA, qSOFA, and SIRS for Mortality Among Emergency Department Patients with Infections. Infect Drug Resist [Internet] 2021 [cited 2022 Jun 4];14:2763–75. Available from: <https://pubmed.ncbi.nlm.nih.gov/34321893/>
 26. Isa SE, Iroezindu MO, Awang SK, Simji GS, Onyedibe KI, Mafuka, et al. An audit of diagnosis and treatment of sepsis in north-central Nigeria. Nigerian Journal of Medicine : Journal of the National Association of Resident Doctors of Nigeria [Internet] 2013 [cited 2022 Jun 7];22(4):319–25. Available from: <https://europepmc.org/article/med/24283093>
 27. National Nutrition and Health Survey (NNHS) 2018 | UNICEF Nigeria [Internet]. [cited 2022 Jun 7]; Available from: <https://www.unicef.org/nigeria/reports/national-nutrition-and-health-survey-nnhs-2018>

28. Medugu N, Iregbu K, Tam PYI, Obaro S. Aetiology of neonatal sepsis in Nigeria, and relevance of Group b streptococcus: A systematic review. PLoS ONE [Internet] 2018 [cited 2022 Jun 7];13(7). Available from: [/pmc/articles/PMC6049915/](https://pubmed.ncbi.nlm.nih.gov/30356806/)
29. EconPapers: Neonatal Sepsis in Nigeria 2 - Narrative Review of Epidemiology and Socioeconomic Factors [Internet]. [cited 2022 Jun 7]; Available from: https://econpapers.repec.org/article/abfjournal/v_3a36_3ay_3a2021_3ai_3a2_3ap_3a28304-28310.htm
30. Adejuyigbe EA, Adeodu OO, Ako-Nai KA, Taiwo O, Owa JA. Septicaemia in high risk neonates at a teaching hospital in Ile-Ife, Nigeria. East Afr Med J [Internet] 2001 [cited 2022 Jun 7];78(10):540–3. Available from: <https://pubmed.ncbi.nlm.nih.gov/11921599/>
31. (Danny) Liang L, Kotadia N, English L, Kissoon N, Mark Ansermino J, Kabakyenga J, et al. Predictors of Mortality in Neonates and Infants Hospitalized With Sepsis or Serious Infections in Developing Countries: A Systematic Review. Front Pediatr [Internet] 2018 [cited 2022 Jun 7];6. Available from: <https://pubmed.ncbi.nlm.nih.gov/30356806/>
32. Hatherill M, McIntyre AG, Wattie M, Murdoch IA. Early hyperlactataemia in critically ill children. Intensive Care Med [Internet] 2000 [cited 2022 Jun 7];26(3):314–8. Available from: <https://pubmed.ncbi.nlm.nih.gov/10823388/>
33. Gorgis N, Asselin JM, Fontana C, Heidersbach RS, Flori HR, Ward SL. Evaluation of the Association of Early Elevated Lactate With Outcomes in Children With Severe Sepsis or Septic Shock. Pediatr Emerg Care [Internet] 2019 [cited 2022 Jun 7];35(10):661–5. Available from: <https://pubmed.ncbi.nlm.nih.gov/28072671/>
34. Crippa IA, Subirà C, Vincent JL, Fernandez RF, Hernandez SC, Cavicchi FZ, et al. Impaired cerebral autoregulation is associated with brain dysfunction in patients with sepsis. Critical Care [Internet] 2018 [cited 2022 Jun 7];22(1). Available from: [/pmc/articles/PMC6280405/](https://pubmed.ncbi.nlm.nih.gov/30356806/)
35. Schlapbach LJ, Straney L, Bellomo R, MacLaren G, Pilcher D. Prognostic accuracy of age-adapted SOFA, SIRS, PELOD-2, and qSOFA for in-hospital mortality among children with suspected infection admitted to the intensive care unit. Intensive Care Med

[Internet] 2018 [cited 2022 Jun 7];44(2):179–88. Available from:
<https://pubmed.ncbi.nlm.nih.gov/29256116/>

36. Goulden R, Hoyle MC, Monis J, Railton D, Riley V, Martin P, et al. qSOFA, SIRS and NEWS for predicting inhospital mortality and ICU admission in emergency admissions treated as sepsis. *Emerg Med J* [Internet] 2018 [cited 2022 Jun 7];35(6):345–9. Available from: <https://pubmed.ncbi.nlm.nih.gov/29467173/>