Title: The impact of physical activity and exercise on aerobic capacity in individuals with spinal cord

injury: A systematic review with meta-analysis and meta-regression

Authors: Daniel D. Hodgkiss[1], Gurjeet S. Bhangu[2,3], Carole Lunny[4], Catherine R. Jutzeler[5,6], Shin-Yi

Chiou[1,7,8,9], Matthias Walter[2,10], Samuel J. E. Lucas[1,7], Andrei V. Krassioukov[2,11,12], Tom E.

Nightingale[1,2,9]*

Running head: Exercise and fitness in SCI

Author Affiliations:

[1] School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, UK.

[2] International Collaboration on Repair Discoveries (ICORD), University of British Columbia,

Vancouver, British Columbia, Canada.

[3] MD Undergraduate Program, Faculty of Medicine, University of British Columbia, Vancouver,

Canada.

[4] Knowledge Translation Program, Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto,

and the University of British Columbia, Vancouver, British Columbia, Canada.

[5] Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland.

[6] Schulthess Clinic, Zurich, Switzerland.

[7] Centre for Human Brain Health, University of Birmingham, United Kingdom.

[8] MRC Versus Arthritis Centre for Musculoskeletal Ageing Research, University of Birmingham,

United Kingdom.

[9] Centre for Trauma Science Research, University of Birmingham, United Kingdom.

[10] Department of Urology, University Hospital Basel, University of Basel, Basel, Switzerland.

[11] Department of Medicine, Division of Physical Medicine and Rehabilitation, University of British

Columbia, Vancouver, British Columbia, Canada.

[12] GF Strong Rehabilitation Centre, Vancouver Coastal Health, Vancouver, British Columbia, Canada.

Correspondence and requests should be addressed to:

Corresponding author*: Tom E. Nightingale PhD,

Assistant Professor, School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham,

Edgbaston, Birmingham, UK,

Email: T.E.Nightingale@bham.ac.uk **Tel:** +44 (0)121 414 6977

Word count: 9,299/10,000; **Abstract word count:** 348

Number of tables: 5; **Number of figures:** 1

ABSTRACT

Background A low level of cardiorespiratory fitness [CRF; typically defined as peak oxygen uptake ($\dot{V}O_{2peak}$) or peak power output (PPO)] is a widely reported consequence of spinal cord injury (SCI). This systematic review with meta-analysis and meta-regression aimed to assess whether certain SCI characteristics and specific exercise considerations are moderators of changes in CRF.

Methods Eligible studies included randomised controlled trials (RCTs) and pre-post studies that conducted an exercise intervention lasting >2 weeks. The outcome measures of interest were absolute ($A\dot{V}O_{2peak}$) or relative $\dot{V}O_{2peak}$ ($R\dot{V}O_{2peak}$), and/or PPO. Four databases were searched up to July 2021. The Cochrane Risk of Bias 2 tool and the National Institute of Health Quality Assessment Tool were used to assess bias/quality. The certainty of the evidence was assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach. Random effects meta-analyses and meta-regressions were conducted.

Results Ninety studies (110 independent exercise interventions) with a total of 1,191 participants were included in our primary meta-analysis. There were significant improvements in $A\dot{V}O_{2peak}$ [0.22 (0.17, 0.26) L/min, $p<0.001$)], $R\dot{V}O_{2peak}$ [2.8 (2.2, 3.4) mL/kg/min, $p<0.001$)], and PPO [11 (8, 13) W, $p<0.001$]. There were no subgroup differences in $A\dot{V}O_{2peak}$ or $R\dot{V}O_{2peak}$. There were subgroup differences ($p\leq0.008$) for changes in PPO based on time since injury, neurological level of injury, exercise modality, relative exercise intensity, method of exercise intensity prescription, and frequency. The meta-regression found that increased age was associated with increases in $A\dot{V}O_{2peak}$ and $R\dot{V}O_{2peak}$, and exercise intensity prescription and volume were associated with increases in PPO ($p<0.05$). GRADE assessments indicated a low level of certainty in the estimated effects due to study design, risk of bias, inconsistency, and imprecision.

Conclusion The pooled analysis indicates that performing exercise >2 weeks results in significant improvements in $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$ and PPO in individuals with SCI. Subgroup comparisons identify that upper-body aerobic exercise and resistance training appear the most effective at improving PPO. Furthermore, acutely-injured, individuals with paraplegia, exercising at a moderate-to-vigorous intensity, prescribed via a percentage of oxygen consumption or heart rate, for more than 3 sessions/week will likely experience the greatest change in PPO.

Registration PROSPERO CRD42018104342

KEYWORDS: Cardiorespiratory Fitness, Cardiopulmonary Fitness, Function, Spinal Cord Injuries,

Rehabilitation, Exercise

Key Points

- Exercise interventions >2 weeks can significantly improve cardiorespiratory fitness in

individuals with a spinal cord injury, by a magnitude greater than one spinal cord injury adjusted

metabolic equivalent (i.e., ≥ 2.7 mL/kg/min). A one metabolic equivalent improvement has been

associated with a reduction in cardiovascular related mortality risk in non-injured individuals.

- Our findings support the minimum 40 minutes of weekly moderate-to-vigorous intensity

exercise recommended by the spinal cord injury-specific exercise guidelines to significantly

improve fitness. However, a two-fold greater improvement in peak power output may be

achieved with exercising ≥ 90 min/week in comparison to ≥ 40 min/week.

- Our secondary meta-analysis comparing cohort studies indicates that prolonged exercise

participation benefits cardiorespiratory fitness in the long term. However, these studies are

prone to confounding and are inherently biased.

- Future research should consider following the recommendations published in the exercise

intervention reporting guidelines, investigate the dose-response relationship between exercise

and cardiorespiratory fitness in this population, and identify whether differences in supraspinal

sympathetic cardiovascular impacts changes in cardiorespiratory fitness.

1. INTRODUCTION

Spinal cord injury (SCI) is a complex neurological condition, caused by trauma, disease or degeneration, which results in sensory-motor deficits (i.e., paralysis or paresis) below the level of lesion and autonomic dysfunctions. Progressive physical deconditioning following injury results in increased health care utilisation, reliance on personal assistance services and a greater predisposition towards developing chronic diseases [1,2]. Individuals with SCI are at an increased risk of stroke, cardiovascular disease (CVD), and type-2 diabetes mellitus compared to non-injured counterparts [3–5]. The elevated incidence of these conditions in people with SCI emphasises the need for targeted interventions to address modifiable risk factors for these chronic diseases, such as cardiorespiratory fitness (CRF). In clinical populations cardiorespiratory fitness (CRF) is typically defined as an individual's peak oxygen uptake ($\dot{V}O_{2peak}$) or peak power output (PPO). $\dot{V}O_{2peak}$ and PPO are determined during graded cardiopulmonary exercise testing (CPET) to the point of volitional exhaustion, and represents the integrated functioning of different bodily systems (pulmonary, cardiovascular and skeletal) to uptake, transport and utilise oxygen for metabolic processes [6]. A number of prospective studies have indicated that CRF is at least as important, if not more so, than other traditional CVD risk factors (e.g., obesity, hypertension and smoking) and is strongly associated with mortality [7–12].

Low levels of CRF have been widely reported in the SCI-population [13], with the between-person variability partially explained by the neurological level and severity of injury (i.e., lower CRF reported in individuals with tetraplegia) [14]. SCI can damage somatic pathways involved in the voluntary control of skeletal muscles, but also sympatho-excitatory pathways involved in the autonomic control of the cardiovascular system. In individuals with cervical and upper-thoracic SCI, the diminished supra-spinal control to the heart and blood vessels in major capacitance beds can limit exercise capacity [15,16]. This may explain the minimal returns on investment highlighted in a recent systematic review on the effects of aerobic exercise interventions in individuals with tetraplegia [17]. A large proportion of the variance in CRF is also explained by physical activity [18], which is reduced in the SCI-population [19,20]. SCI is characterised by lower-limb impairments and an ensuing reliance on mobility aids that limits the engagement in sufficient levels of physical activity to achieve meaningful health benefits.

Performing regular physical activity and/or structured exercise has long been promoted for improving

CRF in individuals with SCI [21,22]. In 2011, the first evidence-based exercise guidelines, specifically

for individuals with SCI were developed [23], which stated that *"for important fitness benefits, adults*

with SCI should engage in at least 20 minutes of moderate-to-vigorous-intensity aerobic activity and

strength-training exercises 2 times per week". This guideline has since been updated, yet remains the

same with regards to CRF benefits [24]. Although this implies adults with SCI can accrue fitness benefits

from volumes of activity well below that promoted in the general population, others have advocated that

adults with a physical disability [25,26] and individuals with SCI [27] should aim to perform at least 150

minutes of aerobic exercise per week. For additional health benefits it has been suggested that adults

should perform closer to 300 minutes per week of moderate-intensity physical activity [28,29]. While

the current SCI-specific guidelines likely represent the *"minimum"* threshold required to achieve CRF

benefits, it has been suggested that this creates an impression that individuals with SCI do not need to be

as physically active as the general population [30]. The dose-response relationship between exercise and

CRF improvements in individuals with SCI remains to be elucidated.

It is noteworthy that the aforementioned SCI-specific exercise guidelines utilise the terminology of

"moderate-to-vigorous" to describe the desired exercise intensity. This is in contrast to accepted

guidelines in the general population whereby moderate and vigorous-intensity exercise are distinguished

from one another with specific thresholds (e.g., ≥150 minutes of moderate-intensity or ≥75 minutes of

vigorous-intensity activity per week) [26]. Exercise intervention intensity has been shown to influence

the magnitude of change in CRF in patients undergoing cardiac rehabilitation [31,32]. The

feasibility/effectiveness of higher intensity exercise is also currently a topical area of research in the SCI-

population [33–35]. There is the potential for vigorous-intensity exercise to be more time efficient or

lead to superior health benefits, although its impact on CRF in individuals with SCI compared to

moderate-intensity exercise is yet to be determined. A recent systematic review identified that exercise

interventions of a specific modality yield distinct changes in certain cardiometabolic health outcomes

and not others in individuals with SCI [36]. This provides rationale for wanting to investigate the efficacy

of different exercise modalities on CRF in this population. Consequently, a number of research questions

requiring further attention include:

1) Do injury-specific characteristics (e.g., tetraplegia vs. paraplegia, acute vs. chronic injuries, motor-

complete vs. incomplete) mediate CRF responses to exercise?

2) What is the best intensity, frequency, and volume of weekly exercise?

3) Is there an optimal conditioning modality [e.g., upper-body aerobic exercise, resistance training,

functional electrical stimulation (FES), hybrid or multimodal exercise interventions etc.]?

To address these questions, we performed a systematic review with meta-analysis and meta-regression

to investigate the impact of different exercise interventions on changes in CRF in individuals with SCI.

Moreover, we gathered evidence to determine whether key moderators (e.g., participant/injury

characteristics, intervention/study characteristics and risk of bias) influence these intervention effects.

2. METHODS

This current review followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

(PRISMA) guidelines [37] and was prospectively registered (PROSPERO ID CRD42018104342).

Randomised and non-randomised study designs [randomised controlled trials (RCTs) and pre-post

interventions without a comparison control group] were included in the primary meta-analysis of this

review. Our secondary meta-analyses included cohort, cross-sectional and observational studies.

2.1. Eligibility criteria

Studies met the following inclusion criteria: 1) Adult (≥18 years) participants; 2) any acquired (traumatic,

infection, cancer) SCI (*note, studies were included if >80% of the sample met these two aforementioned*

inclusion criteria); 3) an exercise or physical activity intervention lasting >2 weeks (RCTs and pre-post

trials included in the primary meta-analysis); 4) report a measurable exposure variable (i.e., secondary

meta-analysis cohort studies: athletes vs. non-athletes or sedentary vs. active participants; and cross-

sectional studies: self-reported or objectively measured habitual physical activity level) and; 5) report

CRF-specific outcomes [i.e., absolute or relative $\dot{V}O_{2peak}$, evaluated via analysis of expired air during a

peak (or symptom-limited) CPET or submaximal prediction, or PPO].

Studies were excluded if they met the following criteria: 1) non-human; 2) non-original work (i.e.,

reviews, guideline documents, editorials, viewpoints, letter-to-editor, protocol paper); 3) case-reports and

case series with a number (n) of participants <5 (to increase the robustness of our findings given the inclusion of smaller sample sizes in previous reviews [21,38,39]); 4) non-peer reviewed (i.e., conference proceeding/abstracts/posters); 5) children or adolescents (<18 years) ; 6) non-SCI (non-injured participants or other neurological conditions); 7) does not report a CRF-specific outcome; 8) single exercise sessions or an intervention <2 weeks; 9) no suitable comparison (i.e., control group or baseline data pre-intervention) or exposure variable measured; 10) no full text; and 11) not written in English. Studies with concurrent interventions (i.e., diet, lifestyle or respiratory training) were included only if the effects of exercise could be isolated.

2.2. Search strategy

A search of the following electronic databases: MEDLINE (via Pubmed), Excerpta Medica Database (EMBASE; via Ovid), Web of Science and the Cochrane Central Register of Controlled Trials (CENTRAL) was conducted from their respective inception through to July 18, 2021. Search terms were developed by the corresponding author (TN) and agreed upon by co-authors (AK, MW). The search strategy combined key words describing the following: 1) condition (e.g., SCI); 2) 'intervention or exposure variable' (e.g., rehabilitation, exercise and physical activity); and 3) 'outcome' (e.g., $\dot{V}O_{2peak}$ or PPO). Details of the complete search strategy can be found as online supplementary material (S1). Search results were collated using Endnote software (Thomson Reuters, NY) and duplicates removed.

2.3. Study selection and data extraction

The citations retrieved from the search strategy were screened by title, abstract, and full text by two independent reviewers (DH, GB). At each stage of the evaluation, studies were excluded if the inclusion criteria were not satisfied. A conservative approach was taken, whereby if insufficient information was available to warrant study exclusion during the title and abstract stages of the screening, studies were retained in the sample for full text screening. TN resolved any disagreement with regards to study inclusion.

Two authors (DH, GB) independently extracted data in duplicate using Microsoft Excel. Any disagreements were resolved via mutual consensus. Where more than one publication was apparent for the same participants, data were extracted from the study with the largest sample size to avoid

duplication. Author, year, study design, sample size, participant demographics/injury characteristics, exercise parameters (including the type, frequency, duration, intensity and weekly volume), or physical activity exposure details (training history, objective wearable device or validated self-report questionnaire) and adverse events were extracted. For RCTs, pre-post interventions and observational studies, mean ± standard deviation (SD) for $\dot{V}O_{2peak}$ and PPO outcomes at baseline and post-intervention/control or observation period were extracted to assess change in CRF. For cross-sectional studies, mean ± SD outcomes were extracted for the unique cohorts, along with the significance and magnitude of associations between CRF and habitual physical activity. Where possible, $\dot{V}O_{2peak}$ values were extracted in relative (mL/kg/min) and absolute (L/min) terms or calculated using pre- and post-intervention body mass values when provided. PPO values were extracted in watts (W) only. If there was insufficient information, the authors were contacted via email (N=12) and given a two-week window to provide additional data (responses received, N=8). Detailed notes were recorded outlining the reasons for study inclusion/exclusion and the number of studies included and excluded at each stage.

2.4. Data synthesis and analysis

A variety of methods [i.e., indices of heart rate (HR), $\dot{V}O_2$ or ratings of perceived exertion (RPE)] have been utilised in the literature to establish, prescribe and regulate exercise intensity in the SCI-population, which creates complexity when classifying the intensity of exercise. Each intervention was classified as having prescribed either light, moderate, vigorous or supramaximal-intensity aerobic exercise, based on thresholds proposed by the American College of Sports Medicine (ACSM) [40] (S2). If a study reported a progression in intensity that spanned the moderate and vigorous-intensity categories (e.g., 60-65% $\dot{V}O_{2peak}$), it was classified as 'moderate-to-vigorous'. If insufficient data were provided, studies were classified as 'mixed-intensity/cannot determine'. Furthermore, where a study reported frequency of sessions or length of interventions as a range (e.g., 6-8 weeks), the midpoint was extracted and if a study reported duration as a range (e.g., 40-45 min), the greater value was extracted. Descriptions of adverse events in the included studies were also collated. These were categorised into the following subgroups: 1) bone, joint or muscular pain, 2) autonomic or cardiovascular function, 3) skin irritation or pressure sores, and 4) other.

Means ± SD were estimated from median and interquartile range (IQR) [41] or median and range [42],

where required. Where CRF data was only presented in figures, data were extrapolated using Photoshop

(Adobe Inc). To combine within-study subgroups and to estimate SD of the delta (Δ) change in CRF

using correlation factors, we followed guidance from the Cochrane handbook [41]. Correlation factors

were calculated for $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$ and PPO using studies that reported pre-post SD and SD of the

Δ change using the following equation:

$$Corr = \frac{(SD_{Pre})^2 + (SD_{Post})^2 - (SD_{Change})^2}{2 \times SD_{Pre} \times SD_{Post}}$$

The specific correlation factors that were calculated for each study were averaged across each study

design (S3) and applied in the following equation to calculate SD of the change for studies where these

values were not reported:

$$SD_{Change} = \sqrt{(SD_{Pre})^2 + (SD_{Post})^2 - 2 \times corr \times SD_{Pre} \times SD_{Post}}$$

where *corr* represents the correlation coefficient.

Since $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$, and PPO are continuous variables, expressed using the same units across

studies, we utilised weighted mean differences (WMDs) and 95% confidence intervals (CI) as summary

statistics. A primary meta-analysis was carried out in R (Version 3.5.1, R Foundation for Statistical

Computing, Vienna, Austria) describing Δ in CRF outcomes in response to prospective, well-

characterised exercise interventions lasting >2 weeks (e.g., combining exercise intervention-arms from

RCTs and pre-post studies). Nine separate primary meta-analyses were performed to describe Δ in each

CRF outcome with studies categorised into subgroups based on the following: 1) time since injury [(TSI),

e.g., Acute (<1-year), chronic (≥1-year)]; 2) neurological level of injury (e.g., tetraplegia, paraplegia); 3)

injury severity [e.g., grading in accordance with the American Spinal Injury Association Impairment

Scale (AIS): motor-complete (AIS A-B), motor-incomplete (AIS C-D)]; 4) exercise modality [e.g.,

aerobic volitional upper-body, resistance training, FES, gait training, behaviour change]; 5) relative

exercise intensity (e.g., light, moderate, moderate-to-vigorous, vigorous, supramaximal); 6) method used

to prescribe exercise intensity (e.g., $\dot{V}O_2$, HR, RPE, workload); 7) frequency of exercise sessions (<3, ≥3

to <5, ≥5); 8) exercise volume [e.g., SCI-specific exercise guidelines for fitness (40 - 89 min/wk) [24],

SCI-specific exercise guidelines for cardiometabolic health (90 - 149 min/wk) [24], achieving general

population exercise guidelines (≥150 min/wk) [26], and 9) length of intervention (≤6 weeks, >6 to ≤12

weeks, >12 weeks). Studies were also classified as 'mixed' or 'not reported/cannot determine' subgroups

based on the aforementioned categories. Four secondary meta-analyses were also conducted for different

trial designs: 1. comparing inactive vs active participants (e.g., cross-sectional cohort studies); 2.

describing Δ in CRF outcomes with standard of care inpatient rehabilitation or free-living follow up (e.g.,

observational studies); 3. comparing Δ in CRF outcomes relative to control groups (RCTs only), and 4.

head-to-head comparison of different exercise intensities (RCTs with exercise interventions of differing

intensities). Statistical heterogeneity was assessed using the I^2 and accompanying p-value from the chi-

squared test. A fixed-effect model was used when no significant heterogeneity was detected among

studies ($P>0.10$, $I^2<50\%$), otherwise, a random effect model was used. Evidence for differences in effects

between the subgroups was explored by comparing effects in the subgroups and the corresponding *p*-

values for interaction. To assess the effect of potential outlier studies, we conducted a sensitivity analysis

where studies were removed, and pooled WMD recalculated, when their CIs did not overlap with the CIs

of the pooled effect. Sensitivity analyses were also conducted by comparing the WMDs of low and high

risk of bias studies, as well as studies with and without imputed data (i.e., extracted from figures or where

mean ± SD were calculated from median, IQR or range), to confirm the robustness of our findings.

Potential publication bias in the dataset was assessed using funnel plots and Egger's tests in R. Data is

visualised in R (see Github for scripts: https://github.com/jutzca/Exercise-and-fitness-in-SCI). A 2.7

mL/kg/min, and thus 1 metabolic equivalent in SCI (1 SCI-MET) [43], change in $R\dot{V}O_{2peak}$ was

considered clinically meaningful.

To explore potential sources of heterogeneity, a random-effects meta-regression was performed using

preselected moderator variables in Stata (Version 13, StataCorp LLC, College Station, TX, USA),

adjusted for multiple testing. As per Cochrane recommendations [44], for each included covariate in the

model a minimum of 10 studies were required. To achieve this, and to also overcome the issue of

collinearity between moderators, some moderators were not included in the analysis. Moderators were

selected *a priori*, based on their potential to influence CRF responses. Exercise intensity prescription was

later added as a moderator in the meta-regression in light of a recent study challenging strategies for

prescribing exercise intensity in individuals with SCI [45]. Moderators fell into two categories: model 1) participant/injury characteristics [continuous variables: age, TSI and baseline CRF; categorical variables: sex (n=male), neurological level of injury (n=PARA), severity (n=motor-complete)]; or model 2) intervention/study characteristics [continuous variables: exercise session duration, frequency, weekly exercise volume, intervention length; categorical variables: exercise modality, exercise intensity, method of exercise intensity prescription, and risk of bias classification]. Any potential covariates of the effect of $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$, and PPO with $p \leq 0.10$ identified via univariate meta-regression were subsequently included in multivariate meta-regression modelling. The level of significance for multivariate meta-regression was set at $p \leq 0.10$. Because meta-regression can result in inflated false-positive rates when heterogeneity is present, or when there are few studies, a permutation test described by Higgins and Thompson [46] was used to verify the significance of the predictors in the final model, whereby 10,000 permutations were generated.

2.5. Risk of bias

Study quality was appraised by at least two independent reviewers in duplicate (DH, GB, SYC), with any conflicts resolved by a third reviewer (TN). The Cochrane Risk of Bias 2 (RoB 2) was used to assess the risk of bias of the RCTs [47]. Reviewers determined the level of bias for each domain using the RoB 2 algorithms and is presented visually using robvis [48]. Non-randomised designs were assessed using assessment tools generated by the National Institutes of Health (NIH) and National Heart, Lung and Blood Institute (NHLBI, Bethesda, MD). Pre-post studies were rated using the Quality Assessment Tool for Before-After (Pre-Post) Studies with No Control Group (12 items) and observational and cross-sectional studies were rated using the Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies (14 items). Studies were subsequently classified as good, fair or poor quality using the guidance provided within each tool and is presented visually in online supplementary material.

2.6. Certainty on the body of the evidence assessment using the GRADE approach

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach [49] was used to evaluate the certainty of the evidence for $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$ and PPO. Two authors (DH, SYC) independently assessed the certainty of evidence for each outcome, with any conflicts resolved by the corresponding author (TN). The certainty of the evidence was graded from 'High' to 'Moderate',

'Low' or 'Very Low'. GRADE certainty in the evidence was downgraded if one or more of the following

criteria were present: 1) risk of bias, 2) inconsistency in the results for a given outcome, 3) indirectness,

4) imprecision, and 5) publication bias.

3. RESULTS

The initial database search identified 12,885 articles after removal of duplicates. Further, 11,029 studies

were removed following the screening of titles and abstracts. The remaining 1,856 articles were selected

for full-text review based on inclusion and exclusion criteria (S1). Of these, a total of 110 eligible studies,

across each specific study design (RCT = 27, pre-post = 63, observational = 5, cross-sectional cohort =

9, cross-sectional association = 6), were included in this review. Ninety studies, comprising the RCTs

and pre-post studies, were included in the primary meta-analysis. Summaries of the pooled cohorts and

descriptions of the individual studies included within each secondary meta-analysis are provided as

supplementary material.

[PLEASE INSERT FIGURE 1 HERE]

3.1. Primary meta-analysis: Effects of prescribed, prospective exercise intervention studies

CRF responses were pooled across 90 studies, comprising 110 exercise interventions in total, taken from

76 pre-post exercise interventions and 34 independent exercise intervention arms from RCTs. Some

studies included multiple exercise intervention arms/phases, hence the greater total number of exercise

interventions than studies. A summary of the demographic/injury characteristics and intervention

parameters for the pooled cohort included in the primary analyses for $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$, and PPO are

presented in Tables 1-2.

[PLEASE INSERT TABLE 1 HERE]

3.1.1. Participants

Across the 110 exercise interventions, there were a total of 1,191 participants. Most interventions

included both males and females (64% of studies), where females made up between 6-80% of the mixed

cohorts. There were no female-only cohorts. Mean age ranged between 24 to 58 years and the majority

of participants had chronic injuries (69% >1-year), with mean TSI ranging between 56 days to 24 years. Sixty-three interventions included a mixed cohort of paraplegia and tetraplegia, of which individuals with paraplegia made up between 10-88% of the mixed cohorts. Four interventions recruited individuals with tetraplegia-only, 34 paraplegia-only, and nine did not specify. Participants across all AIS groups were included, of which 39 interventions were motor-complete-only, 19 were motor-incomplete-only, and 17 did not report. Thirty-five interventions recruited both motor-complete and incomplete individuals, of which 32% were motor-incomplete. Mean $A\dot{V}O_{2peak}$ and $R\dot{V}O_{2peak}$ at baseline was 1.26 (0.51-3.50) L/min and 18.0 (7.3-36.9) mL/kg/min, respectively, and PPO was 49 (0-168) W.

[PLEASE INSERT TABLE 2 HERE]

3.1.2. Exercise intervention characteristics

Length of interventions ranged from 4 to 52 weeks, and whilst most studies reported a specific, predetermined intervention length, some reported a range [50–52], a total or targeted number of sessions [51,53–57], or provided an average [56,58,59]. Exercise sessions were completed between two to seven times per week. Eleven studies reported a range (e.g., "two to three sessions") or maximum frequency (e.g., "up to three sessions/week") [51,54,57,60–67], and frequency was either not reported or could not be determined in five studies [68–72]. The remainder reported an exact frequency (e.g., three sessions per week). The duration of exercise sessions ranged from 5 to 90 minutes, with four studies reporting a range (e.g., 20-30 min) [51,73–75] and six studies reporting a progression to a target duration [54,76–80]. Duration was not reported or could not be determined in 13 studies. Based on current exercise guidelines, 22 interventions prescribed exercise within the SCI-specific exercise guidelines for fitness (40-89 min/week), 44 interventions targeted the SCI-specific exercise guidelines for cardiometabolic health (90-149 min/week), and 26 were greater than general population exercise guidelines (≥150 min/week).

Forty-one interventions utilised aerobic upper-body exercise, 5 upper-body resistance training/circuits, 22 FES, 15 gait training, 4 behaviour change, and 23 mixed/multimodal interventions. Following the ACSM thresholds, one intervention prescribed light-intensity (<1%), 15 prescribed moderate-intensity (14%), 33 prescribed moderate-to-vigorous-intensity (30%), 25 prescribed vigorous-intensity (23%), and

2 prescribed supramaximal-intensity exercise (2%). Intensity could not be determined from 34

interventions (31%). With regards to exercise intensity prescription methods, 32 interventions used HR,

regulated either via HR_{peak} (%HR_{peak}, i.e., determined via a CPET; N=8), HR_{max} (%HR_{max}, i.e., age-

predicted; N=11), or HR reserve (%HRR; N=13). Fourteen interventions established intensity using

$\dot{V}O_{2peak}$ (%$\dot{V}O_{2peak}$; N=13) or $\dot{V}O_2$ reserve (%$\dot{V}O_{2reserve}$; N=1) calculated from the pre-intervention CPET.

Thirteen interventions utilised RPE, using either the Borg CR10 scale (N=7) or the Borg 6-20 scale

(N=6). Workload was used to prescribe intensity in 10 interventions, via a percentage of PPO (%PPO;

N=5), one repetition maximum (%1RM; N=4), or maximal tolerated power (%MTP; N=1). Forty-one

interventions either used a mixture of prescription methods or intensity could not be classified. Detail for

the specific studies is presented in the forest plots in online supplementary material (S4).

3.1.3. Adverse events

Adverse events were described in 17 interventions, comprising 49/1,191 (4.1%) participants (S10). These

events were related to: 1) bone, joint or muscular pain (n=10 participants), 2) autonomic or cardiovascular

function (n=8 participants), 3) skin irritation or pressure sores (n=18 participants), and 4) other events

including anxiety, nausea, dizziness and issues with testing equipment (n=3 participants). Adverse events

were reported in three other pre-post studies. Beillot et al. [68] stated that participants experienced

"spontaneous fractures of lower limbs, occurrence of a syringomyelia and pressure sores at the foot and

ankle" (n=10), but did not define the number of participants who sustained each event. Likewise, Janssen

and Pringle [61] reported *"lightheadedness in some subjects"*, and Gibbons et al. [81] stated that *"a*

number of participants showed some level of autonomic dysreflexia during the FES response test", but

both studies did not quantify further.

3.1.4. Change in CRF outcomes

The summary statistics for the nine primary meta-analyses are presented in Tables 3-4 and their

corresponding forest plots can be found in supplementary material (S4).

[PLEASE INSERT TABLES 3-4 HERE]

Sixty-nine exercise interventions assessed the change in $\text{A}\dot{\text{V}}\text{O}_{2\text{peak}}$, revealing a significant increase of

0.22 [0.17, 0.26] L/min (p<0.001). There were no significant subgroup differences for any of the nine

meta-analyses. Seventy-four exercise interventions assessed the change in $\text{R}\dot{\text{V}}\text{O}_{2\text{peak}}$, revealing a

significant increase of 2.8 [2.2, 3.4] mL/kg/min (p<0.001). There were no significant subgroup

differences for any of the nine meta-analyses. Sixty-one exercise interventions assessed the change in

PPO, revealing a significant increase of 11 [8, 13] W (p<0.001). There were significant subgroup

differences for TSI (p<0.001), neurological level of injury (p<0.001), exercise modality (p=0.003),

relative exercise intensity (p=0.003), method of exercise intensity prescription (p<0.001), and frequency

(p<0.001) (Tables 3-4).

Sensitivity analyses

The removal of potential outliers resulted in no meaningful changes to the overall pooled effects for any

outcome. A sensitivity analysis for risk of bias revealed no differences in the pooled effects for low and

high risk of bias studies (S11). A sensitivity analysis for imputed data revealed a greater $\text{R}\dot{\text{V}}\text{O}_{2\text{peak}}$ in

studies with imputed data (3.9 mL/kg/min) compared to studies without (2.5 mL/kg/min). Yet, there were

no differences in the pooled effects for $\text{A}\dot{\text{V}}\text{O}_{2\text{peak}}$ or PPO (S11). An additional analysis grouped

interventions into those that matched the CPET modality to the exercise intervention and those that did

not. Following the adjustment for subgroup comparisons, there was a significantly greater $\text{R}\dot{\text{V}}\text{O}_{2\text{peak}}$ in

studies with matched CPET and intervention modalities (p = 0.02). There were no significant differences

in $\text{A}\dot{\text{V}}\text{O}_{2\text{peak}}$ or PPO (S12). A sub-analysis on gait training CPETs alone also revealed no subgroup

differences in any outcome (S13).

3.1.5. Meta-regression

Model 1 - Participant and injury characteristics

Increased age was associated with increases in $\text{A}\dot{\text{V}}\text{O}_{2\text{peak}}$ (p = 0.045) and $\text{R}\dot{\text{V}}\text{O}_{2\text{peak}}$ (p = 0.025). There

were no associations between other moderator variables included in this model and CRF outcomes. There

were also no associations between PPO and the other moderator variables (Table 5).

Model 2 - Exercise intervention and study characteristics

There was no evidence that the exercise intervention and study characteristics included in model 2 were

associated with increases in $A\dot{V}O_{2peak}$ or $R\dot{V}O_{2peak}$. However, there was evidence for an association

between the method of exercise intensity prescription and increases in PPO ($p<0.01$). Additionally, there

was evidence for an association between exercise volume and increases in PPO ($p = 0.04$) (Table 5).

[PLEASE INSERT TABLE 5 HERE]

3.1.6. Publication bias

There was no significant publication bias for $A\dot{V}O_{2peak}$ ($Z = -1.23$, $p = 0.22$), $R\dot{V}O_{2peak}$ ($Z = -0.54$, $p =$

0.59), or PPO ($Z = 0.73$, $p = 0.46$). Funnel plots are provided in supplementary material (S4).

3.2. Secondary Meta-Analyses

3.2.1. Cross-sectional studies

Nine studies included cross-sectional data comparing CRF outcomes in active (n=129 participants) vs.

inactive (n=115 participants) individuals with SCI. Inactive participants were mainly classified as

sedentary, whereas active participants varied from recreationally active wheelchair sport players to

paralympic athletes. A meta-analysis of cross-sectional cohort studies revealed significantly ($p<0.001$)

higher $A\dot{V}O_{2peak}$ [0.54 (0.44, 0.63) L/min], $R\dot{V}O_{2peak}$, [9.4 (7.0, 11.8) mL/kg/min] and PPO [37 (29, 44)

W] in active compared to inactive individuals with SCI (S5). Given the significant heterogeneity in

$R\dot{V}O_{2peak}$, a sensitivity analysis was conducted to compare inactive individuals with either 'active' or

'elite athletes'. There was a significantly higher $R\dot{V}O_{2peak}$ [5.4 (3.0, 7.7) mL/kg/min, $p<0.001$] in 'active'

compared to inactive individuals, but an even higher $R\dot{V}O_{2peak}$ [11.2 (9.6, 12.9) mL/kg/min, $p<0.001$] in

'elite athletes' compared to inactive.

Six studies (n=380 participants) included cross-sectional data and assessed associations between habitual

physical activity level (as a continuous variable) and CRF outcomes. Five studies assessed physical

activity exposure using self-report methods [82–86], whereas one study used a validated wearable device

[87]. The measurement period used to capture physical activity dimensions ranged from 3 to 7 days.

There was considerable variability across studies with regards to the physical activity dimensions

captured: hours per week of exercise/sport, minutes per day or week of mild, moderate, heavy-intensity

for the subcategories of leisure time physical activity (LTPA), lifestyle or household activity or cumulative activity (S6). Collectively, data indicates significant positive correlations of a larger magnitude between CRF/PPO outcomes and the volume of sport, exercise or LTPA rather than household activity. The only study to use a validated wearable device indicated that participants performing ≥150 min/wk of moderate-to-vigorous physical activity (MVPA) had a significantly higher CRF relative to a low activity group (performing <40 min/wk). Whereas, there was no significant difference in CRF between the low activity group and participants achieving the SCI fitness specific exercise guidelines (40 - 149 min/wk) [87]. Significant, positive correlations were reported for the amount of moderate-to-vigorous LTPA or cumulative activity with CRF/PPO outcomes, which was not the case for mild or light-intensity activity.

3.2.2. Observational inpatient rehabilitation or community free-living studies

Five studies (n=343 participants) included observational longitudinal data and assessed changes in CRF outcomes following either standard of care inpatient rehabilitation [88–90] or a period of community free-living [88,91,92]. The duration between assessments for standard of care varied, ranging from 5 to 28 weeks, whereas the follow-up period for community observations ranged from 1 to 2.9 years. Reporting on the therapies used within standard of care was poor and only one study included a measurement of physical activity during the community-based free-living follow-up (self-reported mean sport activity) [91]. There were significant improvements following standard of care, but not following community-based free-living, in absolute [0.12 (0.07, 0.17) L/min, $p<0.001$ vs. 0.09 (0.00, 0.19) L/min, $p=0.06$] and relative $\dot{V}O_{2peak}$ [2.1 (1.0, 3.2) mL/kg/min, $p<0.001$ vs -0.1 (-2.9, 2.7) mL/kg/min, $p=0.94$] (S7). Significant improvements in PPO were identified following both standard of care [6 (3, 9) W, $p<0.001$] and community-based free-living [7 (2, 12) W, $p=0.006$] (S7).

3.2.3. RCTs

Twenty RCTs assessed changes in CRF outcomes between exercise intervention (n=255 participants) and control (n=229 participants) groups. A meta-analysis of RCTs revealed a significantly higher A$\dot{V}O_{2peak}$ [0.15 (0.06, 0.24) L/min, $p=0.001$], R$\dot{V}O_{2peak}$ [2.9 (1.7, 4.0) mL/kg/min, $p<0.001$], and PPO [10 (5, 14) W, $p<0.001$] following an exercise intervention relative to SCI controls (S8).

Seven RCTs compared changes in CRF outcomes between moderate (n=52 participants) and vigorous

(n=51 participants) exercise intensity groups. These studies utilised upper-body aerobic exercise and gait

training. A meta-analysis revealed no significant differences between moderate and vigorous-intensity

in $A\dot{V}O_{2peak}$ (p=0.67), $R\dot{V}O_{2peak}$ (p=0.88) or PPO (p=0.62) (S9). There were also no significant subgroup

differences between studies that matched exercise volume between intensity groups and those that did

not.

3.3. Risk of Bias

Full risk of bias assessments for pre-post and RCT interventions can be found in supplementary material

(S4, S8, S9). Twenty-six pre-post studies were rated as having good, 25 as having fair, and 12 as having

poor methodological quality. Six RCTs were rated as having a low risk of bias, 8 as having some

concerns, and 13 as having a high risk of bias. The most common domains in the RCTs with either some

concerns or high risk were 'bias in the measurement of the outcome' and 'bias in selection of the reported

result'. Reporting was inadequate in many of the included studies, which made the assessment of risk of

bias challenging. Notably, reporting of blinding, eligibility or selection criteria, as well as the enrollment

of participants (i.e., a lack of CONSORT flow diagrams) was poor. Individual risk of bias assessments

for each study design are provided in supplementary material (S4-9).

3.4. Evidence appraisal using GRADE

Overall, the GRADE assessment revealed a 'Low' certainty in the body of evidence for improvements

in all CRF outcomes (Table 6). The certainty rating for $A\dot{V}O_{2peak}$ was downgraded due to imprecision

and a lack of high quality study designs, whereas $R\dot{V}O_{2peak}$ was downgraded as a result of imprecision

and a high risk of bias in the RCTs. The confidence rating for PPO was downgraded due to imprecision

and inconsistency, resulting from considerable heterogeneity in the included exercise interventions.

562   *[PLEASE INSERT TABLE 6 HERE]*

564   **4. DISCUSSION**

565   This review provides a large evidence-based summary and appraisal on the effects of prescribed and

566   prospective exercise interventions >2 weeks on CRF in individuals with SCI. The results from the meta-

analysis support the role of exercise in improving CRF in this population by 0.22 L/min and 11W in $A\dot{V}O_{2peak}$ and PPO, respectively. The meta-analysis also indicates a clinically meaningful change in $R\dot{V}O_{2peak}$ of 2.8 mL/kg/min. However, the GRADE assessment revealed 'Low' certainty in the evidence for significant improvements in $A\dot{V}O_{2peak}$, $R\dot{V}O_{2peak}$, and PPO. Subgroup analyses revealed no effects of injury characteristics or exercise intervention parameters on $A\dot{V}O_{2peak}$ or $R\dot{V}O_{2peak}$. However, there were significant subgroup differences for PPO based on TSI, neurological level of injury, exercise modality, exercise intensity, method of exercise intensity prescription, and frequency of sessions.

4.1. Impact of injury characteristics

4.1.1. Time since injury

Following exercise interventions $\dot{V}O_{2peak}$ improves in individuals with both acute and chronic SCI. However, this review highlights the need for more exercise interventions in the acute phase post-SCI. Indeed, a recent review by Van der Scheer et al. [38] rated the confidence in the evidence base for exercise in acute SCI as 'Very Low', and called for more RCTs to control for the deteriorations in fitness and health occurring almost immediately following SCI. With regards to PPO in the current review, subgroup analysis based on TSI reveals that individuals with acute SCI exhibit a greater change than individuals with chronic SCI. This could be due to spontaneous motor recovery in the first few months following SCI [93], or speculatively, a familiarisation effect to novel modalities of exercise or additive upper-limb physiological adaptations in response to concurrent inpatient rehabilitation. To support this point, the secondary meta-analysis with longitudinal observational studies indicates a 6W improvement in PPO with standard of care inpatient rehabilitation during the subacute period. Ultimately, more rigorous RCTs are required in the subacute phase post-SCI that compare standard of care versus standard of care plus a specific exercise intervention to truly quantify improvements in CRF outcomes.

4.1.2. Neurological level of injury

Exercise results in improved $\dot{V}O_{2peak}$ regardless of the neurological level of injury. In particular, this review reveals a pooled improvement of 5.9 mL/kg/min in studies that included only individuals with tetraplegia (N=3). For comparison, there is a considerably larger evidence-base for studies including only individuals with paraplegia (N=28). A recent systematic review suggested that aerobic exercise results in minimal returns on investment in individuals with tetraplegia, with $\dot{V}O_{2peak}$ improving on average only

9% following 10-37 weeks of training [17]. However, their review excluded studies with a sample size

<10. Consequently, the Dicarlo study [94], which reported a 94% increase in $R\dot{V}O_{2peak}$ was excluded

from their analysis. Whilst the inclusion of this study in the current analysis may have augmented the

overall effect, our findings indicate that exercise improves CRF in individuals with tetraplegia and that

the magnitude of change is not significantly different to individuals with paraplegia. However, this meta-

analysis highlights that individuals with paraplegia (16W) are likely to accrue greater absolute changes

in PPO than those with tetraplegia (9W). Typically, higher neurological levels of injury result in a loss

of trunk control, motor impairments in the upper-limbs and reduced mechanical efficiency, compared to

lower levels of injury [95,96]. Therefore, individuals with tetraplegia may not have the physical or motor

capacity to adapt as effectively as individuals with paraplegia, and thus could experience a ceiling effect

with training. Indeed, a recent study identified lesion level as a significant predictor of PPO in a group

of handcyclists with SCI [97]. To account for baseline motor function differences between individuals

with tetraplegia and paraplegia, we determined relative percentage change for studies that included

upper-body aerobic exercise interventions only. The relative percentage change was similar between

neurological level of injury classifications: 46% tetraplegia (N=1) vs. 53% paraplegia (N=9). While only

one tetraplegia-only intervention was included in this subgroup analysis [98], normalising for baseline

values seems to indicate similar relative magnitudes of change in PPO.

Williams et al. [99] demonstrated that individuals with a lower level of injury (<T6) significantly

improved PPO compared to individuals with a higher level of injury (≥T6), suggesting a potential role

of disrupted cardiovascular control in mediating changes in PPO. Whilst methods for ameliorating the

reduction in sympathetic cardiovascular control typically associated with injuries ≥T6 have been

investigated (e.g., abdominal binding [100], lower-body positive pressure [101], and midodrine [102]),

the evidence for an improved CRF is still mixed. A recent case-report has indicated that epidural spinal

cord stimulation (SCS) can safely and effectively restore cardiovascular control and improve CRF [103].

With an explosion in SCS studies over the last few years [104], particularly including transcutaneous

SCS, the pairing of exercise with novel and non-invasive neuromodulatory approaches will likely

continue to receive considerable research attention. Future, adequately powered, research may want to

consider separating participants into paraplegia and tetraplegia groups or dichotomize by injuries above

and below T6 to account for differences in sympathetic cardiovascular control. Currently, there is a

paucity of studies analysing data in this fashion, which limits our understanding of how neurological

level of injury and the degree of impaired sympathetic cardiovascular control influences the magnitude

of change in CRF following an exercise intervention. Researchers may want to consider conducting a

battery of autonomic nervous system stress tests at baseline (e.g., Valsalva manoeuvre, head-up tilt,

sympathetic skin responses etc. [105]), to determine the degree of supraspinal sympathetic disruption

rather than relying on a neurological level of injury derived from a motor-sensory examination. This is

important as recent research has indicated that cardiovascular instability cannot be predicted by motor-

sensory level and completeness of SCI [106].

4.1.3. Injury severity

There were no significant subgroup differences in CRF. However, the subgroup analysis suggests that

individuals with a motor-incomplete SCI may not yield PPO improvements of the same magnitude as

individuals with a motor-complete SCI. This is most likely due to the majority of motor-incomplete

studies implementing gait training as its exercise modality, which we reveal is the least effective modality

for improving CRF. The gait training studies that measured PPO (N=2) used arm-crank ergometry (ACE)

as the CPET modality, demonstrating no transfer effect from lower-body to upper-body exercise. During

data extraction, reviewers noted a poor reporting of injury severity in a number of studies. Whilst this

may be due to older studies having used now outdated severity scales (e.g., International Stoke

Mandeville Games Federation or Frankel), researchers should endeavour to perform an International

Standards for Neurological Classification of SCI (ISNCSCI) exam during screening, and subsequently

report an AIS grade, to enable better comparisons to be made between injury severities in the future.

4.2. Impact of exercise intervention parameters

4.2.1. Exercise modality

Despite a number of recent reviews summarising the effects of specific exercise modalities on the change

in CRF following SCI, including aerobic ACE [107], FES-cycling [39], and aerobic plus muscle strength

training (mixed multimodal) interventions [108], this meta-analysis is the first to directly compare the

effects of a wide range of exercise modalities on the change in CRF in individuals with SCI.

This review revealed there were no significant subgroup differences between exercise modalities in

A$\dot{V}O_{2peak}$ or R$\dot{V}O_{2peak}$, indicating that improvements can be gained from any form of exercise

intervention. The change in R$\dot{V}O_{2peak}$ in the current review (21%) is equivalent to the average 21%

improvement reported in a recent systematic review on the effects of ACE in chronic SCI [107]. Whilst

the current review did not exclusively investigate ACE, it is evident that aerobic, volitional upper-body

exercise training can improve CRF in individuals with SCI. Activating larger amounts of skeletal muscle

mass via FES exercise interventions also appears to improve $\dot{V}O_{2peak}$, yet it is noteworthy that more

accessible and less expensive training modalities such as aerobic and resistance training may yield similar

or even greater increases in $\dot{V}O_{2peak}$, despite utilising less muscle mass. Additionally, $\dot{V}O_{2peak}$ improves

following multimodal/hybrid exercise interventions, which challenges a 2015 review reporting

inconclusive findings on the effects of combined upper-body aerobic and muscle strength training on

CRF [108]. Yet, as the current review included a wide range of interventions not restricted to the upper-

body (e.g., aquatic treadmill [54], hybrid cycling [55,60,109], multimodal exercises [110,111], etc.), it is

recommended that more research is conducted to delineate whether the improvements in $\dot{V}O_{2peak}$ with

multimodal/hybrid exercise interventions are due to the combination of upper- and lower-body exercise

modalities, or due to concurrent training modalities that predominantly use the upper-body (e.g., aerobic

plus muscle strength training). Finally, both gait training and behaviour change interventions appear less

effective at improving $\dot{V}O_{2peak}$ and PPO.

Aerobic, upper-body exercise and resistance training modalities demonstrate the greatest improvements

in PPO, by 15W and 20W, respectively. It is perhaps unsurprising that resistance training resulted in the

largest change in PPO given that these interventions included upper-body exercises prescribed to increase

muscular strength, as shown by Jacobs et al. [112]. Ultimately, improvements in PPO have important

ramifications for individuals with SCI that are dependent on performing explosive upper-body

movements during transfers or wheelchair propulsion [88,92], and may lead to increased quality of life

with more functional independence [113].

Several studies directly compared the effects of specific exercise modalities on the change in CRF

[54,76,114]. Notably, Gorman et al. [54] demonstrated that there were no transfer effects from a robotic

treadmill exercise intervention to ACE performance in a CPET. This review also demonstrates that

greater changes in $R\dot{V}O_{2peak}$ are likely achieved when the CPET modality is matched to the intervention

(S12). Therefore, researchers should endeavour to match the CPET modality to their exercise

intervention, or at the very least be careful when interpreting changes in CRF when using different

modalities.

4.2.2. Exercise intensity

The current SCI-specific exercise guidelines recommend that exercise should be performed at a

moderate-to-vigorous intensity [24]. A recent overview of systematic reviews also advocated the use of

moderate-to-vigorous intensity for improving aerobic fitness [115]. The current meta-analysis

demonstrates robust improvements across all CRF outcomes for interventions prescribing exercise at this

particular intensity. Furthermore, the secondary meta-analysis including cross-sectional studies reveals

significant associations of a greater magnitude between MVPA and CRF, as compared to lower-intensity

activity. Despite this, our classification of moderate-to-vigorous exercise intensity spans two of the

ACSM exercise intensity thresholds (S2). There may be considerable variation in the actual intensity

performed by participants given the noticeable range across thresholds (e.g., 46-90% $\dot{V}O_{2peak}$, 64-95%

HR_{peak}, 12-17 RPE etc.). Therefore, individuals with SCI and exercise practitioners should be cautious

when prescribing such a broad exercise intensity.

The secondary meta-analysis comparing RCT exercise intensities reveals similar changes in CRF

outcomes between moderate- and vigorous-intensity interventions. This is in agreement with a previous

review [33] and supports the viewpoint from a special communication on high-intensity interval training

(HIIT) [34], which suggested that vigorous-intensity exercise is more time efficient and may result in

similar if not superior CRF and skeletal muscle oxidative capacity improvements in comparison to

moderate-intensity exercise. Interestingly, in a response to a Letter-to-the-Editor [30], the SCI-specific

exercise guideline developers acknowledge the need for shorter, effective protocols to be documented in

the literature [116]. In the current review, a number of HIIT-based studies result in an improved CRF

[60,109,117–121]. Furthermore, recent evidence has suggested that HIIT may be more enjoyable than

moderate-intensity exercise for individuals with SCI [122]. Therefore, this form of training may offer a

more time efficient and readily available alternative to moderate-intensity protocols. However, in

echoing the thoughts of Astorino et al. [35], research must first corroborate its safety and feasibility in the SCI population before it can be recommended as an exercise strategy to improve CRF.

4.2.3. Exercise intensity prescription methods

This review reveals that $\dot{V}O_{2peak}$ improves regardless of the method used to prescribe exercise intensity. With regards to PPO, the subgroup difference indicates that the magnitude of change is greater when prescribing intensity via indices of HR (i.e., $\%HR_{peak}$, $\%HR_{max}$, $\%HRR$) or $\dot{V}O_2$ (i.e., $\%\dot{V}O_{2peak}$, $\%\dot{V}O_{2reserve}$), compared to RPE and workload. Previous research has revealed that RPE results in inter-individual responses to exercise, with the potential for two individuals to perform the same bout of exercise above or below lactate threshold despite being prescribed the same intensity, which prevents the development of SCI-specific RPE recommendations [123]. The difference in PPO may also be due to individuals with SCI being unaccustomed to subjective measures of exertion. Accordingly, recent systematic reviews have called for better reporting of the standardisation and familiarisation procedures used for RPE [124] and have only tentatively recommended its use before the evidence base is expanded [125]. Therefore, it seems plausible to suggest that the blunted improvements in PPO with intensity prescribed via RPE, as compared to other prescription methods, may have resulted from insufficient familiarisation before an exercise intervention.

Although HR and $\dot{V}O_2$ have long been used to prescribe exercise intensity, these approaches can result in large training ranges and ignore individual metabolic responses. Particularly, issues may arise with using HR for individuals with a neurological level of injury $\geq$T6, given that these individuals typically exhibit a lower HR_{peak} [126]. The use of fixed percentages (i.e, $\%HR_{peak}$, $\%\dot{V}O_{2peak}$) in the non-injured population has been questioned [127] and has recently been investigated in individuals with SCI, whereby Hutchinson et al. [45] showed that fixed $\%HR_{peak}$ and $\%\dot{V}O_{2peak}$ could not guarantee a homogenous domain-specific exercise intensity prescription. Notably, individuals were spread across moderate, heavy and severe domains at the "moderate" and "vigorous" intensity classifications; thereby questioning whether the "moderate-to-vigorous" terminology used in the SCI-specific exercise guidelines is suitable for adults with SCI.

Given that prescribing exercise intensity via HR and $\dot{V}O_2$ can typically be resource and cost-intensive,

there is some scope for using RPE as a cheaper and more practical method for community-based exercise

prescription. However, this may not be as effective as other objective methods. Future research should

aim to identify the optimal methods of exercise intensity prescription, as well as consider revisiting the

current "moderate-to-vigorous intensity" recommendations.

4.2.4. Frequency and exercise volume

Subgroup analyses based on frequency of sessions and exercise volume reveal no differences in $\dot{V}O_{2peak}$,

thereby supporting the minimal volume of exercise required to attain CRF benefits in individuals with

SCI. Furthermore, although there are no subgroup differences in PPO, the meta-regression identifies that

a greater volume of exercise is associated with greater changes in PPO. Indeed, there is a greater

magnitude of change observed for individuals exercising 90-149 min/wk in comparison to 40-89 min/wk

(12W vs 6W change, respectively). A greater weekly exercise volume may therefore accrue greater

changes in PPO and, as already described, may be important in improving the capacity to perform daily

tasks such as bed or wheelchair transfers [88,92].

Although changes in CRF are similar between each exercise volume subgroup, and thus exercise

guideline, the secondary meta-analysis on cross-sectional cohorts indicates a significant cumulative

impact of prolonged participation in physical activity and exercise. To support this point, a sensitivity

analysis revealed a larger difference in $R\dot{V}O_{2peak}$ between inactive individuals and elite athletes,

compared to between inactive and active individuals, suggesting that those who exercise more exhibit a

greater CRF. Indeed, a cross-sectional association study [87], using a wearable device to objectively

monitor habitual physical activity, reported a significantly higher CRF in those performing the general

population exercise guidelines (≥150 min/wk) compared to the SCI-specific fitness guidelines (40-89

min/wk). In fact, a recent study by Hoevenaars et al. [128] explored whether meeting the guidelines

proposed by Tweedy et al. [27] ("≥150 min/wk of moderate or ≥60 min/wk of vigorous exercise"), which

are nearly consistent with the general population exercise guidelines, is associated with greater health

and fitness benefits than the current SCI-specific guidelines by Martin-Ginis et al. [24]. Individuals

meeting the Tweedy guidelines had a significantly greater $A\dot{V}O_{2peak}$ and PPO than those meeting the

guidelines developed by Martin-Ginis et al. [24]. Looking forward, longitudinal RCTs with multiple

intervention arms would be the best way to explore dose-response changes with regards to differing

volumes of exercise, as has been done in the non-injured population [129–132].

4.3. Adverse events

Adverse events were reported for 4.1% of the total included participants, with the majority of events

related to skin sores, pressure sores or ulcers. Qualitatively, there was no particular exercise modality

that suggested an increased risk for an adverse event, but higher-intensity exercise appeared to reveal

more adverse events, albeit being swayed by one study in particular [111]. Reporting was poor in a

number of studies with reviewers at times unable to determine the exact number of events per participant.

Furthermore, there is generally a lack of follow-up assessments following exercise interventions, so it is

currently unknown whether there are any detrimental long-term effects of exercise in SCI.

4.4. Strengths and limitations of the review and future directions

4.4.1. Limitations of the included studies

Poor reporting of injury characteristics and exercise parameters prevented a perfect comparison of

exercise interventions. Overall, studies could have provided more precise descriptions of training

parameters to aid with any future refinements to the SCI-specific exercise guidelines. Reporting of

adherence to interventions was also poor and should be encouraged to provide an indication of the

feasibility or applicability of specific exercise interventions for individuals with SCI. Moreover, adverse

events should be transparently reported, even if none occur so that practitioners are able to identify forms

of exercise that are most likely to be safe for this population. Additionally, studies typically failed to

utilise the training principle of progression, which during prolonged exercise interventions is essential

for preventing a plateau in training adaptations and perhaps particularly important in this population for

supporting the transition from an inactive lifestyle to higher levels of activity, and ultimately achieving

greater CRF benefits [27]. On the whole, the reporting of $\dot{V}O_{2peak}$ attainment criteria was poor, with only

16% of the included exercise interventions using at least three criterion methods for identifying when an

individual had reached peak capacity. Thus, the magnitude of change in these studies could be inflated

or underestimated. Furthermore, to the best of our knowledge, only 30% of interventions had a

prospectively registered clinical trial entry and only 6.4% had a protocol manuscript published. To

sustain the integrity and transparency of reporting in this field, researchers are encouraged to
prospectively register any planned clinical trials using publicly available repositories.

The risk of bias assessments on pre-post studies revealed that no study conducted multiple baseline or
follow-up assessments. Whilst often time-consuming and impractical with larger sample sizes, multiple
assessments ensure reproducibility by accounting for any technical or biological variation, as shown
previously in non-injured individuals at risk for type-2 diabetes [133]. In the SCI population, individuals
are typically deconditioned and often exhibit variable responses to a CPET. This variance may be
explained by profound blood pressure instability [134], including unintentional 'boosting' via episodes
of autonomic dysreflexia [135]. Researchers should therefore consider performing multiple CPETs at
baseline and follow-up to attain reliable assessments of CRF.

There are also several limitations with regards to the studies included in the secondary meta-analyses for
this review. First, there is only one cross-sectional study using a wearable device to investigate the
association between physical activity and CRF [87]. Whilst self-report questionnaires are valid tools for
estimating levels of physical activity [86,136–138], there are important drawbacks including the
difficulty of accurately capturing intensity, lack of questionnaires measuring activities of daily living,
and recall bias. Secondly, there is a lack of RCTs comparing near-maximal, maximal or supramaximal
exercise intensities to moderate-intensity exercise. The only supramaximal intervention included in this
review demonstrated a 17W improvement in PPO [120]. The inclusion of more RCTs comparing
vigorous-intensity to lower intensity exercise could identify whether there are, in fact, benefits to
performing shorter but more vigorous-intensity exercise bouts, in comparison to longer continuous forms
of exercise.

4.4.2. Strengths and limitations of the review
A major strength of the current study is that we pre-planned and prospectively registered (PROSPERO
ID CRD42018104342) our systematic review. We used GRADE to assess the certainty in the body of
evidence and used quality appraisal tools for the specific study designs included in this review. Our
GRADE assessment demonstrates generalisability within the SCI population, through the inclusion of
participants across the lifespan and with a wide range of injury characteristics. Yet, the 'Low' confidence

in the evidence across all CRF outcomes emphasises the need for more rigorous exercise interventions

to address current gaps in the literature [38].

As there were not enough RCTs to perform a meta-regression on this study design specifically, we pooled

pre-post and RCT exercise interventions. The changes in $R\dot{V}O_{2peak}$ and PPO in the primary meta-analysis

(2.8 mL/kg/min and 11W, respectively) are somewhat similar to those reported with RCT interventions

relative to controls (2.9 mL/kg/min and 10W, respectively), and thus confirms the robustness of our

overall findings. Furthermore, our rigorous approach of adjusting for multiple comparisons minimises

any erroneous interpretations of subgroup differences and therefore strengthens our conclusions on the

available evidence.

Despite this, the categorisation of interventions within each subgroup could be considered a limitation of

the current review. Whilst this was done to directly compare the effects of different subgroups (i.e., acute

vs chronic, tetraplegia vs paraplegia, aerobic vs resistance vs FES etc.), it resulted in an unequal number

of interventions within each classification and likely underpowered the subgroup comparisons. For

example, the subgroup analysis based on exercise intensity reveals an effect of exercise intensity on PPO,

yet this may be influenced by the small number of interventions for light- and supramaximal-intensity.

Despite reporting some significant subgroup differences across dichotomised studies, these variables

were not identified as significant moderator variables in the random-effects meta-regression, meaning

these findings should be viewed with caution. It is perhaps more of a limitation of the evidence-base *per*

se, rather than our meta-analysis, in that more studies should be conducted to increase the power of these

subgroups and to ascertain whether there would be any significant improvements with a greater study

sample size.

Another limitation is that despite our comprehensive search strategy we may have missed relevant studies

as we did not search the grey literature and abstracts were not included. Finally, this review excluded

studies that were not published in English, introducing a source of language bias. However, of the full

texts screened for eligibility only 0.6% were excluded for being unavailable in English and is therefore

highly unlikely to have influenced the overall findings.

4.4.3. Implications and future directions

Our results support the current guidelines regarding the minimal weekly volume of exercise necessary to improve CRF in the SCI population. However, our pooled analysis indicates subgroup differences for PPO based on certain exercise intervention parameters. To the best of our knowledge, there are no large scale epidemiological studies investigating the dose-response relationship between physical activity and CRF in this population using sensitive and validated methods to quantify the exposure variable (e.g. free-living physical activity). Such studies have been performed in non-injured individuals [139,140]. To identify the optimal stimulus for beneficial CRF responses in this population, dose-ranging studies, akin to those that are used in the pharmaceutical industry, should be conducted. A recent overview of systematic reviews [141] highlighted the poor reporting in exercise interventions in health and disease and called upon the inclusion of checklists [e.g., the Consensus on Exercise Reporting Template (CERT) [142] or the Template for Intervention Description and Replication (TIDieR) [143]] to improve study quality. This would ultimately lead to a better understanding of the 'dose' of exercise as medicine required to optimise CRF outcomes in this population.

Exercise interventions >2 weeks result in an overall pooled increase in $R\dot{V}O_{2peak}$ of 2.8 mL/kg/min, which is roughly equivalent to 1 MET-SCI [metabolic equivalent in SCI (2.7 mL/kg/min)] [43]. An increase in maximal aerobic capacity (an estimate of CRF) by 1 MET (3.5 mL/kg/min) in non-injured individuals is associated with a 13% and 15% reduction in all-cause and cardiovascular mortality, respectively [144]. The current review shows that individuals meeting the SCI-specific guidelines for cardiometabolic health [24] can improve $R\dot{V}O_{2peak}$ to a similar magnitude to the overall pooled effect (~1 MET-SCI), highlighting that these guidelines may offer a reduction in CVD risk, and therefore mortality. Nonetheless, an association between an improvement in CRF and a reduction in mortality is yet to be established specifically in the SCI population, and remains an important avenue of research for the future.

5. CONCLUSION

This systematic review with meta-analysis provides an updated, evidence-based summary of the effects of exercise interventions on CRF in individuals with SCI. It reveals that exercise interventions >2 weeks are associated with significant improvements to CRF, and in particular, a clinically meaningful change in $R\dot{V}O_{2peak}$. Subgroup comparisons identified that upper-body aerobic exercise and resistance training

appear the most effective at improving PPO. Furthermore, acutely-injured, paraplegic individuals,

exercising at a moderate-to-vigorous intensity, prescribed via $\dot{V}O_2$ or HR, for more than 3 sessions/week

will likely experience the greatest change in PPO. Importantly, there is an ever-growing need for studies

to establish a dose-response relationship between exercise and CRF in the SCI population to determine

the most optimal form of exercise prescription to reduce the wide-ranging consequences typically

associated with SCI.

6. ACKNOWLEDGEMENTS

We would like to thank Mr. Sajjad Tavassoly (Faculty of Medicine, UBC, Vancouver, Canada) for his assistance with piloting the initial database search. We would also like to thank Dr. Matthew Querée (Department of Physical Therapy, UBC, Spinal Cord Injury Research Evidence Team, GF Strong Rehabilitation Centre, Vancouver, Canada) for his assistance with the risk of bias assessments. We appreciate the assistance of Mr Adrian Cheng (ICORD) and Dr. Alex Williams (ICORD), who helped pilot the inclusion/exclusion of articles and supported the use of Photoshop to extract data from figures, respectively. We thank Dr. Sam Weaver (University of Birmingham) for support with the meta-analysis using R.

7. DECLARATIONS

7.1. Author Contributions

Conceptualisation and study design were conducted by MW, AK and TN. Literature searches were completed by DH, GB and TN. Risk of bias and GRADE assessments were completed by DH, GB, SYC and TN. Statistical analysis and data interpretation were performed by DH, CL, CJ and TN. All authors contributed to the drafting and critical revision of the work.

7.2. Funding

Daniel Hodgkiss is supported by a Nathalie Rose Barr PhD studentship (#NRB123) from the International Spinal Research Trust (ISRT), a registered charity in the UK. Gurjeet Bhangu was supported by the Faculty of Medicine within the University of British Columbia (UBC) via the Flexible and Enhanced Learning Program. Carole Lunny's postdoctoral work is supported by a CIHR project grant (2021-2023). Catherine Jutzeler was supported by the Swiss National Science Foundation (Ambizione Grant [PZ00P3_186101]) and Wings for Life (Wings for Life Research Foundation [#ID 2020_118]). Tom Nightingale (grant number 17767) and Matthias Walter (grant number 17110) were recipients of Michael Smith Foundation for Health Research Trainee Awards in conjunction with the International Collaboration on Repair Discoveries and Rick Hansen Foundation, respectively.

7.3. Availability of data and materials

All data included in this systematic review can be provided upon request to the corresponding author.

7.4. Ethics approval and consent to participate

Not applicable.

7.5. Consent for publication

Not applicable.

7.6. Code availability

R scripts can be found on the Github repository: https://github.com/jutzca/Exercise-and-fitness-in-SCI

7.7. Competing interests

Daniel Hodgkiss, Gurjeet Bhangu, Carole Lunny, Catherine Jutzeler, Shin-Yi Chiou, Matthias Walter, Samuel Lucas, Andrei Krassioukov and Tom Nightingale declare that they have no competing interests relevant to the content of this article.

8. LIST OF ABBREVIATIONS

1RM	One repetition maximum
	Arm-crank ergometry
ACSM	American College of Sports Medicine
AIS	American Spinal Injury Association Impairment Scale
A$\dot{V}O_{2peak}$	Absolute peak oxygen uptake
CENTRAL	Cochrane Central Register of Controlled Trials
CERT	Consensus on Exercise Reporting Template
CI	Confidence interval
CPET	Cardiopulmonary exercise test
CRF	Cardiorespiratory fitness
CVD	Cardiovascular disease
EMBASE	Excerpta Medica Database
FES	Functional electrical stimulation
GRADE	Grading of Recommendations, Assessment, Development and Evaluation
HIIT	High-intensity interval training
HR	Heart rate
HR$_{max}$	Maximum heart rate (age-predicted)
HR$_{peak}$	Peak heart rate
HRR	Heart rate reserve
IQR	Interquartile range
ISNCSCI	International Standards for Neurological Classification of Spinal Cord Injury
LTPA	Leisure time physical activity
MET	Metabolic equivalent
MTP	Maximal tolerated power
MVPA	Moderate-to-vigorous physical activity
PPO	Peak power output
PRISMA	Preferred Reporting Items for Systematic Reviews

RCT	Randomised-controlled trial	
RoB 2	The Cochrane Risk of Bias 2 tool	
RPE	Rating of perceived exertion	
R$\dot{V}O_{2peak}$	Relative peak oxygen uptake	
SCI	Spinal cord injury	
SCS	Spinal cord stimulation	
SD	Standard deviation	
TIDieR	Template for Intervention Description and Replication	
TSI	Time since injury	
$\dot{V}O_2$	Oxygen uptake	
$\dot{V}O_{2peak}$	Peak oxygen uptake	
$\dot{V}O_{2reserve}$	Reserve oxygen uptake	
W	Watts	
WMD	Weighted mean difference	

9. REFERENCES

1   Miller LE, Herbert WG. Health and economic benefits of physical activity for patients with spinal

cord injury. *Clinicoecon Outcomes Res* 2016;**8**:551–8.

2   Rimmer JH, Schiller W, Chen M-D. Effects of disability-associated low energy expenditure

deconditioning syndrome. *Exerc Sport Sci Rev* 2012;**40**:22–9.

3   Cragg JJ, Noonan VK, Dvorak M, *et al.* Spinal cord injury and type 2 diabetes: results from a

population health survey. *Neurology* 2013;**81**:1864–8.

4   Cragg JJ, Noonan VK, Krassioukov A, *et al.* Cardiovascular disease and spinal cord injury:

results from a national population health survey. *Neurology* 2013;**81**:723–8.

5   Wu J-C, Chen Y-C, Liu L, *et al.* Increased risk of stroke after spinal cord injury: a nationwide 4-

year follow-up cohort study. *Neurology* 2012;**78**:1051–7.

6   Ross R, Blair SN, Arena R, *et al.* Importance of Assessing Cardiorespiratory Fitness in Clinical

Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the American

Heart Association. *Circulation* 2016;**134**:e653–99.

7   Clausen Johan S.R., Marott Jacob L., Holtermann Andreas, *et al.* Midlife Cardiorespiratory

Fitness and the Long-Term Risk of Mortality. *J Am Coll Cardiol* 2018;**72**:987–95.

8   Lee D-C, Sui X, Artero EG, *et al.* Long-term effects of changes in cardiorespiratory fitness and

body mass index on all-cause and cardiovascular disease mortality in men: the Aerobics Center

Longitudinal Study. *Circulation* 2011;**124**:2483–90.

9   Myers J, Prakash M, Froelicher V, *et al.* Exercise capacity and mortality among men referred for

exercise testing. *N Engl J Med* 2002;**346**:793–801.

10   Laukkanen JA, Kurl S, Salonen R, *et al.* The predictive value of cardiorespiratory fitness for

cardiovascular events in men with various risk profiles: a prospective population-based cohort

study. *Eur Heart J* 2004;**25**:1428–37.

11   Erikssen G, Liestøl K, Bjørnholt J, *et al.* Changes in physical fitness and changes in mortality.

1022  *Lancet* 1998;**352**:759–62.

1023 12 Lee DC, Sui X, Ortega FB, *et al.* Comparisons of leisure-time physical activity and

1024   cardiorespiratory fitness as predictors of all-cause mortality in men and women. *Br J Sports Med*

1025   2011;**45**:504–10.

1026 13 Haisma JA, van der Woude LHV, Stam HJ, *et al.* Physical capacity in wheelchair-dependent

1027   persons with a spinal cord injury: a critical review of the literature. *Spinal Cord* 2006;**44**:642–52.

1028 14 Simmons OL, Kressler J, Nash MS. Reference fitness values in the untrained spinal cord injury

1029   population. *Arch Phys Med Rehabil* 2014;**95**:2272–8.

1030 15 Cruz S, Blauwet CA. Implications of altered autonomic control on sports performance in athletes

1031   with spinal cord injury. *Auton Neurosci* 2018;**209**:100–4.

1032 16 West CR, Gee CM, Voss C, *et al.* Cardiovascular control, autonomic function, and elite

1033   endurance performance in spinal cord injury. *Scand J Med Sci Sports* 2015;**25**:476–85.

1034 17 Figoni SF, Dolbow DR, Crawford EC, *et al.* Does aerobic exercise benefit persons with

1035   tetraplegia from spinal cord injury? A systematic review. *J Spinal Cord Med* 2021;**44**:690–703.

1036 18 Stofan JR, DiPietro L, Davis D, *et al.* Physical activity patterns associated with cardiorespiratory

1037   fitness and reduced mortality: the Aerobics Center Longitudinal Study. *Am J Public Health*

1038   1998;**88**:1807–13.

1039 19 Nightingale TE, Williams S, Thompson D, *et al.* Energy balance components in persons with

1040   paraplegia: daily variation and appropriate measurement duration. *Int J Behav Nutr Phys Act*

1041   2017;**14**:132.

1042 20 van den Berg-Emons RJ, Bussmann JB, Stam HJ. Accelerometry-based activity spectrum in

1043   persons with chronic physical conditions. *Arch Phys Med Rehabil* 2010;**91**:1856–61.

1044 21 Hicks AL, Martin Ginis KA, Pelletier CA, *et al.* The effects of exercise training on physical

1045   capacity, strength, body composition and functional performance among adults with spinal cord

1046   injury: a systematic review. *Spinal Cord* 2011;**49**:1103–27.

22　Valent L, Dallmeijer A, Houdijk H, *et al.* The effects of upper body exercise on the physical

capacity of people with a spinal cord injury: a systematic review. *Clin Rehabil* 2007;**21**:315–30.

23　Ginis KAM, Hicks AL, Latimer AE, *et al.* The development of evidence-informed physical

activity guidelines for adults with spinal cord injury. *Spinal Cord* 2011;**49**:1088–96.

24　Martin Ginis KA, van der Scheer JW, Latimer-Cheung AE, *et al.* Evidence-based scientific

exercise guidelines for adults with spinal cord injury: an update and a new guideline. *Spinal Cord*

2018;**56**:308–21.

25　Smith B, Kirby N, Skinner B, *et al.* Infographic. Physical activity for disabled adults. *Br J Sports*

Med 2019;**53**:335–6.

26　Bull FC, Al-Ansari SS, Biddle S, *et al.* World Health Organization 2020 guidelines on physical

activity and sedentary behaviour. *Br J Sports Med* 2020;**54**:1451–62.

27　Tweedy SM, Beckman EM, Geraghty TJ, *et al.* Exercise and sports science Australia (ESSA)

position statement on exercise and spinal cord injury. *J Sci Med Sport* 2017;**20**:108–15.

28　Piercy KL, Troiano RP, Ballard RM, *et al.* The Physical Activity Guidelines for Americans.

JAMA 2018;**320**:2020–8.

29　Powell KE, Paluch AE, Blair SN. Physical activity for health: What kind? How much? How

intense? On top of what? *Annu Rev Public Health* 2011;**32**:349–65.

30　Tweedy SM, Beckman EM, Connick MJ, *et al.* Correspondence re 'Evidence-based scientific

exercise guidelines for adults with spinal cord injury: an update and new guideline'. Spinal Cord.

2018;**56**:406–8.

31　Yue T, Wang Y, Liu H, *et al.* Effects of High-Intensity Interval vs. Moderate-Intensity

Continuous Training on Cardiac Rehabilitation in Patients With Cardiovascular Disease: A

Systematic Review and Meta-Analysis. *Front Cardiovasc Med* 2022;**9**:845225.

32　Hannan AL, Hing W, Simas V, *et al.* High-intensity interval training versus moderate-intensity

continuous training within cardiac rehabilitation: a systematic review and meta-analysis. *Open*

Access J Sports Med 2018;**9**:1–17.

33 Peters J, Abou L, Rice LA, *et al.* The effectiveness of vigorous training on cardiorespiratory

fitness in persons with spinal cord injury: a systematic review and meta-analysis. *Spinal Cord*

2021;**59**:1035–44.

34 Nightingale TE, Metcalfe RS, Vollaard NB, *et al.* Exercise Guidelines to Promote

Cardiometabolic Health in Spinal Cord Injured Humans: Time to Raise the Intensity? *Arch Phys*

Med Rehabil 2017;**98**:1693–704.

35 Astorino TA, Hicks AL, Bilzon JLJ. Viability of high intensity interval training in persons with

spinal cord injury-a perspective review. *Spinal Cord* 2021;**59**:3–8.

36 Farrow M, Nightingale TE, Maher J, *et al.* Effect of Exercise on Cardiometabolic Risk Factors in

Adults With Chronic Spinal Cord Injury: A Systematic Review. *Arch Phys Med Rehabil*

2020;**101**:2177–205.

37 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-

analyses: the PRISMA statement. *BMJ* 2009;**339**:b2535.

38 van der Scheer JW, Martin Ginis KA, Ditor DS, *et al.* Effects of exercise on fitness and health of

adults with spinal cord injury: A systematic review. *Neurology* 2017;**89**:736–45.

39 van der Scheer JW, Goosey-Tolfrey VL, Valentino SE, *et al.* Functional electrical stimulation

cycling exercise after spinal cord injury: a systematic review of health and fitness-related

outcomes. *J Neuroeng Rehabil* 2021;**18**:99.

40 American College Medicine. *ACSM's guidelines for exercise testing and prescription*. 9th ed.

Lippincott Williams & Wilkins 2013. https://shop.lww.com/ACSM-s-Guidelines-for-Exercise-

Testing-and-Prescription/p/9781975150181 (accessed 3 Dec 2021).

41 Higgins JPT, Li T, Deeks JJ. Choosing effect measures and computing estimates of effect.

Cochrane Handbook for Systematic Reviews of Interventions. 2019;:143–76.

doi:10.1002/9781119536604.ch6

42 Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and

the size of a sample. *BMC Med Res Methodol* 2005;**5**:13.

43   Collins EG, Gater D, Kiratli J, *et al.* Energy cost of physical activities in persons with spinal cord

1100   injury. *Med Sci Sports Exerc* 2010;**42**:691–700.

1101   44   Higgins JPT, Thomas J, Chandler J, *et al. Cochrane Handbook for Systematic Reviews of*

1102   *Interventions*. John Wiley & Sons 2019.

1103   45   Hutchinson MJ, Goosey-Tolfrey VL. Rethinking aerobic exercise intensity prescription in adults

1104   with spinal cord injury: time to end the use of 'moderate to vigorous' intensity? *Spinal Cord*

1105   2021;:1–7.

1106   46   Higgins JPT, Thompson SG. Controlling the risk of spurious findings from meta-regression. *Stat*

1107   *Med* 2004;**23**:1663–82.

1108   47   Higgins JPT, Sterne JAC, Savovic J, *et al.* A revised tool for assessing risk of bias in randomized

1109   trials. *Cochrane Database Syst Rev* 2016;**10**:29–31.

1110   48   McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web

1111   app for visualizing risk-of-bias assessments. *Res Synth Methods* 2021;**12**:55–61.

1112   49   Guyatt GH, Oxman AD, Vist GE, *et al.* GRADE: an emerging consensus on rating quality of

1113   evidence and strength of recommendations. *BMJ* 2008;**336**:924–6.

1114   50   Han D-S, Hsiao M-Y, Wang T-G, *et al.* Association of serum myokines and aerobic exercise

1115   training in patients with spinal cord injury: an observational study. *BMC Neurol* 2016;**16**:142.

1116   51   Hoekstra F, van Nunen MPM, Gerrits KHL, *et al.* Effect of robotic gait training on

1117   cardiorespiratory system in incomplete spinal cord injury. *J Rehabil Res Dev* 2013;**50**:1411–22.

1118   52   Pollack SF, Axen K, Spielholz N, *et al.* Aerobic training effects of electrically induced lower

1119   extremity exercises in spinal cord injured people. *Arch Phys Med Rehabil* 1989;**70**:214–9.

1120   53   Barstow TJ, Scremin AM, Mutton DL, *et al.* Changes in gas exchange kinetics with training in

1121   patients with spinal cord injury. *Med Sci Sports Exerc* 1996;**28**:1221–8.

1122   54   Gorman PH, Scott W, VanHiel L, *et al.* Comparison of peak oxygen consumption response to

1123   aquatic and robotic therapy in individuals with chronic motor incomplete spinal cord injury: a

randomized controlled trial. *Spinal Cord* 2019;**57**:471–81.

55 Heesterbeek PJC, Berkelmans HWA, Thijssen DHJ, *et al.* Increased physical fitness after 4-week

training on a new hybrid FES-cycle in persons with spinal cord injury. *Technol Disabil*

2005;**17**:103–10.

56 Mutton DL, Scremin AM, Barstow TJ, *et al.* Physiologic responses during functional electrical

stimulation leg cycling and hybrid exercise in spinal cord injured subjects. *Arch Phys Med*

Rehabil 1997;**78**:712–8.

57 Ozturk ED, Lapointe MS, Kim D-I, *et al.* Effect of 6-Month Exercise Training on Neurovascular

Function in Spinal Cord Injury. *Med Sci Sports Exerc* 2021;**53**:38–46.

58 Hooker SP, Scremin AM, Mutton DL, *et al.* Peak and submaximal physiologic responses

following electrical stimulation leg cycle ergometer training. *J Rehabil Res Dev* 1995;**32**:361–6.

59 Hooker SP, Figoni SF, Rodgers MM, *et al.* Physiologic effects of electrical stimulation leg cycle

exercise training in spinal cord injured persons. *Arch Phys Med Rehabil* 1992;**73**:470–6.

60 Hasnan N, Engkasan JP, Husain R, *et al.* High-Intensity Virtual-reality Arm plus FES-leg Interval

Training in Individuals with Spinal Cord Injury. *Biomed Tech* 2013;**58 Suppl 1**.

doi:10.1515/bmt-2013-4028

61 Janssen TWJ, Pringle DD. Effects of modified electrical stimulation-induced leg cycle ergometer

training for individuals with spinal cord injury. *J Rehabil Res Dev* 2008;**45**:819–30.

62 Jeon JY, Hettinga D, Steadward RD, *et al.* Reduced plasma glucose and leptin after 12 weeks of

functional electrical stimulation-rowing exercise training in spinal cord injury patients. *Arch Phys*

Med Rehabil 2010;**91**:1957–9.

63 Midha M, Schmitt JK, Sclater M. Exercise effect with the wheelchair aerobic fitness trainer on

conditioning and metabolic function in disabled persons: a pilot study. *Arch Phys Med Rehabil*

1999;**80**:258–61.

64 Milia R, Roberto S, Marongiu E, *et al.* Improvement in hemodynamic responses to metaboreflex

activation after one year of training in spinal cord injured humans. *Biomed Res Int*

2014;**2014**:893468.

65   Solinsky R, Mercier H, Picard G, *et al.* Cardiometabolic Effects of High-Intensity Hybrid

Functional Electrical Stimulation Exercise after Spinal Cord Injury. *PM R* 2021;**13**:937–44.

66   Lavado EL, Cardoso JR, Silva LGA, *et al.* Effectiveness of aerobic physical training for treatment

of chronic asymptomatic bacteriuria in subjects with spinal cord injury: a randomized controlled

trial. *Clin Rehabil* 2013;**27**:142–9.

67   Brazg G, Fahey M, Holleran CL, *et al.* Effects of Training Intensity on Locomotor Performance

in Individuals With Chronic Spinal Cord Injury: A Randomized Crossover Study. *Neurorehabil*

Neural Repair 2017;**31**:944–54.

68   Beillot J, Carré F, Le Claire G, *et al.* Energy consumption of paraplegic locomotion using

reciprocating gait orthosis. *Eur J Appl Physiol Occup Physiol* 1996;**73**:376–81.

69   Bombardier CH, Dyer JR, Burns P, *et al.* A tele-health intervention to increase physical fitness in

people with spinal cord injury and cardiometabolic disease or risk factors: a pilot randomized

controlled trial. *Spinal Cord* 2021;**59**:63–73.

70   Kooijmans H, Post MWM, Stam HJ, *et al.* Effectiveness of a Self-Management Intervention to

Promote an Active Lifestyle in Persons With Long-Term Spinal Cord Injury: The HABITS

Randomized Clinical Trial. *Neurorehabil Neural Repair* 2017;**31**:991–1004.

71   Ma JK, West CR, Martin Ginis KA. The Effects of a Patient and Provider Co-Developed,

Behavioral Physical Activity Intervention on Physical Activity, Psychosocial Predictors, and

Fitness in Individuals with Spinal Cord Injury: A Randomized Controlled Trial. *Sports Med*

2019;**49**:1117–31.

72   Nooijen CF, Stam HJ, Sluis T, *et al.* A behavioral intervention promoting physical activity in

people with subacute spinal cord injury: secondary effects on health, social participation and

quality of life. *Clin Rehabil* 2017;**31**:772–80.

73   Bakkum AJT, de Groot S, Stolwijk-Swüste JM, *et al.* Effects of hybrid cycling versus

handcycling on wheelchair-specific fitness and physical activity in people with long-term spinal

cord injury: a 16-week randomized controlled trial. *Spinal Cord* 2015;**53**:395–401.

74   Capodaglio P, Grilli C, Bazzini G. Tolerable exercise intensity in the early rehabilitation of
paraplegic patients. A preliminary study. *Spinal Cord* 1996;**34**:684–90.

75   Gorgey AS, Lai RE, Khalil RE, *et al.* Neuromuscular electrical stimulation resistance training
enhances oxygen uptake and ventilatory efficiency independent of mitochondrial complexes after
spinal cord injury: a randomized clinical trial. *J Appl Physiol* 2021;**131**:265–76.

76   Alrashidi AA, Nightingale TE, Currie KD, *et al.* Exercise Improves Cardiorespiratory Fitness, but
Not Arterial Health, after Spinal Cord Injury: The CHOICES Trial. *J Neurotrauma*
2021;**38**:3020–9.

77   Berry HR, Perret C, Saunders BA, *et al.* Cardiorespiratory and power adaptations to stimulated
cycle training in paraplegia. *Med Sci Sports Exerc* 2008;**40**:1573–80.

78   Duffell LD, Paddison S, Alahmary AF, *et al.* The effects of FES cycling combined with virtual
reality racing biofeedback on voluntary function after incomplete SCI: a pilot study. *J Neuroeng*
Rehabil 2019;**16**:149.

79   Durán FS, Lugo L, Ramírez L, *et al.* Effects of an exercise program on the rehabilitation of
patients with spinal cord injury. *Arch Phys Med Rehabil* 2001;**82**:1349–54.

80   Gorman PH, Scott W, York H, *et al.* Robotically assisted treadmill exercise training for
improving peak fitness in chronic motor incomplete spinal cord injury: A randomized controlled
trial. *J Spinal Cord Med* 2016;**39**:32–44.

81   Gibbons RS, Stock CG, Andrews BJ, *et al.* The effect of FES-rowing training on cardiac structure
and function: pilot studies in people with spinal cord injury. *Spinal Cord* 2016;**54**:822–9.

82   Janssen TWJ, Dallmeijer AJ, Veeger DJHEJ, *et al.* Normative values and determinants of
physical capacity in individuals with spinal cord injury. *J Rehabil Res Dev* 2002;**39**:29–39.

83   Lannem AM, Sørensen M, Lidal IB, *et al.* Perceptions of exercise mastery in persons with
complete and incomplete spinal cord injury. *Spinal Cord* 2010;**48**:388–92.

84   Latimer AE, Ginis KAM, Craven BC, *et al.* The physical activity recall assessment for people

with spinal cord injury: validity. *Med Sci Sports Exerc* 2006;**38**:208–16.

85   Manns PJ, McCubbin JA, Williams DP. Fitness, inflammation, and the metabolic syndrome in

men with paraplegia. *Arch Phys Med Rehabil* 2005;**86**:1176–81.

86   Martin Ginis KA, Úbeda-Colomer J, Alrashidi AA, *et al.* Construct validation of the leisure time

physical activity questionnaire for people with SCI (LTPAQ-SCI). *Spinal Cord* 2021;**59**:311–8.

87   Nightingale TE, Walhin J-P, Thompson D, *et al.* Biomarkers of cardiometabolic health are

associated with body composition characteristics but not physical activity in persons with spinal

cord injury. *J Spinal Cord Med* 2019;**42**:328–37.

88   Haisma JA, Post MW, van der Woude LH, *et al.* Functional independence and health-related

functional status following spinal cord injury: a prospective study of the association with physical

capacity. *J Rehabil Med* 2008;**40**:812–8.

89   Leving MT, de Groot S, Woldring FAB, *et al.* Motor learning outcomes of handrim wheelchair

propulsion during active spinal cord injury rehabilitation in comparison with experienced

wheelchair users. *Disabil Rehabil* 2021;**43**:1429–42.

90   Stewart MW, Melton-Rogers SL, Morrison S, *et al.* The measurement properties of fitness

measures and health status for persons with spinal cord injuries. *Arch Phys Med Rehabil*

2000;**81**:394–400.

91   Dallmeijer AJ, van der Woude LH, Hollander PA, *et al.* Physical performance in persons with

spinal cord injuries after discharge from rehabilitation. *Med Sci Sports Exerc* 1999;**31**:1111–7.

92   Janssen TW, van Oers CA, Rozendaal EP, *et al.* Changes in physical strain and physical capacity

in men with spinal cord injuries. *Med Sci Sports Exerc* 1996;**28**:551–9.

93   Steeves JD, Kramer JK, Fawcett JW, *et al.* Extent of spontaneous motor recovery after traumatic

cervical sensorimotor complete spinal cord injury. *Spinal Cord* 2011;**49**:257–65.

94   DiCarlo SE. Effect of arm ergometry training on wheelchair propulsion endurance of individuals

with quadriplegia. *Phys Ther* 1988;**68**:40–4.

95 de Groot S, Dallmeijer AJ, Kilkens OJ, *et al.* Course of gross mechanical efficiency in handrim

wheelchair propulsion during rehabilitation of people with spinal cord injury: a prospective cohort

study. *Arch Phys Med Rehabil* 2005;**86**:1452–60.

96 Sprigle S, Maurer C, Holowka M. Development of valid and reliable measures of postural

stability. *J Spinal Cord Med* 2007;**30**:40–9.

97 Kouwijzer I, Valent L, Osterthun R, *et al.* Peak power output in handcycling of individuals with a

chronic spinal cord injury: predictive modeling, validation and reference values. *Disabil Rehabil*

2020;**42**:400–9.

98 Hjeltnes N, Wallberg-Henriksson H. Improved work capacity but unchanged peak oxygen uptake

during primary rehabilitation in tetraplegic patients. *Spinal Cord* 1998;**36**:691–8.

99 Williams AM, Ma JK, Martin Ginis KA, *et al.* Effects of a Tailored Physical Activity

Intervention on Cardiovascular Structure and Function in Individuals With Spinal Cord Injury.

Neurorehabil Neural Repair 2021;**35**:692–703.

100 West CR, Campbell IG, Goosey-Tolfrey VL, *et al.* Effects of abdominal binding on field-based

exercise responses in Paralympic athletes with cervical spinal cord injury. *J Sci Med Sport*

2014;**17**:351–5.

101 Pitetti KH, Barrett PJ, Campbell KD, *et al.* The effect of lower body positive pressure on the

exercise capacity of individuals with spinal cord injury. *Med Sci Sports Exerc* 1994;**26**:463–8.

102 Nieshoff EC, Birk TJ, Birk CA, *et al.* Double-blinded, placebo-controlled trial of midodrine for

exercise performance enhancement in tetraplegia: a pilot study. *J Spinal Cord Med* 2004;**27**:219–

25.

103 Nightingale TE, Walter M, Williams AMM, *et al.* Ergogenic effects of an epidural

neuroprosthesis in one individual with spinal cord injury. *Neurology* 2019;**92**:338–40.

104 Laskin JJ, Waheed Z, Thorogood NP, *et al.* Spinal Cord Stimulation Research in the Restoration

of Motor, Sensory, and Autonomic Function for Individuals Living With Spinal Cord Injuries: A

Scoping Review. *Arch Phys Med Rehabil* Published Online First: 23 February 2022.

doi:10.1016/j.apmr.2022.01.161

105 Berger MJ, Kimpinski K, Currie KD, *et al.* Multi-Domain Assessment of Autonomic Function in
Spinal Cord Injury Using a Modified Autonomic Reflex Screen. *J Neurotrauma* 2017;**34**:2624–
33.

106 Wang S, Wecht JM, Legg Ditterline B, *et al.* Heart rate and blood pressure response improve the
prediction of orthostatic cardiovascular dysregulation in persons with chronic spinal cord injury.
Physiol Rep 2020;**8**:e14617.

107 Chiou SY, Clarke E, Lam C, *et al.* Effects of Arm-Crank Exercise on Fitness and Health in
Adults With Chronic Spinal Cord Injury: A Systematic Review. *Front Physiol* 2022;**13**:831372.

108 Bochkezanian V, Raymond J, de Oliveira CQ, *et al.* Can combined aerobic and muscle strength
training improve aerobic fitness, muscle strength, function and quality of life in people with
spinal cord injury? A systematic review. *Spinal Cord* 2015;**53**:418–31.

109 Brurok B, Helgerud J, Karlsen T, *et al.* Effect of aerobic high-intensity hybrid training on stroke
volume and peak oxygen consumption in men with spinal cord injury. *Am J Phys Med Rehabil*
2011;**90**:407–14.

110 Gant KL, Nagle KG, Cowan RE, *et al.* Body System Effects of a Multi-Modal Training Program
Targeting Chronic, Motor Complete Thoracic Spinal Cord Injury. *J Neurotrauma* 2018;**35**:411–
23.

111 Lotter JK, Henderson CE, Plawecki A, *et al.* Task-Specific Versus Impairment-Based Training on
Locomotor Performance in Individuals With Chronic Spinal Cord Injury: A Randomized
Crossover Study. *Neurorehabil Neural Repair* 2020;**34**:627–39.

112 Jacobs PL, Nash MS, Rusinowski JW. Circuit training provides cardiorespiratory and strength
benefits in persons with paraplegia. *Med Sci Sports Exerc* 2001;**33**:711–7.

113 Sweet SN, Martin Ginis KA, Tomasone JR. Investigating intermediary variables in the physical
activity and quality of life relationship in persons with spinal cord injury. *Health Psychol*
2013;**32**:877–85.

114 Farkas GJ, Gorgey AS, Dolbow DR, *et al.* Energy Expenditure, Cardiorespiratory Fitness, and

Body Composition Following Arm Cycling or Functional Electrical Stimulation Exercises in

Spinal Cord Injury: A 16-Week Randomized Controlled Trial. *Top Spinal Cord Inj Rehabil*

2021;**27**:121–34.

115 Eitivipart AC, de Oliveira CQ, Arora M, *et al.* Overview of Systematic Reviews of Aerobic

Fitness and Muscle Strength Training after Spinal Cord Injury. *J Neurotrauma* 2019;**36**:2943–63.

116 Martin Ginis KA, van der Scheer JW, Latimer-Cheung AE, *et al.* Response to correspondence

from the ESSA Statement authors. Spinal Cord. 2018;**56**:409–11.

117 Tordi N, Dugue B, Klupzinski D, *et al.* Interval training program on a wheelchair ergometer for

paraplegic subjects. *Spinal Cord* 2001;**39**:532–7.

118 de Groot PCE, Hjeltnes N, Heijboer AC, *et al.* Effect of training intensity on physical capacity,

lipid profile and insulin sensitivity in early rehabilitation of spinal cord injured individuals. *Spinal*

Cord 2003;**41**:673–9.

119 Gauthier C, Brosseau R, Hicks AL, *et al.* Feasibility, Safety, and Preliminary Effectiveness of a

Home-Based Self-Managed High-Intensity Interval Training Program Offered to Long-Term

Manual Wheelchair Users. *Rehabil Res Pract* 2018;**2018**:8209360.

120 Mcleod JC, Diana H, Hicks AL. Sprint interval training versus moderate-intensity continuous

training during inpatient rehabilitation after spinal cord injury: a randomized trial. *Spinal Cord*

2020;**58**:106–15.

121 Graham K, Yarar-Fisher C, Li J, *et al.* Effects of High-Intensity Interval Training Versus

Moderate-Intensity Training on Cardiometabolic Health Markers in Individuals With Spinal Cord

Injury: A Pilot Study. *Top Spinal Cord Inj Rehabil* 2019;**25**:248–59.

122 Astorino TA, Thum JS. Within-session responses to high-intensity interval training in spinal cord

injury. *Disabil Rehabil* 2018;**40**:444–9.

123 Hutchinson MJ, Kouwijzer I, de Groot S, *et al.* Comparison of two Borg exertion scales for

monitoring exercise intensity in able-bodied participants, and those with paraplegia and

tetraplegia. *Spinal Cord* 2021;**59**:1162–9.

124 Valentino SE, Hutchinson MJ, Goosey-Tolfrey VL, *et al.* Effects of Perceptually Regulated

Exercise Training on Cardiorespiratory Fitness and Peak Power Output in Adults With Spinal

Cord Injury: A Systematic Review and Meta-analysis. *Arch Phys Med Rehabil* Published Online

First: 6 April 2022. doi:10.1016/j.apmr.2022.03.008

125 van der Scheer JW, Hutchinson MJ, Paulson T, *et al.* Reliability and Validity of Subjective

Measures of Aerobic Intensity in Adults With Spinal Cord Injury: A Systematic Review. *PM R*

2018;**10**:194–207.

126 Cowley KC. A new conceptual framework for the integrated neural control of locomotor and

sympathetic function: implications for exercise after spinal cord injury. *Appl Physiol Nutr Metab*

2018;**43**:1140–50.

127 Jamnick NA, Pettitt RW, Granata C, *et al.* An Examination and Critique of Current Methods to

Determine Exercise Intensity. *Sports Med* 2020;**50**:1729–56.

128 Hoevenaars D, Holla JFM, Postma K, *et al.* Associations between meeting exercise guidelines,

physical fitness, and health in people with spinal cord injury. *Disabil Rehabil* 2022;:1–8.

129 Tjønna AE, Leinan IM, Bartnes AT, *et al.* Low- and high-volume of intensive endurance training

significantly improves maximal oxygen uptake after 10-weeks of training in healthy men. *PLoS*

One 2013;**8**:e65382.

130 Stensvold D, Viken H, Steinshamn SL, *et al.* Effect of exercise training for five years on all cause

mortality in older adults-the Generation 100 study: randomised controlled trial. *BMJ*

2020;**371**:m3485.

131 Church TS, Earnest CP, Skinner JS, *et al.* Effects of different doses of physical activity on

cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with

elevated blood pressure: a randomized controlled trial. *JAMA* 2007;**297**:2081–91.

132 Foulds HJA, Bredin SSD, Charlesworth SA, *et al.* Exercise volume and intensity: a dose-response

relationship with health benefits. *Eur J Appl Physiol* 2014;**114**:1563–71.

133   Phillips BE, Kelly BM, Lilja M, *et al.* A Practical and Time-Efficient High-Intensity Interval

Training Program Modifies Cardio-Metabolic Risk Factors in Adults with Risk Factors for Type

II Diabetes. *Front Endocrinol* 2017;**8**:229.

134   Katzelnick CG, Weir JP, Jones A, *et al.* Blood Pressure Instability in Persons With SCI: Evidence

From a 30-Day Home Monitoring Observation. *Am J Hypertens* 2019;**32**:938–44.

135   Nightingale TE, Eginyan G, Balthazaar SJT, *et al.* Accidental boosting in an individual with

tetraplegia - considerations for the interpretation of cardiopulmonary exercise testing. *J Spinal*

Cord Med 2021;:1–6.

136   Martin Ginis KA, Phang SH, Latimer AE, *et al.* Reliability and validity tests of the leisure time

physical activity questionnaire for people with spinal cord injury. *Arch Phys Med Rehabil*

2012;**93**:677–82.

137   Ginis KAM, Latimer AE, Hicks AL, *et al.* Development and evaluation of an activity measure for

people with spinal cord injury. *Med Sci Sports Exerc* 2005;**37**:1099–111.

138   de Groot S, van der Woude LHV, Niezen A, *et al.* Evaluation of the physical activity scale for

individuals with physical disabilities in people with spinal cord injury. *Spinal Cord* 2010;**48**:542–

7.

139   Healy GN, Wijndaele K, Dunstan DW, *et al.* Objectively measured sedentary time, physical

activity, and metabolic risk: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab).

Diabetes Care 2008;**31**:369–71.

140   O'Donovan G, Hillsdon M, Ukoumunne OC, *et al.* Objectively measured physical activity,

cardiorespiratory fitness and cardiometabolic risk factors in the Health Survey for England. *Prev*

Med 2013;**57**:201–5.

141   Hansford HJ, Wewege MA, Cashin AG, *et al.* If exercise is medicine, why don't we know the

dose? An overview of systematic reviews assessing reporting quality of exercise interventions in

health and disease. *Br J Sports Med* 2022;**56**:692–700.

142   Slade SC, Dionne CE, Underwood M, *et al.* Consensus on Exercise Reporting Template (CERT):

Explanation and Elaboration Statement. *Br J Sports Med* 2016;**50**:1428–37.

143 Hoffmann TC, Glasziou PP, Boutron I, *et al.* Better reporting of interventions: template for

intervention description and replication (TIDieR) checklist and guide. *BMJ* 2014;**348**:g1687.

144 Kodama S, Saito K, Tanaka S, *et al.* Cardiorespiratory fitness as a quantitative predictor of all-

cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*

2009;**301**:2024–35.

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

 FIGURE/TABLE LEGEND

 Figure 1. PRISMA flow diagram. Abbreviation: PRISMA, Preferred Reporting Items for Systematic

 Reviews and Meta-Analyses

 Table 1. Participant demographics and injury characteristics reported within the included studies of the

 primary meta-analysis.

 Table 2. Exercise intervention parameters reported within the included studies of the primary meta-

 analysis.

 Table 3. Summary statistics of the three subgroup analyses on injury characteristics describing Δ in CRF

 outcomes.

 Table 4. Summary statistics of the three subgroup analyses on exercise parameters describing Δ in CRF

 outcomes.

 Table 5. Meta-regression models with adjusted values for each cardiorespiratory fitness outcome.

 Table 6. Grading of recommendations assessment, development and evaluation analysis for each

 cardiorespiratory fitness outcome.

1393

SUPPLEMENTARY MATERIAL

Supplementary file 1 (S1): Systematic Review Search Strategy

Supplementary file 2 (S2): Description of Exercise Intensity Classifications as per the American

College of Sports Medicine (ACSM) guidelines

Supplementary file 3 (S3): Calculated Correlation Factors

Supplementary file 4 (S4): Change in CRF outcomes in response to prospective, well-

characterised exercise interventions lasting >2 weeks (Primary meta-analysis)

1.   Summary of the individual studies included in the review

2.   Forest and funnel plots for change in each CRF outcome for each subgroup comparison (time since

injury, neurological level of injury, injury severity, exercise modality, length of intervention,

relative exercise intensity, method of exercise intensity prescription, frequency of exercise sessions,

and exercise volume)

3.   Quality assessment ratings for each pre-post study included in the primary meta-analysis

4. Risk of bias for each RCT intervention arm included in the primary meta-analysis

5. References

Supplementary file 5 (S5): Cross-sectional cohort comparisons summary (secondary meta-

analysis 1)

1. Overview of participant demographics and injury characteristics for the pooled cohort

comparisons

2. Summary of the individual studies included in the review

3. Quality assessment rating for each study using the NIH tool for observational cohort and cross-

sectional studies

4. Forest plots and funnel plots for each CRF outcome

5. Sensitivity analysis on $\dot{R}VO_{2peak}$

6. References

Supplementary file 6 (S6): Cross-sectional associations between physical activity and CRF

outcomes

1. Overview of participant demographics and injury characteristics for the pooled association

comparisons

2. Summary of the individual studies included in the review

3. Quality assessment rating for each study using the NIH tool for observational cohort and cross-

sectional studies

4. Visualisation of correlation coefficients between physical activity dimensions and CRF outcomes

across included studies

5. References

Supplementary file 7 (S7): Observational studies (secondary meta-analysis 2)

1. Overview of participant demographics and injury characteristics for the pooled observational

studies

2. Summary of the individual studies included in the review

Table 1. Participant demographics and injury characteristics reported within the included studies of the primary meta-analysis.

	A$\dot{V}O_{2peak}$ (L/min)	R$\dot{V}O_{2peak}$ (mL/kg/min)	PPO (W)
Baseline CRF			
Total number of interventions [sum of participants] Mean (range)	69 [766] 1.26 (0.51 – 3.50)	74 [768] 18.0 (7.3 – 36.9)	61 [662] 49 (0 – 168)
Participant demographics			
Age (years)	38 (24 - 54)	39 (24 - 58)	39 (25 - 57)
Sex			
Male	22 [181]	21 [150]	20 [157]
Female	-	-	-
Mixed (% F)	43 [559] (22%)	44 [535] (24%)	39 [492] (28%)
Not reported/cannot determine	4 [26]	9 [83]	2 [13]
Injury characteristics			
Time since injury (years)	8 (0 - 21)	6 (0 - 24)	7 (0 - 21)
Acute (<1-year)	7 [111]	8 [95]	9 [117]
Chronic (>1-year)	47 [472]	48 [443]	38 [367]
Mixed (% acute)	7 [89] (13.5%)	6 [64] (17%)	7 [84] (24%)
Not reported/cannot determine	8 [94]	12 [166]	7 [94]
Neurological level of injury (TETRA/PARA)			
TETRA	2 [18]	3 [23]	3 [23]
PARA	19 [176]	27 [264]	22 [220]
Mixed (% PARA)	41 [488] (59%)	36 [398] (51%)	32 [382] (62%)
Not reported/cannot determine	7 [84]	8 [83]	4 [37]
Severity			
Motor-complete (AIS A-B)	27 [248]	29 [253]	24 [219]
Motor-incomplete (AIS C-D)	8 [102]	13 [142]	2 [14]
Mixed (% motor-incomplete)	22 [303] (32%)	21 [270] (34%)	25 [344] (35%)
Not reported/cannot determine	12 [113]	11 [103]	10 [85]

Total number of studies (N) and participants (Σ), along with descriptive characteristics for the primary meta-analysis included in this systematic review that describes Δ in CRF outcomes in response to prospective, well-characterised exercise interventions lasting >2 weeks (e.g., combining exercise intervention-arms from RCTs and pre-post studies). Continuous variables are displayed as weighted means (range: lowest – highest mean values reported from studies). Categorical variables are displayed as n (%). Weighted means were calculated to account for differences in sample size between studies using the following formula: $\Sigma n*\bar{x}/\Sigma n$, where Σ = the sum of, n = number of participants in each study, and $\bar{x}$ = mean CRF outcome of each study. AIS, American Spinal Injury Association Impairment Scale; F, females; M, males; NR, not reported; PARA, paraplegia; PPO, peak power output; TETRA, tetraplegia; $\dot{V}O_{2peak}$, peak oxygen consumption; W, watts.

Table 2. Exercise intervention parameters reported within the included studies of the primary meta-analysis.

	A$\dot{V}O_{2peak}$ (L/min)	R$\dot{V}O_{2peak}$ (mL/kg/min)	PPO (W)
Baseline CRF			
Total number of interventions [sum of participants] Mean (range)	69 [766] 1.26 (0.51 – 3.50)	74 [768] 18.0 (7.3 – 36.9)	61 [662] 49 (0 – 168)
Exercise intervention parameters			
Modality			
Upper-body aerobic exercise	25 [272]	33 [299]	26 [259]
Upper-body resistance training/circuits	4 [33]	3 [29]	3 [25]
Functional electrical stimulation	17 [170]	8 [66]	14 [140]
Gait/locomotor training	10 [130]	10 [126]	2 [28]
Mixed/multimodal	10 [94]	18 [227]	12 [136]
Behaviour change	3 [67]	2 [21]	4 [74]
Relative intensity			
Light	1 [14]	-	1 [14]
Moderate	8 [58]	12 [94]	10 [73]
Moderate-to-vigorous	21 [270]	24 [305]	16 [183]
Vigorous	14 [119]	20 [194]	11 [104]
Supramaximal	-	1 [4]	1 [10]
Mixed/cannot determine	25 [305]	17 [171]	22 [278]
Relative intensity prescription method			
VO_2 (%peak, %reserve)	8 [61]	12 [112]	9 [93]
Heart rate (%HRR, %HR_{peak}, %HR_{max})	16 [156]	26 [285]	14 [113]
RPE	9 [144]	8 [111]	6 [90]
Workload (%PPO, %MTP, %1RM)	9 [71]	6 [49]	7 [53]
Mixed/cannot determine	27 [334]	22 [211]	25 [313]
Session duration (min)	41 (20 - 90)	41 (15 - 90)	39 (5 - 90)
Frequency (sessions/week)	3 (2 - 7)	3 (2 - 7)	3 (2 - 7)
< 3	19 [230]	13 [168]	13 [156]
≥ 3 and < 5	35 [339]	49 [500]	38 [387]
≥ 5	11 [116]	9 [65]	5 [31]
Not reported	4 [81]	3 [35]	5 [88]
Volume (min/week)	113 (40 - 450)	116 (40 - 330)	107 (15 - 330)
SCI-specific exercise guidelines [fitness (40 – 89 min/wk)]	13 [140] 45 (40 - 84)	14 [156] 47 (40 - 88)	15 [146] 48 (15 - 88)
SCI-specific exercise guidelines [cardiometabolic (90 – 149 min/wk)]	30 [309] 99 (90 - 135)	31 [336] 102 (90 - 135)	26 [290] 113 (90 - 135)
Achieving general population exercise guidelines (≥150 min/wk)	13 [135] 229 (150 - 450)	21 [197] 206 (150 - 330)	13 [117] 212 (171 - 330)
Cannot classify	13 [182]	8 [79]	7 [109]
Length (weeks)	17 (6 - 52)	12 (4 - 52)	16 (4 - 52)
≤ 6 weeks	10 [85]	23 [215]	18 [175]
> 6 and ≤ 12 weeks	33 [368]	36 [371]	21 [223]
> 12 weeks	26 [313]	15 [182]	22 [264]
Adverse events reported			
Bone, joint or muscular pain	5 [5] [a]	6 [9] [a]	4 [4] [a]

Autonomic or cardiovascular function	3 [5]	2 [1]	4 [3]
Skin irritation or pressure sores	2 [2] [a]	5 [18] [a]	2 [2] [a]
Other [d]	2 [NR] [a,c]	4 [3] [a]	3 [1] [a,c]

Total number of studies (N) and participants, (Σ) along with descriptive characteristics for the primary meta-analysis included in this systematic review that describes Δ in CRF outcomes in response to prospective, well-characterised exercise interventions lasting >2 weeks (e.g., combining exercise intervention-arms from RCTs and pre-post studies). Continuous variables are displayed as weighted means (range: lowest – highest mean values reported from studies). Categorical variables are displayed as n (%). Weighted means were calculated to account for differences in sample size between studies using the following formula: $\Sigma n*\bar{x}/\Sigma n$, where Σ = the sum of, n = number of participants in each study, and $\bar{x}$ = mean CRF outcome of each study. F, females; HR $_{max}$, maximal heart rate; HR $_{peak}$, peak heart rate; HRR, heart rate reserve; 1RM, one repetition maximum; M, males; MTP, maximal tolerated power; NR, not reported; PPO, peak power output; $\dot{V}O_{2\ peak}$, peak oxygen consumption; W, watts. [a] Beillot et al. [68] (pre-post intervention study) reported n=10 suffered major complications including spontaneous fractures of lower limbs, occurrence of syringomyelia and pressure sores but did not specify the sum of participants for each adverse event. [b] Gibbons et al. [81] reported that some individuals experienced autonomic dysreflexia during the FES response test but did not quantify further. [c] Sum of participants experiencing adverse events were not reported by Janssen and Pringle [61]. [d] Other adverse events included: anxiety, nausea, dizziness and issues with testing equipment.

Table 3. Summary statistics of the three subgroup analyses on injury characteristics describing Δ in CRF outcomes.

	$A\dot{V}O_{2peak}$ (L/min)		$R\dot{V}O_{2peak}$ (mL/kg/min)		PPO (W)	
	N [Σ] (%)	WMD (95% CIs) *p*-values	N [Σ] (%)	WMD (95% CIs) *p*-values	N [Σ] (%)	WMD (95% CIs) *p*-values
Main effect	69 [696]	0.22 [0.17, 0.26] ***p* < 0.001**	74 [716]	2.8 [2.2, 3.4] ***p* < 0.001**	61 [602]	11 [8, 13] ***p* < 0.001**
Heterogeneity (I²)	74% (*p* < 0.001)		52% (*p* < 0.001)		78% (*p* < 0.001)	
Time since injury						
Acute (<1-year)	7 [86] (10.4%)	0.23 [0.11, 0.35] ***p* < 0.001**	8 [70] (10.9%)	3.4 [1.5, 6.1] ***p* = 0.002**	9 [95] (13.6%)	16 [11, 22] ***p* < 0.001**
Chronic (≥1-year)	47 [461] (62.6%)	0.20 [0.14, 0.27] ***p* < 0.001**	48 [431] (61.8%)	2.7 [1.9, 3.5] ***p* < 0.003**	38 [343] (61.8%)	9 [6, 12] ***p* < 0.001**
Mixed	7 [79] (14%)	0.25 [0.10, 0.39] ***p* < 0.001**	6 [54] (5.9%)	1.9 [0.1, 3.7] ***p* = 0.03**	7 [75] (12.8%)	6 [5, 7] ***p* < 0.001**
Not reported/cannot determine	8 [70] (13%)	0.25 [0.11, 0.38] ***p* < 0.001**	12 [161] (21.4%)	2.6 [2.0, 3.3] ***p* < 0.003**	7 [89] (11.8%)	16 [9, 23] ***p* < 0.001**
Subgroup differences	-	*p* = 0.87	-	*p* = 0.64	-	***p* < 0.001**
Neurological level of injury						
Tetraplegia	2 [18] (5.1%)	0.45 [-0.28, 1.19] *p* = 0.23	3 [23] (8.5%)	5.9 [0.2, 11.7] ***p* = 0.04**	3 [23] (6.8%)	9 [6, 13] ***p* < 0.002**
Paraplegia	20 [174] (28.4%)	0.24 [0.17, 0.32] ***p* < 0.002**	28 [262] (45.2%)	2.8 [2.2, 3.4] ***p* < 0.003**	22 [216] (42.2%)	16 [12, 19] ***p* < 0.002**
Mixed	44 [470] (58.9%)	0.20 [0.15, 0.25] ***p* < 0.002**	41 [418] (42.5%)	2.2 [1.5, 2.8] ***p* < 0.003**	34 [350] (48.6%)	6 [4, 8] ***p* < 0.002**
Not reported/cannot determine	3 [34] (7.6%)	0.19 [0.11, 0.27] ***p* < 0.002**	2 [13] (3.8%)	2.8 [0.7, 4.8] ***p* = 0.02**	2 [13] (2.4%)	17 [7, 27] ***p* = 0.001**
Subgroup differences	-	*p* = 0.65	-	*p* = 0.34	-	***p* < 0.001**
Injury severity						
Motor-complete (AIS A-B)	27 [235] (40%)	0.21 [0.14, 0.27] ***p* < 0.002**	29 [241] (47.4%)	2.7 [2.0, 3.4] ***p* < 0.002**	24 [210] (49.2%)	11 [8, 15] ***p* < 0.002**
Motor-incomplete (AIS C-D)	8 [103] (9%)	0.10 [-0.01, 0.21] *p* = 0.08	13 [139] (12.5%)	1.6 [0.2, 2.9] ***p* = 0.02**	2 [14] (3.1%)	4 [-3, 12] *p* = 0.25
Mixed (AIS A-D)	22 [247] (26.2%)	0.18 [0.13, 0.24] ***p* < 0.002**	21 [244] (23.3%)	2.7 [1.7, 3.6] ***p* < 0.002**	25 [296] (31.1%)	10 [6, 14] ***p* < 0.002**
Not reported/cannot determine	12 [111] (24.8%)	0.32 [0.20, 0.44] ***p* < 0.002**	11 [92] (16.8%)	3.9 [1.7, 6.1] ***p* < 0.002**	10 [82] (16.6%)	11 [5, 17] ***p* < 0.002**
Subgroup differences	-	*p* = 0.06	-	*p* = 0.28	-	*p* = 0.43

Total number of interventions (N), sum of participants analysed at post-intervention (Σ), weighting of subgroups (%). Thresholds for statistically significant subgroup differences were adjusted for the number of subgroup comparisons and are highlighted in bold: time since injury (*p*<0.0125), neurological level of injury (*p*<0.0125) and injury severity (*p*<0.0125). Individual subgroup p-values were adjusted for multiple comparisons via the Bonferroni correction method. AIS, American Spinal Injury Association Impairment Scale; $A\dot{V}O_{2peak}$, absolute peak oxygen consumption; CIs, confidence intervals; PPO, peak power output; $R\dot{V}O_{2peak}$, relative peak oxygen consumption; WMD, weighted mean difference.

1498

Table 4. Summary statistics of the six subgroup analyses on exercise parameters describing Δ in CRF outcomes.

	A$\dot{V}O_{2peak}$ (L/min)		R$\dot{V}O_{2peak}$ (mL/kg/min)		PPO (W)	
	N [Σ] (%)	WMD (95% CIs) *p*-values	N [Σ] (%)	WMD (95% CIs) *p-values*	N [Σ] (%)	WMD (95% CIs) *p-values*
Main effect	69 [696]	0.22 [0.17, 0.26] **_p < 0.001_**	74 [716]	2.8 [2.2, 3.4] **_p < 0.001_**	61 [602]	11 [8, 13] **_p < 0.001_**
Heterogeneity (I²)	74% (*p* < 0.001)		52% (*p* < 0.001)		78% (*p* < 0.001)	
Exercise modality						
Aerobic, volitional upper-body	25 [235] (33.5%)	0.25 [0.16, 0.34] **_p < 0.004_**	33 [264] (43.1%)	3.4 [2.4, 4.4] **_p < 0.004_**	26 [223] (32.5%)	15 [11, 19] **_p < 0.003_**
Resistance training	4 [31] (3.5%)	0.33 [0.13, 0.52] **_p = 0.003_**	3 [27] (3.9%)	5.0 [2.9, 7.0] **_p < 0.004_**	3 [25] (4.6%)	20 [12, 28] **_p < 0.003_**
Functional electrical stimulation	17 [168] (33.9%)	0.22 [0.15, 0.29] **_p < 0.004_**	8 [66] (14.4%)	2.4 [0.9, 3.9] **_p = 0.006_**	14 [138] (37.1%)	6 [3, 10] **_p < 0.003_**
Gait training	10 [127] (12%)	0.07 [-0.02, 0.17] *p = 0.14*	10 [120] (9.7%)	1.0 [-0.5, 2.6] *p = 0.40*	2 [24] (2.2%)	4 [-9, 18] *p = 0.54*
Behaviour change	3 [49] (2.5%)	0.22 [0.00, 0.44] *p = 0.10*	2 [21] (3.3%)	1.1 [-1.2, 3.5] *p = 0.35*	4 [56] (4.6%)	12 [-1, 24] *p = 0.12*
Mixed	10 [86] (14.6%)	0.21 [0.14, 0.27] **_p < 0.004_**	18 [218] (25.6%)	2.4 [1.7, 3.2] **_p < 0.004_**	12 [136] (19%)	10 [5, 16] **_p < 0.003_**
Subgroup differences	-	*p = 0.07*	-	*p = 0.02*	-	**_p = 0.003_**
Length of intervention						
≤6 weeks	10 [79] (14.5%)	0.26 [0.19, 0.39] **_p < 0.001_**	23 [206] (23.1%)	2.9 [1.9, 3.9] **_p < 0.001_**	17 [159] (26.7%)	10 [6, 14] **_p < 0.001_**
>6 – ≤12 weeks	32 [327] (46.6%)	0.21 [0.14, 0.29] **_p < 0.001_**	36 [337] (54.6%)	3.2 [2.3, 4.1] **_p < 0.001_**	22 [202] (34.5%)	13 [8, 17] **_p < 0.001_**
>12 weeks	27 [290] (38.9%)	0.22 [0.15, 0.28] **_p < 0.001_**	15 [173] (22.3%)	1.8 [1.0, 2.6] **_p < 0.001_**	22 [241] (38.8%)	9 [5, 13] **_p < 0.001_**
Subgroup differences	-	*p = 0.59*	-	*p = 0.05*	-	*p = 0.49*
Relative exercise intensity						
Light	1 [10] (0.6%)	-0.05 [-0.57, 0.47] *p = 0.85*	-	-	1 [10] (1%)	-1 [-22, 20] *p = 0.92*
Moderate	8 [58] (9.5%)	0.32 [0.09, 0.54] **_p = 0.01_**	12 [92] (18.1%)	3.2 [1.1, 5.3] **_p = 0.006_**	10 [71] (15.9%)	13 [4, 21] **_p = 0.009_**
Moderate-to-vigorous	21 [247] (33.7%)	0.21 [0.14, 0.27] **_p < 0.003_**	24 [279] (32.5%)	2.7 [2.0, 3.5] **_p < 0.003_**	16 [161] (24.1%)	17 [13, 21] **_p < 0.004_**
Vigorous	14 [109] (15.2%)	0.19 [0.14, 0.25] **_p < 0.003_**	20 [183] (21.6%)	2.2 [1.4, 3.0] **_p < 0.003_**	11 [96] (12%)	10 [7, 16] **_p < 0.004_**
Supramaximal	-	-	1 [4] (0.4%)	1.1 [-8.2, 10.4] *p = 0.82*	1 [10] (0.6%)	17 [-12, 46] *p = 0.50*
Mixed/cannot determine	25 [272] (41%)	0.21 [0.14, 0.29] **_p < 0.003_**	17 [158] (27.4%)	2.6 [1.5, 3.8] **_p < 0.003_**	22 [254] (46.4%)	8 [5, 10] **_p < 0.004_**
Subgroup differences	-	*p = 0.71*	-	*p = 0.67*	-	**_p = 0.003_**

Exercise intensity prescription					
Oxygen consumption	8 [57] (13.1%)	0.19 [0.07, 0.32] *p* = 0.003	12 [107] (17.9%)	2.3 [1.5, 3.2] *p* < 0.003	9 [89] (12.6%)	20 [15, 25] *p* < 0.003
Heart rate	16 [156] (20.2%)	0.28 [0.15, 0.40] *p* < 0.002	26 [284] (37.7%)	3.1 [2.0, 4.3] *p* < 0.003	14 [113] (21.8%)	14 [8, 19] *p* < 0.003
Rating of perceived exertion	9 [121] (12.1%)	0.18 [0.09, 0.26] *p* < 0.002	8 [84] (8.3%)	3.5 [1.2, 5.07] *p* = 0.002	6 [66] (5.7%)	9 [1, 17] *p* = 0.03
Workload	9 [65] (9.9%)	0.23 [0.13, 0.33] *p* < 0.002	6 [43] (6.5%)	3.0 [1.2, 4.8] *p* = 0.002	7 [49] (11.9%)	11 [4, 19] *p* = 0.01
Mixed/cannot determine	27 [297] (44.7%)	0.21 [0.14, 0.27] *p* < 0.002	22 [198] (29.6%)	2.4 [1.4, 3.3] *p* < 0.003	25 [285] (48%)	7 [5, 10] *p* < 0.003
Subgroup differences	-	*p* = 0.72	-	*p* = 0.71	-	*p* < 0.001
Frequency of exercise sessions					
<3 sessions/wk	19 [213] (24%)	0.18 [0.12, 0.25] *p* < 0.002	13 [155] (15.2%)	2.8 [1.5, 4.0] *p* < 0.002	13 [148] (21.4%)	4 [1, 6] *p* = 0.003
≥3 – <5 sessions/wk	35 [306] (57.6%)	0.26 [0.19, 0.33] *p* < 0.002	49 [465] (65.9%)	2.8 [2.1, 3.6] *p* < 0.002	38 [357] (61.8%)	13 [10, 15] *p* < 0.004
≥5 sessions/wk	11 [118] (14.4%)	0.15 [0.07, 0.23] *p* < 0.002	9 [65] (14.2%)	3.6 [1.7, 5.4] *p* < 0.002	5 [31] (10.9%)	10 [-2, 23] *p* = 0.10
Not reported/cannot determine	4 [59] (4%)	0.13 [-0.05, 0.31] *p* = 0.15	3 [31] (4.7%)	0.8 [-1.2, 2.8] *p* = 0.42	5 [66] (5.9%)	9 [0, 18] *p* = 0.10
Subgroup differences	-	*p* = 0.14	-	*p* = 0.24	-	*p* < 0.001
Exercise volume					
SCI-specific exercise guidelines for fitness (40 - 89 min/wk)	13 [132] (13.1%)	0.23 [0.13, 0.33] *p* < 0.001	14 [151] (13.9%)	3.2 [2.0, 4.5] *p* < 0.002	15 [138] (19.1%)	6 [2, 10] *p* = 0.002
SCI-specific exercise guidelines for cardiometabolic health (90 - 149 min/wk)	30 [269] (52.3%)	0.23 [0.16, 0.30] *p* < 0.001	31 [295] (48.3%)	2.8 [1.9, 3.8] *p* < 0.002	26 [260] (48.7%)	12 [9, 16] *p* < 0.002
Achieving general population exercise guidelines (≥150 min/wk)	13 [133] (18%)	0.18 [0.11, 0.25] *p* < 0.001	21 [195] (28.1%)	2.8 [1.8, 3.9] *p* < 0.002	13 [117] (23.9%)	11 [5, 17] *p* < 0.002
Not reported/cannot determine	13 [162] (16.6%)	0.24 [0.12, 0.36] *p* < 0.001	8 [75] (9.7%)	2.1 [0.7, 3.5] *p* = 0.003	7 [87] (8.3%)	10 [4, 17] *p* = 0.002
Subgroup differences	-	*p* = 0.67	-	*p* = 0.70	-	*p* = 0.17

Total number of interventions (N), sum of participants analysed at post-intervention (Σ), weighting of subgroups (%). Thresholds for statistically significant subgroup differences were adjusted for the number of subgroup comparisons and are highlighted in bold: exercise modality ($p<0.008$), length of intervention ($p<0.017$), relative exercise intensity [$A\dot{V}O_{2peak}$ and $R\dot{V}O_{2peak}$ ($p<0.01$), PPO ($p<0.008$)], exercise intensity prescription ($p<0.01$), frequency of exercise sessions ($p<0.025$), and exercise volume ($p<0.025$). Individual subgroup *p*-values were adjusted for multiple comparisons via the Bonferroni correction method. $A\dot{V}O_{2peak}$, absolute peak oxygen consumption; CIs, confidence intervals; PPO, peak power output; $R\dot{V}O_{2peak}$, relative peak oxygen consumption; WMD, weighted mean difference.

1499

1500

Table 5: Meta-regression models with adjusted values for each cardiorespiratory fitness outcome.

Covariate	Coef.	Std.Err.	t	Unadjusted P>t	95% CI	Adjusted P>t
AV̇O$_{2peak}$ Model 1 (N = 69)[1] (# covariates = 5)						
Male	0.006	0.004	1.610	0.084	-0.001 to 0.013	0.322
Mean age	-0.003	0.001	-2.580	0.010	-0.006 to -0.001	0.045
TSI	0.000	0.001	0.260	0.714	-0.002 to 0.003	0.999
Neurological level of injury	-0.002	0.004	-0.470	0.584	-0.010 to 0.006	0.984
Severity	-0.005	0.003	-1.340	0.153	-0.012 to 0.002	0.534
AV̇O$_{2peak}$ Model 2 (N = 69)[2] (# covariates = 6)						
Exercise modality	-0.015	0.016	-0.940	0.281	-0.048 to 0.017	0.865
Exercise intensity	-0.006	0.016	-0.390	0.666	-0.038 to 0.026	0.999
Length of intervention	0.000	0.002	0.050	0.828	-0.004 to 0.004	1.000
Duration (mins)	-0.005	0.006	-0.790	0.327	-0.018 to 0.008	0.917
Frequency	0.002	0.011	0.180	0.902	-0.019 to 0.023	1.000
Volume	-0.000	0.003	-0.040	0.929	-0.006 to 0.006	1.000
RV̇O$_{2peak}$ Model 1 (N = 74)[3] (# covariates = 5)						
Male	0.084	0.047	1.790	0.84	-0.010 to 0.177	0.395
Mean age	-0.041	0.013	-3.010	0.004	-0.068 to -0.014	0.025
TSI	-0.012	0.013	-0.940	0.386	-0.039 to 0.014	0.932
Neurological level of injury	-0.043	0.042	-1.020	0.356	-0.128 to 0.042	0.907
Severity	-0.030	0.047	-0.650	0.521	-0.124 to 0.063	0.983
RV̇O$_{2peak}$ Model 2 (N = 74)[4] (# covariates = 6)						
Exercise modality	-0.276	0.164	-1.680	0.110	-0.603 to 0.051	0.511
Exercise intensity	-0.076	0.200	-0.380	0.718	-0.474 to 0.323	1.00
Length of intervention	-0.024	0.033	-0.740	0.437	-0.089 to 0.041	0.982
Risk of bias	-0.057	0.350	-0.160	0.866	-0.756 to 0.642	1.000
Duration (mins)	-0.043	0.063	-0.690	0.494	-0.169 to 0.082	0.986
Frequency	0.143	0.139	1.030	0.308	-0.133 to 0.420	0.908
PPO Model 1 (N = 61)[5] (# covariates = 6)						
Male	-0.090	0.236	-0.380	0.712	-0.562 to 0.383	0.997
Mean age	-0.035	0.082	-0.420	0.694	-0.199 to 0.130	0.995
TSI	-0.149	0.077	-1.940	0.075	-0.303 to 0.005	0.296
Neurological level of injury	-0.122	0.193	-0.630	0.556	-0.509 to 0.265	0.972
Severity	-0.051	0.182	-0.280	0.796	-0.416 to 0.314	1.000
PPO Model 2 (N = 61)[6] (# covariates = 5)						
Exercise modality	-0.685	0.733	-0.930	0.068	-2.156 to 0.786	0.266
Exercise intensity prescription	-1.465	0.749	-1.960	0.001	-2.967 to 0.036	0.002
Duration (mins)	-0.170	0.361	-0.470	0.476	-0.893 to 0.554	0.945
Frequency	0.254	0.643	0.390	0.997	-1.036 to 1.543	1.00
Volume	-0.217	0.178	-1.220	0.009	-0.574 to 0.140	0.041

* Permutations = 10,000

[1] tau^2 = 0.02339; I^2 res = 98.61%; Adj R^2 = 13.00%; Model F (5,63) = 2.77; Prob > F = 0.0252

[2] tau^2 = 0.2932; I^2 res = 98.52%; Adj R^2 = -9.04%; Model F (7,61) = 0.31; Prob > F = 0.9446

[3] tau^2 = 3.639; I^2 res = 97.98%; Adj R^2 = 11.98%; Model F (5,68) = 2.89; Prob > F = 0.0201

Adj R^2, proportion of between-study variance explained; A$\dot{V}O_{2peak}$, absolute peak oxygen consumption; Coef, coefficient of variation; I^2 res, I^2 residual variation due to heterogeneity; Model F, joint test for all covariates; PP Prob > F, with Knapp-Hartung modification; R$\dot{V}O_{2peak}$, relative peak oxygen consumption; Std.Err, standard error; TSI, time since injury.

1501

Table 6. Grading of recommendations assessment, development and evaluation analysis for each cardiorespiratory fitness outcome.

	A$\dot{V}O_{2peak}$ (L/min)	R$\dot{V}O_{2peak}$ (mL/kg/min)	PPO (W)
Summary of findings according to GRADE analysis			
GRADE	LOW	LOW	LOW
Comments	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect. Study design, imprecision, an unclear dose response and residual confounding reduced the Grade to Low. The evidence supporting improvements in A$\dot{V}O_{2peak}$ is predominantly in young and middle-aged males that had been injured for >1-year (chronic TSI). Participants were mostly paraplegic (70%) but there were a mixture of injury severities (AIS A-D). There were no subgroup differences in exercise intervention characteristics to suggest the optimal training parameters.	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect. High risk of bias, imprecision, an unclear dose response and residual confounding reduced the Grade to Low. The evidence supporting improvements in R$\dot{V}O_{2peak}$ is predominantly in young and middle-aged males that had been injured for >1-year (chronic TSI). Participants were mostly paraplegic (70.5%) but there were a mixture of injury severities (AIS A-D). There were no subgroup differences in exercise intervention characteristics to suggest the optimal training parameters.	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect. Inconsistency, imprecision, an unclear dose response and residual confounding reduced the Grade to Moderate. The evidence supporting improvements in PPO is predominantly in young and middle-aged males that had been injured for >1-year (chronic TSI). Participants were mostly paraplegic (76%) but there were a mixture of injury severities (AIS A-D). Subgroup differences suggest that upper-body aerobic exercise and resistance training appear the most effective at improving PPO. Furthermore, acutely-injured, individuals with paraplegia, exercising for >3 sessions/week at a moderate-to-vigorous-intensity, prescribed via $\dot{V}O_2$ or heart rate, will likely experience the greatest change in PPO.
Lower quality criteria			
Study design	Mixture of RCTs and pre-post studies with no control groups.	Mixture of RCTs and pre-post studies with no control groups.	Mixture of RCTs and pre-post studies with no control groups.

	Overall WMDs for RCT interventions relative to controls and pre-post interventions only: RCTs (0.15 L/min) and pre-post studies (0.23 L/min). **DOWNGRADE**	Overall WMDs for RCT interventions relative to controls and pre-post interventions only: RCTs (2.9 mL/kg/min) and pre-post studies (2.9 mL/kg/min). **NO DOWNGRADE**	Overall WMDs for RCT interventions relative to controls and pre-post interventions only: RCTs (10 W) and pre-post studies (11 W). **NO DOWNGRADE**
Risk of bias (RoB)	28% of pre-post studies were rated as good, 56% as fair, and 16% as poor. 31% of RCTs had low RoB, 23% had some concerns, and 46% had high RoB. **NO DOWNGRADE**	38% of pre-post studies were rated as good, 48% as fair, and 14% as poor. 15% of RCTs had low RoB, 25% had some concerns, and <u>60% had high RoB</u>. **DOWNGRADE**	26% of pre-post studies were rated as good, 59% as fair, and 15% as poor. 23% of RCTs had low RoB, 38.5% had some concerns, and 38.5% had high RoB. **NO DOWNGRADE**
Inconsistency of results	Effect estimates were consistent, with 91% of the included exercise interventions favouring an increase in $A\dot{V}O_{2peak}$, but most had a low effect estimate. $I^2 = 74\%$ **NO DOWNGRADE**	Effect estimates were consistent, with 91% of the included exercise interventions favouring an increase in $R\dot{V}O_{2peak}$, and most had a large effect estimate. $I^2 = 52\%$ **NO DOWNGRADE**	Effect estimates were consistent, with 93% of the included exercise interventions favouring an increase in PPO, and most had a large effect estimate. $I^2 = 78\%$ **DOWNGRADE**
Indirectness	Most studies (83%) included $A\dot{V}O_{2peak}$ in their main outcome measures, across a range of participant characteristics. **NO DOWNGRADE**	Most studies (72%) included $R\dot{V}O_{2peak}$ in their main outcome measures, across a range of participant characteristics. **NO DOWNGRADE**	Most studies (82%) included PPO in their main outcome measures, across a range of participant characteristics. **NO DOWNGRADE**
Imprecision	Large sample size (N=696), however, 62% of the included exercise interventions had 95% CI overlap 0. **DOWNGRADE**	Large sample size (N=716), however, 76% of the included exercise interventions had 95% CI overlap 0. **DOWNGRADE**	Large sample size (N=601), however, 67% of the included exercise interventions had 95% CI overlap 0. **DOWNGRADE**
Publication bias	An exhaustive approach was used during the search strategy (i.e., scientific databases and grey literature search). Egger's test: Z = -1.23 (p = 0.22). Visual inspection of the funnel plots, data extraction sheets and Tables 3-4 revealed no noticeable publication bias. **NO DOWNGRADE**	An exhaustive approach was used during the search strategy (i.e., scientific databases and grey literature search). Egger's test: Z = -0.54 (p = 0.59). Visual inspection of the funnel plots, data extraction sheets and Tables 3-4 revealed no noticeable publication bias. **NO DOWNGRADE**	An exhaustive approach was used during the search strategy (i.e., scientific databases and grey literature search). Egger's test: Z = 0.73 (p = 0.46). Visual inspection of the funnel plots, data extraction sheets and Tables 3-4 revealed no noticeable publication bias. **NO DOWNGRADE**
Higher quality criteria			
Large effect	Yes Z = 9.5 ($p < 0.001$) **NO UPGRADE**	Yes Z = 9.4 ($p < 0.001$) **NO UPGRADE**	Yes Z = 8.7 ($p < 0.001$) **NO UPGRADE**

Dose response	No clear dose response. **NO UPGRADE**	No clear dose response. **NO UPGRADE**	No clear dose response. **NO UPGRADE**
Residual confounding	Mixture of exercise modalities, levels of injury, etc. **NO UPGRADE**	Mixture of exercise modalities, levels of injury, etc. **NO UPGRADE**	Mixture of exercise modalities, levels of injury, etc. **NO UPGRADE**

GRADE certainty in the evidence can be 'High', 'Moderate', 'Low' or 'Very Low' according to published guidelines [34]. Risk of bias was downgraded where >50% of RCTs had a high risk of bias. Heterogeneity was also included as a measure of inconsistency, whereby an outcome with I^2 >75% was classed as considerable and resulted in a downgrade. Imprecision was downgraded where >50% of studies had confidence intervals overlap the no effect line. Indirectness would have been downgraded where <50% of studies did not include the appropriate main outcome measure or assess a range of participant characteristics. Overall effect sizes are presented as Z-scores. Statistical significance accepted as $p < 0.05$. AIS, American Spinal Injury Association Impairment Scale; $A\dot{V}O_{2peak}$, absolute peak oxygen consumption; CI, confidence intervals; CRF, cardiorespiratory fitness; PPO, peak power output; RCTs, randomised-controlled trials; RoB, risk of bias; $R\dot{V}O_{2peak}$, relative peak oxygen consumption; $\dot{V}O_2$, peak oxygen consumption.