SARS-CoV-2 genomic diversity in households highlights the challenges of sequence-based transmission inference

Authors: Emily Bendall[1], Gabriela Paz-Bailey[2], Gilberto A. Santiago[2], Christina A. Porucznik[3], Joseph B. Stanford[3], Melissa S. Stockwell[4], Jazmin Duque[5], Zuha Jeddy[5], Vic Veguilla[2], Chelsea Major[2], Vanessa Rivera-Amill, PhD[6], Melissa A. Rolfes[2], Fatimah S. Dawood[2], Adam S. Lauring[1,7] *

Affiliations: [1]Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA; [2]Centers for Disease Control and Prevention; [3]Division of Public Health, Department of Family and Preventive Medicine, University of Utah School of Medicine; [4]Division of Child and Adolescent Health, Department of Pediatrics, Columbia University Vagelos College of Physicians and Surgeons, and Department of Population and Family Health, Mailman School of Public Health, Columbia University Irving Medical Center, New York, NY; [5]Abt Associates; [6]Ponce Health Sciences University/Ponce Research Institute; [7]Division of Infectious Diseases, Department of Internal Medicine, University of Michigan, Ann Arbor, MI, USA;

* Corresponding Author: Adam S. Lauring, MS2 4742C, SPC 1621, 1137 Catherine Street, Ann Arbor, MI 48109; (734) 764-7731; alauring@med.umich.edu.

Disclaimer: The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Keywords: SARS-CoV-2; genomic epidemiology; transmission; household
Running Title: SARS-CoV-2 diversity in households
Abstract Word Count: 200
Main Text Word Count: 3009
Summary: High depth of coverage whole genome sequencing can identify SARS-CoV-2 transmission chains in settings where there is strong epidemiologic linkage but is not reliable as a stand-alone method for transmission inference.

ABSTRACT

Background: The reliability of sequence-based inference of SARS-CoV-2 transmission is not

clear. Sequence data from infections among household members can define the expected

genomic diversity of a virus along a defined transmission chain.

Methods: SARS-CoV-2 cases were identified prospectively among 2,369 participants in 706

households. Specimens with an RT-PCR cycle threshold ≤30 underwent whole genome

sequencing. Intrahost single nucleotide variants (iSNV) were identified at ≥5% frequency.

Phylogenetic trees were used to evaluate the relationship of household and community

sequences.

Results: There were 178 SARS-CoV-2 cases in 706 households. Among 147 specimens

sequenced, 106 yielded a whole genome consensus with coverage suitable for identifying iSNV.

Twenty-six households had sequences from multiple cases within 14 days. Consensus

sequences were indistinguishable among cases in 15 households, while 11 had $\geq$1 consensus

that differed by 1-2 mutations. Sequences from households and the community were often

interspersed on phylogenetic trees. Identification of iSNV improved inference in 2 of 15

households with indistinguishable consensus sequences and 6 of 11 with distinct ones.

Conclusions: In multiple infection households, whole genome consensus sequences differed by

0-1 mutations. Identification of shared iSNV occasionally resolved linkage, but the low genomic

diversity of SARS-CoV-2 limits the utility of "sequence-only" transmission inference.

INTRODUCTION

INTRODUCTION

RNA viruses evolve rapidly and accumulate mutations as outbreaks grow [1]. As a result, the

evolutionary relationships among sequenced cases hold important information about the

processes that drive epidemics [2]. For example, sequence data can help define transmission

chains and outbreaks [3–5], the timing and location of viral introductions into communities [6–

8], and larger patterns of spread [9–12]. Over the course of the COVID-19 pandemic, SARS-CoV-

2 sequences have been used to infer transmission linkage in hospitals and other congregate

settings [13–18]. Inferring these linkages with high confidence is necessary for subsequent

studies of the biology of transmission and effectiveness of mitigation strategies.

To infer transmission, one can ask whether the sequences within a group of close contacts,

such as a household, are more similar than sequences in the broader community. This approach

depends on both the granularity of the sequence data and the amount of genomic diversity in

the underlying community or meta-population. The relatedness of viral sequences identified

from potential transmission chains versus community virologic surveillance is compared using

phylogenetic trees of whole genome consensus sequences or clustering of transmission-

associated sequences [2]. In the setting of insufficient community sampling and/or low genomic

diversity, consensus trees can miss true linkages and identify false ones. Greater coverage

sequencing can improve resolution by identifying intrahost single nucleotide variants (iSNV) in

host-derived viral populations that have yet to achieve consensus levels, or >50% within-host

frequency, along a transmission chain [19,20]. While these approaches have proven useful for

influenza and other viruses, the reliability of sequence-based inference of SARS-CoV-2

transmission is less clear. For example, we and others have found that participants without

known epidemiologic linkage can share indistinguishable consensus sequences and even

minority (<50%) iSNV [21–23].

Households are ideal settings for studies of the biology and epidemiology of viral transmission.

Documentation of close contact and concurrent symptoms or test positivity provide strong

epidemiologic evidence of within-household transmission. Sequence data from infected

participants can therefore define the expected genomic diversity of a virus along a transmission

chain and inform sequence-based studies in other transmission settings, where epidemiologic

linkage may be uncertain. Here, we use whole genome sequencing of SARS-CoV-2 populations

from participants in two prospective household studies of COVID-19 that were conducted at

three sites. To assess the utility of SARS-CoV-2 sequence data as a tool for inferring

transmission, we used phylogenetic analysis of sequences from households with at least two

SARS-CoV-2 infection cases to assess the clustering of within-household sequences relative to

contemporaneous community sequences. We used iSNV to further resolve transmission

linkages in selected households.

METHODS

Cohorts

The Coronavirus Household Evaluation and Respiratory Testing (C-HEaRT) study enrolled

households in Utah (Salt Lake, Weber, Davis, Box Elder, Cache, Tooele, Wasatch, Summit, Utah,

and Iron Counties) and New York City [24] during August 2020 through February 2021 and

followed them with surveillance for SARS-CoV-2 infection during September 2020 through

August 2021. The Communities Organized for the Prevention of Arboviruses (COPA) was

expanded to include investigation of the epidemiology of COVID-19, creating the COCOVID

study, and recruited households in Ponce, Puerto Rico. For C-HEaRT, household eligibility

criteria included: ≥1 child aged 0-17 years, ≥75% of household members met individual level

eligibility (all members if a 2- or 3-person household), one adult member was willing to

complete monthly questionnaires, and adult members could communicate in English or

Spanish. Individual eligibility criteria included: anticipated residence in the household for ≥3

consecutive months, and willingness to complete study surveys, weekly symptom assessments,

and self-collect respiratory specimens. For COCOVID, household members were eligible if they

were aged ≥1 year, slept in the house ≥4 nights per week, had no definite plans to move in the

next year, and were willing and able to comply with study requirements. For both studies,

written informed consent (paper or electronic) was obtained from adults (aged >18 years in C-

HEaRT and >20 years in COCOVID). Parents or legal guardians of minor children provided

written informed consent on behalf of their children; older children (aged 12–17 years in C-

HEaRT and 7–20 years in COCOVID) also provided assent to study participation. The C-HEaRT

study protocol was reviewed and approved by the University of Utah Institutional Review Board

(IRB) as the single IRB for all collaborators. The COCOVID study protocol was reviewed and

approved by the Ponce Medical School Foundation, Inc. IRB.

Sample Collection and Testing

Participants were asked to self-collect (or parent/guardian-collect for children) mid-turbinate

nasal swabs every week, regardless of illness symptoms, and place the swabs in viral transport

media. Participants were also contacted by text message or email every week to ascertain if

they had COVID-19-like (CLI) or any other illness symptoms; they were asked to self-collect an

additional mid-turbinate flocked nasal swab once with onset of CLI symptoms. CLI was defined

as 1 or more of the following: fever or feverishness, cough, shortness of breath, sore throat,

diarrhea, muscle aches, chills, or change in taste or smell. Respiratory specimens were shipped

overnight to a central lab and tested using either the Quidel Lyra SARS-CoV-2 Assay or the

ThermoFisher Combo Kit platform. The assays were approved under Emergency Use

Authorization for the diagnosis of SARS-CoV-2 infection prior to use in this study. Test-positive

infections in the same household that were first detected by RT-PCR within 14 days of each

other (including those detected on the same date) were considered epidemiologically linked

and likely to have resulted from within-household transmission.

SARS-CoV-2 Genomic Sequencing

SARS-CoV-2 genomic sequencing was attempted on all specimens with an RT-PCR cycle

threshold (Ct) ≤30 on either the nucleocapsid protein 1 or 2 target. SARS-CoV-2 genomes were

sequenced as described previously [10]. Briefly, RNA was extracted from midturbinate nasal

swab specimens with the MagMax MVPII viral nucleic acid isolation kit on a Kingfisher Flex

(Thermofisher) and reverse transcribed with Lunascript (NEB). We amplified SARS-CoV-2 cDNA

in two pools using the ARTIC Network v3 primers and protocol. Amplicon pools were combined

in equal volumes for a given sample and purified with magnetic beads. Barcoded sequencing

libraries were prepared using the NEBNext ARTIC SARS-CoV-2 Library Prep Kit with magnetic

bead size selection. Individual barcoded sample libraries were pooled (up to 96) and sequenced

on an Illumina MiSeq (v2 chemistry, 2x250 cycles).

Reads were mapped to the Wuhan/Hu-1/2019 reference genome (GenBank MN908947.3) with

BWA-MEM [25]. We used iVar 1.2.1 [26] to trim ARTIC amplification primer sequences and to

determine consensus sequences using bases with >50% frequency and placing a designated

unknown base N at positions covered by fewer than 10 reads. Genomes with 29,000 or more

unambiguous bases (> 97% completeness) were used in downstream analysis. We identified

iSNV with iVar using the following parameters: sample with a minimum consensus genome

length of 29,000 bases; sample with an average genome sequencing coverage depth of greater

than 200 reads per position; iSNV frequency of 5–95%; read depth of 400 at iSNV sites with a

Phred score of >30; iVar p-value of <0.00001. We masked sites commonly affected by

sequencing errors in both consensus sequences and iSNV calls [27].

Phylogenetic Analysis

Consensus sequences for each household were placed on the global SARS-CoV-2 phylogenetic

tree using UShER [28]. The tree versions used were from the week of 14 February 2022 and

included over 7.8 million genome sequences from GISAID, GenBank, COG-UK and CNCB. The

level of genomic sampling of the state or territory of each study site (Figure 1) was estimated

with subsampler [10] using case data and GISAID submission data. Subtrees were initially

constructed with 30 samples and then reconstructed with additional samples as needed to

visualize all genomes from a household in a single subtree (e.g., when samples existed within

large clusters of indistinguishable samples). The JSON files for each master tree and subtree are

available in Supplemental Dataset 1 and can be visualized in the auspice viewer at

https://auspice.us/. Trees were annotated and edited in FigTree using the subtree.nwk files

generated by UShER.

Data and materials availability

The consensus genomes that we generated for this study are publicly available on

https://github.com/lauringlab/SARS-CoV-2_Household_Diversity Accessions. Those for the

community sequences (largely from GISAID) can be found in the pdf tree files

("<Household_ID>.pdf") in Supplemental Dataset 1. Laboratories responsible for submissions

are acknowledged in Supplemental Table 1. Analysis code for the generation of consensus

sequences and phylogenetic analysis are available at https://github.com/lauringlab/SARS-CoV-

2_Household_Diversity.

RESULTS

The C-HEaRT (Utah and New York City) and COCOVID (Puerto Rico) studies performed active

surveillance for SARS-CoV-2 infection and CLI in 706 households with 2,369 participants (Table

1). During September 2020 through August 2021, the cumulative incidence of SARS-CoV-2

infection was 11% [96/842 participants in 41/190 (22%) households under surveillance] at the

Utah site and 7% [33/499 participants in 13/135 (10%) households] at the New York City site;

during June 2020 through September 2021, cumulative incidence was 5% at the Puerto Rico site

(49/1028 infections detected in 28/381 households).

Of the 191 participants with SARS-CoV-2 infections in these households, 147 (77%) in 70

households had samples with a Ct value <30 that were processed for whole genome

sequencing, of whom, 106 (72%) had samples that were successfully sequenced to sufficient

breadth and depth of coverage (see Methods). Of the 706 households at the three sites, 56

included ≥2 participants who were test-positive within a 14-day period, suggestive of within-

household transmission (Table 1). Twenty-six households had high quality sequence data on ≥2

of these contemporaneous infections. The SARS-CoV-2 clades and lineages identified (Table 2)

were the same among participants of the same household and reflect viruses circulating in the

corresponding time periods in the United States (www.outbreak.info)(Table 2).

We first used phylogenetic analysis of whole genome sequences to infer transmission linkage

within these 26 households. We used Usher [28] to obtain local sequences for each household

and to place household sequences simultaneously on a phylogenetic tree. Over 7.8 million

whole genome SARS-CoV-2 sequences were available at the time of this analysis. Because most

of these contextual sequences were from GISAID, we estimated the level of sampling at each

study site over time by dividing the number of GISAID sequences by the number of reported

cases [10]. Sampling of locally circulating viruses was low in 2020 (<2% cases sequenced) and

increased at all three sites beginning in early 2021 (>5% cases sequenced, Figure 1). In 2022,

Utah was generally better sampled (~10-30%) than New York or Puerto Rico (5-15%).

In 15 out of the 26 households that we studied, the consensus sequences of all cases were

indistinguishable and grouped together on their respective trees (representative trees in Figure

2, additional trees in Supplemental Figures 1-4). Given the epidemiologic linkage in the same

household, these can be considered sequence-confirmed transmission events. However, we

also found several trees in which these monophyletic groupings also included indistinguishable,

contemporaneous sequences from non-household members within the community of the same

locality (see trees in Figure 2). In two of the New York households, there were many such

sequences during a B.1.526 (Iota) variant wave (Supplemental Figures 1 and 2). Therefore, even

with a modest level of sampling (Figure 1), it is not uncommon to find indistinguishable viral

sequences from participants at the same region and time who presumably lack a documented

epidemiologic linkage.

In 11 households, the consensus sequences of the virus from one or more household members

differed at 1-2 positions over the nearly 30kb genome. This is not uncommon in transmission

chains, particularly ones that are longer or in which the samples are collected 7-14 days apart

(see Table 2 for time span). In nearly all cases, the trees from these households demonstrated

linkage and/or an ancestor/descendant relationship for the viral sequences (representative

trees in Figure 3, additional trees in Supplemental Figure 5). In some cases, the household

lineages were phylogenetically distinct from contemporaneous local sequences (e.g., UT2, UT4).

These tree structures supported transmission linkage, but low sampling of community cases

makes it hard to rule out missed linkages between members of the household and the larger

community. Indeed, there were some households in which there were sequences from the

larger community included among the same branches of the within-household sequences (e.g.,

UT10, PR3). As above, the low genomic diversity of SARS-CoV-2 and modest sampling of

community cases made it difficult to define a threshold to effectively rule in or rule out

transmission.

We next determined whether transmission inference could be improved by identifying iSNV

that were shared among members of a household. These would manifest as polymorphic sites

where the alternative allele, or mutation, is present but not fixed in the transmission chain.

While there was just one household where two participants shared a minority iSNV (PR5, Figure

4), several had iSNV in at least one individual at a consensus level (i.e., frequency >0.5), but that

had not yet achieved fixation (i.e., frequency >0.95). In the 15 households with

indistinguishable consensus sequences, each of the two participants in households NY5 and PR5

shared an iSNV that was consensus level, but not fixed. In three (UT12, UT5, UT3) out of the 11

households with distinct consensus sequences (Table 2), the consensus differences were due to

one or more participants having a non-reference iSNV that achieved consensus level, but not

fixation. Household UT4 had two participants with consensus level iSNV (Figure 4). In household

PR3, there was one site where one out of four members had a consensus level iSNV and

another member had this as a fixed mutation.

DISCUSSION

We evaluated the utility of SARS-CoV-2 sequence data in transmission inference using data

from two studies of household cohorts at three sites. In the household setting, where at least

two incident infections occurring within 14 days of one another are strongly suggestive of

transmission, we found that sequencing generally confirmed transmission linkage. The whole

genome consensus sequences of participants within a household were nearly always

indistinguishable or differed by one mutation. In some cases, these links were further

supported by the identification of iSNV shared among members of the household. Out of the 26

households evaluated, there was just one (UT10) in which the high average number of

consensus differences (two) and absence of shared iSNV called linkage into doubt. Importantly,

we frequently found multiple sequences from the community that were indistinguishable to

those within the household with even modest sampling (<5%) over the course of the pandemic.

This highlights the limits of "sequence-only" inference of transmission in hospitals or other

congregate settings where epidemiologic linkage is less certain.

Strengths of the study include our reliance on samples from active surveillance of longitudinal

cohorts and our use of quality controlled, deep sequencing. With weekly sampling of all

participants from a household, we were able to identify asymptomatic or mildly symptomatic

cases and avoid some of the bias of case-ascertainment studies, in which cases are recruited

based on a test-positive index. Together with our use of contemporaneous community

specimens collected from participants not in the households but from the same site, our data

provide a valuable benchmark for the expected SARS-CoV-2 diversity in households relative to

that in the community. The cohorts are also drawn from diverse geographic areas with varied

household sizes and composition [21,29]. Our assessment of viral diversity is strengthened by

our criteria for identifying consensus and minority iSNV [22]. The low observed diversity in this

study, in part, reflects the stringent thresholds applied to the sequence data . This conservative

approach reduces sequencing errors, which can be systematic and lead to incorrect

ascertainment of shared iSNV among unrelated participants [21–23,30].

This study had several notable limitations. First, we were relatively stringent in our criteria for

identifying iSNV and therefore may have underascertained shared diversity in the rare (<5%)

variant fraction within households. Second, given the limited number of households with

sequenced cases, we were unable to formulate a statistically robust approach to sequence-

based inference with clear cut-offs and associated positive and negative predictive values. Case-

ascertained cohorts or contact tracing studies offer a more efficient way to capture and

sequence many putative transmission pairs and will be useful as a setting in which to further

develop this approach. Third, while we believe that our data provide an important framework

for interpreting sequence data in studies of SARS-CoV-2 transmission, data from households

may not translate completely to hospitals and other congregate living settings, which may differ

in case density, contact frequency, and force of infection. Fourth, we assumed that household

cases testing positive for SARS-CoV-2 within 14 days of one another were linked by

transmission. If these cases represented distinct introductions into the household, we could

overestimate expected within-household diversity. Fifth, it is possible, but in our opinion

unlikely, that some of the community cases in our analysis actually had an epidemiologic

linkage to participants in these households.

Despite the limitations identified in this study, integration of sequence and epidemiologic data

can be a powerful approach to studies of SARS-CoV-2 transmission. In settings where there is

strong epidemiologic linkage among cases (e.g., known exposure or clear temporal and spatial

association), indistinguishable consensus sequences with or without shared iSNV should be

confirmatory. In these situations, single mutation differences among consensus sequences in a

cluster are not uncommon; mutations can fix along a transmission chain, particularly longer

ones over a greater timespan. However, if epidemiologic linkage is less certain, sequence

identity can only confirm transmission if the metapopulation is highly sampled and genetically

diverse. For example, early in the pandemic when circulating SARS-CoV-2 diversity was low,

many inpatients and employees in hospitals were found to share indistinguishable consensus

sequences, and even iSNV, without any apparent epidemiologic linkage [22,31]. This contrasts

with other studies of hospital outbreaks where the combination of contact tracing and

sequence data confirmed suspected transmission chains and identified new ones. We expect

that future studies of transmission in households, hospitals, and other congregate settings will

benefit from Bayesian methods, which can integrate epidemiologic and sequence data for

improved inference [32].

ACKNOWLEDGMENTS

We thank the participants in the C-HEaRT and COCOVID cohorts and all GISAID submitting

laboratories. We acknowledge Joseph Fauver for developing and suggesting the subsampler

tool used to generate Figure 1.

REFERENCES

1.   Lauring AS. Within-Host Viral Diversity: A Window into Viral Evolution. Annu Rev Virol

287   **2020**; 7:63–81.

288   2.   Kao RR, Haydon DT, Lycett SJ, Murcia PR. Supersize me: how whole-genome sequencing

289   and big data are transforming epidemiology. Trends Microbiol **2014**; 22:282–291.

290   3.   Lemieux JE, Siddle KJ, Shaw BM, et al. Phylogenetic analysis of SARS-CoV-2 in Boston

291   highlights the impact of superspreading events. Science **2021**; 371:eabe3261.

292   4.   Siddle KJ, Krasilnikova LA, Moreno GK, et al. Transmission from vaccinated individuals in a

293   large SARS-CoV-2 Delta variant outbreak. Cell **2022**; 185:485-492.e10.

294   5.   Zeller M, Gangavarapu K, Anderson C, et al. Emergence of an early SARS-CoV-2 epidemic in

295   the United States. Cell **2021**; 184:4939-4952.e15.

296   6.   Worobey M, Pekar J, Larsen BB, et al. The emergence of SARS-CoV-2 in Europe and North

297   America. Science **2020**; 370:564–570.

298   7.   du Plessis L, McCrone JT, Zarebski AE, et al. Establishment and lineage dynamics of the

299   SARS-CoV-2 epidemic in the UK. Science **2021**; 371:708–712.

300   8.   Candido DS, Claro IM, de Jesus JG, et al. Evolution and epidemic spread of SARS-CoV-2 in

301   Brazil. Science **2020**; 369:1255–1260.

302   9.   Viana R, Moyo S, Amoako DG, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron

303   variant in southern Africa. Nature **2022**; 603:679–686.

10.   Alpert T, Brito AF, Lasek-Nesselquist E, et al. Early introductions and transmission of SARS-

CoV-2 variant B.1.1.7 in the United States. Cell **2021**; 184:2595-2604.e13.

11.   Valesano AL, Fitzsimmons WJ, Blair CN, et al. SARS-CoV-2 Genomic Surveillance Reveals

Little Spread From a Large University Campus to the Surrounding Community. Open Forum

Infect Dis **2021**; 8:ofab518.

12.   Aggarwal D, Warne B, Jahun AS, et al. Genomic epidemiology of SARS-CoV-2 in a UK

university identifies dynamics of transmission. Nat Commun **2022**; 13:751.

13.   Lucey M, Macori G, Mullane N, et al. Whole-genome Sequencing to Track Severe Acute

Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Transmission in Nosocomial Outbreaks.

Clin Infect Dis **2021**; 72:e727–e735.

14.   Francis RV, Billam H, Clarke M, et al. The Impact of Real-Time Whole-Genome Sequencing

in Controlling Healthcare-Associated SARS-CoV-2 Outbreaks. J Infect Dis **2022**; 225:10–18.

15.   Meredith LW, Hamilton WL, Warne B, et al. Rapid implementation of SARS-CoV-2

sequencing to investigate cases of health-care associated COVID-19: a prospective

genomic surveillance study. Lancet Infect Dis **2020**; 20:1263–1271.

16.   Hamilton WL, Fieldman T, Jahun A, et al. Applying prospective genomic surveillance to

support investigation of hospital-onset COVID-19. Lancet Infect Dis **2021**; 21:916–917.

17.   Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 Infections and

Transmission in a Skilled Nursing Facility. N Engl J Med **2020**; 382:2081–2090.

18.   Addetia A, Crawford KHD, Dingens A, et al. Neutralizing Antibodies Correlate with

324   Protection from SARS-CoV-2 in Humans during a Fishery Vessel Outbreak with a High

325   Attack Rate. J Clin Microbiol **2020**; 58:e02107-20.

326   19.   McCrone JT, Woods RJ, Martin ET, Malosh RE, Monto AS, Lauring AS. Stochastic processes

327   constrain the within and between host evolution of influenza virus. eLife **2018**; 7:e35962.

328   20.   Worby CJ, Lipsitch M, Hanage WP. Shared Genomic Variants: Identification of Transmission

329   Routes Using Pathogen Deep-Sequence Data. Am J Epidemiol **2017**; 186:1209–1216.

330   21.   Braun KM, Moreno GK, Wagner C, et al. Acute SARS-CoV-2 infections harbor limited

331   within-host diversity and transmit via tight transmission bottlenecks. PLOS Pathog **2021**;

332   17:e1009849.

333   22.   Valesano AL, Rumfelt KE, Dimcheff DE, et al. Temporal dynamics of SARS-CoV-2 mutation

334   accumulation within and across infected hosts. PLOS Pathog **2021**; 17:e1009499.

335   23.   Tonkin-Hill G, Martincorena I, Amato R, et al. Patterns of within-host genetic diversity in

336   SARS-CoV-2. eLife **2021**; 10:e66857.

337   24.   Dawood FS, Porucznik CA, Veguilla V, et al. Incidence Rates, Household Infection Risk, and

338   Clinical Characteristics of SARS-CoV-2 Infection Among Children and Adults in Utah and

339   New York City, New York. JAMA Pediatr **2022**; 176:59.

25.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

ArXiv13033997 Q-Bio **2013**; Available at: http://arxiv.org/abs/1303.3997. Accessed 31 July

2020.

26.   Grubaugh ND, Gangavarapu K, Quick J, et al. An amplicon-based sequencing framework

for accurately measuring intrahost virus diversity using PrimalSeq and iVar. Genome Biol

2019; 20:8.

27.   De Maio N, Walker C, Borges R, Weilguny L, Slodkowicz G, Goldman N. Masking strategies

for SARS-CoV-2 alignments. Virological. 2020; Available at:

https://virological.org/t/masking-strategies-for-sars-cov-2-alignments/480. Accessed 10

March 2022.

28.   Turakhia Y, Thornlow B, Hinrichs AS, et al. Ultrafast Sample placement on Existing tRees

(UShER) enables real-time phylogenetics for the SARS-CoV-2 pandemic. Nat Genet **2021**;

53:809–816.

29.   Walter KS, Kim E, Verma R, et al. Shared within-host SARS-CoV-2 variation in households.

medRxiv **2022**; :2022.05.26.22275279.

30.   Lythgoe KA, Hall M, Ferretti L, et al. SARS-CoV-2 within-host diversity and transmission.

Science **2021**; :eabg0821.

31.   Braun KM, Moreno GK, Buys A, et al. Viral Sequencing to Investigate Sources of SARS-CoV-

2 Infection in US Healthcare Personnel. Clin Infect Dis **2021**; 73:e1329–e1336.

32.   Lindsey BB, Villabona-Arenas ChJ, Campbell F, et al. Characterising within-hospital SARS-

CoV-2 transmission events using epidemiological and viral genomic data across two

pandemic waves. Nat Commun **2022**; 13:671.

Table 1: Distribution of SARS-CoV-2 test-positive cases across cohorts and households

	New York City	Utah	Puerto Rico
Number of Households	135	190	381
Number of Participants	499	842	1028
Median household size (range)	4 (2-9)	4 (2-10)	2 (1-6)
Number of unique SARS-CoV-2-positive cases[a]	33	96	49
Households with 1 case	3	17	3
Households with 2 cases[b]	5	7	5
Households with 3 cases[b]	3	5	8
Households with 4 cases[b]	0	4	6
Households with 5 cases[b]	1	4	5
Households with 6 cases[b]	1	0	1
Households with 7 cases[b]	0	1	0
# of cases with sequence data / # sequenced	28/29	52/86	26/32

[a] total number over study period
[b] only households with cases testing positive within 14 days of each other

367 **Table 2:** Households with two or more incident SARS-CoV-2 infections within a 14- day period
368

Household	Number of Specimens Sequenced	Date First Specimen	Date Last Specimen	Nextclade Clade[a]	PANGO lineage[b]	Mean Consensus Diff[c]
PR1	3	9/2/20	9/8/20	20C	B.1.426	0
UT1	3	10/14/20	10/15/20	20G	B.1.2	0
PR2	4	10/24/20	10/30/20	20C	B.1.588	0.5
UT2	4	11/17/20	11/24/20	20B	B.1.1	0.5
UT3	6	11/24/20	12/2/20	20G	B.1.2	1.4
UT4	4	11/30/20	12/7/20	20B	B.1.1	1.5
UT5	2	12/2/20	12/3/20	20A	B.1.400	1
UT6	3	12/3/20	12/10/20	20G	B.1.2	0.67
PR3	4	12/3/20	12/11/20	20B	B.1.1.486	1.17
UT7	3	12/15/20	12/28/20	20A	B.1.596	0
UT8	2	12/28/20	1/1/21	21C (Epsilon)	B.1.429	0
UT9	2	1/12/21	1/14/21	20A	B.1.400	0
NY1	3	1/29/21	2/4/21	21F (iota)	B.1.526	0.67
PR4	2	2/8/21	2/8/21	20A	B.1.240	0
NY2	3	2/9/21	2/9/21	21F (Iota)	B.1.526	0
NY3	3	2/11/21	2/18/21	20C	B.1.582	0
NY4	2	2/21/21	3/5/21	21F (iota)	B.1.526	0
NY5	2	2/23/21	2/23/21	21F (Iota)	B.1.526	0
NY6	6	2/24/21	3/11/21	20C	B.1.637	1.13
UT10	2	3/1/21	3/15/21	21C (Epsilon)	B.1.427	2
NY7	3	3/3/21	3/16/21	20C	B.1.637	0
NY8	2	3/22/21	4/4/21	21F (Iota)	B.1.526	0
PR5	2	3/23/21	3/30/21	20B	R.1	0
PR6	3	4/23/21	4/23/21	20I (Alpha,V1)	Q.4	0
UT11	3	7/26/21	8/10/21	21J (Delta)	AY.44	0
UT12	2	8/4/21	8/4/21	21J (Delta)	AY.44	1

a Defined using nextclade, https://clades.nextstrain.org

b Defined using pango, https://cov-lineages.org/resources/pangolin.html

c Total number of pairwise unambiguous consensus differences between
sequences divided by total number of sequences in a household

369
370

    FIGURE LEGENDS

Fig. 1: **Cases and sampling density.** Columns show the number of SARS-CoV-2 infections (left y-

axis) in households from New York (NY, top), Puerto Rico (PR, middle), and Utah (UT, bottom)

cohorts by epiweek (x-axis). The sampling density (line) for community genomes in each state

or territory (right y-axis) was estimated as the proportion of cases with sequences available on

GISAID.

Fig. 2: Phylogenetic trees of sequences from households where all participants had

indistinguishable consensus sequences. Shown are four representative trees. Trees from 11

other households are shown in Supplemental Figures 1-4. Each tree is labeled with the

household identifier (NY = New York, UT = Utah, PR = Puerto Rico). The tips of household

sequences are colored cyan and those from non-household participants in the same community

in the same state or territory (2 letter abbreviation) are colored magenta. All other tips are

colored black. The collection date for each specimen is indicated. Genetic distance is

represented by the bar and corresponds to one mutation.

Fig. 3: Phylogenetic trees of sequences from households where participants had distinct

consensus sequences. Shown are four representative trees. Trees from 7 other households are

shown in Supplemental Figure 5. Each tree is labeled with the household identifier (NY = New

York, UT = Utah, PR = Puerto Rico). The tips of household sequences are colored cyan and those

from the same state or territory (2 letter abbreviation) are colored magenta. All other tips are

colored black. The collection date for each sample is indicated. Genetic distance is represented

by the bar and corresponds to one mutation.

Fig. 4: Shared single nucleotide polymorphisms within households. Each panel shows one of

the 8 households in which members shared a polymorphic site. The frequency (5-95%) of the

indicated mutation (relative to the Wuhan/Hu-1 reference) is shown on the y-axis and the

individual/sequence identifier is shown on the x-axis. Shared variants that did and did not lead

to a consensus level difference between household members are shown as circles and

diamonds, respectively. Mutations that are fixed (>95% frequency) are shown as squares.

Supplemental Figures 1-4: Phylogenetic trees of sequences from households where all

participants had indistinguishable consensus sequences. Each tree is labeled with the

household identifier (NY = New York, UT = Utah, PR = Puerto Rico). The tips of household

sequences are colored cyan and those from the same state or territory (2 letter abbreviation)

are colored magenta. All other tips are colored black. The collection date for each sample is

indicated. Genetic distance is represented by the bar and corresponds to one mutation.

Supplemental Figure 5: Phylogenetic trees of sequences from seven households where

participants had distinct consensus sequences. Each tree is labeled with the household

identifier (NY = New York, UT = Utah, PR = Puerto Rico). The tips of household sequences are

colored cyan and those from the same state or territory (2 letter abbreviation) are colored

magenta. All other tips are colored black. The collection date for each sample is indicated.

Genetic distance is represented by the bar and corresponds to one mutation.

Supplemental Table 1: Submitting laboratories for GISAID sequences used in this study. GISAID

identifiers can be found in the unedited trees in the Supplemental Dataset.

Figure 1

Figure 2

Figure 3

Figure 4

Supplemental Figure 1

NY1

Supplemental Figure 2

NY2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

UT2

UT6

UT12

NY6

UT5

UT8

PR2