

Two years of COVID-19: Excess mortality by age, region, gender, and race/ethnicity in the United States during the COVID-19 pandemic, March 1, 2020, through February 28, 2022.

Jeremy Samuel Faust, MD MS*

Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts

Chengan Du, PhD

Center for Outcomes Research and Evaluation, Yale University School of Medicine, New Haven, Connecticut

Benjamin Renton, BA

Ariadne Labs, Brigham and Women's Hospital, Boston, Massachusetts

Chenxue Liang, MS, MPH

Center for Outcomes Research and Evaluation, Yale University School of Medicine, New Haven, Connecticut

Alexander Junxiang Chen

Harvard University, Cambridge, Massachusetts

Shu-Xia Li, PhD

Center for Outcomes Research and Evaluation, Yale University School of Medicine, New Haven, Connecticut

Zhenqiu Lin, PhD

Center for Outcomes Research and Evaluation, Yale University School of Medicine, New Haven, Connecticut

Marcella Nunez-Smith, MD, MHS

Yale University School of Medicine, New Haven, Connecticut

Harlan M. Krumholz, MD, SM

Center for Outcomes Research and Evaluation, Yale University School of Medicine, New Haven, Connecticut

*corresponding author

Brigham and Women's Hospital, Harvard Medical School

Jsf Faust@BWH.Harvard.edu

Abstract

Introduction: Excess mortality does not depend on labeling the cause of death and is an accurate representation of the pandemic population-level effects. A comprehensive evaluation of all-cause excess mortality in the United States during the first two years of the COVID-19 pandemic, stratified by age, sex, region, and race/ethnicity can provide insight into the extent and variation in harm.

Methods: With Centers for Disease Control and Prevention (CDC)/National Center for Health Statistics (NCHS) data from 2014-2022, we use seasonal autoregressive integrated moving averages (sARIMA) to estimate excess mortality during the pandemic, defined as the difference between the number of observed and expected deaths. We continuously correct monthly expected deaths to reflect the decreased population owing to cumulative pandemic-associated excess deaths recorded. We calculate excess mortality for the total US population, and by age, sex, US census division, and race/ethnicity.

Results: From March 1, 2020, through February 28, 2022, there were 1.17 million excess deaths in the United States. Overall, mortality was 20% higher than expected during the study period. Of the excess deaths, 799,477 (68%) were among residents aged 65 and older. The largest relative increase in all-cause mortality was 27% among adults ages 18-49 years. Males comprised most of the excess mortality (57%), but this predominance declined with age. A higher relative mortality occurred among non-Hispanic American Indian/Alaskan Native, non-Hispanic Black, non-Hispanic Native Hawaiian and Other Pacific Islander, Hispanic people. Excess mortality differed by region; the highest rates were in the South, including in the population ages ≥ 65 years. Excess mortality rose and fell contemporaneously with COVID-19 waves.

Conclusion: In the first two years of the pandemic, the US experienced 1.17 million excess deaths, with greater relative increases in all-cause mortality among men, in American Indian/Alaskan Native, Black and Hispanic people, and the South.

Main body

Introduction

Disease-specific mortality determinations are subjective, leading to overcounts and undercounts.^{1,2} For a new cause of death, such as SARS-CoV-2 infection, the subjectivity may be amplified because of variations in testing and changes in other causes of death.^{3,4} All-cause excess mortality provides a better estimate of the overall burden of the pandemic as it does not depend on the cause of death determinations.⁵ Also, excess mortality measurement by subgroups enables comparisons without the need to adjust for health differences among populations or historical differences.

The COVID-19 outbreak in the United States produced all-cause excess mortality.^{6,7} Several reports noted marked disparities in all-cause excess mortality or COVID-19-attributed mortality by age, gender, and race/ethnicity.^{8–11} Here, we provide an excess mortality update two years into the COVID-19 pandemic in the United States. These estimates, unlike others, reflect the differences in the denominator by region, age, and gender, race/ethnicity owing to variation in pandemic-associated excess mortality. Thus, the present analysis, which accounts for these changes, provides a more accurate (and higher) excess mortality estimate for the first two years of the US COVID-19 outbreak than official estimates. We also provide period and overall sub-analyses by age group, sex, geographic location, and race/ethnicity.

Methods

Data Sources

We used publicly available Centers for Disease Control and Prevention (CDC)/National Center for Health Statistics (NCHS) population data from 2014-2019, and monthly mortality data from January 2015-February 2022.^{12,13} Based on 2014-2019 census data, we used autoregressive integrated moving averages (ARIMA) to project the expected populations for 2020-2022, and then smoothed the yearly increases by dividing the yearly change by 12 and adding 1/12 of the growth to each successive month. We further adjusted (i.e., lowered) the population during the pandemic period to in proportion to the cumulative excess mortality recorded during the pandemic, so as to capture the smaller than expected population, owing to pandemic-associated all-cause excess mortality, as described previously.^{14–16}

Excess Mortality

Excess mortality is defined as the difference between the number of observed and expected deaths.

Statistical Analysis

We divided the two-year study period (March 1, 2020, through February 28, 2022) into 6 periods: Wave 1 (March 1, 2020-May 31, 2020), Wave 2 (June 1, 2020-September 30, 2020), Wave 3 (October 1, 2020-February 28, 2021), Spring 2021 (March 1, 2020-June 30, 2021), Delta (July 1, 2021-December 31, 2021), and Omicron (January 1, 2022-February 28, 2022).

Then, using CDC Wonder data, we apply seasonal ARIMA (sARIMA) to monthly mortality data from January 1, 2015-February 29, 2020 to project the monthly number of expected deaths for the period of March 1, 2020-February 28, 2022, normalizing for difference in days in each month.¹² Expected deaths for each of the four US Census Bureaus is determined by summing estimated from the nine US Census Divisions, which in turn are determined by age and gender subgroup projections (ages 0-17, 18-49, 50-64, and ≥ 65 by sex). These 72 components are then summed to create US-level estimates. Separately, state-level estimates are also determined. For 37 states, data were adequate to sum age-specific components models; for the 14 states with inadequate age-specific data, models are determined without age-stratification.

For the race and ethnicity analysis, a separate model is used because of recent changes in how CDC/NCHS reports relevant data and due to the smaller population of one group. US level race/ethnicity estimates are determined by summing 3 age-stratified sARIMA components parts (ages 0-24, 25-64, and ≥ 65) for 6 groups. The 7th group (Native Hawaiian and Other Pacific Islander), sARIMA by age group was not possible due to the small size of the population and therefore the entire population is modelled together. Regional data are not available.

As we have described previously, we continuously corrected the number of monthly expected deaths to reflect the decreased population owing to cumulative pandemic-associated excess deaths recorded. The raw number of excess deaths, observed-to-expected ratios, and incident rates per 100,000 person-periods are reported for excess mortality.^{14,15} Raw and incident rates for COVID-19-specific mortality (ICD U07.1) is also reported. Analyses were conducted with R version 4.1.2. Statistical significance was defined as a 95% CI that did not include 0.

This study used publicly available data and was not subject to institutional review board approval according to the National Center for Health Statistics policy.

Results

During the first two years of the COVID-19 pandemic in the US (March 1, 2020 – February 28, 2022), there were 1,173,156 all-cause excess deaths (5,817,974 expected, 6,991,130 observed, 337.1 per 100,000 pandemic period population), and observed-to-expected ratio of 1.20 (Figure 1A, Table 1). All-cause excess mortality varied by region (Figure 1A, Table 1), and rose and fell contemporaneously with COVID-19-specific mortality in all states and regional jurisdictions (Supplemental Appendix Figure 1A).

Regional analysis. During the overall study period, there were notable differences in excess mortality by Census Bureau region, (Figure 1A), by Census Division, and by state (Supplemental Appendix Figure 1). There was excess mortality in all four Census Bureau regions during each of the 6 periods (Table 2). Census bureau contribution (Figure 1B-C) to and share of excess mortality (Supplemental Appendix Figure 1D) changed over time. During the study period, the highest cumulative excess mortality was observed in the South (rate and number), and the lowest overall excess mortality was observed in the West (rate) or Northeast (number), (Figure 1C, Table 2, Supplemental Appendix Figure 1B-C). Overall, 43.3% of the excess mortality was in the South despite comprising 38.5% of the US population; this accounted for 507,454 excess deaths (398.9 per 100,000), corresponding to an observed-to-expected ratio of 1.22. By comparison, the least excess mortality was observed in the Midwest where 230,098 excess

deaths occurred (337.1 per 100,000), corresponding to an observed-to-expected ratio of 1.17, representing 20.7% of US excess mortality despite comprising 19.6% of the population. After the initial wave, the lowest rates of excess mortality were in the Northeast (220.7 per 100,000) while rates were higher in the other regions (range: 291.0-380.0 per 100,000, Table 2).

Age analysis. Excess mortality was observed among all adult age groups during the overall study period and varied by age (Figure 2A-D, Supplemental Appendix Figure 2). In adults, there was excess mortality in each of the 6 periods, except for US residents ages 65 and older during the spring 2021 period (Table 1-2, Figure 2A)

Numerically, the highest number of excess deaths occurred in residents ages 65 and older (Table 1). However, the largest relative increase in mortality occurred among residents ages 18-49 (Table 1).

Statistically significant excess mortality was not observed among children ages 0-17 during the pandemic months of 2020 but was observed during the Delta wave (1,532 excess deaths, IR 2.1 per 100,000 population, observed-to-expected ratio of 1.09) and 2021 overall (2,228, IR. 3.1 per 100,000 population, observed-to-expected ratio of 1.07) (Table 1, Table 3) for both males and females. Most excess deaths among children occurred in the South. Census Divisions and states large enough to provide adequate data indicated similar regional and subregional trends. (Supplemental Appendix Figure 2).

Among residents ages 65 years or older, the highest overall rates of excess mortality during the pandemic period occurred in the South, even though during Wave 1, the highest rate of excess mortality recorded in any period among any age group occurred in Northeast residents ages 65 years and over (Supplemental Appendix Table 1).

Sex analysis. Males accounted for 57% of excess mortality during the study period (Table 1). There were 663,228 excess deaths in men and 509,928 in women during the study period. Males comprised most of the excess deaths in all age groups, although the male share of excess mortality decreased with age (Table 3, Supplemental Figure 3, Figure 3A, 3B). The rate of excess death was 419.5 per 100,000 in men and 305.4 per 100,000 in women. Increased disparities became apparent during the vaccine period (Table 3, Figure 3C).

Race/Ethnicity analysis. There were marked differences in excess mortality by race, and ethnicity (Table 4, Figure 4A-B, Supplemental Appendix Figure 4). The highest increase in mortality was observed among American Indian/Alaska Native people, followed by Hispanic people, followed by non-Hispanic Black people and non-Hispanic Native Hawaiian and Other Pacific Islander (Table 4). There were marked differences by age group and wave. The highest incident rate of excess death was 661.1 per 100,000 in American Indian/Alaska Native people, followed by 361.2 per 100,000 in non-Hispanic White people. There were 708,740 excess deaths in non-Hispanic White people, 201,667 excess deaths in non-Hispanic Black people, and 187,654 excess deaths in Hispanic people.

Discussion

Using methodology that accounts for population changes across the pandemic and employs seasonal autoregressive integrated moving averages, we find that the United States had 1,173,156 all-cause excess deaths. This number of deaths represents a 20% higher number than expected based on the pattern in the five years immediately before the pandemic. This study is the first to show the distinct regional and age-specific changes in all-cause excess mortality during the initial two years of the pandemic.

The number of deaths was 43% higher than expected in American Indian/Alaska Native people and 41% higher for Hispanic people, and 28% higher for Black people, compared with 16% for White people. The South suffered the greatest burden, with 507,454 excess deaths, representing 22% higher than expected, compared with the Midwest, where mortality was 17% higher than expected. The Northeast experienced a 20% increase in mortality. Although experiencing the brunt of the initial wave, smaller deviations from historical mortality rates occurred after the first wave (14%) as compared to elsewhere, where after the initial wave, deviations from expected deaths were 18% (Midwest), 22% (West), and 24% (South).

The regional differences show that even accounting for baseline differences in expected mortality and the variation in the population, the South suffered the most mortality burden from the pandemic, with the share increasing later in the pandemic period. While the South historically has higher age-adjusted mortality than the rest of the nation, the excess mortality metric conveys increases beyond the usual mortality patterns. Moreover, the pattern reveals that while the Northeast bore the brunt of the early pandemic period, the South exceeded the Northeast especially as vaccines became available, consistent with national data about vaccine adoption. However, other COVID-19 mitigation efforts, which varied by region may have contributed to these findings. The tracking the excess mortality with the infection waves indicates that these excess deaths are related to the virus and not a by-product of other societal changes.

This study puts a number on the excess mortality toll of the pandemic and provides an assessment that is not dependent on the cause of death labels. The study also illuminates the disproportionate toll of excess mortality on American Indian/Alaska natives, Black and Hispanic people, as we and others have reported in the past, and which continued during the vaccine and variant era.^{8,17} Of note, the numerical majority of excess mortality in American Indian/Alaska Native people occurred in people under the age of 65, a finding unique to this demographic, and consistent with recent reports of life expectancy changes during the pandemic.¹⁸ These findings are consistent with other data describing how infection, hospitalization and death varied by social factors, including race and ethnicity.^{19–22} Differences in mortality by race/ethnicity reflect not genetic differences, but social identity and race/ethnicity-based injustices derived from the organization of the United States and the racism that has spanned centuries and persists today, including access to primary care.²³ The excess deaths are beyond the higher historic baseline death rates of these groups, indicating that the impact of structural racism on health outcomes was augmented during the pandemic. Considering this, the US will need to reckon with the magnitude of the loss of life and the inherent unfairness of how the excess deaths are distributed among the population and seek solutions for the future.

The age-related findings also warrant attention. First, the vaccine rollout period (Winter and Spring 2021) was associated with a precipitous decrease in the share (and, in fact, the temporary elimination) of excess mortality contributed by the ≥ 65 -year-old demographic. In fact, during 2021, the only age group with less excess mortality than 2020, was the ≥ 65 -year-old group, likely reflecting vaccine uptake. Remarkably, during April and May of 2021, a numerical majority of excess mortality was derived from adults ages 50-64 years, the only time a non-geriatric demographic represented most of the excess mortality during the pandemic. However, the share of deaths from the ≥ 65 -year-old demographic soon became the majority again, as younger groups began to receive vaccinations, and as the Delta variant became dominant, which decreased vaccine effectiveness against severe disease in older populations.²⁴ Second, this report reveals all-cause excess mortality in US residents ages 0-17 during the COVID-19 period, which occurred during the Delta period, which was sufficient to cause statistically significant all-cause excess mortality for 2021 overall. Reasons for this are unclear, but 69% (1,532 of 2,228) excess deaths in this demographic recorded in 2021 occurred during the Delta wave, and were concentrated in the South, where vaccine uptake among the general population was lower and fewer non-pharmacologic interventions remained in place.

The study also found that men had higher excess mortality, accounting for 57% of those deaths even though on a population basis, there are many more women in older age groups. Sex differences in hospitalizations and deaths have been noted in the past, but the reason why men are more vulnerable remains unexplained.^{10,25} The method used in this study addresses the higher historic mortality rates for men compared with women, and these excess deaths are in addition to a baseline higher risk. However, during the vaccine era, men have been less likely to receive COVID vaccines, which may be one explanation, especially given the increase in this disparity since the vaccine rollout began in early 2021.²⁶ Another explanation is that men are less likely to have existing primary care, and may seek out healthcare less often and later.²³

This study has notable features that strengthen the validity of our findings. By summing component age and sex models to create regional and ultimately US-level figures, we generated the high fidelity excess mortality estimates, avoiding biases that are otherwise introduced due to secular demographic heterogeneity that exists across jurisdictions. While taking age differences and secular changes in demographics across jurisdictions is important, we believe that age-adjustment obscures authentic region-to-region comparisons. In contrast, by age stratifying, our model preserves real-world populations differences, while avoiding Simpson's Paradox.²⁷ Additionally, we corrected for the ongoing population changes (i.e., smaller increases than expected), owing to cumulative excess mortality observed during the pandemic, as described previously.^{8,14-16}

The study has several limitations. First, while all-cause and COVID-19-specific mortality trends were highly aligned, we cannot definitively establish the cause of the excess deaths. We note that the strength of the excess mortality metric is that it encompasses all deaths, thereby more comprehensively capturing the population-level effects of the pandemic. However, analysis of disease-specific causes of death, particularly in younger demographics, may still be fruitful. Second, we used mortality categorized by decedent residence (not occurrence) due to the data availability. Therefore, a small fraction of deaths may have occurred in jurisdictions other than the decedents' home states. Third, the race/ethnicity data depends on what was reported on death

certificates. This is particularly important because of the magnitude of the disparities among American Indian/Alaska Native people described. However, prior research indicates that American Indian/Alaska Native demographic information is the most likely group to have its race/ethnicity data omitted on death certificates (which are most frequently misclassified as White), suggesting that our findings are a lower threshold estimate of this disparity.²⁸

In conclusion, the COVID-19 pandemic was associated with an increase in all-cause deaths in the United States by more than 20% beyond what would have been expected based on recent historic norms, resulting in nearly 1.2 million more deaths than expected in the first two years. Several groups suffered considerably a higher deviation in all-cause deaths compared with the historic norms, including young people, American Indian/Alaska Native, non-Hispanic Black and Hispanic people, men, and those living in the South.

Future pandemic planning must consider strategies to better mitigate the overall harm, and specifically address populations at highest risk. The challenge going forward is to develop pandemic response that mitigates harm, particularly where the identified risk is greatest.

Acknowledgements: The authors wish to thank Katie Dickerson Mayes, MD, PhD, Andrew Marshall, MD, Jacob Aguilar, PhD, Kristen Panthagani, MD, PhD and Lauren Rossen, PhD for their assistance and expertise.

Data Statement: Dr. Faust and Mr. Renton had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Conflict of Interest Disclosures: Dr. Krumholz reported receiving consulting fees from UnitedHealth, Element Science, Aetna, Reality Labs, F-Prime, and Tesseract/4Catalyst; serving as an expert witness for Martin Baughman law firm, Arnold and Porter law firm, and Siegfried and Jensen law firm; being a cofounder of Hugo Health, a personal health information platform; being a cofounder of Refactor Health, an enterprise health-care, artificial intelligence-augmented data management company; receiving contracts from the Centers for Medicare & Medicaid Services through Yale New Haven Hospital to develop and maintain performance measures that are publicly reported; and receiving grants from Johnson & Johnson outside the submitted work. The other authors declare no competing interests.

References:

1. Sehdev AES, Hutchins GM. Problems With Proper Completion and Accuracy of the Cause-of-Death Statement. *Arch Intern Med*. 2001;161(2):277-284. doi:10.1001/archinte.161.2.277
2. Mieno MN, Tanaka N, Arai T, et al. Accuracy of Death Certificates and Assessment of Factors for Misclassification of Underlying Cause of Death. *J Epidemiol*. 2016;26(4):191-198. doi:10.2188/jea.JE20150010
3. Gill JR, DeJoseph ME. The Importance of Proper Death Certification During the COVID-19 Pandemic. *JAMA*. 2020;324(1):27-28. doi:10.1001/jama.2020.9536
4. US Centers for Disease Control and Prevention/NCHS. Guidance for certifying deaths due to coronavirus disease 2019 (COVID-19). Accessed September 3, 2022. <https://stacks.cdc.gov/view/cdc/86344>
5. National Center for Health Statistics. Excess Deaths Associated with COVID-19. Published December 30, 2020. Accessed January 4, 2021. https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm
6. Rossen LM. Excess Deaths Associated with COVID-19, by Age and Race and Ethnicity — United States, January 26–October 3, 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69. doi:10.15585/mmwr.mm6942e2
7. Woolf SH, Chapman DA, Sabo RT, Zimmerman EB. Excess Deaths From COVID-19 and Other Causes in the US, March 1, 2020, to January 2, 2021. *JAMA*. 2021;325(17):1786-1789. doi:10.1001/jama.2021.5199
8. Rossen LM. Disparities in Excess Mortality Associated with COVID-19 — United States, 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70. doi:10.15585/mmwr.mm7033a2
9. Stoto MA, Schlageter S, Kraemer JD. COVID-19 mortality in the United States: It's been two Americas from the start. *PLOS ONE*. 2022;17(4):e0265053. doi:10.1371/journal.pone.0265053
10. Nielsen J, Nørgaard SK, Lanzieri G, Vestergaard LS, Moelbak K. Sex-differences in COVID-19 associated excess mortality is not exceptional for the COVID-19 pandemic. *Sci Rep*. 2021;11(1):20815. doi:10.1038/s41598-021-00213-w
11. Faust JS, Krumholz HM, Du C, et al. All-Cause Excess Mortality and COVID-19–Related Mortality Among US Adults Aged 25–44 Years, March–July 2020. *JAMA*. Published online December 16, 2020. doi:10.1001/jama.2020.24243
12. US Centers for Disease Control and Prevention/NCHS. CDC WONDER. Accessed September 3, 2022. <https://wonder.cdc.gov/>

13. US Centers for Disease Control and Prevention/NCHS. Provisional Mortality Statistics, 2018 through Last Month Request. Accessed September 3, 2022. <https://wonder.cdc.gov/mcd-icd10-provisional.html>
14. Faust JS, Du C, Liang C, et al. Excess Mortality in Massachusetts During the Delta and Omicron Waves of COVID-19. *JAMA*. Published online May 20, 2022. doi:10.1001/jama.2022.8045
15. Faust JS, Du C, Li SX, Lin Z, Krumholz HM. Correcting excess mortality for pandemic-associated population decreases. *medRxiv*. Published online February 12, 2021:2021.02.10.21251461. doi:10.1101/2021.02.10.21251461
16. Faust JS, Renton B, Chen AJ, et al. Uncoupling of all-cause excess mortality from COVID-19 cases in a highly vaccinated state. *Lancet Infect Dis*. 2022;0(0). doi:10.1016/S1473-3099(22)00547-3
17. Shiels MS, Haque AT, Haozous EA, et al. Racial and Ethnic Disparities in Excess Deaths During the COVID-19 Pandemic, March to December 2020. *Ann Intern Med*. 2021;174(12):1693-1699. doi:10.7326/M21-2134
18. Arias E, Ahmad FB. Provisional life expectancy estimates for 2021. August 2022. DOI: <https://dx.doi.org/10.15620/cdc:118999>. *Vital Stat Rapid Release*.:16.
19. Acosta AM, Garg S, Pham H, et al. Racial and Ethnic Disparities in Rates of COVID-19–Associated Hospitalization, Intensive Care Unit Admission, and In-Hospital Death in the United States From March 2020 to February 2021. *JAMA Netw Open*. 2021;4(10):e2130479. doi:10.1001/jamanetworkopen.2021.30479
20. Song Z, Zhang X, Patterson LJ, Barnes CL, Haas DA. Racial and Ethnic Disparities in Hospitalization Outcomes Among Medicare Beneficiaries During the COVID-19 Pandemic. *JAMA Health Forum*. 2021;2(12):e214223. doi:10.1001/jamahealthforum.2021.4223
21. US Centers for Disease Control and Prevention/NCHS. AH Excess Deaths by Sex, Age, and Race and Hispanic Origin | Data | Centers for Disease Control and Prevention. Accessed September 6, 2021. <https://data.cdc.gov/NCHS/AH-Excess-Deaths-by-Sex-Age-and-Race-and-Hispanic-/m74n-4hbs>
22. Romano SD. Trends in Racial and Ethnic Disparities in COVID-19 Hospitalizations, by Region — United States, March–December 2020. *MMWR Morb Mortal Wkly Rep*. 2021;70. doi:10.15585/mmwr.mm7015e2
23. Levine DM, Linder JA, Landon BE. Characteristics of Americans With Primary Care and Changes Over Time, 2002-2015. *JAMA Intern Med*. 2020;180(3):463-466. doi:10.1001/jamainternmed.2019.6282
24. Rosenberg ES. New COVID-19 Cases and Hospitalizations Among Adults, by Vaccination Status — New York, May 3–July 25, 2021. *MMWR Morb Mortal Wkly Rep*. 2021;70. doi:10.15585/mmwr.mm7034e1

25. Nguyen NT, Chinn J, Ferrante MD, Kirby KA, Hohmann SF, Amin A. Male gender is a predictor of higher mortality in hospitalized adults with COVID-19. *PLOS ONE*. 2021;16(7):e0254066. doi:10.1371/journal.pone.0254066
26. Ungar L. The Gender Vaccine Gap: More Women Than Men Are Getting Covid Shots. Kaiser Health News. Published April 12, 2021. Accessed September 8, 2022. <https://khn.org/news/article/gender-vaccine-gap-more-women-than-men-vaccinated-against-covid/>
27. Shiels MS, Almeida JS, García-Closas M, Albert PS, Freedman ND, de González AB. Impact of Population Growth and Aging on Estimates of Excess U.S. Deaths During the COVID-19 Pandemic, March to August 2020. *Ann Intern Med*. Published online December 15, 2020. doi:10.7326/M20-7385
28. Arias E, Heron M, Ni H. The Validity of Race and Hispanic-origin Reporting on Death Certificates in the United States: An Update. *Vital Health Stat*. 2(2016 Aug 1;(172):1-21.):29.

List of Exhibits:

Figure 1. Monthly excess deaths in the United States, March 1, 2020 through February 28, 2022.

Panel A: Observed (yellow line) and expected (blue line) deaths, for the US and by Census Bureau regions per 100,000 person-months.

Panel B: Cumulative excess deaths by Census Bureau region, per 100,000 person-months.

Panel C: Cumulative excess deaths by Census Bureau region, number.

Table 1: Excess and COVID-19–Attributed Deaths in the United States, by sex, or age group, March 1, 2020, through February 28, 2022

Table 2: Excess and COVID-19–Attributed Deaths in the United States, by Census Bureau region, March 1, 2020, through February 28, 2022

Figure 2. Monthly observed (yellow) and expected (blue) deaths in the United States, by age group for each Census Bureau region, and Division, March 1, 2020, through February 28, 2022.

Panel A: Ages ≥ 65 years, per 100,000 person-months.

Panel B: Ages 50–64 years, per 100,000 person-months.

Panel C: Ages 18–49 years, per 100,000 person-months.

Panel D: Ages 0–17 years, per 100,000 person-months.

Panel E: Cumulative excess deaths by Census Bureau division and age, per 100,000 person months.

Panel F: Cumulative excess deaths by Census Bureau region and age, per 100,000 person months.

Table 3: Excess and COVID-19–Attributed Deaths in the United States, by sex and age group, March 1, 2020, through February 28, 2022.

Figure 3. Monthly excess deaths in the United States, by sex, March 1, 2020, through February 28, 2022.

Panel A: Excess mortality, all ages, ages 0–17, ages 18–49, ages 50–64, ages ≥ 65 years, by sex, per 100,000 person-months.

Panel B: Observed (yellow) and expected (blue) deaths, by sex and age, per 100,000 person-months.

Panel C: Cumulative excess deaths by sex and age, per 100,000 person months.

Table 4: Excess and COVID-19–Attributed Deaths in the United States, race/ethnicity, March 1, 2020, through February 28, 2022.

Figure 4. Monthly excess deaths in the United States, by race or ethnicity, March 1, 2020, through February 28, 2022.

Panel A: Excess mortality, all ages, ages 0–25, ages 25–64, ages ≥ 65 years, by race or ethnicity, per 100,000 person-months.

Panel B: Observed (yellow) and expected (blue) deaths, by age and race or ethnicity, per 100,000 person-months.

Panel C: Cumulative excess deaths by age and race or ethnicity, per 100,000 person months.

Figure 1A




Figure 1B

Cumulative All-Cause Excess Mortality by Census Bureau Region, IR
March 1, 2020-February 28, 2022





Figure 1C

Cumulative All-Cause Excess Mortality by Census Bureau Region
March 1, 2020-February 28, 2022



It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Table 1

	Expected deaths (95% CI)	Observed deaths	Ratio of observed to expected deaths (95% CI)	Excess deaths, no (95% CI)	Excess deaths, IR (100,000 person-months)	COVID-19 deaths, no	COVID-19 deaths, IR (100,000 person-months)	Excess Deaths (100,000 person-waves)
All Genders, All Ages								
Wave 1 (March-May 2020, 3 months)	742,778 (707,746 - 777,810)	871,236	1.17 (1.12 - 1.23)	128,458 (93,426 - 163,490)	13.00	110,609	11.20	39.01
Wave 2 (June 2020-September 2020, 4 months)	916,705 (875,502 - 957,907)	1,061,830	1.16 (1.11 - 1.21)	145,125 (103,923 - 186,328)	11.01	97,728	7.42	44.05
Wave 3 (October 2020-February 2021, 5 months)	1,237,280 (1,190,610 - 1,283,950)	1,596,881	1.29 (1.24 - 1.34)	359,601 (312,931 - 406,271)	21.81	329,498	19.98	109.03
Spring 2021 (March 2021-June 2021, 4 months)	974,008 (932,338 - 1,015,677)	1,029,283	1.06 (1.01 - 1.10)	55,276 (13,606 - 96,945)	4.19	64,694	4.90	16.74
Delta (July 2021-December 2021, 6 months)	1,447,202 (1,394,883 - 1,499,520)	1,778,560	1.23 (1.19 - 1.28)	331,359 (279,040 - 383,677)	16.71	242,846	12.24	100.24
Omicron (January 2022-February 2022, 2 months)	500,003 (472,251 - 527,754)	653,340	1.31 (1.24 - 1.38)	153,338 (125,586 - 181,089)	23.18	131,741	19.91	46.35
2020 (March 2020-December 2020, 10 months)	2,401,950	2,875,140	1.20	473,191	14.36	384,189	11.66	143.60
2021 (January 2021-December 2021, 12 months)	2,916,022	3,462,650	1.19	546,628	13.79	461,186	11.63	165.49
2022 (January 2022-February 2022, 2 months)	500,003 (472,251 - 527,754)	653,340	1.31 (1.24 - 1.38)	153,338 (125,817 - 180,858)	23.18	131,741	19.91	46.35
Post Wave 1 (June 2020-February 2022, 21 months)	5,075,196	6,119,894	1.21	1,044,698	15.07	866,507	12.50	316.45
Pandemic period (March 2020-February 2022, 24 months)	5,817,974	6,991,130	1.20	1,173,156	14.81	977,116	12.34	355.48
Male								
Wave 1 (March-May 2020, 3 months)	384,606 (371,662 - 397,550)	453,962	1.18 (1.14 - 1.22)	69,356 (56,412 - 82,300)	14.26	59,430	12.22	42.77
Wave 2 (June 2020-September 2020, 4 months)	480,538 (465,334 - 495,742)	559,107	1.16 (1.11 - 1.20)	78,569 (63,365 - 93,773)	12.11	53,232	8.20	48.43
Wave 3 (October 2020-February 2021, 5 months)	639,923 (622,656 - 657,191)	843,744	1.32 (1.28 - 1.36)	203,821 (186,553 - 221,088)	25.10	181,799	22.39	125.51
Spring 2021 (March 2021-June 2021, 4 months)	507,429 (492,016 - 522,841)	545,186	1.07 (1.04 - 1.11)	37,758 (22,345 - 53,170)	5.81	35,818	5.51	23.23
Delta (July 2021-December 2021, 6 months)	756,602 (737,185 - 776,019)	943,617	1.25 (1.22 - 1.28)	187,015 (167,598 - 206,432)	19.16	137,178	14.05	114.93
Omicron (January 2022-February 2022, 2 months)	259,010 (248,929 - 269,091)	345,719	1.33 (1.28 - 1.39)	86,709 (76,628 - 96,790)	26.63	73,710	22.64	53.26
2020 (March 2020-December 2020, 10 months)	1,249,777	1,508,531	1.21	258,755	15.95	208,541	12.85	159.47
2021 (January 2021-December 2021, 12 months)	1,519,321	1,837,085	1.21	317,764	16.29	258,916	13.27	195.43
2022 (January 2022-February 2022, 2 months)	259,010 (248,929 - 269,091)	345,719	1.33 (1.28 - 1.39)	86,709 (76,648 - 96,770)	26.63	73,710	22.64	53.26
Post Wave 1 (June 2020-February 2022, 21 months)	2,643,502	3,237,373	1.22	593,871	17.40	481,737	14.12	365.41
Pandemic period (March 2020-February 2022, 24 months)	3,028,108	3,691,335	1.22	663,228	17.01	541,167	13.88	408.21
Female								
Wave 1 (March-May 2020, 3 months)	358,172 (340,905 - 375,440)	417,274	1.17 (1.11 - 1.22)	59,102 (41,834 - 76,369)	11.79	51,179	10.21	35.37
Wave 2 (June 2020-September 2020, 4 months)	436,167 (415,802 - 456,532)	502,723	1.15 (1.10 - 1.21)	66,556 (46,191 - 86,921)	9.95	44,496	6.65	39.80
Wave 3 (October 2020-February 2021, 5 months)	597,357 (574,184 - 620,530)	753,137	1.26 (1.21 - 1.31)	155,780 (132,607 - 178,953)	18.61	147,699	17.64	93.04
Spring 2021 (March 2021-June 2021, 4 months)	466,579 (445,875 - 487,283)	484,097	1.04 (0.99 - 1.09)	17,518 (-3,186 - 38,222)	2.61	28,876	4.31	10.45
Delta (July 2021-December 2021, 6 months)	690,600 (664,251 - 716,948)	834,943	1.21 (1.16 - 1.26)	144,343 (117,995 - 170,692)	14.33	105,668	10.49	85.99
Omicron (January 2022-February 2022, 2 months)	240,993 (227,756 - 254,229)	307,621	1.28 (1.21 - 1.35)	66,629 (53,392 - 79,865)	19.83	58,031	17.27	39.66
2020 (March 2020-December 2020, 10 months)	1,152,173	1,366,609	1.19	214,436	12.82	175,648	10.50	128.21
2021 (January 2021-December 2021, 12 months)	1,396,701	1,625,565	1.16	228,864	11.37	202,270	10.05	136.46
2022 (January 2022-February 2022, 2 months)	240,993 (227,756 - 254,229)	307,621	1.28 (1.21 - 1.35)	66,629 (53,395 - 79,862)	19.83	58,031	17.27	39.66
Post Wave 1 (June 2020-February 2022, 21 months)	2,431,695	2,882,521	1.19	450,826	12.81	384,770	10.93	268.97
Pandemic period (March 2020-February 2022, 24 months)	2,789,867	3,299,795	1.18	509,928	12.68	435,949	10.84	304.35
0-17 All Genders								
Wave 1 (March-May 2020, 3 months)	8,728 (8,332 - 9,123)	8,356	0.96 (0.92 - 1.00)	-372 (-767 - 24)	-0.17	0	0.00	-0.51
Wave 2 (June 2020-September 2020, 4 months)	11,501 (11,020 - 11,982)	11,691	1.02 (0.98 - 1.06)	190 (-291 - 671)	0.07	0	0.00	0.26
Wave 3 (October 2020-February 2021, 5 months)	13,670 (13,106 - 14,234)	13,656	1.00 (0.96 - 1.04)	-14 (-578 - 550)	0.00	0	0.00	-0.02
Spring 2021 (March 2021-June 2021, 4 months)	11,370 (10,880 - 11,861)	11,954	1.05 (1.01 - 1.10)	584 (93 - 1,074)	0.20	0	0.00	0.80
Delta (July 2021-December 2021, 6 months)	16,677 (16,030 - 17,324)	18,209	1.09 (1.05 - 1.14)	1,532 (885 - 2,179)	0.35	134	0.03	2.11
Omicron (January 2022-February 2022, 2 months)	5,263 (4,989 - 5,537)	5,432	1.03 (0.98 - 1.09)	169 (-105 - 443)	0.12	118	0.08	0.23
2020 (March 2020-December 2020, 10 months)	28,603	28,295	0.99	-308	-0.04	0	0.00	-0.42
2021 (January 2021-December 2021, 12 months)	33,343	35,571	1.07	2,228	0.26	134	0.02	3.07
2022 (January 2022-February 2022, 2 months)	5,263 (4,989 - 5,537)	5,432	1.03 (0.98 - 1.09)	169 (-103 - 440)	0.12	118	0.08	0.23
Post Wave 1 (June 2020-February 2022, 21 months)	58,482	60,942	1.04	2,460	0.16	252	0.02	3.39
Pandemic period (March 2020-February 2022, 24 months)	67,209	69,298	1.03	2,089	0.12	252	0.01	2.88
18-49 All Genders								
Wave 1 (March-May 2020, 3 months)	60,938 (57,994 - 63,881)	71,069	1.17 (1.11 - 1.23)	10,131 (7,188 - 13,075)	2.44	4,905	1.18	7.33
Wave 2 (June 2020-September 2020, 4 months)	82,369 (78,445 - 86,293)	99,869	1.21 (1.16 - 1.27)	17,500 (13,576 - 21,424)	3.17	5,756	1.04	12.66
Wave 3 (October 2020-February 2021, 5 months)	100,400 (95,282 - 105,519)	125,035	1.25 (1.18 - 1.31)	24,635 (19,516 - 29,753)	3.56	12,677	1.83	17.82
Spring 2021 (March 2021-June 2021, 4 months)	82,523 (78,532 - 86,513)	99,771	1.21 (1.15 - 1.27)	17,248 (13,258 - 21,239)	3.12	5,094	0.92	12.48
Delta (July 2021-December 2021, 6 months)	125,761 (119,323 - 132,199)	174,771	1.39 (1.32 - 1.46)	49,010 (42,572 - 55,448)	5.91	29,970	3.61	35.44
Omicron (January 2022-February 2022, 2 months)	39,902 (37,935 - 41,870)	52,280	1.31 (1.25 - 1.38)	12,378 (10,410 - 14,345)	4.47	7,993	2.89	8.95
2020 (March 2020-December 2020, 10 months)	204,782	245,129	1.20	40,347	2.92	16,940	1.23	29.19
2021 (January 2021-December 2021, 12 months)	247,208	325,386	1.32	78,178	4.71	41,462	2.50	56.54
2022 (January 2022-February 2022, 2 months)	39,902 (37,935 - 41,870)	52,280	1.31 (1.25 - 1.38)	12,378 (10,411 - 14,344)	4.47	7,993	2.89	8.95
Post Wave 1 (June 2020-February 2022, 21 months)	430,955	551,726	1.28	120,771	4.16	61,490	2.12	87.35
Pandemic period (March 2020-February 2022, 24 months)	491,893	622,795	1.27	130,902	3.95	66,395	2.00	94.69
50-64 All Genders								
Wave 1 (March-May 2020, 3 months)	120,495 (116,780 - 124,210)	141,376	1.17 (1.14 - 1.21)	20,881 (17,166 - 24,596)	11.08	16,628	8.82	33.24
Wave 2 (June 2020-September 2020, 4 months)	152,829 (148,448 - 157,211)	181,660	1.19 (1.16 - 1.22)	28,831 (24,449 - 33,212)	11.48	16,897	6.73	45.94
Wave 3 (October 2020-February 2021, 5 months)	199,988 (195,014 - 204,962)	257,037	1.29 (1.25 - 1.32)	57,049 (52,075 - 62,023)	18.19	47,172	15.04	90.97
Spring 2021 (March 2021-June 2021, 4 months)	157,917 (153,475 - 162,360)	180,127	1.14 (1.11 - 1.17)	22,210 (17,767 - 26,652)	8.86	15,217	6.07	35.45
Delta (July 2021-December 2021, 6 months)	235,794 (230,195 - 241,393)	319,916	1.36 (1.33 - 1.39)	84,122 (78,523 - 89,721)	22.41	63,321	16.87	134.43
Omicron (January 2022-February 2022, 2 months)	79,887 (77,004 - 82,769)	107,482	1.35 (1.30 - 1.40)	27,595 (24,713 - 30,478)	22.08	24,562	19.65	44.16
2020 (March 2020-December 2020, 10 months)	393,677	472,481	1.20	78,804	12.55	56,480	9.00	125.55
2021 (January 2021-December 2021, 12 months)	473,346	607,635	1.28	134,289	17.87	102,755	13.67	214.45
2022 (January 2022-February 2022, 2 months)	79,887 (77,004 - 82,769)	107,482	1.35 (1.30 - 1.40)	27,595 (24,717 - 30,474)	22.08	24,562	19.65	44.16
Post Wave 1 (June 2020-February 2022, 21 months)	826,415	1,046,222	1.27	219,807	16.71	167,169	12.71	350.84
Pandemic period (March 2020-February 2022, 24 months)	946,910	1,187,598	1.25	240,688	16.00	183,797	12.22	384.05
65+ All Genders								
Wave 1 (March-May 2020, 3 months)	552,618 (530,797 - 574,439)	650,435	1.18 (1.13 - 1.23)	97,817 (75,996 - 119,638)	58.89	89,076	53.63	176.68
Wave 2 (June 2020-September 2020, 4 months)	670,005 (645,149 - 694,861)	768,610	1.15 (1.11 - 1.19)	98,605 (73,749 - 123,461)	44.23	75,075	33.68	176.92
Wave 3 (October 2020-February 2021, 5 months)	923,222 (895,156 - 951,289)	1,201,153	1.30 (1.26 - 1.34)	277,931 (249,864 - 305,997)	98.93	269,649	95.98	494.66
Spring 2021 (March 2021-June 2021, 4 months)	722,197 (696,823 - 747,571)	737,431	1.02 (0.99 - 1.06)	15,234 (-10,140 - 40,608)	6.73	44,383	19.60	26.90
Delta (July 2021-December 2021, 6 months)	1,068,970 (1,037,701 - 1,100,238)	1,265,664	1.18 (1.15 - 1.22)	196,695 (165,426 - 227,963)	57.29	149,421	43.52	343.71
Omicron (January 2022-February 2022, 2 months)	374,950 (357,170 - 392,731)	488,146	1.30 (1.24 - 1.37)	113,196 (95,415 - 130,976)	98.22	99,068	85.96	196.44
2020 (March 2020-December 2020, 10 months)	1,774,887	2,129,235	1.20	354,349	63.57	310,769	55.76	635.75
2021 (January 2021-December 2021, 12 months)	2,162,125	2,494,058	1.15	331,933	48.64	316,835	46.43	583.67
2022 (January 2022-February 2022, 2 months)	374,950 (357,170 - 392,731)	488,146	1.30 (1.24 - 1.37)	113,196 (95,280 - 131,111)	98.22	99,068	85.96	196.44
Post Wave 1 (June 2020-February 2022, 21 months)	3,759,344	4,461,004	1.19	701,660	59.01	637,596	53.63	1239.30
Pandemic period (March 2020-February 2022, 24 months)	4,311,962	5,111,439	1.19	799,477	59.00	726,672	53.63	1415.98

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Table 2

	Expected deaths (95% CI)	Observed deaths	Ratio of observed to expected deaths (95% CI)	Excess deaths, no (95% CI)	Excess deaths, IR (100,000 person-months)	COVID-19 deaths, no	COVID-19 deaths, IR (100,000 person-months)	Excess Deaths (100,000 person-waves)
Midwest								
Wave 1 (March-May 2020, 3 months)	169,210 (163,064 - 175,357)	190,830	1.13 (1.09 - 1.17)	21,620 (15,473 - 27,766)	10.55	19,287	9.41	31.65
Wave 2 (June 2020-September 2020, 4 months)	208,820 (201,819 - 215,822)	229,752	1.10 (1.06 - 1.14)	20,932 (13,930 - 27,933)	7.66	13,530	4.95	30.65
Wave 3 (October 2020-February 2021, 5 months)	281,819 (273,913 - 289,725)	361,646	1.28 (1.25 - 1.32)	79,827 (71,921 - 87,733)	23.39	77,914	22.83	116.94
Spring 2021 (March 2021-June 2021, 4 months)	220,908 (213,760 - 228,056)	227,088	1.03 (1.00 - 1.06)	6,180 (-968 - 13,328)	2.26	13,744	5.04	9.06
Delta (July 2021-December 2021, 6 months)	330,397 (321,589 - 339,205)	397,703	1.20 (1.17 - 1.24)	67,306 (58,498 - 76,114)	16.44	52,610	12.85	98.64
Omicron (January 2022-February 2022, 2 months)	113,180 (108,171 - 118,189)	147,414	1.30 (1.25 - 1.36)	34,234 (29,225 - 39,243)	25.10	30,605	22.44	50.21
2020 (March 2020-December 2020, 10 months)	547,878 (536,781 - 558,975)	650,335	1.19 (1.16 - 1.21)	102,457 (91,360 - 113,554)	15.00	87,931	12.87	150.02
2021 (January 2021-December 2021, 12 months)	663,277 (650,965 - 675,589)	756,684	1.14 (1.12 - 1.16)	93,407 (81,095 - 105,719)	11.41	89,154	10.89	136.89
2022 (January 2022-February 2022, 2 months)	113,180 (108,134 - 118,227)	147,414	1.30 (1.25 - 1.36)	34,234 (29,187 - 39,280)	25.10	30,605	22.44	50.21
Post Wave 1 (June 2020-February 2022, 21 months)	1,155,125 (1,137,275 - 1,172,975)	1,363,603	1.18 (1.16 - 1.20)	208,478 (190,628 - 226,328)	14.55	188,403	13.15	305.47
Pandemic period (March 2020-February 2022, 24 months)	1,324,335 (1,304,869 - 1,343,802)	1,554,433	1.17 (1.16 - 1.19)	230,098 (210,631 - 249,564)	14.05	207,690	12.68	337.11
Northeast								
Wave 1 (March-May 2020, 3 months)	128,432 (122,948 - 133,915)	203,194	1.58 (1.52 - 1.65)	74,762 (69,279 - 80,246)	44.62	62,271	37.16	133.85
Wave 2 (June 2020-September 2020, 4 months)	156,560 (150,314 - 162,807)	164,027	1.05 (1.01 - 1.09)	7,467 (1,220 - 13,713)	3.35	7,737	3.47	13.39
Wave 3 (October 2020-February 2021, 5 months)	215,604 (208,551 - 222,657)	263,371	1.22 (1.18 - 1.26)	47,767 (40,714 - 54,820)	17.14	51,173	18.37	85.72
Spring 2021 (March 2021-June 2021, 4 months)	166,293 (159,917 - 172,670)	174,572	1.05 (1.01 - 1.09)	8,279 (1,902 - 14,655)	3.72	14,478	6.50	14.87
Delta (July 2021-December 2021, 6 months)	247,093 (239,235 - 254,951)	282,253	1.14 (1.11 - 1.18)	35,160 (27,302 - 43,018)	10.54	25,797	7.73	63.21
Omicron (January 2022-February 2022, 2 months)	86,624 (82,156 - 91,092)	110,851	1.28 (1.22 - 1.35)	24,227 (19,759 - 28,695)	21.80	23,041	20.73	43.60
2020 (March 2020-December 2020, 10 months)	413,705 (403,805 - 423,604)	520,001	1.26 (1.23 - 1.29)	106,296 (96,397 - 116,196)	19.06	94,566	16.95	190.55
2021 (January 2021-December 2021, 12 months)	500,278 (489,294 - 511,261)	567,416	1.13 (1.11 - 1.16)	67,138 (56,155 - 78,122)	10.05	66,890	10.02	120.65
2022 (January 2022-February 2022, 2 months)	86,624 (82,122 - 91,126)	110,851	1.28 (1.22 - 1.35)	24,227 (19,725 - 28,729)	21.80	23,041	20.73	43.60
Post Wave 1 (June 2020-February 2022, 21 months)	872,175 (855,844 - 888,505)	995,074	1.14 (1.12 - 1.16)	122,899 (106,569 - 139,230)	10.51	122,226	10.45	220.74
Pandemic period (March 2020-February 2022, 24 months)	1,000,606 (982,661 - 1,018,552)	1,198,268	1.20 (1.18 - 1.22)	197,662 (179,716 - 215,607)	14.79	184,497	13.80	354.88
South								
Wave 1 (March-May 2020, 3 months)	293,632 (277,758 - 309,506)	317,323	1.08 (1.03 - 1.14)	23,691 (7,817 - 39,565)	6.24	19,699	5.19	18.73
Wave 2 (June 2020-September 2020, 4 months)	366,649 (347,936 - 385,362)	449,941	1.23 (1.17 - 1.29)	83,292 (64,579 - 102,005)	16.43	55,788	11.00	65.72
Wave 3 (October 2020-February 2021, 5 months)	490,516 (469,254 - 511,778)	636,488	1.30 (1.24 - 1.36)	145,972 (124,710 - 167,234)	22.98	125,047	19.69	114.90
Spring 2021 (March 2021-June 2021, 4 months)	387,358 (368,394 - 406,323)	418,757	1.08 (1.03 - 1.14)	31,399 (12,434 - 50,363)	6.16	26,524	5.21	24.66
Delta (July 2021-December 2021, 6 months)	578,259 (554,307 - 602,211)	737,320	1.28 (1.22 - 1.33)	159,061 (135,109 - 183,013)	20.76	117,354	15.32	124.56
Omicron (January 2022-February 2022, 2 months)	197,871 (185,594 - 210,148)	261,911	1.32 (1.25 - 1.41)	64,040 (51,763 - 76,317)	25.02	53,539	20.92	50.05
2020 (March 2020-December 2020, 10 months)	955,700 (924,855 - 986,545)	1,135,033	1.19 (1.15 - 1.23)	179,333 (148,488 - 210,178)	14.15	136,744	10.79	141.49
2021 (January 2021-December 2021, 12 months)	1,160,715 (1,126,436 - 1,194,994)	1,424,796	1.23 (1.19 - 1.26)	264,081 (229,802 - 298,360)	17.26	207,668	13.57	207.14
2022 (January 2022-February 2022, 2 months)	197,871 (185,592 - 210,150)	261,911	1.32 (1.25 - 1.41)	64,040 (51,761 - 76,319)	25.02	53,539	20.92	50.05
Post Wave 1 (June 2020-February 2022, 21 months)	2,020,654 (1,951,554 - 2,089,754)	2,504,417	1.24 (1.20 - 1.28)	483,763 (414,663 - 552,863)	18.09	378,252	14.15	379.96
Pandemic period (March 2020-February 2022, 24 months)	2,314,286 (2,236,338 - 2,392,234)	2,821,740	1.22 (1.18 - 1.26)	507,454 (429,506 - 585,402)	16.62	397,951	13.03	398.89
West								
Wave 1 (March-May 2020, 3 months)	151,504 (146,193 - 156,815)	159,889	1.06 (1.02 - 1.09)	8,385 (3,074 - 13,696)	3.56	9,352	3.97	10.67
Wave 2 (June 2020-September 2020, 4 months)	184,675 (178,625 - 190,725)	218,110	1.18 (1.14 - 1.22)	33,435 (27,385 - 39,485)	10.62	20,673	6.57	42.49
Wave 3 (October 2020-February 2021, 5 months)	249,341 (242,510 - 256,173)	335,376	1.35 (1.31 - 1.38)	86,035 (79,203 - 92,866)	21.84	75,364	19.13	109.19
Spring 2021 (March 2021-June 2021, 4 months)	199,448 (193,272 - 205,624)	208,866	1.05 (1.02 - 1.08)	9,418 (3,242 - 15,594)	2.98	9,948	3.15	11.94
Delta (July 2021-December 2021, 6 months)	291,452 (283,842 - 299,063)	361,284	1.24 (1.21 - 1.27)	69,832 (62,221 - 77,442)	14.73	47,085	9.93	88.37
Omicron (January 2022-February 2022, 2 months)	102,327 (97,999 - 106,655)	133,164	1.30 (1.25 - 1.36)	30,837 (26,509 - 35,165)	19.49	24,556	15.52	38.98
2020 (March 2020-December 2020, 10 months)	484,667 (475,079 - 494,256)	569,771	1.18 (1.15 - 1.20)	85,104 (75,515 - 94,692)	10.82	64,948	8.25	108.15
2021 (January 2021-December 2021, 12 months)	591,753 (581,115 - 602,391)	713,754	1.21 (1.18 - 1.23)	122,001 (111,363 - 132,639)	12.88	97,474	10.29	154.54
2022 (January 2022-February 2022, 2 months)	102,327 (97,967 - 106,688)	133,164	1.30 (1.25 - 1.36)	30,837 (26,476 - 35,197)	19.49	24,556	15.52	38.98
Post Wave 1 (June 2020-February 2022, 21 months)	1,027,243 (1,007,954 - 1,046,533)	1,256,800	1.22 (1.20 - 1.25)	229,557 (210,267 - 248,846)	13.86	177,626	10.72	290.99
Pandemic period (March 2020-February 2022, 24 months)	1,178,747 (1,157,162 - 1,200,332)	1,416,689	1.20 (1.18 - 1.22)	237,942 (216,357 - 259,527)	12.57	186,978	9.88	301.76

Figure 2A




Figure 2B



Figure 2C



Figure 2D




Figure 2E

Cumulative All-Cause Excess Mortality by Census Bureau Division, IR March 1, 2020-February 28, 2022





Figure 2F

Cumulative All-Cause Excess Mortality by Census Bureau Region, IR March 1, 2020-February 28, 2022



It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Table 3

	Expected deaths (95% CI)	Observed deaths	Ratio of observed to expected deaths (95% CI)	Excess deaths, no (95% CI)	Excess deaths, IR (100,000 person-months)	COVID-19 deaths, no	COVID-19 deaths, IR (100,000 person-months)	Excess Deaths (100,000 person-waves)
0-17 Male								
Wave 1 (March-May 2020, 3 months)	5,007 (4,829 - 5,185)	4,997	1.00 (0.96 - 1.03)	-10 (-188 - 168)	-0.01	0	0.00	-0.03
Wave 2 (June 2020-September 2020, 4 months)	6,733 (6,530 - 6,936)	6,897	1.02 (0.99 - 1.06)	164 (-39 - 367)	0.11	0	0.00	0.44
Wave 3 (October 2020-February 2021, 5 months)	7,929 (7,700 - 8,159)	7,959	1.00 (0.98 - 1.03)	30 (-200 - 259)	0.02	0	0.00	0.08
Spring 2021 (March 2021-June 2021, 4 months)	6,570 (6,363 - 6,777)	6,928	1.05 (1.02 - 1.09)	358 (151 - 565)	0.24	0	0.00	0.97
Delta (July 2021-December 2021, 6 months)	9,733 (9,477 - 9,988)	10,586	1.09 (1.06 - 1.12)	853 (598 - 1,109)	0.38	63	0.03	2.30
Omicron (January 2022-February 2022, 2 months)	3,049 (2,904 - 3,194)	3,143	1.03 (0.98 - 1.08)	94 (-51 - 239)	0.13	55	0.07	0.25
2020 (March 2020-December 2020, 10 months)	16,610	16,756	1.01	146	0.04	0	0.00	0.39
2021 (January 2021-December 2021, 12 months)	19,362	20,611	1.06	1,249	0.28	63	0.01	3.37
2022 (January 2022-February 2022, 2 months)	3,049 (2,904 - 3,194)	3,143	1.03 (0.98 - 1.08)	94 (-53 - 240)	0.13	55	0.07	0.25
Post Wave 1 (June 2020-February 2022, 21 months)	34,014	35,513	1.04	1,499	0.19	118	0.02	4.04
Pandemic period (March 2020-February 2022, 24 months)	39,021	40,510	1.04	1,489	0.17	118	0.01	4.01
0-17 Female								
Wave 1 (March-May 2020, 3 months)	3,720 (3,508 - 3,932)	3,359	0.90 (0.85 - 0.96)	-361 (-573 - -149)	-0.34	0	0.00	-1.01
Wave 2 (June 2020-September 2020, 4 months)	4,768 (4,516 - 5,021)	4,794	1.01 (0.95 - 1.06)	26 (-227 - 278)	0.02	0	0.00	0.07
Wave 3 (October 2020-February 2021, 5 months)	5,741 (5,451 - 6,030)	5,697	0.99 (0.94 - 1.05)	-44 (-333 - 246)	-0.02	0	0.00	-0.12
Spring 2021 (March 2021-June 2021, 4 months)	4,800 (4,544 - 5,056)	5,026	1.05 (0.99 - 1.11)	226 (-30 - 482)	0.16	0	0.00	0.64
Delta (July 2021-December 2021, 6 months)	6,944 (6,616 - 7,272)	7,623	1.10 (1.05 - 1.15)	679 (351 - 1,007)	0.32	71	0.03	1.91
Omicron (January 2022-February 2022, 2 months)	2,214 (2,051 - 2,377)	2,289	1.03 (0.96 - 1.12)	75 (-88 - 238)	0.11	63	0.09	0.21
2020 (March 2020-December 2020, 10 months)	11,993	11,539	0.96	-454	-0.13	0	0.00	-1.28
2021 (January 2021-December 2021, 12 months)	13,981	14,960	1.07	979	0.23	71	0.02	2.76
2022 (January 2022-February 2022, 2 months)	2,214 (2,051 - 2,377)	2,289	1.03 (0.96 - 1.12)	75 (-87 - 237)	0.11	63	0.09	0.21
Post Wave 1 (June 2020-February 2022, 21 months)	24,467	25,429	1.04	962	0.13	134	0.02	2.71
Pandemic period (March 2020-February 2022, 24 months)	28,188	28,788	1.02	600	0.07	134	0.02	1.69
18-49 Male								
Wave 1 (March-May 2020, 3 months)	40,662 (38,646 - 42,677)	47,937	1.18 (1.12 - 1.24)	7,275 (5,260 - 9,291)	3.48	3,490	1.67	10.44
Wave 2 (June 2020-September 2020, 4 months)	55,600 (52,822 - 58,378)	67,550	1.21 (1.16 - 1.28)	11,950 (9,172 - 14,728)	4.29	3,748	1.34	17.14
Wave 3 (October 2020-February 2021, 5 months)	66,650 (62,967 - 70,334)	83,136	1.25 (1.18 - 1.32)	16,486 (12,802 - 20,169)	4.73	8,043	2.31	23.64
Spring 2021 (March 2021-June 2021, 4 months)	55,570 (52,770 - 58,370)	67,353	1.21 (1.15 - 1.28)	11,783 (8,983 - 14,583)	4.22	3,079	1.10	16.89
Delta (July 2021-December 2021, 6 months)	84,737 (80,037 - 89,436)	116,308	1.37 (1.30 - 1.45)	31,571 (26,872 - 36,271)	7.54	18,423	4.40	45.25
Omicron (January 2022-February 2022, 2 months)	26,638 (25,322 - 27,953)	34,367	1.29 (1.23 - 1.36)	7,729 (6,414 - 9,045)	5.54	4,780	3.42	11.08
2020 (March 2020-December 2020, 10 months)	137,158	165,026	1.20	27,868	4.00	11,208	1.61	39.98
2021 (January 2021-December 2021, 12 months)	166,061	217,258	1.31	51,197	6.12	25,575	3.06	73.39
2022 (January 2022-February 2022, 2 months)	26,638 (25,322 - 27,953)	34,367	1.29 (1.23 - 1.36)	7,729 (6,416 - 9,042)	5.54	4,780	3.42	11.08
Post Wave 1 (June 2020-February 2022, 21 months)	289,195	368,714	1.27	79,519	5.43	38,073	2.60	114.01
Pandemic period (March 2020-February 2022, 24 months)	329,857	416,651	1.26	86,794	5.19	41,563	2.48	124.46
18-49 Female								
Wave 1 (March-May 2020, 3 months)	20,276 (19,242 - 21,310)	23,132	1.14 (1.09 - 1.20)	2,856 (1,822 - 3,890)	1.39	1,415	0.69	4.17
Wave 2 (June 2020-September 2020, 4 months)	26,769 (25,441 - 28,096)	32,319	1.21 (1.15 - 1.27)	5,550 (4,223 - 6,878)	2.03	2,008	0.73	8.10
Wave 3 (October 2020-February 2021, 5 months)	33,750 (32,078 - 35,422)	41,899	1.24 (1.18 - 1.31)	8,149 (6,477 - 9,821)	2.38	4,634	1.35	11.89
Spring 2021 (March 2021-June 2021, 4 months)	26,953 (25,604 - 28,302)	32,418	1.20 (1.15 - 1.27)	5,465 (4,116 - 6,814)	1.99	2,015	0.74	7.98
Delta (July 2021-December 2021, 6 months)	41,024 (38,976 - 43,072)	58,463	1.43 (1.36 - 1.50)	17,439 (15,391 - 19,487)	4.24	11,547	2.81	25.45
Omicron (January 2022-February 2022, 2 months)	13,264 (12,540 - 13,988)	17,913	1.35 (1.28 - 1.43)	4,649 (3,925 - 5,373)	3.39	3,213	2.34	6.78
2020 (March 2020-December 2020, 10 months)	67,625	80,103	1.18	12,478	1.82	5,732	0.84	18.21
2021 (January 2021-December 2021, 12 months)	81,147	108,128	1.33	26,981	3.28	15,887	1.93	39.38
2022 (January 2022-February 2022, 2 months)	13,264 (12,540 - 13,988)	17,913	1.35 (1.28 - 1.43)	4,649 (3,926 - 5,371)	3.39	3,213	2.34	6.78
Post Wave 1 (June 2020-February 2022, 21 months)	141,760	183,012	1.29	41,252	2.87	23,417	1.63	60.21
Pandemic period (March 2020-February 2022, 24 months)	162,036	206,144	1.27	44,108	2.68	24,832	1.51	64.38
50-64 Male								
Wave 1 (March-May 2020, 3 months)	72,990 (70,958 - 75,022)	87,982	1.21 (1.17 - 1.24)	14,992 (12,960 - 17,024)	16.36	11,145	12.16	49.08
Wave 2 (June 2020-September 2020, 4 months)	93,570 (90,982 - 95,759)	112,236	1.20 (1.17 - 1.23)	18,666 (16,477 - 21,254)	15.46	10,729	8.79	61.83
Wave 3 (October 2020-February 2021, 5 months)	121,228 (118,534 - 123,921)	159,443	1.32 (1.29 - 1.35)	38,215 (35,522 - 40,909)	25.08	30,097	19.75	125.41
Spring 2021 (March 2021-June 2021, 4 months)	95,695 (93,283 - 98,107)	111,051	1.16 (1.13 - 1.19)	15,356 (12,944 - 17,768)	12.62	9,246	7.60	50.47
Delta (July 2021-December 2021, 6 months)	143,092 (140,066 - 146,117)	196,065	1.37 (1.34 - 1.40)	52,973 (49,948 - 55,999)	29.07	38,553	21.16	174.41
Omicron (January 2022-February 2022, 2 months)	48,275 (46,659 - 49,891)	65,725	1.36 (1.32 - 1.41)	17,450 (15,834 - 19,066)	26.78	14,787	24.39	57.55
2020 (March 2020-December 2020, 10 months)	239,372	292,573	1.22	53,201	17.44	36,568	11.98	174.35
2021 (January 2021-December 2021, 12 months)	287,003	374,204	1.30	87,201	23.90	63,202	17.32	286.80
2022 (January 2022-February 2022, 2 months)	48,275 (46,659 - 49,891)	65,725	1.36 (1.32 - 1.41)	17,450 (15,848 - 19,052)	26.78	14,787	24.39	57.55
Post Wave 1 (June 2020-February 2022, 21 months)	501,660	644,520	1.28	142,860	22.36	103,412	16.18	469.50
Pandemic period (March 2020-February 2022, 24 months)	574,650	732,502	1.27	157,852	21.61	114,557	15.68	518.52
50-64 Female								
Wave 1 (March-May 2020, 3 months)	47,505 (45,703 - 49,306)	53,394	1.12 (1.08 - 1.17)	5,889 (4,088 - 7,691)	6.08	5,483	5.66	18.25
Wave 2 (June 2020-September 2020, 4 months)	59,459 (57,339 - 61,579)	69,424	1.17 (1.13 - 1.21)	9,965 (7,845 - 12,085)	7.72	6,168	4.78	30.90
Wave 3 (October 2020-February 2021, 5 months)	78,760 (76,346 - 81,175)	97,594	1.24 (1.20 - 1.28)	18,834 (16,419 - 21,248)	11.68	17,075	10.59	58.42
Spring 2021 (March 2021-June 2021, 4 months)	62,223 (60,064 - 64,381)	69,076	1.11 (1.07 - 1.15)	6,853 (4,695 - 9,012)	5.32	5,971	4.63	21.26
Delta (July 2021-December 2021, 6 months)	92,702 (89,949 - 95,456)	123,851	1.34 (1.30 - 1.38)	31,149 (28,395 - 33,902)	16.12	24,768	12.82	96.73
Omicron (January 2022-February 2022, 2 months)	31,611 (30,238 - 32,985)	41,757	1.32 (1.27 - 1.38)	10,146 (8,772 - 11,519)	15.77	9,775	15.19	31.54
2020 (March 2020-December 2020, 10 months)	154,306	179,908	1.17	25,602	7.94	19,912	6.17	79.38
2021 (January 2021-December 2021, 12 months)	186,343	233,431	1.25	47,088	12.18	39,553	10.23	146.16
2022 (January 2022-February 2022, 2 months)	31,611 (30,238 - 32,985)	41,757	1.32 (1.27 - 1.38)	10,146 (8,775 - 11,516)	15.77	9,775	15.19	31.54
Post Wave 1 (June 2020-February 2022, 21 months)	324,756	401,702	1.24	76,946	11.37	63,757	9.42	238.80
Pandemic period (March 2020-February 2022, 24 months)	372,260	455,096	1.22	82,836	10.71	69,240	8.95	257.03
65+ Male								
Wave 1 (March-May 2020, 3 months)	265,947 (255,256 - 276,638)	313,046	1.18 (1.13 - 1.23)	47,099 (36,408 - 57,790)	63.61	44,795	60.50	190.82
Wave 2 (June 2020-September 2020, 4 months)	324,834 (312,230 - 337,438)	372,424	1.15 (1.10 - 1.19)	47,590 (34,986 - 60,194)	47.87	38,755	38.98	191.47
Wave 3 (October 2020-February 2021, 5 months)	444,116 (429,808 - 458,425)	593,206	1.34 (1.29 - 1.38)	149,090 (134,781 - 163,398)	118.98	143,659	114.64	594.89
Spring 2021 (March 2021-June 2021, 4 months)	349,594 (336,819 - 362,369)	359,854	1.03 (0.99 - 1.07)	10,260 (-2,515 - 23,035)	10.16	23,493	23.26	40.63
Delta (July 2021-December 2021, 6 months)	519,041 (502,938 - 535,144)	620,658	1.20 (1.16 - 1.23)	101,617 (85,514 - 117,720)	66.33	80,139	52.31	397.97
Omicron (January 2022-February 2022, 2 months)	181,048 (172,746 - 189,350)	242,484	1.34 (1.28 - 1.40)	61,436 (53,134 - 69,738)	119.46	54,088	105.17	238.93
2020 (March 2020-December 2020, 10 months)	856,637	1,034,176	1.21	177,539	71.43	160,765	64.68	714.26
2021 (January 2021-December 2021, 12 months)	1,046,895	1,225,012	1.17	178,117	58.51	170,076	55.87	702.12
2022 (January 2022-February 2022, 2 months)	181,048 (172,746 - 189,350)	242,484	1.34 (1.28 - 1.40)	61,436 (53,151 - 69,721)	119.46	54,088	105.17	238.93
Post Wave 1 (June 2020-February 2022, 21 months)	1,818,633	2,188,626	1.20	369,993	69.76	340,134	64.13	1465.00
Pandemic period (March 2020-February 2022, 24 months)	2,084,579	2,501,672	1.20	417,093	69.01	384,929	63.69	1656.19
65+ Female								
Wave 1 (March-May 2020, 3 months)	286,671 (272,173 - 301,169)	337,389	1.18 (1.12 - 1.24)	50,718 (36,220 - 65,216)	55.10	44,281	48.11	165.30
Wave 2 (June 2020-September 2020, 4 months)	345,174 (328,085 - 362,257)	396,186	1.15 (1.09 - 1.21)	51,015 (33,929 - 68,101)	41.30	36,320	29.41	165.21
Wave 3 (October 2020-February 2021, 5 months)	479,106 (459,646 - 498,566)	607,947	1.27 (1.22 - 1.32)	128,841 (109,381 - 148,301)	82.79	125,990	80.96	413.95
Spring 2021 (March 2021-June 2021, 4 months)	377							

Figure 3A




Figure 3B





Figure 3C

Excess Deaths Cumulative IR: Gender



It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Table 4

	Expected deaths (95% CI)	Observed deaths	Ratio of observed to expected deaths (95% CI)	Excess deaths, no (95% CI)	Excess deaths, IR (100,000 person-months)	COVID-19 deaths, no	COVID-19 deaths, IR (100,000 person-months)	Excess Deaths (100,000 person-waves)
American Indian or Alaska Native								
Wave 1 (March-May 2020, 3 months)	4,693 (3,921 - 5,464)	5,601	1.19 (1.03 - 1.43)	908 (137 - 1,680)	12.45	761	10.43	37.34
Wave 2 (June 2020-September 2020, 4 months)	5,869 (5,158 - 6,579)	8,329	1.40 (1.25 - 1.60)	2,360 (1,650 - 3,071)	24.26	1,394	14.33	97.05
Wave 3 (October 2020-February 2021, 5 months)	7,923 (7,182 - 8,664)	12,708	1.60 (1.47 - 1.77)	4,785 (4,044 - 5,526)	39.34	4,119	33.87	196.71
Spring 2021 (March 2021-June 2021, 4 months)	6,214 (5,561 - 6,868)	7,548	1.21 (1.10 - 1.36)	1,334 (680 - 1,987)	13.70	534	5.49	54.81
Delta (July 2021-December 2021, 6 months)	9,180 (8,490 - 9,869)	14,359	1.56 (1.45 - 1.69)	5,179 (4,490 - 5,869)	35.46	2,799	19.16	212.76
Omicron (January 2022-February 2022, 2 months)	3,298 (2,651 - 3,945)	4,816	1.46 (1.22 - 1.82)	1,518 (871 - 2,165)	31.19	1,243	25.53	62.37
2020 (March 2020-December 2020, 10 months)	15,225	21,474	1.41	6,249	25.69	4,596	18.90	256.92
2021 (January 2021-December 2021, 12 months)	18,654	26,971	1.45	8,317	28.48	5,011	17.16	341.77
2022 (January 2022-February 2022, 2 months)	3,298 (2,651 - 3,945)	4,816	1.46 (1.22 - 1.82)	1,518 (871 - 2,165)	31.19	1,243	25.53	62.37
Post Wave 1 (June 2020-February 2022, 21 months)	32,484	47,660	1.47	15,176	10.089	10,089	19.75	623.72
Pandemic period (March 2020-February 2022, 24 months)	37,177	53,261	1.43	16,084	27.54	10,850	18.58	661.07
Asian								
Wave 1 (March-May 2020, 3 months)	18,754 (15,345 - 22,164)	25,816	1.38 (1.16 - 1.68)	7,062 (3,652 - 10,471)	12.20	5,553	9.60	36.61
Wave 2 (June 2020-September 2020, 4 months)	23,234 (20,130 - 26,339)	27,156	1.17 (1.03 - 1.35)	3,922 (817 - 7,026)	5.05	2,998	3.86	20.22
Wave 3 (October 2020-February 2021, 5 months)	32,607 (29,343 - 35,870)	45,966	1.41 (1.28 - 1.57)	13,359 (10,096 - 16,623)	13.69	12,012	12.31	68.47
Spring 2021 (March 2021-June 2021, 4 months)	25,279 (22,452 - 28,106)	27,711	1.10 (0.99 - 1.23)	2,432 (-395 - 5,259)	3.10	2,416	3.08	12.40
Delta (July 2021-December 2021, 6 months)	37,602 (34,548 - 40,656)	43,727	1.16 (1.08 - 1.27)	6,125 (3,071 - 9,179)	5.17	4,177	3.53	31.01
Omicron (January 2022-February 2022, 2 months)	13,953 (11,136 - 16,770)	17,771	1.27 (1.06 - 1.60)	3,818 (1,001 - 6,635)	9.62	3,480	8.77	19.23
2020 (March 2020-December 2020, 10 months)	61,161	77,946	1.27	16,785	8.66	13,496	6.96	86.56
2021 (January 2021-December 2021, 12 months)	76,315	92,430	1.21	16,115	6.83	13,657	5.79	81.92
2022 (January 2022-February 2022, 2 months)	13,953 (11,136 - 16,770)	17,771	1.27 (1.06 - 1.60)	3,818 (1,001 - 6,635)	9.62	3,480	8.77	19.23
Post Wave 1 (June 2020-February 2022, 21 months)	132,675	162,331	1.22	29,656	7.20	25,083	6.09	151.22
Pandemic period (March 2020-February 2022, 24 months)	151,429	188,147	1.24	36,718	7.82	30,636	6.52	187.62
Black or African American								
Wave 1 (March-May 2020, 3 months)	90,201 (76,393 - 104,010)	127,266	1.41 (1.22 - 1.67)	37,065 (23,256 - 50,873)	29.86	26,382	21.26	89.59
Wave 2 (June 2020-September 2020, 4 months)	113,447 (100,783 - 126,111)	143,606	1.27 (1.14 - 1.42)	30,159 (17,495 - 42,823)	18.21	17,403	10.51	72.83
Wave 3 (October 2020-February 2021, 5 months)	153,311 (140,082 - 166,541)	201,389	1.31 (1.21 - 1.44)	48,078 (34,848 - 61,307)	23.19	36,490	17.60	115.97
Spring 2021 (March 2021-June 2021, 4 months)	119,686 (108,181 - 131,192)	136,887	1.14 (1.04 - 1.27)	17,201 (5,695 - 28,706)	10.36	11,674	7.03	41.44
Delta (July 2021-December 2021, 6 months)	179,373 (166,913 - 191,833)	227,843	1.27 (1.19 - 1.37)	48,470 (36,010 - 60,930)	19.43	31,376	12.58	116.60
Omicron (January 2022-February 2022, 2 months)	63,379 (52,078 - 74,680)	84,084	1.33 (1.13 - 1.61)	20,705 (9,404 - 32,006)	24.88	17,811	21.40	49.76
2020 (March 2020-December 2020, 10 months)	294,843	387,236	1.31	92,393	22.31	61,395	14.83	223.12
2021 (January 2021-December 2021, 12 months)	361,176	449,755	1.25	88,579	17.77	61,930	12.43	213.28
2022 (January 2022-February 2022, 2 months)	63,379 (52,078 - 74,680)	84,084	1.33 (1.13 - 1.33)	20,705 (20,705 - 20,705)	24.88	17,811	21.40	49.76
Post Wave 1 (June 2020-February 2022, 21 months)	629,197	793,809	1.26	164,612	18.89	114,754	13.17	396.62
Pandemic period (March 2020-February 2022, 24 months)	719,398	921,075	1.28	201,677	20.25	141,136	14.17	486.12
Hispanic								
Wave 1 (March-May 2020, 3 months)	56,266 (47,240 - 65,292)	77,452	1.38 (1.19 - 1.64)	21,186 (12,160 - 30,212)	11.54	18,328	9.99	34.63
Wave 2 (June 2020-September 2020, 4 months)	71,340 (63,025 - 79,654)	103,665	1.45 (1.30 - 1.64)	32,325 (23,911 - 40,540)	13.13	25,557	10.41	52.51
Wave 3 (October 2020-February 2021, 5 months)	95,887 (87,368 - 104,405)	156,635	1.63 (1.50 - 1.79)	60,748 (52,330 - 69,267)	19.72	55,117	17.89	96.60
Spring 2021 (March 2021-June 2021, 4 months)	75,778 (68,303 - 83,253)	89,027	1.17 (1.07 - 1.30)	13,259 (5,784 - 20,734)	5.36	10,877	4.40	21.44
Delta (July 2021-December 2021, 6 months)	113,339 (105,260 - 121,417)	156,038	1.38 (1.29 - 1.48)	42,699 (34,621 - 50,778)	11.45	32,090	8.61	68.73
Omicron (January 2022-February 2022, 2 months)	40,562 (33,240 - 47,884)	58,098	1.43 (1.21 - 1.75)	17,536 (10,214 - 24,858)	14.06	15,933	12.78	28.13
2020 (March 2020-December 2020, 10 months)	184,176	267,065	1.45	82,889	13.51	69,060	11.25	135.06
2021 (January 2021-December 2021, 12 months)	228,433	315,662	1.38	87,229	11.73	72,909	9.81	140.79
2022 (January 2022-February 2022, 2 months)	40,562 (33,240 - 47,884)	58,098	1.43 (1.21 - 1.75)	17,536 (10,214 - 24,858)	14.06	15,933	12.78	28.13
Post Wave 1 (June 2020-February 2022, 21 months)	396,905	563,373	1.42	166,468	12.82	139,574	10.75	269.25
Pandemic period (March 2020-February 2022, 24 months)	453,171	640,825	1.41	187,654	12.66	157,902	10.66	303.91
More than one race								
Wave 1 (March-May 2020, 3 months)	3,437 (2,061 - 4,812)	3,806	1.11 (0.79 - 1.85)	369 (-1,006 - 1,745)	1.64	261	1.16	4.92
Wave 2 (June 2020-September 2020, 4 months)	4,645 (3,451 - 5,839)	5,058	1.09 (0.87 - 1.47)	413 (-781 - 1,607)	1.36	324	1.07	5.44
Wave 3 (October 2020-February 2021, 5 months)	5,905 (4,652 - 7,158)	7,415	1.26 (1.04 - 1.59)	1,510 (257 - 2,763)	3.94	1,122	2.93	19.71
Spring 2021 (March 2021-June 2021, 4 months)	4,887 (3,747 - 6,028)	5,264	1.08 (0.87 - 1.41)	377 (-764 - 1,517)	1.22	257	0.83	4.87
Delta (July 2021-December 2021, 6 months)	7,414 (6,212 - 8,615)	8,971	1.21 (1.04 - 1.44)	1,557 (356 - 2,759)	3.32	1,151	2.45	19.91
Omicron (January 2022-February 2022, 2 months)	2,516 (1,427 - 3,606)	3,066	1.22 (0.85 - 2.15)	550 (-540 - 1,639)	3.48	594	3.76	6.96
2020 (March 2020-December 2020, 10 months)	11,612	13,198	1.14	1,586	2.09	1,128	1.49	20.93
2021 (January 2021-December 2021, 12 months)	14,676	17,316	1.18	2,640	2.83	1,987	2.13	33.97
2022 (January 2022-February 2022, 2 months)	2,516 (1,427 - 3,606)	3,066	1.22 (0.85 - 2.15)	550 (-540 - 1,639)	3.48	594	3.76	6.96
Post Wave 1 (June 2020-February 2022, 21 months)	25,368	29,774	1.17	4,406	2.71	3,448	2.12	57.01
Pandemic period (March 2020-February 2022, 24 months)	28,804	33,580	1.17	4,776	2.58	3,709	2.01	62.01
Native Hawaiian or Other Pacific Islander								
Wave 1 (March-May 2020, 3 months)	962 (685 - 1,240)	1,014	1.05 (0.82 - 1.48)	52 (-226 - 329)	2.83	79	4.31	8.49
Wave 2 (June 2020-September 2020, 4 months)	1,237 (986 - 1,487)	1,481	1.20 (1.00 - 1.50)	244 (-6 - 495)	9.93	298	12.12	39.72
Wave 3 (October 2020-February 2021, 5 months)	1,634 (1,382 - 1,886)	2,203	1.35 (1.17 - 1.59)	569 (317 - 821)	18.40	589	19.03	91.98
Spring 2021 (March 2021-June 2021, 4 months)	1,325 (1,089 - 1,561)	1,496	1.13 (0.96 - 1.37)	171 (-65 - 407)	6.85	139	5.58	27.42
Delta (July 2021-December 2021, 6 months)	2,006 (1,768 - 2,244)	2,822	1.41 (1.26 - 1.60)	816 (578 - 1,054)	21.68	755	20.06	130.06
Omicron (January 2022-February 2022, 2 months)	687 (449 - 925)	1,002	1.46 (1.08 - 2.23)	315 (77 - 553)	24.95	240	19.03	49.90
2020 (March 2020-December 2020, 10 months)	3,168	3,793	1.20	625	10.17	685	11.15	101.70
2021 (January 2021-December 2021, 12 months)	3,996	5,223	1.31	1,227	16.37	1,175	15.68	196.43
2022 (January 2022-February 2022, 2 months)	687 (449 - 925)	1,002	1.46 (1.08 - 2.23)	315 (77 - 553)	24.95	240	19.03	49.90
Post Wave 1 (June 2020-February 2022, 21 months)	6,889	9,004	1.31	2,115	16.18	2,021	15.46	339.78
Pandemic period (March 2020-February 2022, 24 months)	7,851	10,018	1.28	2,167	14.54	2,100	14.09	348.93
White								
Wave 1 (March-May 2020, 3 months)	564,686 (436,634 - 692,739)	626,994	1.11 (0.91 - 1.44)	62,308 (-65,745 - 190,360)	10.55	58,177	9.85	31.65
Wave 2 (June 2020-September 2020, 4 months)	695,108 (581,076 - 809,139)	770,420	1.11 (0.95 - 1.33)	75,312 (-38,719 - 189,344)	9.58	49,709	6.32	38.30
Wave 3 (October 2020-February 2021, 5 months)	946,915 (823,812 - 1,070,019)	1,167,257	1.23 (1.09 - 1.42)	220,342 (97,238 - 343,445)	22.44	219,710	22.38	112.22
Spring 2021 (March 2021-June 2021, 4 months)	739,144 (632,406 - 845,882)	759,279	1.03 (0.90 - 1.20)	20,135 (-86,603 - 126,873)	2.57	38,822	4.95	10.27
Delta (July 2021-December 2021, 6 months)	1,087,270 (971,541 - 1,202,999)	1,321,643	1.22 (1.10 - 1.36)	234,373 (118,644 - 350,102)	19.95	170,668	14.53	119.70
Omicron (January 2022-February 2022, 2 months)	388,053 (284,546 - 491,561)	484,324	1.25 (0.99 - 1.70)	96,271 (-7,237 - 199,778)	24.62	93,297	23.86	49.25
2020 (March 2020-December 2020, 10 months)	1,820,855	2,096,818	1.15	275,963	14.03	232,531	11.83	140.34
2021 (January 2021-December 2021, 12 months)	2,212,268	2,548,775	1.15	336,507	14.31	304,555	12.95	171.72
2022 (January 2022-February 2022, 2 months)	388,053 (284,546 - 491,561)	484,324	1.25 (0.99 - 1.70)	96,271 (-7,237 - 199,778)	24.62	93,297	23.86	49.25
Post Wave 1 (June 2020-February 2022, 21 months)	3,856,491	4,502,923	1.17	646,432	15.70	572,206	13.89	329.63
Pandemic period (March 2020-February 2022, 24 months)	4,421,177	5,129,917	1.16	708,740	15.05	630,383	13.39	361.23

Figure 4A

Excess Deaths IR: Race




Figure 4B




Figure 4C

