Epidemiology of leprosy identified through active case detection in six districts of Nepal

Ram Kumar Mahato[1*], Uttam Ghimire[1], Madhav Lamsal[1], Bijay Bajracharya[2], Mukesh

Poudel[1], Prashanna Naapit[3], Krishna Lama[4], Gokarna Dahal[1], David TS Hayman[5], Chuman

Lal Das[1], Krishna Prasad Paudel[1*]

Affiliations:

[1] Epidemiology and Disease Control Division, Department of Health Services, Ministry of Health

and Population, Kathmandu, Nepal

[2] Epidemiology and Disease Control Division- Malaria Program Management Unit- SCI-GF,

Nepal

[3] Nepal Academy of Medical Sciences, Kathmandu, Nepal

[4] Lalgadh Leprosy Hospital & Service center, Nepal Leprosy Trust, Lalgadh, Nepal

[5] Molecular Epidemiology and Public Health Laboratory, Infectious Disease Research Centre,

Hopkirk Research Institute, Massey University, Palmerston North, New Zealand

[*]**Corresponding author**

kpkalyan@gmail.com

mahatoram811@gmail.com

Abstract

Background

Nepal has achieved and sustained elimination of leprosy as a public health problem since 2009, but 17 districts and 3 provinces have yet to eliminate the disease. Pediatric cases and grade-2 disabilities (G2D) indicate recent transmission and late diagnosis respectively, which necessitate active and early case detection. This operational research was performed to identify approaches best suited for early case detection, determine community-based leprosy epidemiology, and identify hidden leprosy cases early and respond with prompt treatment.

Methods

Active case detection was performed by: house-to-house visits among vulnerable populations (*n*=26,469), contact examination and tracing (*n*=7,608) and screening prison populations (*n*=4,428) in Siraha, Bardiya, Rautahat, Banke, Lalitpur and Kathmandu districts of Nepal.

Results

New case detection rates were highest for contact tracing (250), followed by house-to-house visits (102) and prison screening (45) per 100,000 population screened. However, cost per case identified was cheapest for house-to-house visits (Nepalese rupee (NPR) 76,500/case), then contact tracing (NPR90,286/case) and prison screening (NPR298,300/case). House-to-house and contact tracing case paucibacillary/multibacillary (PB:MB) ratios were 59:41 and 68:32; female/male ratios 63:37 and 57:43; pediatric cases 11% in both approaches; and G2D 11% and 5% respectively. Developing leprosy was similar among household and neighbor contacts (Odds ratios (OR)=1.4, 95% confidence interval (CI), 0.24-5.85) and for contacts of MB versus PB cases (*OR*=0.7, 0.26-2.0). Attack rates were similar among household contacts of MB cases (0.32%, 0.07-0.94%) and

PB cases (0.13%, 0.03-0.73) and neighbor contacts of MB cases (0.23%, 0.1-0.46) and PB cases

(0.48%, 0.19-0.98). BCG vaccination with scar presence had a significant protective effect against

leprosy (*OR*=0.42, 0.22-0.81).

Conclusions

The most effective case identification approach here is contact tracing, followed by house-to-house

visits in vulnerable populations and screening in prisons. The findings suggest hidden cases, recent

transmission, and late diagnosis in the community exist and highlight the importance of early case

detection.

Introduction

Leprosy is a contagious, but low pathogenic and chronic infectious disease caused by *Mycobacterium leprae*. *Mycobacterium leprae* mainly affects peripheral nerves and skin, which results in progressive physical, psychological and social disability in some cases (1,2). Disability affects social and working life of the people; social stigma is a significant consequence of leprosy. The first and prime objective of leprosy control programs is to focus on early case detection so treatment can begin as early as possible after symptoms appear and disability prevented (3). In 2019, 202,256 new leprosy cases were reported from 118 countries (26.0 per million population), in which 16 countries reported more than 1,000 new cases; the World Health Organization (WHO) South-East Asia Region (SEAR) accounted for 71% of the cases. The new cases (0.23/10000 population) and child (7.4%), grade 2 disability (G2D, 5.4%) and female (39%) case proportions in 2019 indicate ongoing transmission, late diagnosis and under reported cases in females (4,5).

Nepal has maintained and sustained elimination as a public health problem level at country level from 2009. However, in 2018, Nepal reported more than 3,200 cases with a registered prevalence of 0.99/10,000 population. Seventeen districts and 3 provinces have registered leprosy prevalence >1/10,000 population, where Madhesh Province (40%) and Lumbini Province (18%) accounted for most cases. The proportion of child, female and G2D cases in Nepal in 2018 were 7.92, 42% and 4.75 (6). The pediatric cases indicate recent transmission, lower female proportions indicate underreporting, and G2D suggests late diagnosis, all threatening the elimination status that Nepal achieved in 2009.

The WHO Global Leprosy Strategy 2016–2020 launched in 2016 envisioned accelerating action towards a leprosy-free world. The indicators for this vision were zero children diagnosed with leprosy and visible deformities, the rate of newly diagnosed leprosy patients with visible

deformities <1 per million, and no countries with legislation allowing discrimination on basis of

leprosy. The promotion of voluntary self-reporting is crucial to case detection and to achieve the

desired target. The Global Leprosy Strategy 2016–2020 recommends targeting high-risk and

vulnerable groups with increasing active case detection (7). Active case detection is a more

effective strategy which enables early diagnosis and treatment and prevents disability and

potentially the spread of the disease (8,9).

House-to-house visits of high-risk and vulnerable populations, such as Dalit, Mushhar and other

marginalized communities, could identify hidden cases that might transmit the disease in favorable

conditions. Contact tracing is a recognized form of undertaking active case detection in a group

which is significantly more likely to have leprosy than the general population in high- and low-

endemic disease burden countries. Among different types of contacts, household contacts

reportedly have 3.5 times greater likelihood of having leprosy than social contacts and almost

double that of neighbors; however, even social contacts are 2.5 to 3 times more likely to have

leprosy than the general population (10). Studies suggest the most susceptible populations include

family contacts of multibacillary (MB) cases, followed by neighboring contacts and the contacts

of paucibacillary (PB) cases (11). However, overcrowding within prisons also make the

environment conducive for disease spread. Poor diet, lack of hygiene and physical inactivity are

enabling factors and hence prisoners are more at risk of transmission compared to general

population (12,13). Finally, BCG vaccination, the attenuated Bacillus Calmette-Guérin strain of

the related *Mycobacterium bovis* bacteria, also reduces leprosy transmission (14,15).

Here, we use three active case detection methods: 1) house-to-house visits of high-risk and

vulnerable populations in Nepali districts with leprosy public health problems; 2) house-to-house

visits and examination of contacts of leprosy cases identified between 2-5 years ago; and 3)

examination of prisoners to identify early cases in a cross-sectional study. The study also assessed

the cost effectiveness of methods to identify active cases and measures of association to highlight

key epidemiological features of leprosy in risk areas relevant for control.

Methods

The general study objective is to determine the epidemiology of leprosy with its protective and

risk factors through active and early case detection approach using three active case detection

approaches.

Approach 1. House-to-house visits in communities with high-risk groups and vulnerable

populations, such as marginalized habitants of Dalit, Mushhar, and Chamar groups, were

undertaken in Rautahat District of Madhesh Province and Banke District of Lumbini Province. In

Rautahat, four rural municipalities (Palika): Dewahi Gonahi, Rajpur, Ishnath and Rajdevi were

selected in close coordination with district health authorities. These municipalities were considered

to have inhabitants from more vulnerable populations. From the four municipalities, 24 sites

(wards) covered by 24 health facilities were selected. The same process was followed in Banke,

where 27 sites (wards) covered by 27 health facilities from four municipalities, Baijnath,

Narainapur, Janaki and Nepalgunj, were selected. A total of 60 to 100 households with inhabitants

of marginalized people living in overcrowded houses made of soil or mud, which favored leprosy

transmission, were used for the census. Trained local health workers and local female community

health volunteers (FCHV) visited the selected sites and performed house-to-house visits,

examining all the members present in the household for any signs of leprosy. In total 13,420 and

13,049 individuals were examined respectively in Rautahat and Banke (Table S1). Local trained

health workers examined male and FCHV examined female individuals present in the household.

Simultaneously, demographic and epidemiological variables were collected by the trained local

health worker.

Members of the households were informed 2 days before the survey to be present at their own

household at the time of the survey via the local FCHV. All suspected cases identified by local

health workers and FCHV were invited to health facilities and cases confirmed by a dermatologist.

After diagnosis confirmation, leprosy cases were treated as per the national protocol.

Approach 2. Household and neighboring contacts of previously identified confirmed leprosy cases

in the previous 2-5 years were examined in Siraha district of Madhesh Province and Bardiya of

Lumbini Province by trained local health workers. The cases diagnosed between recent 2-5 years

in the respective districts were selected randomly in planning meetings conducted before the

implementation of field work. Local trained health workers and FCHVs examined 106 and 177

confirmed leprosy case contacts correspondingly in Siraha and Bardiya. A total of 7608 contacts

were screened during the case-contact survey (Table 1).

Table 1. Leprosy cases and their contacts screened during a case-contact survey

District	Total index cases	Clinical disease	Index cases	Household contacts*	Neighboring contacts**	Total screened population
Siraha	106	MB	53	327	1353	3170
		PB	53	372	118	
Bardiya	177	MB	107	599	2102	4438
		PB	70	2102	1352	
Total						7608

*Household contacts comprised all members >2 years old residing in the index case household.

**Neighboring contacts comprised all individuals >2 years old residing in the nearest 4-6

neighboring houses of index case.

Approach 3. Siraha (*n=449*), Rautahat (*n=360*), Banke (*n=826*), Bardiya (*n=319*), Lalitpur

(*n=251*) and Kathamandu (*n=2223*) prisons were used as screening sites for active case detection

using convenience sampling among 4428 prisoners to assess the transmission status of leprosy in

prisons.

The whole study was carried out between October 2020 and December 2021.

Informed consent

In all approaches, participants were requested to give verbal informed consent. As this study was

part of regular surveillance of epidemiology and disease control division (EDCD), written

informed consent was not taken. Approval for data publication was obtained from EDCD and

exemption from ethical review (347/2022) was obtained from ethical board of Nepal Health

Research council.

Statistical analysis

Data collected on paper-based questionnaires, developed by the Leprosy Control and Disability

Management Section (LCDMS)/EDCD, were entered in Excel® spreadsheets. Consistency was

checked and data analysis done in IBM SPSS statistics 22 and R version 4.2.0. New case detection

rate, attack rate (AR) with respect to different demographic variables and types of leprosy cases

were calculated with 95% confidence intervals (CI) when appropriate, e.g., using binomial models

for the attack rates. Chi-squared tests (χ^2) and odds ratio (OR) with 95% CIs were calculated for

associations between attack rates with respect to different demographic variables and between

BCG scar presence and leprosy. To adjust for screened population at risk and index cases present

in the case-contact survey, we use simple Poisson regression with an offset for index cases per

population present to assess risk (16), where:

$$\log{(E(new\ cases_i))} = \beta_0 + \beta_{district}x_i + \log\left(\frac{index\ cases_i}{population\ screened_i}\right)$$

Finally, for discussion, we converted costs from national currencies to US dollars for comparison.

We used the date in publications and adjust to 25 December 2021 rates using Google's default

currency convertor provided by Morningstar.

Results

Comparison of different approaches of active case detection

New leprosy cases were identified during house-to-house visits (n=27), contact tracing (n=19) and

prison screening (n=2) from a total of 38,505 screened people (Table 4). New case detection rates

were highest in contact tracing (250 per 100,000 population), followed by house-to-house visits

(102 per 100,000) and prison screening (45 per 100,000). However, house-to-house visits were the

cheapest cost per case identified at Nepalese rupee (NPR) 76,500/case, followed by contact tracing

(NPR 90,286/case) and prison screening (NPR 298,300/case).

Just two MB cases were discovered in adult male prisoners. House-to-house and contact tracing

case PB:MB ratios were 59:41 and 68:32; female/male ratios 63:37 and 57:43; pediatric cases 11%

in both approaches; and G2D 11% and 5% respectively.

Table 2: Comparison of different approaches of active case detection

Approach	Screened	Suspected cases	Confirmed cases	New case detection rate (/100,000)	PB:MB (%)	F:M (%)	Pediatric cases (%)	G2D (%)	Cost/case (USD$)
House-to-house visits	26469	365	27	102	16:11 (59:41)	17:10 (63:37)	3 (11)	3(11)	76,500 ($654)
Contact tracing	7608	214	19	250	13:6 (68:32)	11:8 (57:43)	2 (11)	1 (5)	90,286 ($772)
Prison screening	4428	185	2	45	0:2 (0:100)	0:2 (0:100)	0 (0)	0 (0)	298,300 ($2550)

175 Screened - total numbers screened; PB/MB - paucibacillary/multibacillary ratio; F/M -

176 female/male ratio; G2D – Grade 2 deformity

Age and gender of leprosy cases

178 In aggregate, the number of females among confirmed leprosy cases (28/48) was higher than

179 males, but not significantly different (χ^2 = 1.3, df = 1, p-value = 0.25) (Figure 1, Table S2).

186   Figure 1. Age and gender distributions of cases

Attack rate and associations of contacts and leprosy

Individuals with the history of two to five years of proximate contact with leprosy confirmed cases

were examined, in which attack rates were higher among household contacts of MB cases (0.32%,

95% CI 0.07-0.94) compared to neighboring contacts of the same cases (0.23%, 0.1-0.46), but this

was not significant (χ^2 = 0.07, df = 1, p-value = 0.9). Neighboring contacts of the PB cases were

found to have higher attack rate (0.48%, 0.19-0.98) compared to the household contact (0.13%,

0.03-0.73) of the same cases, but this was again not significant (χ^2 = 0.8, df = 1, p-value = 0.7)

(Table S3, Figure S1).

Differences in related associations, such as household contacts of MB case odds of being a case

compared to neighbours (*OR*=1.4, 95% CI 0.24-5.85) or PB household cases (*OR*=2.5, 95% CI 0.2-

129.1) and other variations of these associations were not significantly different. Further data and tables

are provided in the supplementary information (Table S4, Figure S2).

The contact tracing of 106 and 177 leprosy index cases in Siraha ($n = 3170$) and Bardiya ($n =$

4438) after 2-5 years of proximate contact with index cases identified 14 new cases Siraha (0.13

cases per index case, 0.07-0.21) and 5 in Bardiya (0.03, 0.01-0.06). This was statistically

significant ($\chi^2 = 9.8$, df = 1, p-value = 0.002), including adjusting for index cases and screened

contacts ($\beta = 1.2$, standard error = 0.5, p-value = 0.02).

Gender, case classification and disability

206 We found no statistical differences between genders and either having advanced (MB) leprosy

207 (*OR* = 0.47, 0.11-1.78, p-value = 0.24) or G2D (*OR* = 0.76, 0.05-11.4, p-value = 1) using all 48

208 new confirmed cases.

BCG and leprosy transmission

211 Among the total participants, 569 visited a health facility for confirmation of leprosy and were

212 also inspected for a BCG vaccination scar by a dermatologist. Those participants with the presence

213 of a BCG scar were found to have a significantly lower odds of having leprosy (18 of 341) than

214 without (27 of 201), with an OR = 0.42 (0.22 – 0.81, p-value = 0.007).

Discussion

We report 48 new leprosy cases from 38,505 screened people, comprising 29 from house-to-house

screening among vulnerable populations, 19 from case-contact tracing and 2 from prisoner

screening. House-to-house screening and contact tracing discovered 11% pediatric cases in both

approach and 11% and 5% G2D cases respectively, indicating new transmission events and late

diagnosis, highlighting the gaps in leprosy control programs.

The new case detection rate was highest in contact tracing (250) followed by house-to-house visits

(102) and prison screening (45) in per 100,000 population screened, whereas the most cost-

efficient approach here was house-to-house visits (NPR 76,500/case), followed by contact tracing

(NPR 90286/case) and prison screening (NPR 298,300/case). These costs per case detected are

similar to those reported for other countries, for example case contact tracing was approximately

US$529 (vs ~US$758) in similar study of Nigeria (17), scaled by inflation and using December

2021 exchange rates. The effectiveness and cost efficiency suggest implementing both approaches

in parallel may be optimal.

The epidemiological and clinical features of the identified confirmed cases were not significantly

different. This is possibly because of the small sample of confirmed cases. The PB:MB ratio

differed to the global status (35:65), but greater sample sizes might alter this. Similarly, the female:

male ratio differed but with great sample sizes might change to match the national (42:58) and

global (40:60) ratios. However, if the findings here are true but simply lacking statistical power

due to smaller sample sizes, then these altered ratios could be due to active case detection versus

passive case detection and suggest females with leprosy are often hidden with passive surveillance

(4). Future efforts should aim for earlier detection through active case detection and to reduce the

stigma attached to leprosy, so cases are not hidden.

The transmission attack rates observed in household (0.32%, 95% CI 0.07-0.94) and neighboring

(0.13%, 95% CI 0.03-0.73) contacts of MB cases, and household (0.23%, 95% CI 0.1-0.46) and

neighboring (0.48%, 95% CI 0.19-0.98) contacts of PB cases were not statistically different. The

rates, however, are lower than some other reports, such as 2% in Brazil in 2008 (15). This requires

additional studies in more districts to confirm, but the lower attack rate in the current Nepalese

situation also indicates progress towards elimination of the disease. The lack of a significant

difference in household contacts of cases developing leprosy compared to neighboring contacts

(0.78, 95% CI 0.19-2.45) differs from other findings where households contacts may have twice

the risk of developing disease compared to neighboring contacts (10). The reasons for this could

be sample size and statistical power, or that other factors are either reducing the within-household

transmission or increasing the neighbor-case transmission. Again, further work is needed to

determine which is occurring, but if reduced within-household transmission, this could be a sign

of successful case management.

The contact tracing of leprosy index cases was conducted after 2-5 years of proximate contact in

both Siraha (Madhesh Province) and Bardiya (Lumbini Province), where more new cases per index

case in Siraha (14 , 0.13 cases per index case, 0.07-0.21) than in Bardia (5, 0.013 per case, 0.01-

0.06). The difference was significant, including adjusting for index cases and screened

populations. It was reported that leprosy post exposure prophylaxis (LPEP) has been implemented

in Bardia for several years.

A further encouraging finding was that BCG vaccination with the presence of a scar has a

significant protective effect against leprosy (OR = 0.42, 95% CI 0.22 − 0.81). The finding is

consistent with other findings, such as in Brazil, where the OR = 0.27 (95% CI 0.13-0.59) (15).

The findings suggest BCG immunization programs will successfully contribute to leprosy

elimination. For Nepal, this is encouraging, because BCG is given at birth and national coverage

is high at 97.8% (95% CI 95.8–98.7) for BCG (18,19). However, like all immunization programs,

there are often small pockets of people where there is lower vaccine coverage, and lower BCG is

reported for at risk populations such as Madhesi, Dalit, and some religious minorities, who were

targeted for screening here (20). Future immunization programs should aim to ensure at risk

communities are reached to achieve the leprosy elimination goals.

Conclusion

The new case detection rates identified in this study suggest sustained levels of transmission in the communities screened. The proportion of pediatric cases (>10%) is evidence of recent transmission and the proportion of G2D confers evidence of late diagnosis and inadequate surveillance in the community. Though not significant, the female: male case ratio being the reverse of the global and national reports from passive case surveillance systems indicates hidden cases in the community, suggesting active surveillance is required to hasten leprosy elimination. The reduced attack rate compared to earlier studies, however, suggest some progress towards disease elimination and BCG vaccination should be given more attention as a tool for elimination, as it reduces transmission. Together, active case detection, through house-to-house visits and contact tracing to detect early and hidden cases, along with the optimal use of BCG, might help block transmission, prevent disabilities, and move Nepal closer towards elimination.

Contributors

RKM conceived and designed the manuscript; RKM, UG, ML, PN, KL KPP, CLD managed and supervise the data collection; RKM, DTSH analyzed the data; RKM, BB, MKP wrote the original draft KPP, CLD, DTSH reviewed the manuscript. All authors read, reviewed, and approved the final manuscript.

Declarations of interests

The authors declare that they have no competing interests and views does not represent the organization's views.

ACKNOWLEDGMENTS

We are grateful to all the provincial health directors, section chiefs and focal people of the

municipalities, health facilities, health offices, where active case detection was carried out for their

kind cooperation, direct involvement in case screening, monitoring and supervision. We are more

grateful to EDCD LCDMS staff who contributed to coordination and monitoring of the activities.

We sincerely express our gratitude to Dr Amrita Shrestha, Consultant Dermatologist of Badegaun

PHC, Dr Badri Chapagain, Chief Consultant Dermatologist of Nepalgunj Hospital, Dr Sumit

Pandey, Consultant Dermatologist of Nepalgunj Medical college, Dr. Niraj Parajuli, Dr Mahesh

Sah, Senior Consultant Dermatologist of Anandban Hospital, Dr Abhisek Bilash Panta, Consultant

Dermatologist of Janakpur Hospital and Mr Dipak Bam, Leprosy Expert who worked on the

confirmation and validation of suspected cases. This work was supported by grants from WHO

through Red book of the Ministry of Health and Population. We are especially thankful to Dr Usha

Kiran, coordinator of NTDs section of WHO Nepal who continuously supported us in technical

design. DTSH is funded by New Zealand Royal Society Te Apārangi, grant number MAU1701

and a Percival Carmine Chair in Epidemiology and Public Health.

Data availability

The datasets generated and/or analyzed during the current study are not publicly available due to individual privacy of patients could be compromised but are available from the corresponding author on reasonable request.

Ethics approval, consent to participate, and consent for publication.

Ethical approval was waived by the board of Nepal Health Research Council Nepal as the data collection is a part of regular surveillance activities. However informed consent was taken from individual participants. The data was generated through contact tracing and house to house visit among vulnerable population as a part of regular program in EDCD and approved for the analysis by the Epidemiology and Disease Control Division of Department of Health, Ministry of Health and Population of Nepal. No personal information (anonymized) was obtained or used in this manuscript writing.

References

1.   Bhat RM, Prakash C. Leprosy: An overview of pathophysiology. Interdiscip Perspect Infect Dis. 2012;2012.

2.   Joseph GA, Sundar Rao PSS. Impact of leprosy on the quality of life. Int J Lepr Other Mycobact Dis. 1999;67(4 SUPPL.):518.

3.   Leprosy/Hansen Disease: Management of reactions and prevention of disabilities.

4.   Harrell GT. Epidemiology of leprosy. J Am Med Assoc. 1947;135(11):732.

5.   Strategy GL. Global consultation of National Leprosy Programme managers , partners and affected persons on Global Leprosy Strategy. 2021.

6.   Department of Health services(DoHS). Annual Report: Department of Health Services 2075/76 (2018/19) [Internet]. Vol. 76, Departement of Health Services, Ministry of Health and Population, Government of Nepal. 2019. Available from: https://publichealthupdate.com/department-of-health-services-dohs-annual-report-2075-

334   76-2018-19/

7.   WHO. Global Leprosy Strategy 2016-2020: accelerating towards a leprosy-free world. [Internet]. Vol. 1, Weekly Epidemiological record. 2016. 349–360 p. Available from: http://apps.who.int/iris/bitstream/10665/205149/1/B5233.pdf?ua=1

8.   Moura MLN, Dupnik KM, Sampaio GAA, Nóbrega PFC, Jeronimo AK, do Nascimento-Filho JM, et al. Active Surveillance of Hansen's Disease (Leprosy): Importance for Case Finding among Extra-domiciliary Contacts. PLoS Negl Trop Dis. 2013;7(3):1–7.

9.   Tiendrebéogo A, Sow SO, Traore M, Sissoko K, Coulibaly B. Comparison of two methods of leprosy case finding in the circle of Kita in Mali. Int J Lepr Other Mycobact Dis. 1999;67(3):237–42.

10.   WHO. Leprosy/Hansen disease: Contact tracing and post-exposure prophylaxis.

11.   de Campos DCC, Dutra APB, Suares VL, de Carvalho PAC, Camargo LMA. New strategies for active finding of leprosy cases in the Amazonian region. Rev Soc Bras Med Trop. 2015;48(4):488–90.

12.   Vijayakumaran P, Mahipathy P V, Misra RK, Petro TS, Ramanujan R, Karunakaran S, et al. Hidden cases of leprosy (in prison). Indian J Lepr [Internet]. 1997;69(3):271—274. Available from: http://europepmc.org/abstract/MED/9394177

13.   Bernardes Filho F, Santana JM, de Almeida RCP, Voltan G, de Paula NA, Leite MN, et al. Leprosy in a prison population: A new active search strategy and a prospective clinical analysis. PLoS Negl Trop Dis. 2020;14(2):1–17.

14.   Merle CS, Cunha SS, Rodrigues LC. BCG vaccination and leprosy protection: Review of

current evidence and status of BCG in leprosy control. Expert Rev Vaccines.

2010;9(2):209–22.

15. Goulart IMB, Bernardes Souza DO, Marques CR, Pimenta VL, Gonçalves MA, Goulart

LR. Risk and protective factors for leprosy development determined by epidemiological

surveillance of household contacts. Clin Vaccine Immunol. 2008;15(1):101–5.

16. Tim CE. Epidemiologic programs for computers and calculators: Decision-tree analysis

using a microcomputer. Am J Epidemiol. 1986;124(5):843–50.

17. Ezenduka C, Post E, John S, Suraj A, Namadi A, Onwujekwe O. Cost-Effectiveness

Analysis of Three Leprosy Case Detection Methods in Northern Nigeria. PLoS Negl Trop

Dis. 2012;6(9).

18. Government of Nepal M of H and P. National Immunization Programme [Internet]. [cited

2022 Jul 18]. Available from: https://www.mohp.gov.np/eng/program/child-health-

services/nip

19. Rauniyar SK, Iwaki Y, Yoneoka D, Hashizume M, Nomura S. Age-appropriate

vaccination coverage and its determinants in children aged 12–36 months in Nepal: a

national and subnational assessment. BMC Public Health. 2021;21(1):1–12.

20. Shrestha S, Shrestha M, Wagle RR, Bhandari G. Predictors of incompletion of

immunization among children residing in the slums of Kathmandu valley, Nepal: A case-

control study. BMC Public Health [Internet]. 2016;16(1):1–9. Available from:

http://dx.doi.org/10.1186/s12889-016-3651-3

Figure