Perfusion imaging heterogeneity during NO inhalation distinguishes pulmonary arterial
hypertension (PAH) from healthy subjects and has potential as an imaging biomarker

Tilo Winkler PhD[1], Puja Kohli MD MMSc[2], Vanessa J Kelly PhD[2], Ekaterina G. Kehl MD[2], Alison S.
Witkin, MD[2], Josanna Rodriguez-Lopez MD[2], Kathryn A. Hibbert MD[2], Mamary Kone MD MPH[2],
David M. Systrom MD[3], Aaron B. Waxman MD[3], Jose G. Venegas PhD[1], Richard Channick MD[2], R.
Scott Harris MD[2]

[1] Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital and
Harvard Medical School, Boston, Massachusetts
[2] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Massachusetts General
Hospital and Harvard Medical School, Boston, Massachusetts
[3] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and Women's
Hospital and Harvard Medical School, Boston, Massachusetts

ORCID
Tilo Winkler   https://orcid.org/0000-0002-7276-5550
Kathryn A. Hibbert   https://orcid.org/0000-0001-7908-5446
David M. Systrom   https://orcid.org/0000-0002-9610-6330
Aaron B. Waxman   https://orcid.org/0000-0001-7797-0361

Corresponding author:
Tilo Winkler, PhD
Massachusetts General Hospital
Dept of Anesthesia, Critical Care and Pain Medicine
55 Fruit Street, Boston, MA 02114
USA
Email address: twinkler@mgh.harvard.edu
Phone: ++1-617-724-4083

ABSTRACT

Background: Without aggressive treatment, pulmonary arterial hypertension (PAH) has a 5-year mortality

of approximately 40%. A patient's response to vasodilators at diagnosis impacts the therapeutic options

and prognosis. We hypothesized that analyzing perfusion images acquired before and during vasodilation

could identify characteristic differences between PAH and control subjects.

Methods: We studied 5 controls and 4 subjects with PAH using HRCT and ^{13}NN PET imaging of

pulmonary perfusion and ventilation. The total spatial heterogeneity of perfusion (CV^2_{Qtotal}) and its

components in the vertical (CV^2_{Qvgrad}) and cranio-caudal (CV^2_{Qzgrad}) directions, and the residual

heterogeneity (CV^2_{Qr}), were assessed at baseline and while breathing oxygen and nitric oxide (O_2+iNO).

The length scale spectrum of CV^2_{Qr} was determined from 10 to 110 mm, and the response of regional

perfusion to O_2+iNO was calculated as the mean of absolute differences. Vertical gradients in perfusion

(Q_{vgrad}) were derived from perfusion images, and ventilation-perfusion distributions from images of ^{13}NN

washout kinetics.

Results: O_2+iNO significantly enhanced perfusion distribution differences between PAH and controls,

allowing differentiation between PAH subjects from controls. During O_2+iNO, CV^2_{Qvgrad} was significantly

higher in controls than in PAH (0.08 (0.055-0.10) vs. 6.7 x 10^{-3} (2x10^{-4}-0.02), p<0.001) with a

considerable gap between groups. Q_{vgrad} and CV^2_{Qtotal} showed smaller differences: –7.3 vs. -2.5, p = 0.002,

and 0.12 vs. 0.06, p = 0.01. CV^2_{Qvgrad} had the largest effect size among the primary parameters during

O_2+iNO. CV^2_{Qr}, and its length scale spectrum were similar in PAH and controls. Ventilation-perfusion

distributions showed a trend towards a difference between PAH and controls at baseline, but it was not

statistically significant.

Conclusions: Perfusion imaging during O2+iNO showed a significant difference in the heterogeneity

associated with the vertical gradient in perfusion, distinguishing PAH subjects from controls.

Trial registration: Not applicable

KEYWORDS

Pulmonary Circulation

Vascular Physiology

Functional Imaging

Positron Emission Tomography

Perfusion distribution

Ventilation/Perfusion distribution

Inhaled nitric oxide

INTRODUCTION

Pulmonary arterial hypertension (PAH) is a progressive disease with a 5-year mortality, if not aggressively treated, of approximately 40% [1]. Wide variations in patients' response to vasodilator treatment is a major challenge to improving outcomes [2]. Treatment guidelines define a positive response to the vasodilator challenge as a reduction of the mean pulmonary arterial pressure (mPAP) $\geq$ 10 mmHg to reach an absolute value of mPAP $\leq$ 40 mmHg with an increased or unchanged cardiac output [3]. But, only about 10% of patients with idiopathic PAH meet these clinical criteria for a response to vasodilators [3] that allow greater treatment options and have better survival [2,3]. Interestingly, it has been shown that the magnitude of responses in pulmonary vascular resistance (PVR) or mean PAP to inhaled nitric oxide with oxygen predicts long-term survival even among patient that are non-responders [4]. Overall, however, we have little insight to date into the pulmonary vascular function that is associated with increased risk.

In contrast to global parameters such as mPAP or pulmonary vascular resistance (PVR), imaging of pulmonary perfusion provides insights into regional changes associated with the disease process and its heterogeneity affecting the perfusion distribution inside the lungs. However, it is unclear if pulmonary perfusion imaging can detect substantial differences between PAH and healthy controls or assess regional changes in response to inhaled nitric oxide with oxygen. In healthy lungs, there is a vertical gradient in perfusion [5,6] including reduced lung inflation of dependent lung regions due to gravity [7]. Additionally, there is heterogeneity in perfusion within isogravitational slices that might be related to the fractal nature of the distribution of alveoli and the structure of the pulmonary vascular tree [8–10]. In addition, super-imposed disease processes can lead to regional changes in lung structure or function with characteristic sizes reflected in the length-scale spectrum of perfusion heterogeneity. It is however unclear if changes in the vascular structure in PAH, such as plexiform lesions [11], increase the regional heterogeneity of perfusion associated with the specific size of affected anatomical structures. We showed that length-scale analysis of pulmonary perfusion images could detect exercise PAH at rest [12]. Other imaging studies found associations between PAH and changes in the redistribution of pulmonary perfusion in response to changes

in body position [13,14], regional pulmonary blood volume [15–17], pulmonary impedance associated with

perfusion [18], and the vertical gradient in pulmonary perfusion [19]. Though early detection of PAH could

improve the survival of patients [20], no pulmonary imaging parameter has been identified yet to allow a

precise differentiation between PAH patients and healthy individuals,

Here, we used positron-emission tomography and high-resolution computer tomography (PET-CT)

imaging for quantitatively characterize the spatial distributions of pulmonary perfusion and ventilation in

patients with PAH. We compared the patterns of perfusion while breathing air and during inhalation of

nitric oxide (iNO) and oxygen (O_2) (O_2+iNO) to differentiate active effects of vasoconstriction [21],

anatomical structure of the vascular tree and the vertical gradient in hydrostatic pressure due to gravity. We

hypothesized that differences in pulmonary perfusion images acquired at baseline and during vasodilation

could identify characteristic perfusion patterns between controls and PAH subjects, even in PAH subjects

deemed "non-responders" to acute vasodilation during cardiac catheterization.

METHODS

Subjects

The study was approved by the Institutional Review Board of the Massachusetts General Hospital. Informed

consent was obtained from each subject before the study. We previously published an imaging study of

exercise PAH [12], using the same control group as in the present study. However, there is no overlap in

primary parameters and findings between both studies. Subjects enrolled with PAH at the time of

recruitment, as defined by the World Health Organization Group 1 classification, were required to have a

right heart catheterization (RHC) within 12 months of study entry demonstrating mPAP >25 mmHg and

pulmonary capillary wedge pressure (PCWP) < 15 mm Hg.

Spirometry data were collected for subjects with PAH. Trends in cardiac output were non-invasively

recorded in all subjects using impedance cardiography (ICON, Cardiotronic, LaJolla, CA). Due to the

known limitations of impedance cardiography, a trend rather than an absolute change in cardiac output, was

monitored. For assessment of within-subject changes in perfusion and perfusion heterogeneity compared to

baseline and for the comparison of regional perfusion between PAH and controls, we used values of mean-

normalized regional perfusion obtained directly from imaging data. Pulse oximetry and heart rate were

continuously monitored from a fingertip pulse oximeter.

PAH subjects receiving vasodilators as part of their treatment were maintained on those therapies during

the day of study.

Image Acquisition

A catheter was placed in an antecubital vein for radioisotope injection. Subjects were positioned supine and

with arms abducted on the table of the PET-CT scanner (Biograph 64, Siemens Healthcare, Malvern PA,

USA). Relative lung volume was continuously measured using impedance plethysmography (SomnoStar

PT, SensorMedics Corp., Yorba Linda, CA, USA). After calibration of the plethysmograph, mean lung

volume (MLV) was determined during 30 seconds of steady tidal breathing prior to imaging. Image

acquisition started a 'topogram' CT scan to determine the field of view for PET and HRCT scans. In

preparation for an HRCT scan, the subject was asked to inhale to total lung capacity (TLC), followed by

slow exhalation, and to hold the breath when the impedance plethysmography signal showed that MLV was

reached. During the breath-hold the HRCT scan was acquired. Following this, a dynamic ^{13}NN-saline PET

scan was acquired starting with an equivalent breath-hold at MLV. Simultaneously with the beginning of

the breath-hold we injected a bolus of ^{13}NN dissolved in saline (25mL) and started the acquisition of the

dynamic PET scan. After 30 seconds of breath-hold, we asked the subject to resume breathing while the

PET scan acquisition continued. Total acquisition time of the PET scan was 6 minutes and 40 seconds.

During this first PET-CT imaging sequence the subject was breathing air. After it was completed, subjects

were administered a gas mixture (iNO, iNOMax Delivery System) of balance gas oxygen with 30 ppm inhaled nitric oxide (O_2+iNO) through a one-way valve with a mouthpiece and nose clip. After at least 5 minutes of breathing O_2+iNO, we repeated the PET-CT imaging sequence while the subject continued to breath the O_2+iNO gas mixture. Given the sample size and the desire to prevent carry over effects from the inhaled nitric oxide, all subjects underwent baseline imaging followed by imaging with iNO without randomization of order of imaging.

HRCT scans (helical mode with 64 slices per rotation, 120 kVp, and 80 mAs) were reconstructed with filtered back-projection (convolution kernel B31f, 0.5 mm slice increment, voxel size: 0.7324 x 0.7324 x 0.5 mm^3). The PET field of view included on average 75% of the CT volume of the thorax. Dynamic PET scans were reconstructed with filtered back-projection and attenuation corrections using the HRCT scan. The dynamic PET scans yielded a stack of 19 images (8x5s, 3x10s, 7x30s, 60s, 2x30s), each image a 3D matrix of 128x128x81 voxels (voxel size 5.3456 x 5.3456 x 2.0250 mm^3). The PET scans have a full-width at half maximum (FWHM) resolution of 6 mm (Biograph 64, Siemens Healthcare, Malvern PA, USA) so that the effective resolution with a contrast recovery coefficient of 80% at two times FWHM is approximately 12 mm. PET images were filtered before image analysis using a moving average filter of five slices (~10mm) in the axial direction and a filter with 10 mm diameter within slices, which is comparable to the effective resolution of the scans.

Image Analysis

Image segmentation software Apollo (VIDA Diagnostics, IA, USA) was used to identify the lung region of interest (ROI) in the HRCT scans. Using custom software tools written in Matlab (Mathworks, Natick, MA), the lung ROIs were converted to ROI masks for PET scan analysis. Additionally, HRCT scans were converted into gas fraction (F_{gas}) images using the equation:

F_{gas} = (tissue density – density value of the voxel) / (tissue density – density of air),

where tissue density = 0 Hounsfield units (HU) and density of gas = -1000 HU. Tissue fraction (F_{tis}) images

were calculated using $F_{tis} = 1 - F_{gas}$.

Perfusion and ventilation images at baseline and after O_2+iNO were generated from ^{13}NN-saline PET scans.

The analysis of dynamic PET scans of ^{13}NN kinetics following a bolus injection have been described

previously [22,23]. Briefly, nitrogen gas has a very low solubility, and intravenously injected ^{13}NN,

dissolved in saline, diffuses during its passage through the pulmonary capillaries rapidly out of the blood

into the alveolar gas space, where it accumulates during the breath hold reaching a plateau value that is

directly proportional to the local fraction of perfusion. Thus, the voxel values of a PET image of ^{13}NN

activity during the plateau, divided by the mean activity within the lung represents those of a mean-

normalized perfusion distribution, allowing straightforward comparisons among images taken at different

conditions. After the breath hold, the subject resumes tidal breathing resulting in the washout of the ^{13}NN

by ventilation with the tracer kinetics during this period yielding an estimate of specific ventilation ($s\dot{V}$)

defined as: $s\dot{V}$ = alveolar ventilation ($\dot{V}_A$) / alveolar gas volume.

Assessment of regional ventilation and $\dot{V}/\dot{Q}$ ratio using ^{13}NN tracer kinetics modelling was performed as

previously described [23]. Briefly, the tracer kinetics curves of each voxel were analysed to determine $s\dot{V}$,

the perfusion associated with the $s\dot{V}$ kinetics, and $\dot{V}/\dot{Q}$ ratio. Four models were used for quantification of

tracer washout kinetics: 1) single-exponential washout describing homogenous voxels, 2) complete gas-

trapping for voxels with no change in activity during washout, 3) a double-exponential washout model

describing heterogenous voxels with two compartments that have different $s\dot{V}$, and 4) a partial gas trapping

model for heterogeneous voxels with one washout and one gas trapping compartment. The perfusion

associated with the $s\dot{V}$ of a voxel or compartment was proportional to the initial activity of the tracer

kinetics. After fitting all four models for each voxel, we used Akaike Information Criterion (AIC) (32) to

select for each individual voxel the optimal model yielding $s\dot{V}$ and perfusion of the voxel or sub-voxel

compartments. $\dot{V}_A$ was calculated as $s\dot{V}/V$, where V = Fgas * voxel volume. For voxels with two

compartments, the gas volume of each compartment was assumed to be proportional to the fraction of blood

flow in this compartment relative to the whole voxel: V_i = Fgas * voxel volume * Q_i / Q_{voxel}, were i is

compartment 1 or 2, and $Q_{voxel} = Q_1 + Q_2$. Mean normalized $\dot{V}/\dot{Q}$ distributions were computed from the $\dot{V}/\dot{Q}$

ratios of the individual voxels and sub-voxel compartments using $\dot{V}_A = \dot{V}_A$ / (sum of $\dot{V}_A$ / number of voxels)

and Q = Q / (sum of Q / number of voxels). $\dot{V}/\dot{Q}$ plots were constructed by binning of the typically 20,000-

50,000 individual voxel and sub-voxel $\dot{V}/\dot{Q}$ values using 100 bins over the $\dot{V}/\dot{Q}$ range from 0.001 to 100.

Note that the voxels with two-compartment washout kinetics were included as independent compartments

[24,25]. The $\dot{V}/\dot{Q}$ dispersions of perfusion SD Q log ($\dot{V}/\dot{Q}$) and ventilation SD V log ($\dot{V}/\dot{Q}$) were calculated

as the standard deviations of the perfusion-weighted and ventilation-weighted natural logarithm (Peter

Wagner, personal communication, July 11, 2020) of the $\dot{V}/\dot{Q}$ distribution, respectively [26,27]. Note that

other authors have used the term log SD Q, but we wanted to avoid a misinterpretation as log(SD(Q)).

Perfusion Heterogeneity and Length Scale Spectrum Analysis

Heterogeneity of perfusion was determined as the square of the coefficient of variation (CV^2 = (standard

deviation/mean)^2) using a previously developed and validated method for assessment of noiseless CV^2 of

PET scans based on theoretical properties of measurement errors and empirical assessment of imaging noise

[28]. In short, random errors in the individual voxels of PET images, referred to as imaging noise, are

superimposed to the true voxel values (signal) of the PET image and uncorrelated with the imaging signal

so that the heterogeneity of an image including noise (CV^2_{Qimage}) is the sum of the variance of the true signal

(perfusion) and the variance of the imaging noise $CV^2_{Qimage} = CV^2_{Qtotal} + CV^2_{noise}$ [28]. We estimated CV^2_{noise}

using multiple PET images during the tracer plateau at the end of the breath-hold period and calculated the

total spatial heterogeneity of ^{13}NN images of relative perfusion, excluding the imaging noise, as CV^2_{Qtotal} =

$CV^2_{Qimage} - CV^2_{noise}$. In the supine position, gravitational forces typically cause a gradient in perfusion in

the dorsoventral direction with higher perfusion in the dependent (dorsal) regions than in the non-dependent

(ventral) regions, which can be a substantial component of CV^2_{Qtotal}. We separated the total spatial

heterogeneity of perfusion into components caused by systematic gradients in the vertical direction

(CV^2_{Qvgrad}), the axial (craniocaudal) direction (CV^2_{Qzgrad}), and the remaining, or residual heterogeneity

(CV^2_{Qr}). Thus, $CV^2_{Qtotal} = CV^2_{Qvgrad} + CV^2_{zgrad} + CV^2_{Qr}$. The value of CV^2_{Qr} was further split into its

components due to spatial variations in perfusion across the length scale spectrum [23,25]. Such a length

scale spectrum quantifies the components of heterogeneity originating from spatial patterns in regional

perfusion measured at different image resolutions.

For each condition, the length scale spectrum of perfusion was quantified as follows: First, the vertical and

the axial gradients (z direction) in perfusion (CV^2_{Qvgrad} and CV^2_{Qzgrad}) were determined. Then the gradients

were subtracted voxel-by-voxel from the 3D perfusion scan, and each of the resulting transverse slices of

the scan were low-pass filtered at 10, 30, 50, 70, 90 and 110 mm with correction for edge effects [25], CV^2

was calculated for each filter size, and the contribution of each length scale ranges were calculated as

differences between successive filter sizes, e.g. $CV^2_{Q10-30} = CV^2(10mm) - CV^2(30mm)$. Overall, small

length scales of a spectrum starting at 10 mm correspond to variation among volumes of approximately 1

mL, which corresponds to about 5 acinar units; whereas large length scales of the spatial spectrum

correspond to heterogeneity due to variation in structural, functional and disease processes among larger

regions such as segments or subsegments.

Voxel Mapping and Changes in Regional Perfusion

For the assessment of changes in perfusion at the voxel level, we performed elastic image registration using

the Advanced Normalization Tools (ANTs, free and open-source software, https://stnava.github.io/ANTs/)

to correct for potential tissue displacement between the two perfusion images. First, we identified the

transformation field that mapped the HRCT scan taken at MLV while breathing O_2+iNO to the

corresponding HRCT scan taken at MLV while breathing air. The same method has been successfully used

before to determine the regional volumetric deformation and regional lung strain during mechanical

ventilation [29,30]. In principle, this elastic registration method uses the density information of the HRCT

scans for the identification of the transformation field T(x) describing the change in tissue location from

one HRCT scan to the other. Image acquisition for both CT and PET scans at MLV allows for the

application of the transformation field T(x) to the O_2+iNO PET scan to map its voxels to the corresponding

voxels of the air PET scan correcting differences in tissue location. This correction is essential for an

assessment of changes between PET scans at the voxel level. Images representing the relative changes in

regional perfusion between baseline and O_2+iNO were calculated as voxel-by-voxel differences between

the mean-normalized perfusion at air and the mean-normalized perfusion at O_2+iNO.

Perfusion-height maps were generated as two-dimensional (2D) histograms of the number of voxels in

relation to relative perfusion and relative height from the most dorsal point of the lung mask and the most

ventral point. Difference maps of the perfusion-height distributions between air and O_2+iNO were

calculated as the difference between the perfusion-height distribution at O_2+iNO and the distribution at air.

Values in the difference map are visualized with a grey scale: no change equals 50% grey, decrease in

perfusion is lighter, and an increase is darker.

Mean absolute changes in regional perfusion between air and O_2+iNO were calculated at the voxel level as

mean absolute deviations. Contributions of imaging noise [28] to the mean deviations were removed using

the statistical properties of the difference image. Specifically, total variance of the difference image of

perfusion between air and O_2+iNO includes the sum of the variances of imaging noise in the two original

images. The variance of the imaging PET noise, which has a normal distribution, was determined, using a

previously described method for noise assessment using multiple time frames of a PET scan [28], and

subtracted from the total variance of the difference image. The mean absolute deviation was then calculated

using the relationship to the variance for a normal distribution as square root of two over pi times the

variance. In order to rule out that imaging noise could explain the variations in the difference image, we

tested in each individual study if the overall variance of the difference image was higher than the upper

limit of the 99.9% confidence interval of the variance from noise.

Statistical Analysis

Our primary outcomes of interest were the noise corrected values of total spatial heterogeneity of perfusion (CV^2_{Qtotal}) and its components, including the heterogeneity generated by the vertical gradient in perfusion (CV^2_{Qvgrad}) and the residual perfusion heterogeneity (CV^2_{Qr}) and its length scale components. In secondary analyses, we assessed various regional perfusion and ventilation metrics. We performed two-sided T-tests to evaluate whether there was a difference in means of each metric between study groups and for baseline vs. O_2+iNO within study groups. Additionally, we obtained Cohen's d effect size estimates for the primary parameters. All analyses were performed using R Statistical Software (v4.1.3)[31]. Effect sizes were estimated using the effsize R package (v0.8.1)[32]. Statistical significance was set at $p < 0.05$. Data are presented as mean (range) unless otherwise stated.

RESULTS

Study Participants

Nine subjects were studied: 5 controls, and 4 subjects with PAH (**Table 1**). Subjects were free of infections or rapid progression of their disease for at least one month prior to the study and did not have any other known cardiac or pulmonary disease. No subjects required oxygen therapy at the time of imaging. All controls were non-smokers with subjectively normal exercise tolerance and normal pulmonary function tests. All PAH subjects were initially found to have normal pulmonary function testing and normal chest imaging but two of them were subsequently found to have mildly abnormal spirometry on a repeat testing after imaging. PAH subjects had elevated resting mPAP (mean 43; range 33-63 mmHg) with normal PCWP (mean 11; range 7-18 mmHg). Two PAH subjects were "non-responders" as demonstrated by absence of significant reduction in mean pulmonary artery pressure and increased cardiac output after administration of a short-acting vasodilator, one was a responder, and one did not have a standardized test. Cardiac outputs

254 were lower while breathing O_2+iNO for both groups. However, there were no significant differences in

255 cardiac output trends between groups at rest or while breathing O_2+iNO. All PAH subjects were on

256 combination therapy with a PDE5 inhibitor and endothelin receptor antagonist, and two PAH subjects were

257 also on an inhaled prostacyclin or a prostacyclin receptor agonist.

Regional Perfusion Distribution: Effect of iNO and Oxygen

Total Spatial Heterogeneity in Perfusion

260 At baseline, the total spatial heterogeneity of perfusion (CV^2_{Qtotal}) for the PAH subjects was similar to that

261 of the controls (p=0.51). However, while breathing O_2+iNO, the PAH subjects had a lower CV^2_{Qtotal} (2-

262 fold, p=0.01) compared to control counterparts (Figure 1, **Table 2**).

Vertical and Axial Gradients in perfusion and the associated heterogeneity

A vertical gradient in regional perfusion was observed in both groups, with higher perfusion in the

dependent regions (dorsal) relative to the non-dependent (ventral) regions (Figure 1, **Table 2**). At baseline

breathing air, the magnitude of the vertical gradient (Q_{vgrad}) and the heterogeneity generated by that gradient

(CV^2_{Qvgrad}) were significantly greater in the control group compared with the PAH group (p=0.04 and 0.05,

respectively) (**Table 2**). During O_2+iNO, these differences between groups were enhanced: CV^2_{Qvgrad} was

6.1-fold and Q_{vgrad} 2.9-fold, greater in the controls compared to the PAH subjects (p<0.001 and p=0.013,

respectively) (Figure 1, **Table 2**). Notably, during O_2+iNO, both CV^2_{Qvgrad} and Q_{vgrad} distinguished PAH

subjects from controls with a considerable gap between groups compared to the overlap at baseline

breathing air (Figure 1). Quantifying the effect size using Cohen's d, CV^2_{Qvgrad} during O_2+iNO showed the

largest estimate with a 95%-confidence interval not crossing zero, followed by Q_{vgrad} and CV^2_{Qtotal}: 4.30

(1.42, 7.17), –3.25 (–5.65, –0.84) and 2.23 (0.22, 4.25), respectively (Figure 2). In the individual subjects,

the average response of CV^2_{Qvgrad} to breathing O_2+iNO was an increase of 0.0272 in the controls compared

to a minimal decrease of -0.0017 in PAH subjects. Our study was not designed to assess differences between

responders and non-responders in PAH. Nevertheless, we noted that the only confirmed responder among

the PAH subjects showed an increase in CV^2_{Qvgrad} during O_2-iNO in the contrast to decreases in CV^2_{Qvgrad}

in the other three PAH subjects (Figure 1B). PAH subjects generally displayed low vertical gradients in

perfusion at baseline and during O_2+iNO without a significant change in the gradient (**Figures 3**). Also, the

differences in the vertical gradient in perfusion between groups persisted after adjusting the regional blood

flow by regional lung density determined from HRCT.

The contribution of the axial (craniocaudal) gradient in perfusion to total heterogeneity was small, and there

was no significant difference between the groups regardless of O_2+iNO administration (p=0.17 at baseline,

p=0.39 after O_2+iNO) (Figure 1C).

Residual Perfusion Heterogeneity and the Length Scale Spectrum

After removing the vertical and axial gradients in perfusion from the scans, PAH subjects had similar

residual heterogeneity (CV^2_{Qr}) when compared to controls (p=0.41) at baseline. These similarities in CV^2_{Qr}

remained consistent while breathing O_2+iNO. To further analyze the residual spatial heterogeneity in

perfusion, we assessed the variations in perfusion across the length scale spectrum. At baseline and while

breathing O_2+iNO, PAH subjects had similar residual heterogeneity across the spectrum of length scales of

10-110 mm (p>0.05 for all included length scales), when compared to control subjects (**Tables 3 & 4**).

Change in Perfusion: Results of voxel mapping

PAH and control subjects demonstrated significant changes in regional perfusion between air and O_2+iNO

at the voxel level of the two registered images (**Figures 4 & 5**) without a significant difference between

PAH (mean 0.19, range 0.18-0.20) and controls (mean 0.17, range 0.13-0.20). Also, the variation in regional

perfusion at the voxel level in each individual subject was higher than the 99.9% confidence interval

expected for random differences due to imaging noise, demonstrating that the perfusion distributions during

O_2+iNO at the voxel level were different from those while breathing air.

Regional Ventilation: Effect of iNO and Oxygen

Alveolar ventilation ($\dot{V}_A$) was similar between groups before and during O_2+iNO (p=0.25, p=0.49 after O_2+iNO). There was no significant difference observed in the overall mean specific ventilation ($s\dot{V}$) or spatial heterogeneity of $\dot{V}_A$ (CV^2_{VA}) between controls and PAH subjects, reflecting a lack of parenchymal lung disease (**Table 5**). Additionally, lung parenchymal density (estimated by F_{tis}) was similar between control and PAH subjects.

Ventilation-Perfusion Distributions: Effect of iNO and Oxygen

The distributions of $\dot{V}/\dot{Q}$ ratios at baseline in PAH subjects were overall broader than in controls at baseline, approaching statistical significance, consistent with moderate degrees of vascular disease-causing mismatch (SD Q log ($\dot{V}/\dot{Q}$), controls 0.51 vs PAH 0.70, p=0.11). There was no difference in the $\dot{V}/\dot{Q}$ distribution between groups breathing O_2+iNO, in large part due to widening of the $\dot{V}/\dot{Q}$ distribution in controls (SD Q log ($\dot{V}/\dot{Q}$), controls 0.65 vs PAH 0.74, p=0.58). The distributions at baseline were unimodal in all subjects, except 2 PAH subjects who had bimodal distributions. $\dot{V}/\dot{Q}$ distributions during administration of iNO+O_2 were unimodal in all subjects including those who had a bimodal distribution prior to the intervention (**Figure 6, Table 5**).

DISCUSSION

We report a detailed analysis of regional perfusion and ventilation imaging data in PAH and control subjects obtained with PET-CT, revealing unique perfusion characteristics in PAH subjects that may not be reflected in routine PA pressure measurements. We found that: 1. Breathing O_2+iNO during imaging resulted in significant differences in the regional distribution of perfusion between PAH subjects and controls, distinguishing the subjects of the two groups. 2. CV^2_{Qvgrad} during O_2+iNO had the highest effect size for PAH vs. controls. 3. All subjects in our study responded to O_2+iNO with changes in regional

perfusion, including the two PAH subjects who were non-responders according to vasoreactivity testing.

4. PAH subjects had significantly lower vertical gradients in perfusion than controls and such gradients in

3 of the 4 subjects were unaffected by O_2+iNO. 5. PAH subjects had residual perfusion heterogeneity

across the length-scale spectrum that was similar to that of controls.

An unexpected finding was the magnitude of the vasodilation in healthy controls during O_2+iNO leading

to perfusion redistributions. The significant difference in CV^2_{Qvgrad} between PAH subjects and controls

suggests that perfusion imaging during O_2+iNO has the potential to be as a biomarker for PAH, that could

lead to earlier detection and thus improve patient survival [20]. It remains unclear if the change between

O_2+iNO and air allows identifying responders among PAH patients. The only confirmed responder in our

study showed a greater response in CV^2_{Qvgrad} compared to the other PAH subjects, but it is a single case.

Prior studies using SPECT, MRI and PET have observed that patients with primary precapillary pulmonary

hypertension display a pronounced attenuation in the normal gravity-dependent distribution of lung

perfusion [13,14,17,19] consistent with our findings. However, these studies reported overlaps between

groups, reported only group means, or had not investigated the perfusion distribution during pulmonary

vasodilation. One study by Jones et al. [19] using adenosine demonstrated limited change in perfusion

redistribution and near absence of a vertical perfusion gradient in PAH even after vasodilation. Based on

physiologic principles, studies have hypothesized that since gravity-dependent perfusion redistribution is

reliant on low arterial pressures at rest and the ability to recruit and distend vessels with increased cardiac

output, higher pulmonary arterial pressures and loss of vascular reserves from obstructive vasculopathies

may inhibit vessel recruitment and distension [14,33]. Our study confirmed a lack of significant vertical

perfusion redistribution during exposure to O_2+iNO in those with PAH, indicating that despite measured

changes in regional perfusion with vasodilation, PAH subjects did not show normal vertical perfusion

patterns. In fact, the vertical gradient in perfusion was in some subjects nearly absent in the dependent part

of the lungs, as demonstrated with perfusion-height mapping (**Figure 3**), indicating that vessels at the high

pulmonary arterial pressures in PAH may in these subjects already be maximally recruited and distended

[34], and therefore are minimally affected by hydrostatic pressures.

An unexpected finding was that the two PAH subjects that were "non-responsive" to vasodilator in terms

of the definition of the guidelines for an acute response during vasoreactivity testing [3] had mean values

of absolute changes in voxel-by-voxel perfusion during exposure to O_2+iNO that were similar to those of

controls, despite having no change in their vertical gradient in perfusion. Interestingly, there is clinical

evidence that patients with a decrease of $\geq 30\%$ in PVR in response to O_2+iNO benefit from vasodilator

therapy although they may not meet the guidelines' definition for a response during vasoreactivity testing

[4]. The hydrostatic pressure gradient caused by gravity would be expected to mainly effect blood vessel

distension, depending on the vessel compliance and tone. In this scenario, any loss of vertical gradient could

be thought of as a reflection of a loss of overall pulmonary vessel compliance or increase in vessel tone.

For instance, any mechanism narrowing the inner lumen, while the outside of the vessel is maximally

distended, could explain a high pulmonary vascular resistance, loss of the vertical gradient in perfusion

because the hydrostatic pressure gradient cannot expand the vessel, and the lack of a significant change in

mPAP to vasodilators. Our data suggests that some degree of vascular tone must be present since we see a

change in blood flow during exposure to O_2+iNO that is similar to controls. Perhaps the increase in tone

could be related to the smooth muscle hypertrophy that is seen in PAH [35], and the changes in regional

perfusion in our data could explain why some PAH patients can have symptomatic improvement with

vasodilators despite little change in mPAP [36].

Our study revealed that PAH subjects had relatively normal values of residual perfusion heterogeneity

across the length scale spectrum (**Tables 3 & 4**). This similarity to controls suggests that the vascular

disease process may not lead to increased heterogeneity among the resistances of different pathways of

blood flow throughout the pulmonary vascular tree. We found the lack of differences in the length scale

spectrum very surprising since histopathologic evaluations of the pulmonary vascular tree in patients with

PAH have found significant heterogeneity in the type and extent of pulmonary vascular remodelling [37–

39] that could have increased the heterogeneity in perfusion at some length scales linked to the anatomical

size of the affected sections of the vascular tree, and the length scale spectrum of perfusion heterogeneity

has shown differences in other studies, including exercise pulmonary arterial hypertension [12] and COPD

[25]. However, histopathologic samples showing structural changes cannot provide complete insights into

functional aspects in vivo, such as the vascular resistance along complete pathways through the vessel tree,

and systemic consequences of endothelial injury in pulmonary hypertension, such as alterations in the

coagulation system, abnormal platelet aggregation and altered production of various endothelial vasoactive

mediators, which may be contributing to greater uniformity in the perfusion pattern even in parts of the

vascular tree without histopathologic changes. Indeed, although local vascular remodelling and obstruction

could be expected to result in patches of low perfusion in PAH subjects, there was no evidence for

substantial areas of very low perfusion or significant total spatial heterogeneity of perfusion in our imaging

data (Figure 3 & 5). This is relevant because the ^{13}NN PET imaging technique measures the regional blood

flow through the alveolar capillaries, which is determined by the pressure difference between pulmonary

artery and pulmonary vein and the pathway resistance to blood flow. Therefore, the spatial heterogeneity

in perfusion includes the overall effects of vascular properties (structural and functional) as well as

abnormalities of the vessels along the pathways. In control subjects, O_2-iNO had no effect on the residual

heterogeneity of perfusion CV^2_{Qr} (Table 2), which is consistent with previous findings that hypoxic

pulmonary vasoconstriction had no effect on perfusion heterogeneity in healthy subjects [40].

The relative contribution of gravity and arterial geometry in determining pulmonary perfusion distribution

is still subject to debate. Hughes and West argue that gravity contributes to 24-61% of the overall variance

in blood flow in the human supine lung [41]. Glenny et al, on the other hand, argue that gravity accounts

for a minority of the variability (at most 28%) in perfusion distribution while the geometry of the vascular

tree, much of which is genetically predetermined, is claimed to be the primary determinate of pulmonary

blood flow [42]. Very few have studied the effects of vasodilation on redistribution of this vertical

pulmonary blood flow in healthy subjects. One prior study by Henderson et al. did not find gross

redistribution of pulmonary blood flow during hyperoxia in healthy subjects [43] while other studies have

found redistribution of blood flow [21] and increased inequality in $\dot{V}/\dot{Q}$ with hyperoxia [44] and iNO [45]

in healthy subjects. In our study, vasodilation with O_2+iNO significantly augmented the vertical gradient

of perfusion and widened the $\dot{V}/\dot{Q}$ distribution in control subjects. These findings suggest that the controls

have baseline pulmonary vasomotor tone enhancing $\dot{V}/\dot{Q}$ matching which responds to interventions such as

hyperoxia and iNO.

The SD Q log ($\dot{V}/\dot{Q}$) values for controls in our study (0.31-0.81) are consistent with the range of values

previously reported in healthy adults using MIGET (0.22-0.72) [46]. Additionally, MRI-based $\dot{V}/\dot{Q}$

distributions are in good agreement with MIGET measurements [47]. However, MIGET using micropore

membrane inlet mass spectrometry (MMIMS) to measure the tracer gases resulted in narrower $\dot{V}/\dot{Q}$

distributions compared to MIGET using gas chromatography [48]. In contrast to MIGET, which involves

a smoothing function to estimate the 50 values of a $\dot{V}/\dot{Q}$ distribution from measurements of six tracer gases,

dynamic PET imaging of ^{13}NN tracer kinetics allows the estimations of ventilation and perfusion for up to

two compartments per voxel and binning of the ventilation and perfusion values to derive $\dot{V}/\dot{Q}$ distributions

with 100 discrete values without a smoothing function. A validation of the method using heterogenous $\dot{V}/\dot{Q}$

distributions has shown a good agreement between predicted and measured blood gases [49]. In PAH, $\dot{V}/\dot{Q}$

distributions can be substantially wider than we have found [50–53], but our results are consistent with

examples of narrower distributions in some PAH subjects [50,51] and the average SD Q log ($\dot{V}/\dot{Q}$) for PAH

in our study was slightly higher compared to controls (0.70 vs. 0.51, respectively). We cannot rule out that

the exclusion of regions close to the diaphragm where the tidal breathing would cause partial volume effects

and axial limitations in the field of view of the PET scanner had some effect on the $\dot{V}/\dot{Q}$ distribution.

Additionally, our PET imaging method measures relative pulmonary perfusion and the conversion to blood

flow, e.g., in ml/min, would require reliable cardiac output measurements during imaging, which were not

available in this study. Our $\dot{V}/\dot{Q}$ distributions are therefore based on mean normalized ventilation and

perfusion.

Despite our study's strengths, some important limitations should be considered. Limitations of the [13]NN

imaging technique has been discussed in detail [5,27]. Briefly, although PET scans have lower resolution

than CT scans, they allow assessment of functional parameters and PET resolution is more than sufficient

to quantify functional imaging parameters such as vertical gradients [5]. The effective spatial resolution of

the PET perfusion scans is approximately 12 mm, motion blurring from tidal breathing results in lower

spatial resolution, but our method of sub-voxel compartments can capture functional heterogeneity at this

level. Second, [13]NN tracer kinetics during the breath hold may affect the accuracy of noise-free CV^2

parameters of the perfusion heterogeneity, and errors in image registration causing small misalignments

could affect the assessment of voxel-by-voxel changes in perfusion. However, such errors only diminish

differences among groups and would bias our results toward the null hypothesis of statistical tests.

Additionally, an analysis of the reproducibility of CV^2_{Qtotal}, excluding imaging noise, in our previous

validation showed a standard deviation of ±8.4% in repeated imaging [28]. Third, despite notable

differences observed with this small cohort, future studies that include other types of patients with

pulmonary hypertension and larger studies with matched cohorts will be necessary to assess whether these

findings are more generalizable. Furthermore, additional associations may be present that we are

underpowered to detect in this small cohort.

CONCLUSIONS

Perfusion imaging during O_2-iNO inhalation showed a large effect size for the heterogeneity associated

with the vertical gradient in perfusion, $CV^2_{Qvgrad,}$ for PAH followed in magnitude by the vertical gradient

in perfusion. PAH subjects could be discriminated from healthy controls showing the potential of

CV^2_{Qvgrad} measured during O_2-iNO inhalation to be a biomarker for PAH. Further studies are needed to

investigate NO inhalation in clinically available imaging modalities such as dual-energy CT to translate

our findings into clinics.

ABBREVIATIONS

446		
447	^{13}NN	Isotope nitrogen-13 in a molecule bound to a stable nitrogen atom
448	2D	two-dimensional
449	3D	three-dimensional
450	6MWD	six-minute walk distance
451	AIC	Akaike Information Criterion
452	CV^2	squared coefficient of variation quantifying heterogeneity
453	CV^2_{Q10-30}	heterogeneity of perfusion at the length scale of 10 to 30 millimeter
454	CV^2_{Qr}	residual heterogeneity of perfusion
455	CV^2_{Qtotal}	total spatial heterogeneity of perfusion
456 457	CV^2_{Qvgrad} perfusion	heterogeneity of perfusion associated with the vertical (dorso-ventral) gradient in
458 459	CV^2_{Qzgrad} perfusion	heterogeneity of perfusion associated with the z-axail (cranio-caudal) gradient in
460	FEV_1	forced expiratory volume in first second
461	F_{gas}	gas fraction images using
462	F_{tis}	tissue fraction
463	FWHM	full-width at half maximum
464	FVC	forced vital capacity
465	HRCT	high resolution computed tomography
466	HU	Hounsfield units
467	MLV	mean lung volume
468	mPAP	mean pulmonary arterial pressure
469	MRI	magnetic resonance imaging
470	O_2+iNO	inhalation of nitric oxide (iNO) with balance gas oxygen (O_2)
471	PAH	pulmonary arterial hypertension
472	PCWP	pulmonary capillary wedge pressure

473	PET-CT	positron-emission tomography and high-resolution computer tomography
474	PVR	pulmonary vascular resistance
475	Q_{vgrad}	vertical (dorso-ventral) gradient in perfusion
476	Q_{voxel}	perfusion of a voxel
477	RHC	right heart catheterization
478	ROI	region of interest
479	SD Q log $(\dot{V}/\dot{Q})$	$\dot{V}/\dot{Q}$ dispersions of perfusion
480	SD V log $(\dot{V}/\dot{Q})$	$\dot{V}/\dot{Q}$ dispersions of ventilation
481	SPECT	single-photon emission computed tomography
482	$s\dot{V}$	specific ventilation
483	TLC	total lung capacity
484	$\dot{V}/\dot{Q}$	ventilation/perfusion
485	$\dot{V}_A$	alveolar ventilation
486		

REFERENCES

1. Thenappan T, Ryan JJ, Archer SL. Evolving epidemiology of pulmonary arterial hypertension. Am J Respir Crit Care Med. 2012;186:707–9.

2. Halliday SJ, Hemnes AR, Robbins IM, Pugh ME, Zhao DX, Piana RN, et al. Prognostic value of acute vasodilator response in pulmonary arterial hypertension: beyond the "classic" responders. J Heart Lung Transplant Off Publ Int Soc Heart Transplant. 2015;34:312–8.

3. Galiè N, Humbert M, Vachiery J-L, Gibbs S, Lang I, Torbicki A, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension: The Joint Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS): Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC), International Society for Heart and Lung Transplantation (ISHLT). Eur Respir J. 2015;46:903–75.

4. Malhotra R, Hess D, Lewis GD, Bloch KD, Waxman AB, Semigran MJ. Vasoreactivity to inhaled nitric oxide with oxygen predicts long-term survival in pulmonary arterial hypertension. Pulm Circ. 2011;1:250–8.

5. Musch G, Layfield JDH, Harris RS, Melo MFV, Winkler T, Callahan RJ, et al. Topographical distribution of pulmonary perfusion and ventilation, assessed by PET in supine and prone humans. J Appl Physiol. 2002;93:1841–51.

6. Prisk GK, Yamada K, Henderson AC, Arai TJ, Levin DL, Buxton RB, et al. Pulmonary perfusion in the prone and supine postures in the normal human lung. J Appl Physiol Bethesda Md 1985. 2007;103:883–94.

7. Hopkins SR, Henderson AC, Levin DL, Yamada K, Arai T, Buxton RB, et al. Vertical gradients in regional lung density and perfusion in the supine human lung: the Slinky effect. J Appl Physiol Bethesda Md 1985. 2007;103:240–8.

8. Burrowes KS, Hunter PJ, Tawhai MH. Investigation of the Relative Effects of Vascular Branching Structure and Gravity on Pulmonary Arterial Blood Flow Heterogeneity via an Image-based Computational Model. Acad Radiol. 2005;12:1464–74.

9. Hlastala MP, Glenny RW. Vascular Structure Determines Pulmonary Blood Flow Distribution. Physiology. American Physiological Society; 1999;14:182–6.

10. Nelson TR, West BJ, Goldberger AL. The fractal lung: universal and species-related scaling patterns. Experientia. 1990;46:251–4.

11. Cool CD, Stewart JS, Werahera P, Miller GJ, Williams RL, Voelkel NF, et al. Three-dimensional reconstruction of pulmonary arteries in plexiform pulmonary hypertension using cell-specific markers. Am J Pathol. 1999;155:411–9.

12. Kohli P, Kelly VJ, Kehl EG, Rodriguez-Lopez J, Hibbert KA, Kone M, et al. Perfusion imaging distinguishes exercise pulmonary arterial hypertension at rest. Am J Respir Crit Care Med. 2019;199:1438–41.

13. Horn M, Hooper W, Brach B, Ashburn W, Moser K. Postural changes in pulmonary blood flow in
pulmonary hypertension: a noninvasive technique using ventilation-perfusion scans. Circulation.
1982;66:621–6.

14. Lau EM, Bailey DL, Bailey EA, Torzillo PJ, Roach PJ, Schembri GP, et al. Pulmonary hypertension leads
to a loss of gravity dependent redistribution of regional lung perfusion: a SPECT/CT study. Heart Br Card
Soc. 2014;100:47–53.

15. Lefebvre B, Kyheng M, Giordano J, Lamblin N, de Groote P, Fertin M, et al. Dual-energy CT lung
perfusion characteristics in pulmonary arterial hypertension (PAH) and pulmonary veno-occlusive
disease and/or pulmonary capillary hemangiomatosis (PVOD/PCH): preliminary experience in 63
patients. Eur Radiol. 2022;

16. Giordano J, Khung S, Duhamel A, Hossein-Foucher C, Bellèvre D, Lamblin N, et al. Lung perfusion
characteristics in pulmonary arterial hypertension (PAH) and peripheral forms of chronic
thromboembolic pulmonary hypertension (pCTEPH): Dual-energy CT experience in 31 patients. Eur
Radiol. 2017;27:1631–9.

17. Wollmer P, Rozkovec A, Rhodes CG, Allan RM, Maseri A. Regional pulmonary blood volume in
patients with abnormal blood pressure or flow in the pulmonary circulation. Eur Heart J. 1984;5:924–31.

18. Hovnanian ALD, Costa ELV, Hoette S, Fernandes CJCS, Jardim CVP, Dias BA, et al. Electrical
impedance tomography in pulmonary arterial hypertension. PLOS ONE. Public Library of Science;
2021;16:e0248214.

19. Jones AT, Hansell DM, Evans TW. Quantifying pulmonary perfusion in primary pulmonary
hypertension using electron-beam computed tomography. Eur Respir J. 2004;23:202–7.

20. Lau EMT, Humbert M, Celermajer DS. Early detection of pulmonary arterial hypertension. Nat Rev
Cardiol. Nature Publishing Group; 2015;12:143–55.

21. Asadi AK, Sá RC, Kim NH, Theilmann RJ, Hopkins SR, Buxton RB, et al. Inhaled nitric oxide alters the
distribution of blood flow in the healthy human lung, suggesting active hypoxic pulmonary
vasoconstriction in normoxia. J Appl Physiol Bethesda Md 1985. 2015;118:331–43.

22. Melo MFV, Layfield D, Harris RS, O'Neill K, Musch G, Richter T, et al. Quantification of Regional
Ventilation-Perfusion Ratios with PET. J Nucl Med. 2003;44:1982–91.

23. Wellman TJ, Winkler T, Costa ELV, Musch G, Harris RS, Venegas JG, et al. Effect of regional lung
inflation on ventilation heterogeneity at different length scales during mechanical ventilation of normal
sheep lungs. J Appl Physiol. 2012;113:947–57.

24. Harris RS, Winkler T, Tgavalekos N, Musch G, Melo MFV, Schroeder T, et al. Regional pulmonary
perfusion, inflation, and ventilation defects in bronchoconstricted patients with asthma. Am J Respir Crit
Care Med. 2006;174:245–53.

25. Vidal Melo MF, Winkler T, Harris RS, Musch G, Greene RE, Venegas JG. Spatial Heterogeneity of Lung
Perfusion Assessed with 13N PET as a Vascular Biomarker in Chronic Obstructive Pulmonary Disease. J
Nucl Med. 2010;51:57–65.

26. Kelly VJ, Hibbert KA, Kohli P, Kone M, Greenblatt EE, Venegas JG, et al. Hypoxic pulmonary
vasoconstriction does not explain all regional perfusion redistribution in asthma. Am J Respir Crit Care
Med. 2017;196:834–44.

27. Vidal Melo MF, Layfield D, Harris RS, O'Neill K, Musch G, Richter T, et al. Quantification of regional
ventilation-perfusion ratios with PET. J Nucl Med. 2003;44:1982–91.

28. Winkler T, Melo MFV, Degani-Costa LH, Harris RS, Correia JA, Musch G, et al. Estimation of noise-free
variance to measure heterogeneity. PloS One. 2015;10:e0123417.

29. Motta-Ribeiro G, Winkler T, Hashimoto S, Vidal Melo MF. Spatial heterogeneity of lung strain and
aeration and regional inflammation during early lung injury assessed with PET/CT. Acad Radiol.
2019;26:313–25.

30. Motta-Ribeiro GC, Hashimoto S, Winkler T, Baron RM, Grogg K, Paula LFSC, et al. Deterioration of
regional lung strain and inflammation during early lung injury. Am J Respir Crit Care Med. 2018;198:891–
902.

31. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria;
2022. Available from: https://www.R-project.org/

32. Torchiano M. Effsize - A Package For Efficient Effect Size Computation [Internet]. Zenodo; 2016 [cited
2022 May 4]. Available from: https://zenodo.org/record/1480624

33. West JB, Dollery CT. Distribution of blood flow and ventilation-perfusion ratio in the lung, measured
with radioactive CO_2. J Appl Physiol. 1960;15:405–10.

34. Lai-Fook SJ. A continuum mechanics analysis of pulmonary vascular interdependence in isolated dog
lobes. J Appl Physiol. 1979;46:419–29.

35. Pietra GG, Capron F, Stewart S, Leone O, Humbert M, Robbins IM, et al. Pathologic assessment of
vasculopathies in pulmonary hypertension. J Am Coll Cardiol. 2004;43:S25–32.

36. Barst RJ, Gibbs JSR, Ghofrani HA, Hoeper MM, McLaughlin VV, Rubin LJ, et al. Updated evidence-
based treatment algorithm in pulmonary arterial hypertension. J Am Coll Cardiol. 2009;54:S78-84.

37. Tuder RM, Stacher E, Robinson J, Kumar R, Graham BB. Pathology of pulmonary hypertension. Clin
Chest Med. 2013;34:639–50.

38. Rol N, Timmer EM, Faes TJC, Vonk Noordegraaf A, Grünberg K, Bogaard H-J, et al. Vascular narrowing
in pulmonary arterial hypertension is heterogeneous: rethinking resistance. Physiol Rep. 2017;5.

39. Stacher E, Graham BB, Hunt JM, Gandjeva A, Groshong SD, McLaughlin VV, et al. Modern Age
Pathology of Pulmonary Arterial Hypertension. Am J Respir Crit Care Med. 2012;186:261–72.

40. Arai TJ, Henderson AC, Dubowitz DJ, Levin DL, Friedman PJ, Buxton RB, et al. Hypoxic pulmonary
vasoconstriction does not contribute to pulmonary blood flow heterogeneity in normoxia in normal
supine humans. J Appl Physiol Bethesda Md 1985. 2009;106:1057–64.

41. Hughes M, West JB. Point:Counterpoint: Gravity is/is not the major factor determining the
distribution of blood flow in the human lung. J Appl Physiol. 2008;104:1531–3.

42. Glenny R. Counterpoint: Gravity is not the major factor determining the distribution of blood flow in
the healthy human lung. J Appl Physiol. 2008;104:1533–5.

43. Henderson AC, Sá RC, Theilmann RJ, Buxton RB, Prisk GK, Hopkins SR. The gravitational distribution
of ventilation-perfusion ratio is more uniform in prone than supine posture in the normal human lung. J
Appl Physiol Bethesda Md 1985. 2013;115:313–24.

44. Barberà JA, Roger N, Roca J, Rodriguez-Roisin R, Rovira I, Higenbottam TW. Worsening of pulmonary
gas exchange with nitric oxide inhalation in chronic obstructive pulmonary disease. The Lancet.
1996;347:436–40.

45. Katayama Y, Higenbottam TW, Diaz de Atauri MJ, Cremona G, Akamine S, Barbera JA, et al. Inhaled
nitric oxide and arterial oxygen tension in patients with chronic obstructive pulmonary disease and
severe pulmonary hypertension. Thorax. 1997;52:120–4.

46. Cardús J, Burgos F, Diaz O, Roca J, Barberá JA, Marrades RM, et al. Increase in Pulmonary
Ventilation–Perfusion Inequality with Age in Healthy Individuals. Am J Respir Crit Care Med.
1997;156:648–53.

47. Sá RC, Henderson AC, Simonson T, Arai TJ, Wagner H, Theilmann RJ, et al. Measurement of the
distribution of ventilation-perfusion ratios in the human lung with proton MRI: comparison with the
multiple inert-gas elimination technique. J Appl Physiol Bethesda Md 1985. 2017;123:136–46.

48. Kretzschmar M, Schilling T, Vogt A, Rothen HU, Borges JB, Hachenberg T, et al. Multiple inert gas
elimination technique by micropore membrane inlet mass spectrometry—a comparison with reference
gas chromatography. J Appl Physiol. 2013;115:1107–18.

49. Vidal Melo MF, Harris RS, Layfield JDH, Venegas JG. Topographic basis of bimodal ventilation-
perfusion distributions during bronchoconstriction in sheep. Am J Respir Crit Care Med. 2005;171:714–
21.

50. Bratel T, Lagerstrand L, Brodin L-A, Nowak J, Randmaa I. Ventilation-perfusion relationships in
pulmonary arterial hypertension: effect of intravenous and inhaled prostacyclin treatment. Respir
Physiol Neurobiol. 2007;158:59–69.

51. Dantzker DR, Bower JS. Mechanisms of gas exchange abnormality in patients with chronic
obliterative pulmonary vascular disease. J Clin Invest. 1979;64:1050–5.

52. Mélot C, Naeije R, Mols P, Vandenbossche JL, Denolin H. Effects of nifedipine on
ventilation/perfusion matching in primary pulmonary hypertension. Chest. 1983;83:203–7.

53. Voswinckel R, Reichenberger F, Gall H, Schmehl T, Gessler T, Schermuly RT, et al. Metered dose
inhaler delivery of treprostinil for the treatment of pulmonary hypertension. Pulm Pharmacol Ther.
2009;22:50–6.

 FIGURE LEGENDS

Figure 1: Perfusion heterogeneity profiling distinguished PAH subjects and control subjects. The

total spatial heterogeneity of perfusion (CV^2_{Qtotal})[28] (**A**) is comprised of the heterogeneity of the vertical

(dorsoventral) gradient in perfusion (CV^2_{Qvgrad}) (**B**), the heterogeneity of the craniocaudal (z-axis) gradient

in perfusion (CV^2_{Qzgrad}) (**C**), and the heterogeneity of the residual (or remaining) perfusion (CV^2_{Qr}) (**D**).

PAH subjects demonstrated very low CV^2_{Qvgrad} due to attenuated vertical gradients in perfusion (Q_{vgrad})

(**E**) that were not responsive to O_2+iNO, distinguishing this cohort from their control counterparts. Note

that imaging noise is excluded from all CV^2 parameters.

Figure 2: Cohen's d effect size estimates and 95% confidence intervals of the primary parameters.

The effect sizes for PAH vs. control were substantially larger for O_2+iNO compared to baseline. CV^2_{Qvgrad}

had the most significant effect size and a 95% confidence interval not crossing zero, followed by Q_{vgrad} and

CV^2_{Qtotal}. The relatively large effect sizes for O_2+iNO vs. baseline in the control group are indicators of the

magnitude of the response.

Figure 3: Perfusion-height maps for PAH subjects and controls. The greatest perfusion heterogeneity,

including the largest range of perfusion values at any given height, was present in PAH subjects, followed

by controls. The vertical gradient in perfusion (represented by the slope of the black-and-white dashed line)

increased in magnitude in controls but not PAH subjects breathing O_2+iNO. In contrast to controls, several

PAH subjects had no vertical gradient in perfusion in the lower or dependent part of the lung (dorsal in the

supine position), as if the hydrostatic pressure, which increases towards the dependent regions, had no effect

on perfusion. Surprisingly, there was no substantial fraction of voxels with very low or zero perfusion in

any PAH subject, suggesting there was no severe or complete obstruction of blood vessels. The only

confirmed responder of the PAH group is shown in the second row.

Figure 4: **Difference maps of the perfusion-height distribution visualizing the changes in mean**

normalized perfusion vs. height in response to the administration of O_2+iNO in PAH subjects and

controls. Note that all subjects had regional responses to breathing O2+iNO, including PAH subjects who

had no discernible change in mean pulmonary arterial pressures with vasodilator challenge during cardiac

catheterization. In controls, the changes in perfusion show a substantial spatial heterogeneity with an

overall trend towards an increased vertical gradient during O2+iNO and more uniform perfusion in the

dependent (dorsal) regions (see also Fig. 2). In contrast, there are substantial changes in regional

perfusion in PAH but not associated with a shift in the vertical gradient except for the confirmed

responder among the PAH subjects shown in the second row. All panels use the same scale greyscale for

the magnitude of differences.

Figure 5: 3D renderings and animations of perfusion distributions. (A) Representative 3D renderings of the mean-normalized distribution of perfusion for PAH and healthy control subjects at baseline and after O_2+iNO (animated in Videos V1 (https://doi.org/10.6084/m9.figshare.12442589) and V2 (https://doi.org/10.6084/m9.figshare.12442784)) showing the effects of the administration of O_2+iNO based on PET-CT imaging in supine position. The patchy areas of high perfusion illustrate the size and location of regional variations behind the representations in perfusion-height maps (Fig. 2). **(B)** 3D renderings of the regional changes in perfusion corresponding to the perfusion distributions in panel A(animated in Videos V3 (https://doi.org/10.6084/m9.figshare.12442787) and V4 (https://doi.org/10.6084/m9.figshare.12442790)). The web-like structure of perfusion increases (yellow) and decreases (blue) instead of gradual changes in perfusion among different regions illustrates the spatial variations behind the difference maps of the perfusion-height distribution (Fig. 3). The relatively small sizes suggest the involvement of smaller anatomical structures together with larger anatomical structures linked to the average changes in larger regions. In all 3D renderings, the orientation of the lungs is shown by a small mannequin. Color scales are fully transparent either for average perfusion (A, V1, and V2) or zero change (B, V3, and V4), opaqueness increases with the deviation from these values, and differences in color between PAH and controls are differences in magnitude. Rotating 3D renderings with periodic switching between air and O2+iNO show the location and changes in high and low perfusion areas. This provides a spatial perception of their location and their magnitude lacking in the still images – video download provides the highest resolution and allows playback in loop mode.

Figure 6: V̇/Q̇ distributions in PAH subjects and controls. The distributions of V̇/Q̇ ratios at baseline

in PAH subjects were overall broader than in controls at baseline, approaching statistical significance,

consistent with moderate degrees of vascular disease, causing a mismatch. There was no difference in the

V̇/Q̇ distribution among groups while breathing O_2+iNO, in large part due to the widening of the V̇/Q̇

distribution in controls. V̇/Q̇ distributions while breathing O_2+iNO were unimodal in all subjects,

including those with a bimodal distribution prior to the intervention.

Table 1 Baseline characteristics

Parameter	Control	PAH	p-value
	N=5	N=4	
Age (years)	41.2 (22-60)	47.3 (22-64)	0.64
Gender, Female	1	3	0.14
Race, White	5	3	0.39
Height (inches)	67.3 (63-71.5)	67.2 (66-68.5)	0.91
Weight (pounds)	189.6 (160-296.7)	170.5 (124-264)	0.67
Years of symptoms	NA	8.4 (7-10)	
FEV1 (%predicted)	NA	82.9 (66-95)	
FVC (% predicted)	NA	85.6 (64-110)	
FEV1/FVC	NA	80 (63-86)	
mPAP, rest (mmHg)	NA	41.6 (33-63)	
PVR, rest (dyn·s/cm^5)	NA	442 (322-548)	
PCWP, rest (mmHg)	NA	10 (7-16)	
6MWD (feet)	NA	1519.8 (1253-2198)	

Continuous variables are represented as mean (min-max). Categorical variables are represented as counts.

Definitions of abbreviations: PAH = primary pulmonary arterial hypertension; FEV_1 = forced expiratory volume in first second; FVC= forced vital capacity; mPAP = mean pulmonary arterial pressure; PVR = pulmonary vascular resistance; PCWP = pulmonary capillary wedge pressure; 6MWD = six-minute walk distance.

690 **Table 2 Vertical gradient and heterogeneity in perfusion**

Parameter	Control		PAH	
	Baseline	**O$_2$+iNO**	**Baseline**	**O$_2$+iNO**
Q$_{vgrad}$ (10^{-2} cm^{-1})	-5.46 (-3.32, -7.01)*[†]	-7.37 (-6.43, -8.66)*[‡]	-3.13 (-2.33, -4.78)[†]	-2.51 (-0.039, -0.431) [‡]
CV^2$_{Qtotal}$	0.09 (0.06, 0.16)	0.12 (0.08, 0.16)[‡]	0.08 (0.07, 0.11)	0.06 (0.04, 0.1)[‡]
CV^2$_{Qvgrad}$	0.04 (0.02, 0.08)[†]	0.08 (0.055, 0.10)[‡]	0.01 (0.0, 0.03)[†]	0.01 (0.0002, 0.02)[‡]
CV^2$_{Qr}$	0.04 (0.02, 0.07)	0.04 (0.02, 0.06)	0.05 (0.04, 0.06)	0.03 (0.02, 0.04)

Continuous variables are represented as mean (min, max).

Definitions of abbreviations: PAH = primary pulmonary arterial hypertension; Q$_{vgrad}$ = vertical gradient in regional perfusion (slope); CV^2$_{Qtotal}$ = squared coefficient of variation in perfusion quantifying total spatial heterogeneity of perfusion (noise corrected); CV^2$_{Qvgrad}$ = squared coefficient of variation in perfusion quantifying heterogeneity generated by the dorsal-ventral gradient in perfusion; CV^2$_{Qr}$ = squared coefficient of variation in perfusion quantifying heterogeneity of the residual perfusion.

* significant difference between measures at baseline and during administration of O$_2$+iNO within a group

[†] significant difference between PAH subjects and controls at baseline

[‡] significant difference between PAH subjects and controls while breathing O$_2$+iNO

691   **Table 3 Perfusion Distributions at all Length Scales at Baseline**

Parameter	Controls	PAH	p-value
$CV^2_{Q10\text{-}30}$ (x10^{-3})	13.0 (6.82, 29.77)	19.7 (17.17, 22.25)	0.33
$CV^2_{Q30\text{-}50}$ (x10^{-3})	7.77 (4.06, 16.70)	10.0 (8.05, 11.70)	0.68
$CV^2_{Q50\text{-}70}$ (x10^{-3})	4.46 (2.59, 8.15)	5.61 (4.48, 6.83)	0.45
$CV^2_{Q70\text{-}90}$ (x10^{-3})	2.52 (1.69, 3.57)	3.31 (2.54, 4.67)	0.20
$CV^2_{Q90\text{-}110}$ (x10^{-3})	1.48 (0.96, 2.10)	1.91 (1.41, 3.22)	0.33
$CV^2_{Q>110}$ (x10^{-3})	5.28 (1.19, 16.83)	5.81 (0.80, 21.04)	0.80

692   Definition of abbreviations: $CV^2_{Q(lengthscale)}$ = squared coefficient of variation in perfusion quantifying

693   spatial heterogeneity in regional perfusion in the denoted length scale range (mm)

694   **Table 4 Perfusion Distributions at all Length Scales at During O_2+iNO**

Parameter	Controls	PAH	p-value
$CV^2_{Q10\text{-}30}$ (x10^{-3})	14.4 (6.65, 24.32)	13.4 (8.24, 17.43)	0.62
$CV^2_{Q30\text{-}50}$ (x10^{-3})	8.43 (4.68, 12.80)	7.16 (4.66, 9.59)	0.46
$CV^2_{Q50\text{-}70}$ (x10^{-3})	4.62 (2.44, 6.07)	4.04 (2.50, 6.48)	0.68
$CV^2_{Q70\text{-}90}$ (x10^{-3})	2.61 (1.29, 3.26)	2.32 (1.48, 3.91)	0.72
$CV^2_{Q90\text{-}110}$ (x10^{-3})	1.51 (0.75, 1.98)	1.40 (1.04, 2.06)	0.70
$CV^2_{Q>110}$ (x10^{-3})	9.56 (5.46, 20.68)	4.66 (4.05, 5.23)	0.10

695   Definition of abbreviations: CV2Q(length scale) = squared coefficient of variation in perfusion

696   quantifying spatial heterogeneity in regional perfusion in the denoted length scale range (mm)

697   **Table 5 Regional F_{tis} and Ventilation parameters at baseline and during O_2+iNO**

Parameter	Control		PAH	
	Baseline	**O₂+iNO**	**Baseline**	**O₂+iNO**
F_{tis}	0.35 (0.31, 0.43)	0.39 (0.33, 0.49)	0.34 (0.27, 0.37)	0.37 (0.33, 0.41)
Volume (ml)	2107 (1521, 2646)	1940 (1421, 2347)	2223 (1676, 3209)	2147 (1282, 2997)
Mean s$\dot{V}$ ($10^{-3}s^{-1}$)	2.69 (1.49, 4.13)	2.92 (2.21, 3.51)	4.68 (1.82, 9.72)	3.33 (2.00, 5.91)
$\dot{V}_A$ (ml/min)	5.56 (3.73, 8.53)	5.55 (4.26, 6.72)	9.36 (4.22, 16.30)	6.31 (3.85, 7.84)
$CV^2_{\dot{V}A}$	0.19 (0.07, 0.33)	0.18 (0.12, 0.24)	0.32 (0.07, 0.54)	0.28 (0.09, 0.37)

Continuous variables are represented as mean (min, max).

Definitions of abbreviations: PAH = primary pulmonary arterial hypertension; F_{tis} = tissue fraction; Volume = gas volume; V_A = alveolar ventilation; s$\dot{V}$ = specific ventilation ($\dot{V}_A$ /alveolar volume); $CV^2_{\dot{V}A}$ = squared coefficient of variation quantifying heterogeneity in alveolar ventilation.

DECLARATIONS

Ethics approval and consent to participate

The study was approved by the Institutional Review Board of the Massachusetts General Hospital. Informed

consent was obtained from each subject before the study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests. For full disclosure, ASW has been consulting

for United Therapeutics in the past on topics unrelated to this study, and RC has or had consultant, research,

or speaker relationships with United Therapeutics, Janssen, Bayer, Gossamer, Altavant, Third Pole, and

Aria CV.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on

reasonable request.

Funding

Support for this research was provided by an unrestricted grant from United Therapeutics. The sponsor was

not involved in the study design, analysis, or reporting of study findings.

Author information

Tilo Winkler and Puja Kohli contributed equally as joint first authors.

Affiliations

The following is the list of author affiliations at the time the study was conducted. Changes are listed under acknowledgements.

Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

Tilo Winkler PhD, Jose G. Venegas PhD

Division of Pulmonary and Critical Care Medicine, Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts, USA

Puja Kohli MD MMSc, Vanessa J Kelly PhD, Ekaterina G. Kehl MD, Alison S. Witkin, MD, Josanna Rodriguez-Lopez MD, Kathryn A. Hibbert MD, Mamary Kone MD MPH, Richard Channick MD, R. Scott Harris MD

Division of Pulmonary and Critical Care Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA

David M. Systrom MD, Aaron B. Waxman MD

Contributions

Study design: MK, EGK, RC, RSH; data acquisition: TW, PK, VJK, EGK, JRL, KAH, MK, DMS, JGV, RC, RSH; data analysis: TW, PK, VJK, RSH; data interpretation: TW, PK, RSH; manuscript drafting and editing: PK, TW, VJK, RSH; all authors read and approved the final manuscript.

Corresponding author

Tilo Winkler, PhD – email: twinkler@mgh.harvard.edu

Massachusetts General Hospital

Dept of Anesthesia, Critical Care and Pain Medicine

55 Fruit Street, Boston, MA 02114

USA

Acknowledgements

The authors thank the staff from the Massachusetts General Hospital Departments of Nuclear Medicine
and Nuclear Pharmacy for their assistance with this project, including Steve Weise, Michael Cournoyer,
John Correia, Eugene Lee, Peter Rice, Melissa Bruen, Erin Beloin, Kevin Vernon and Daniel Yokell. In
addition, the authors extend kind thanks to Julian Fischer and Gabriel C. Motta-Ribeiro for their
contributions to the process of image registration, Rebecca Harris for her contributions to mask creation,
Benno F. Rodemann for his contribution to media creation and to Pradeep Natarajan MD (Massachusetts
General Hospital, Division of Cardiology) for helpful discussions and feedback on this work. The present
address of Richard Channick is David Geffen School of Medicine at UCLA, Los Angeles, California,
USA, and the address of Ekaterina G. Kehl is Division of Pulmonary and Critical Care Medicine, Mount
Auburn Hospital, and Harvard Medical School, Boston, Massachusetts, USA.

PAH vs. Control

Baseline

Q_{vgrad}

CV^2_{Qtotal}

CV^2_{Qvgrad}

CV^2_{Qr}

O2+iNO

Q_{vgrad}

CV^2_{Qtotal}

CV^2_{Qvgrad}

CV^2_{Qr}

O$_2$+iNO vs. Baseline

Control

Q_{vgrad}

CV^2_{Qtotal}

CV^2_{Qvgrad}

CV^2_{Qr}

PAH

Q_{vgrad}

CV^2_{Qtotal}

CV^2_{Qvgrad}

CV^2_{Qr}

-5   0   5

Cohen's d effect size

	Air	O$_2$+iNO	Mean absolute change