

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

Coping strategies employed by transgender youth with higher and lower quality of life.

Ítala Raymundo Chinazzo^{1*}¶, Anna Martha Vaitses Fontanari^{1¶}, Bruno de Brito Silva^{3&}, Lucca P. Rodrigues^{3&}, Angelo Brandelli Costa^{2¶}, Maria Inês Rodrigues Lobato^{1¶}

¹ Psychiatry Graduate Program, Hospital de Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

² Psychology Graduate Program, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil.

³Psychology Graduate Program, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil

* Corresponding author

E-mail: italach@gmail.com

¶These authors contributed equally to this work.

&These authors also contributed equally to this work.

25 Abstract

26 Transgender youth are especially susceptible and vulnerable to mental health concerns. Minority
27 Stress Theory may explain these mental health concerns and avoidance to seek healthcare
28 provision in this population. Understanding transgender youth adaptive and maladaptive coping
29 mechanisms may help create strategies that promote quality of life and reduce the negative
30 influence of stress on their mental health. Our study aims to measure the quality of life of Brazilian
31 binary and non-binary transgender youth and analyze the association with their coping strategies
32 used for dealing with general social and gender-related stress. Participants were recruited through
33 Facebook advertisement directed to young Brazilians, aged from 16 to 24 years, who “liked” pages
34 or joined groups related to LGBTQ+ movement. Gender identity was evaluated by the Two Steps
35 Method. Coping was evaluated by the Coping with Stress Checklist; and quality of life, by the
36 Quality of My Life questionnaire. The final sample consisted of 185 participants. Out of the total,
37 95 (46.34%) identified as transgender boys, 45 (21.95%) as transgender girls, and 65 (31.71%) as
38 non-binary youth. The mean age was 18.61 years (SD 2.43). The study found that 53.9% of the
39 sample had low self-perceived quality of life, 48.0% depression, and 68.3% anxiety. Better quality
40 of life was related to socialization ($p=0.02$), whereas a worse quality of life was related to
41 avoidance attitudes ($p=0.05$), concerning coping with general stress. The study found no
42 association between coping strategies for gender-related stress and quality of life. The Brazilian
43 transgender youth may need support to develop adaptive coping skills to deal with both general
44 and gender-related stress. Also, social interventions against prejudice are needed to achieve better
45 results in the quality of life for transgender youth. Mental health professionals should explore the
46 unique needs and stressors of transgender identity and explore adaptive coping strategies.

47
48 **Keywords (3-5):** transgender youth; non-binary youth; coping strategies; quality of life; mental
49 health.

51 Introduction

52 The term transgender denotes individuals whose gender identity does not align with their
53 sex assigned at birth, whereas the expression “cisgender” refers to individuals whose gender
54 identity matches their sex assigned at birth (Giblon & Bauer, 2017). Transgender is an umbrella
55 term that covers a broad continuum of gender identities, including binary and non-binary people.
56 While binary people identify either as a woman or as a man – regardless of whether they are
57 transgender or not – non-binary people recognize themselves as belonging to a third different

58 gender, having more than one gender, or as being agender (Bass et al., 2018). Considering that
59 non-binary people may seek gender-affirming care and that transgender people do not necessarily
60 identify within the gender binaries, there are numerous intersections between non-binary and
61 transgender. Nevertheless, since not all non-binary people identify as transgender, and vice versa,
62 using both expressions is necessary (Bass et al., 2018; Coleman et al., 2012; Richards et al., 2016).

63 Transgender adults face high rates of violence throughout Brazil and the world (Balzer,
64 Lagata, & Berredo, 2016; Benevides & Nogueira, 2019; Silva, 2018; Chinazzo et al. 2020), which
65 not only directly affects their physical and mental health (Magno, Dourado, & Silva, 2018; Zucchi,
66 2019), but also limits their access to health services (Rocon et al. 2016; Rodriguez, 2014; Sousa &
67 Iriart, 2018) and education (Associação Brasileira de Lésbicas, Gays, Bissexuais, Transexuais e
68 Travestis [ABGLT], 2016). For instance, longitudinal studies suggest that transgender people have
69 higher rates of completed suicide than their cisgender peers (Asscherman et al., 2011; Dhejne et
70 al., 2011), and also experience a lower quality of life (Newfield et al.2006; Jones et al., 2019).

71 Transgender binary and non-binary youth are especially susceptible and vulnerable to
72 mental health concerns (Chinazzo et al., 2020; Connolly et al., 2016; Dhejne et al., 2016; Perez-
73 Brumer et al., 2015). Transgender youth from the United States are between 3.2 and 3.6 times
74 more likely to have suicidal ideation and suicidal attempts, respectively, when compared to the
75 general population (Reisner et al., 2015). Furthermore, according to their parents or caregivers,
76 Canadian transgender youth talk 5.1 times more about suicide than their siblings (Aitken et al.,
77 2016). Regarding non-binary transgender youth, Thorne et al. (2019) found evidence of a
78 difference in their mental health (anxiety and depression) when compared with binary transgender
79 youth, whereas Rimes et al. (2019) found no apparent differences when compared with binary
80 transgender youth.

81 Mental health problems are associated with low quality of life in transgender people
82 (Bouman et al., 2016). Quality of life is defined and related to an “individual’s perception of their
83 position in life in the context of the culture and value systems in which they live and in relation to
84 their goals, expectations, standards and concerns” (World Health Organization [WHO], 2021).
85 Studies (Newfield et al., 2006; Nobili, Glazebrook & Arcelus, 2018) have reported that young
86 transgender individuals have a lower perceived quality of life compared to the general population.
87 Moreover, transgender youth have been associated with greater mental health problems (Arcelus
88 et al, 2016; Heylens et al., 2014) and a lower mental health-related quality of life (Newfield et al.,
89 2006). Additionally, the detrimental effects of body dissatisfaction in cisgender groups are
90 observed as a factor for an increasing risk of depression (Paxton et al., 2006), low self-esteem
91 (Davison & McCabe, 2006), and eating disorders (Stice & Shaw, 2002). Reports suggest that body
92 dissatisfaction may be higher among transgender people (Witcomb et al., 2015; Ålgars, Santilla,
93 & Sandnabba, 2010).

94 The Minority Stress Theory helps understanding these health disparities (Hendricks &
95 Testa, 2012; Meyer, 1995, 2003). In addition to the general stress of life, belonging to a minority
96 group increases chronic stress by enacted stigma, expectations of rejection, and internalization of
97 social stigma, resulting in psychological distress and avoidance to seek healthcare provision.
98 However, identification with a minority group may give access to community resources such as
99 support groups, community centers, minority role models, as well as affirmative actions and
100 specialized clinics (Meyer, 2015). Such resources are conceptualized as *minority coping*, which
101 can buffer the effect of minority-specific stressors on the mental health of transgender individuals
102 (DiFulvio, 2015; Stanton, Ali & Chaudhuri, 2017; Zucchi et al., 2019). Specifically, social support
103 from one’s community – a resource derived from minority coping – seems to positively affect

104 psychological well-being (Stanton, Ali, & Chaudhuri, 2017; Zucchi et al., 2019) as well as lessen
105 the consequences of stigma on mental health (DiFulvio, 2015).

106 Stress is the reaction to a situation in which a person’s physical or psychological resources
107 are challenged as aforementioned. Coping consists of the efforts that a person makes to adapt to
108 stress and to manage external and/or internal demands that exceed the resources of the person.
109 However, being able to cope does not necessarily translate as successfully adjusting. A person’s
110 reactions in stressful environments are dependent on the person’s appraisal of a given situation,
111 combined with the physical or psychological resources available (Lazarus & Folkman, 1984). The
112 evaluation of this mechanism is adaptive or non-adaptive, depending on the ability to buffer or
113 not the stresses they experience. Coping strategies can be learned, so it is important knowing which
114 mechanisms are adaptive and, therefore, protective factors for mental health in transgender youth.

115 A protective factor for transgender individuals is the available social support (DiFulvio,
116 2015; Singh, Hays, & Watson, 2011; Puckett et al., 2019; Stanton, Ali, & Chaudhuri, 2017; Zucchi
117 et al., 2019). This may be understood as a coping strategy for patients who have positive social
118 relations, termed as “minority coping.” Social support means the existence of meaningful relations
119 with people, in which there is trust, affection, and acceptance when sharing one’s life experience
120 and feelings. There are two types of social support—informal and formal. The source of informal
121 social support is the daily social network such as family members, friends, and social groups who
122 help in everyday activities; whereas formal social support is related to social assistance
123 organizations, professionals, and health services.

124

125 **Current Study**

126 Most studies investigating coping strategies have focused on gay, lesbian, and transgender
127 adults (Budge, Adelson, & Howard, 2013; Hughto, Pachankis, & Willie, 2017; Lehavot, 2012).
128 Therefore, it is essential to evaluate transgender youth. Better understating of transgender youth
129 coping mechanisms may help create strategies to promote well-being and reduce the negative
130 influence of stress on their mental health. Our study aims to describe the coping behaviors
131 employed by transgender youth who have a higher quality of life in comparison with those with a
132 lower quality of life.

133

134 **Methods**

135

136 **Study Sample**

137 From February to April 2018, participants were recruited via Facebook advertisements
138 directed towards Brazilian adolescents and young adults, aged 16–24, who “liked” pages or joined
139 groups related to the LGBTQ+ movement. Gender identity was evaluated by the Two Steps
140 Method (Bauer et al., 2017; Sausa et al., 2009). Firstly, sex at birth was assessed by the question:
141 “What sex were you assigned at birth on your original birth certificate?”. Participants could choose
142 between the “female” and “male” alternatives. Secondly, gender identity was evaluated as an open
143 question “How do you currently describe your gender?”. All participants responded within the
144 “woman,” “man,” and “non-binary” alternatives.

145 In total, 706 young people started the online protocol. Nevertheless, to guarantee the quality
146 of the data, participants who did not reply to the two-step questions as transgender and non-binary
147 were excluded. Likewise, duplicates and discrepant responses were removed. A total of 165
148 individuals were excluded because they did not complete the Coping with Stress Checklist. The

149 final sample was composed of 185 participants in the data analysis. Fig 1 summarizes the
150 participants' flow.

151 **Fig 1.** Participants' sampling flowchart.

152

153 **Measures**

154 The survey was inspired by the TransYouth CAN! Project (see <http://transyouthcan.ca/>).
155 The cross-cultural adaptation for the Brazilian populations was based on the work by Borsa,
156 Damásio, and Bandeira (2012); it is better described in previous publications (Fontanari et al.,
157 2019).

158 Socioeconomic Status (SES) was measured using the Deprivation Scale, created by the
159 "TransYouth CAN! Team" to access the information about the basic needs of our sample. The
160 Deprivation Scale consisted of five questions asking whether participants had access to school
161 supplies, internet, proper seasonal clothing, other essential clothing, and trustworthy transportation
162 in a one-year period. Each question could be responded using a five-point Likert scale ranging
163 from "never" (one point) to "always" (five points), in which a higher score represents less
164 deprivation. The score was calculated by the arithmetical mean of the five questions (mean = 4.31;
165 SD = 0.72); the Cronbach's alpha coefficient was 0.820.

166 Coping with Stress Checklist, developed by the TransYouth CAN! Team is a questionnaire
167 that assumes that young transgender people deal with a two-tier "stress situation." The first one is
168 the stress that happens in everyday life, identified as General Stress; and the second one is
169 identified as Gender-Related Stress linked to their transgender condition. The questionnaire is
170 composed of a 26-item list containing adaptive and maladaptive coping strategies, which assess
171 the coping style and how each person deals with the General Stress and Gender Related Stress in

172 a 30-day period. The 26-items were divided by the research team into four dimensions for each
173 type of stress: 1. hobby strategies (watching movies, TV or on-line videos, playing videogame,
174 and reading), 2. socialization (talking with LGBT and non-LGBT friends, spending time with
175 family, volunteer work, spending time in social events, and publishing about themselves on on-
176 line social networks), 3. introspection (physical exercises, deep breathing/relaxation exercises,
177 writing a diary, praying, meditation) and 4. avoidance (excessive sleeping, eating, shopping,
178 cosmetic self-care, day dreaming, looking for noncommitted partners, and ignoring or pretending
179 that the stress is not real). For each strategy, participants could choose between “yes” and “no”
180 alternatives. Cronbach’s alpha value of general stress is 0.74, and gender-related stress is 0.88.

181 To assess quality of life, the first item of the Quality of My Life (QoML) Questionnaire
182 states “overall, my life is...” and is answered with a visual analog scale (Bowling, 2005; Feldman
183 et al., 2000). There are five response options on this scale: excellent, very good, good, fair, or poor.
184 Responses were dichotomized into “poor” and “excellent” quality of life, using the median.

185 The Overall Anxiety Severity and Impairment Scale (OASIS) (Norman et al., 2006) is a
186 brief continuous measure of five questions about the level of anxiety and prejudice that participants
187 felt in a seven-day period. Each question was measured on a five-point Likert scale, ranging from
188 0 (no anxiety or impairment from anxiety) to 4 (constant anxiety or impairment from anxiety). The
189 responses were dichotomized and the cut-off point was 8, in which participants who scored more
190 than 8 were more likely to have an anxiety diagnosis.

191 The Modified Depression Scale (MDS) (Dunn, Johnson & Green, 2012) was used to assess
192 the frequency of five depressive symptoms in a 30-day period: “in the last 30 days how often have
193 you... (1) felt very sad, (2) felt very angry or in a bad mood, (3) felt hopeless about the future, (4)
194 slept much more or much less than usual, and (5) had difficulty concentrating on your studies.”

195 Each question was answered on a five-point Likert scale: 1 (never), 2 (rarely), 3 (sometimes), 4
196 (often), and 5 (always). The responses were dichotomized according to the cut-off point of 17, in
197 which participants who scored more than 17 were more likely to have a diagnosis of depression.

198 To evaluate gender dysphoria, a version of the Utrecht Gender Dysphoria Scale was used
199 – adapted by the Trans Youth Can! Team, in which some items were added or deleted, making it
200 more appropriate for non-binary persons. The scale consisted of 15 items, covering dysphoria
201 related to the body (e.g., I wish I had been born in a different body) and social life (e.g., it bothers
202 me to be called by the wrong gender). The items were answered on a five-point Likert scale,
203 ranging from “totally disagree” to “totally agree,” plus the answer option “does not apply to me.”
204 The score was obtained by summing up all items; for analysis the answers were divided into
205 whether the participants identified as being without or almost without gender dysphoria, or with
206 gender dysphoria or with a lot of gender dysphoria.

207

208 **Statistical Analysis**

209 The software SPSS 18.0 was used to perform statistical analysis. Firstly, background
210 characteristics were compared between those reporting in QoML, which was divided into poor and
211 excellent life quality. Secondly, a Poisson Regression in two steps was applied to compare the four
212 dimensions of coping strategy, from the Coping with Stress Checklist, with Quality of My Life,
213 controlled by sociodemographic data. In the first step of the logistic regression, only the four
214 dimensions of coping with stress checklist were included. In the second step, the variables age,
215 SES, anxiety, depression, and gender dysphoria were also included. Both steps were performed
216 for coping strategies with general stress and stress related to gender identity.

217

218 **Ethical Considerations**

219 The Ethical Committee and Research Commission of the Federal University of Rio Grande
220 do Sul Psychology Institute approved the project (14221513.4.0000.5334). All participants signed
221 an online ethical consent form; for participants aged under 18 years, a consent term was provided
222 by their guardians.

223

224 **Results**

225 For this study, 185 participants completed the Coping with Stress Checklist. Among them,
226 90 (48.6%) identified as transgender boys, 37 (20.0%) as transgender girls, and 58 (31.4%) as non-
227 binary persons (Table 1). The mean age was 18.50 years (SD 2.44) and youth with an average
228 quality of life did not differ in sociodemographic characteristics.

229

230 **Table 1. Sample description**

		N	%
Gender identity	Male	90	48.6
	Female	37	20.0
	Non-binary	58	31.4
Education	School	70	37.8
	College preparatory programs	13	7.0
	Technical course	13	7.0
	College	50	27.0
	Not studying	39	21.1
State	São Paulo	51	28.5
	Rio Grande do Sul	37	20.7
	Rio de Janeiro	19	10.6
	Minas Gerais	15	8.4

	Other	57	31.8
Time living in perceived gender	All the time	86	46.5
	Half the time	68	36.8
	Never	31	16.8
Legal identity transition	Social card	14	7.6
	Identity card	2	1.1
	Social and identity card	2	1.1
	It is in process	88	47.6
	Does not plan to change name	18	9.7
	Prefer not to answer	61	33.0
	Public knowledge of your transgender identity	Always	18
Almost always		32	17.3
About half the time		34	18.4
Rarely		58	31.4
Never		23	12.4
I do not know		19	10.3
Hormone use	Yes, regularly	35	19.0
	Yes, once or twice	14	7.6
	Never	125	67.9
	I do not know / I would rather not answer	10	5.5
Hormone use over time	I am not looking for hormonal treatment	85	46.2
	< 1 year	56	30.4
	1 - 2 years	22	12.0
	2 - 3 years	14	7.6
	> 3 years	7	3.8

231

232

233 In relation to the coping strategies used, higher means were perceived in relation to general

234 stress rather than stress related to gender identity (Table 2). Considering the total score of the

235 coping scale (sum), there is no significant relationship with quality of life, both for general stress
 236 and for stress related to gender identity.

237 **Table 2. Mean (SD) coping general stress and gender-related stress**

		Mean (SD)
Coping general stress	Hobby ^a	3.09 (0.09)
	Socialization ^b	3.08 (0.13)
	Introspection ^c	2.80 (0.12)
	Avoidance ^d	3.29 (0.35)
Coping gender-related stress	Hobby ^a	1.22 (0.11)
	Socialization ^b	1.61 (0.12)
	Introspection ^c	1.42 (0.11)
	Avoidance ^d	1.76 (0.13)

238 ^aTotal of 5 strategies.

239 ^bTotal of 8 strategies.

240 ^cTotal of 7 strategies.

241 ^dTotal of 6 strategies.

242
 243
 244 Regarding quality of life, 53.9% of the sample was below the median, representing low
 245 QoL (Table 3). As for the mental health variables, the sample showed 48.0% with significant
 246 presence of depressive symptoms, 68.3% had significant symptoms of anxiety and 49.6% had
 247 some degree of gender dysphoria (GD) – “with GD” and “with a lot of GD”.

248 **Table 3. Mental health variables**

		N	%
Quality of life	Good	95	46.1
	Bad	111	53.9
Depression	Yes	94	48.0
	No	102	52.0
Anxiety	Yes	129	68.3
	No	60	31.7

Gender dysphoria (GD)	Without or almost without GD	174	50.4
	With GD	98	28.4
	With a lot of GD	73	21.2

249
250

251 When grouped together, avoidance strategies were associated with a worse quality of life,
252 in relation to general stress; whereas the other dimensions (hobby, socialization, and introspection)
253 were associated with a better quality of life (Table 4). There was no significant association between
254 coping strategies for stress related to gender identity and quality of life.

255 **Table 4. Correlation quality of life and coping general stress and gender-related stress**

		p
Coping general stress	Hobby	0.036 ^a
	Socialization	0.170 ^b
	Introspection	0.177 ^b
	Avoidance	-0.194 ^a
Coping gender-related stress	Hobby	0.015
	Socialization	-0.001
	Introspection	0.043
	Avoidance	-0.102

256 ^ap<0.001.
257 ^bp<0.05.
258
259

260 Two Poisson regressions were performed in two steps with the quality of life outcome. The
261 first analysis concerned coping strategies with general stress and the second, with coping strategies
262 for stress related to gender identity (Table 5). In the first step, only the four dimensions of coping
263 strategies were used. In the second step, for both analyses, the variables for depressive symptoms,
264 anxiety symptoms, gender dysphoria, age, and socioeconomic status were included.

265 In the first Poisson regression, a worse quality of life was related to avoidance strategies in
 266 dealing with general stress, whereas socialization coping strategies were associated with a better
 267 quality of life. Both relationships were maintained when the other predictor variables were added
 268 to the analysis. In the second step, low socioeconomic status and the presence of anxiety and
 269 depression symptoms were significantly associated with a worse quality of life.

270 **Table 5. Poisson regression quality of life and coping general stress**

Characteristic		Step 1		Step 2	
		PR (95% CI)	p	PR (95% CI)	p
Coping general stress	Hobby	1.04 (0.92; 1.18)	0.50	1.04 (0.93; 1.16)	0.51
	Socialization	0.90 (0.82; 0.97)	0.01 ^b	0.91 (0.83; 0.99)	0.02 ^b
	Introspection	0.92 (0.84; 1.01)	0.09	0.99 (0.91; 1.08)	0.87
	Avoidance	1.23 (1.11; 1.36)	0.00 ^a	1.14 (1.00; 1.30)	0.05 ^b
Gender dysphoria (GD)	With a lot of GD	-	-	1.29 (0.92; 1.81)	0.14
	With some GD	-	-	1.10 (0.81; 1.50)	0.53
	Without or almost without GD	-	-	-	-
Anxiety	Yes	-	-	2.53 (1.41; 4.51)	0.00 ^a
	No	-	-	-	-
Depression	Yes	-	-	1.78 (1.26; 2.52)	0.00 ^a
	No	-	-	-	-
Age		-	-	0.96 (0.91; 1.01)	0.15
Deprivation		-	-	0.80 (0.69; 0.94)	0.00 ^a

271 Abbreviation: PR = Poisson regression.

272 ^ap<0.001.

273 ^bp<0.05.

274

275

276 In the second Poisson regression, there was a significant association between a worse
 277 quality of life and avoidance strategies for coping with gender-related stress (Table 6). This
 278 relationship was not statistically significant when the other predictor variables were added. In the
 279 second step, low socioeconomic status and the presence of anxiety and depression symptoms were
 280 significantly associated with a poorer quality of life.

281 **Table 6. Poisson Regression quality of life and coping gender-related stress**

Characteristic		Step 1		Step 2	
		PR (95% CI)	p	PR (95% CI)	p
Coping gender-related stress	Hobby	0.92 (0.81; 1.05)	0.23	1.00 (0.90; 1.11)	0.98
	Socialization	0.99 (0.89; 1.10)	0.88	1.01 (0.92; 1.11)	0.87
	Introspection	0.95 (0.85; 1.07)	0.38	0.95 (0.85; 1.07)	0.44

	Avoidance	1.15 (1.03; 1.30)	0.01 ^b	1.01 (0.91; 1.12)	0.84
Gender dysphoria	With a lot of GD	-	-	1.27 (0.91; 1.79)	0.16
	With some GD	-	-	1.13 (0.84; 1.53)	0.42
	Without or almost without GD	-	-	-	-
Anxiety	Yes	-	-	2.56 (1.38; 4.74)	0.00 ^a
	No	-	-	-	-
Depression	Yes	-	-	1.85 (1.27; 2.68)	0.00 ^a
	No	-	-	-	-
Age		-	-	0.97 (0.91; 1.03)	0.27
Deprivation		-	-	0.79 (0.68; 0.93)	0.00 ^a

282 Abbreviation: PR = Poisson regression.

283 ^ap<0.001.

284 ^bp<0.05.

285

286

287 Discussion

288 This study highlights the challenges that Brazilian transgender youth face when dealing
 289 with the stress of everyday life and of social prejudice due to their gender status. Our study
 290 confirmed that more than half of the sample showed a self-perceived low quality of life (53.9%).
 291 Other studies have shown similar results (Jones et al., 2019; Newfield et al., 2006; Nobili et al.,
 292 2018). Despite being a subjective concept that involves different social and individual values, this
 293 self-perceived low quality of life raises questions concerning the main challenges that transgender
 294 youth face and what we must do, as a society, to minimize their suffering. Despite the social
 295 advances of recent years regarding sexuality, identifying as transgender is still a stressful and
 296 dysfunctional condition.

297 In accordance with other studies (Budge, Adelson, & Howard, 2013; Lewis et al, 2019;
 298 Müller et al., 2019), we found high rates of depressive symptoms (48.0%) and anxiety symptoms
 299 (68.3%). As predicted, these mental conditions were significantly associated with poor self-
 300 perceived quality of life, in addition to coping with general stress and gender-related stress.
 301 Generally, the Brazilian population shows a higher prevalence of depression (5.8%) and anxiety
 302 (9.3%) disorders (WHO, 2017) among South Americans countries and a probable lower quality of

303 life; the worst prevalence of mental conditions, however, occurs within the transgender youth
304 population. The Brazilian study with trans and gender non-conforming (TGNC) adults found high
305 rates of depressive symptoms (67.2%), indicating that suffering may persist throughout adult life
306 (Chinazzo et al., 2021). Another predictive variable associated with low quality of life in both
307 regression analysis was deprivation, that may harm any population enduring difficult social
308 conditions. Socioeconomical factors are associated with mental health and low quality of life,
309 especially in a stigmatized population.

310 Quality of life from a personal perspective can be improved with appropriate coping
311 strategies to buffer the general stressors of life. Coping strategies can be learned and developed
312 and the concept of adaptive and maladaptive strategies depends on the impact on well-being and
313 mental health of each individual (Lazarus & Folkman, 1984). Transgender people may need more
314 specific coping strategies. A central aspect for our research was to look at strategies that may be
315 adaptive or not for Brazilian transgender youth. In this study, the coping strategies were grouped
316 into four clusters: hobby, socialization, introspection, and avoidance and two types of stress:
317 general and gender-related. These two types of stress were analyzed to understand specific
318 transgender stressors in relation to quality of life in comparison with the general stresses of life
319 that they also experience, and to know which adaptive coping strategies may buffer the
320 dysfunctional consequences of adverse stressful events.

321 Our final findings concerning coping strategies for general stress indicated a significant
322 association between a lower quality of life with less use of socialization ($p=0.02$) and more use of
323 avoidance ($p=0.05$). Socialization coping strategies such as passing time with family, friends, and
324 volunteer work was associated with better self-perception and quality of life, which may indicate

325 that these social relations are positive and protective for transgender youth. Nevertheless, this
326 variable was not related to public health services support.

327 In another study published by our group (Chinazzo et al., 2021), we found that there was
328 an association between supportive affective relationships with a diminished presence of depressive
329 symptoms and suicidal ideation in transgender Brazilian adults. Despite the social difficulties
330 related to prejudice faced by transgender individuals, in our sample we identified socialization as
331 an adaptative coping strategy for general stress. This finding highlights the need for a special social
332 resource and strategies to reduce the chances of serious mental health issues in transgender youth.

333 Avoidance coping strategies for general stress were related with a low self-perceived
334 quality of life. Despite the possible passing pleasure associated with these coping strategies
335 (excessive sleeping, eating, shopping, and superfluous self-care), these methods of coping draw
336 them away from a more mature solution to their challenges in dealing with general stress and as a
337 result, compromise their mental health. This result implies that clinics working with transgender
338 should work on diminishing these maladaptive strategies. The use of maladaptive coping strategies
339 has been previously reported in transgender populations and is implicated in contributing to serious
340 mental health issues (Freese et al., 2017; Hughto et al., 2017).

341 Surprisingly, our study did not find a significant association between quality of life and the
342 four clusters of coping mechanisms to deal with gender-related stress. In the first step of the
343 regression analysis, the use of avoidance coping strategies was significantly associated with a low
344 perceived quality of life ($p=0.01$). However, this association is not apparent in the second step, in
345 which other variables were also included.

346 Avoidance strategy is associated with social isolation, marginalization, and failure to find
347 personal solutions; as cited before, this strategy may lead to anxiety, depression, suicidal behavior,

348 and, as evidenced by our study, poor perceived quality of life. The Brazilian social environment is
349 still full of different expressions of prejudice and aggressive attitudes directed towards transgender
350 people. It is rational to think that avoidance coping strategies could be considered as “protective”
351 from some points of view, as suggested by the Minority Stress Theory. It is a challenge for medical
352 assistance teams to help change this “safe” coping strategy for a more “audacious one.” For these
353 changes in coping attitudes to occur, the transgender youth will need to develop emotional strength
354 and self-assurance, aspects uncommon in many youths.

355 In our study we found a low average use of coping mechanisms when the four clusters and
356 gender-related stress were considered. This low average will need to be studied in the future to
357 better understand the coping mechanisms used by transgender youth in Brazil. Also, a further study
358 should explore the relation between quality of life, coping, and gender-related stress in the
359 transgender population.

360

361 **Limitations**

362 To the best of our knowledge, this is the first study with Brazilian transgender youth from
363 different regions of the country, which includes quality of life and coping strategies for general
364 and gender-related stress. There were some statistical limitations in generalizing the results found
365 in the Brazilian binary and non-binary transgender youth population due to the relatively small
366 sample studied. Nevertheless, the on-line survey gave access to many regions of Brazil which
367 allowed for a heterogeneous sample, albeit a small one.

368 Our study included binary and non-binary transgender youth. However, the sample was
369 not enough to calculate the difference between the two groups, binary and non-binary. Although
370 both groups are included under the transgender umbrella, there is evidence that there may exist

371 different stressors, such as quality of life and mental health issues (Jones, et al., 2019). To better
372 support both groups, we suggest more studies in this area.

373

374 **Acknowledgments**

375 We thank the Health Equity and Epidemiology Research Group (HEER) for multiple, and
376 very useful, insights. We also appreciate TransYouth CAN! Team for sharing their survey.
377 National Council for Scientific and Technological Development (CNPq) and Coordination for the
378 Improvement of Higher Education Personnel (CAPES) provided financial support through
379 scholarships. The authors thank the Research Incentive Fund of the Hospital da Clínicas de Porto
380 Alegre (FIPE/HCPA) for the financial support.

381

382 **References**


- 383 1. Giblon R, Bauer GR. Health care availability, quality, and unmet need: A comparison of
384 transgender and cisgender residents of Ontario, Canada. *BMC Health Services Research*
385 2017;1–10. doi: 10.1186/s12913-017-2226-z
- 386 2. Bass M, Gonzalez LJ, Colip L, Sharon N, Conklin J. Rethinking gender: The nonbinary
387 approach. *American Journal of Health-System Pharmacy* 2018;75,709–711. doi:
388 10.2146/ajhp180236
- 389 3. Coleman E, Bockting W, Botzer M, Cohen-Kettenis P, DeCuypere G, Feldman J, et al.
390 Standards of Care for the Health of Transsexual, Transgender, and Gender-Nonconforming
391 People, Version 7. *International Journal of Transgenderism* 2012;13(4), 165–232. doi:
392 10.1080/15532739.2011.700873
- 393 4. Richards C, Bouman WP, Seal L, Barker MJ, Nieder TO, T’Sjoen G. Non-binary or
394 genderqueer genders. *International Review of Psychiatry* 2016;28(1), 95–102. doi:
395 10.3109/09540261.2015.1106446
- 396 5. Balzer C, Lagata C, Berredo L. 2,190 murders are only the tip of the iceberg – An
397 introduction to the Trans Murder Monitoring project TMM annual report 2016. *TvT*
398 *Publication Series* 2016;14(14), 1–28. Available from: **Erro! A referência de hiperlink**
399 **não é válida.**
- 400 6. Benevides B. Dossiê dos assassinatos e da violência contra travestis e transexuais no Brasil
401 em 2018. Brasília: Distrito Drag, ANTRA; 2022.
- 402 7. Ministério dos Direitos Humanos. Secretaria Nacional de Cidadania. Violência
403 LGTBfóbicas no Brasil: dados da violência. Brasília: Ministério dos Direitos Humanos;

- 404 2018.
- 405 8. Chinazzo IR, Lobato MIR, Nardi HC, Koller SH, Saadeh A, Costa AB. Impact of minority
406 stress in depressive symptoms, suicide idea and suicide attempt in trans people. *Ciência &*
407 *Saúde Coletiva* 2021;26(suppl 3). doi: 10.1590/1413-812320212611.3.28532019
- 408 9. Magno L, Dourado I, Da Silva LAV. Stigma and resistance among travestis and transsexual
409 women in Salvador, Bahia State, Brazil. *Cadernos de Saude Publica* 2018;34(5). doi:
410 10.1590/0102-311x00135917
- 411 10. Zucchi EM, Barros CRS, Redoschi BRL, Deus LFA, Veras MASM. Bem-estar psicológico
412 entre travestis e mulheres transexuais no Estado de São Paulo, Brasil. *Cadernos de Saúde*
413 *Pública* 2019;35(3), e00064618. Epub March 25. doi: 10.1590/0102-311x00064618
- 414 11. Rocon PC, Rodrigues A, Zamboni J, Pedrini MD. Dificuldades vividas por pessoas trans
415 no acesso ao Sistema Único de Saúde. *Ciência & Saúde Coletiva* 2016;21(8), 2517–2526.
416 doi: 10.1590/1413-81232015218.14362015
- 417 12. Rodriguez AMM. Experiências de atenção à saúde e percepções das pessoas transgênero,
418 transexuais e travestis sobre os serviços públicos de saúde em Florianópolis/SC. 2014.
- 419 13. Sousa D, Iriart J. “Viver dignamente”: necessidades e demandas de saúde de homens trans
420 em Salvador, Bahia, Brasil. *Cadernos de Saúde Pública* 2018;34(10). doi: 10.1590/0102-
421 311x00036318
- 422 14. Associação Brasileira de Lésbicas, Gays, Bissexuais, travestis e transexuais [ABGLT]. As
423 experiências de adolescentes e jovens lésbicas, gays, bissexuais, travestis e transexuais em
424 nossos ambientes educacionais. In *Pesquisa nacional sobre o ambiente educacional no*
425 *Brasil*. 2016.
- 426 15. Asscheman H, Giltay EJ, Megens JA, de Ronde WP, van Trotsenburg MA, Gooren LJ. A
427 long-term follow-up study of mortality in transsexuals receiving treatment with cross-sex
428 hormones. *Eur J Endocrinol* 2011;164(4), 635–642. doi: 10.1530/eje-10-1038
- 429 16. Dhejne C, Lichtenstein P, Boman M, Johansson ALV, Langstrom N, Landén M. Long-
430 term follow-up of transsexual persons undergoing sex reassignment surgery: Cohort study
431 in Sweden. *Plos One* 2011;6(2), e16885. doi: 10.1371/journal.pone.0016885
- 432 17. Newfield E, Hart S, Dibble S, Kohler L. Female-to-male transgender quality of life. *Quality*
433 *of Life Research* 2006;15, 1447-457. Doi: 10.1007/s11136-006-0002-3
- 434 18. Jones BA, Bouman WP, Haycraft E, Arcelus J. Mental health and quality of life in non-
435 binary transgender adults: a case control study. *International Journal of Transgenderism*
436 2019;20(2-3), 251-262. doi: 10.1080/15532739.2019.1630346
- 437 19. Connolly MD, Zervos MJ, Barone CJ, Johnson CC, Joseph CLM. The mental health of
438 transgender youth: Advances in understanding. *Journal of Adolescent Health* 2016;59(5),
439 489–495. doi: 10.1016/j.jadohealth.2016.06.012
- 440 20. Dhejne C, Van Vlerken R, Heylens G, Arcelus J. Mental health and gender dysphoria: A
441 review of the literature. *International Review of Psychiatry* 2016;28(1), 44–57. doi:
442 10.3109/09540261.2015.1115753
- 443 21. Perez-Brumer A, Hatzenbuehler ML, Oldenburg CE, Bockting W. Individual- and
444 Structural-Level Risk Factors for Suicide Attempts among Transgender Adults. *Behavioral*
445 *Medicine* 2015;41(3),164–171. doi: 10.1080/08964289.2015.1028322
- 446 22. Reisner SL, Veters R, Leclerc M, Zaslow S, Wolfrum S, Shumer D, Mimiaga MJ. Mental
447 health of transgender youth in care at an adolescent Urban community health center: A
448 matched retrospective cohort study. *Journal of Adolescent Health* 2015;56(3), 274–279.
449 doi: 10.1016/j.jadohealth.2014.10.264

- 450 23. Aitken M, Vanderlaan DP, Wasserman L, Stojanovski S, Zucker KJ. Self-harm and
451 suicidality in children referred for gender dysphoria. *Journal of the American Academy of*
452 *Child & Adolescent Psychiatry* 2016. doi: 10.1016/j.jaac.2016.04.001
- 453 24. Thorne N, Witcomb GL, Nieder T, Nixon E, Yip A, Arcelus J. A comparison of mental
454 health symptomatology and levels of social support in young treatment seeking transgender
455 individuals who identify as binary and non-binary. *International Journal of Transgenderism*
456 2019;20(2-3),241-250. doi: 10.1080/15532739.2018.1452660
- 457 25. Rimes KA, Goodship N, Ussher G, Baker D, West E. Non-binary and binary transgender
458 youth: Comparison of mental health, self-harm, suicidality, substance use and
459 victimization experiences. *International Journal of Transgenderism* 2019;20(2-3),230-240.
460 doi: 10.1080/15532739.2017.1370627
- 461 26. Bouman WP, Davey A, Meyer C, Witcomb GL, Arcelus J. Predictors of psychological
462 well-being among treatment seeking transgender individuals. *Sexual and Relationship*
463 *Therapy* 2016;31(3),359-375.
- 464 27. World Health Organization. WHOQOL: Measuring Quality of Life. 2021. Available from:
465 <https://www.who.int/tools/whoqol>
- 466 28. Nobili A, Glazebrook C, Arcelus J. Quality of life of treatment-seeking transgender adults:
467 A systematic review and meta-analysis. *Reviews in Endocrine and Metabolic Disorders*
468 2018;19,199–220. doi: 10.1007/s11154-018-9459-y
- 469 29. Arcelus J, Claes L, Witcomb G, Marshall E, Bouman WP. Risk factors for non suicidal
470 self injury among trans youth. *The Journal of Sexual Medicine* 2016;13(3),402-412.
- 471 30. Heylens G, Elaut E, Kreukels BP, Paap MC, Cerwenka S, Richter-Appelt H, et al.
472 Psychiatric characteristics in transsexual individuals: multicentre study in four European
473 countries. *The British Journal of Psychiatry* 2014;204(2),151-156.
- 474 31. Paxton SJ, Neumark-Sztainer D, Hannan PJ, Eisenberg ME. Body dissatisfaction
475 prospectively predicts depressive mood and low self-esteem in adolescent girls and boys.
476 *Journal of clinical child and adolescent psychology* 2006;35(4),539-549.
- 477 32. Davison TE, McCabe MP. Adolescent body image and psychosocial functioning. *The*
478 *Journal of social psychology* 2006;146(1),15-30.
- 479 33. Stice E, Shaw HE. Role of body dissatisfaction in the onset and maintenance of eating
480 pathology: A synthesis of research findings. *Journal of psychosomatic research*
481 2002;53(5),985-993.
- 482 34. Witcomb GL, Bouman WP, Brewin N, Richards C, Fernandez-Aranda F, Arcelus J. Body
483 image dissatisfaction and eating-related psychopathology in trans individuals: A matched
484 control study. *European Eating Disorders Review* 2015;23(4),287-293.
- 485 35. Ålgars M, Santilla P, Sandnabba NK. Conflicted gender identity, body dissatisfaction, and
486 disordered eating in adult men and women. *Sex Roles* 2010;63,118–125.
- 487 36. Hendricks ML, Testa RJA. conceptual framework for clinical work with transgender and
488 gender nonconforming clients: An adaptation of the minority stress model. *Professional*
489 *Psychology: Research and Practice* 2012;43(5),460–467. doi: 10.1037/a0029597
- 490 37. Meyer IH. Minority stress and mental-health in gay men. *Journal of Health and Social*
491 *Behavior* 1995;36(1),38–56. doi: 10.2307/2137286
- 492 38. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual
493 populations: Conceptual issues and research evidence. *Psychological Bulletin*
494 2003;129(5),674–697. doi: 10.1037/0033-2909.129.5.674
- 495 39. Meyer IH. Resilience in the study of minority stress and health of sexual and gender

- 496 minorities. *Psychology of Sexual Orientation and Gender Diversity* 2015;2(3),209–213.
- 497 40. DiFulvio GT. Experiencing Violence and Enacting Resilience: The Case Story of a
498 Transgender Youth. *Violence Against Women* 2015;21(11),1385–1405. doi:
499 10.1177/1077801214545022
- 500 41. Stanton MC, Ali S, Chaudhuri S. Individual, social and community-level predictors of
501 wellbeing in a US sample of transgender and gender non-conforming individuals. *Culture,*
502 *health & sexuality* 2017;19(1),32-49.
- 503 42. Lazarus RS, Folkman S. *Stress, appraisal, and coping*. Springer publishing company, 1984.
- 504 43. Singh AA, Hays DG, Watson LS. Strength in the Face of Adversity: Resilience Strategies
505 of Transgender Individuals. *Journal of Counseling & Development* 2011;89:20-27.
506 doi:10.1002/j.1556-6678.2011.tb00057.x
- 507 44. Puckett JA, Matsuno E, Dyar C, Mustanski B, Newcomb ME. Mental health and resilience
508 in transgender individuals: What type of support makes a difference? *Journal of Family*
509 *Psychology* 2019;33(8):954–964. doi: 10.1037/fam0000561
- 510 45. Budge SL, Adelson JL, Howard KA. Anxiety and depression in transgender individuals:
511 the roles of transition status, loss, social support, and coping. *Journal of consulting and*
512 *clinical psychology* 2013;81(3):545.
- 513 46. Hughto J, Pachankis J, Willie T, Reisner S. Victimization and depressive symptomology
514 in transgender adults: The mediating role of avoidant coping. *Journal of Counseling*
515 *Psychology* 2017;64:41-51. doi: 10.1037/cou0000184.
- 516 47. Lehavot K. Coping strategies and health in a national sample of sexual minority women.
517 *American Journal of Orthopsychiatry* 2012;82(4):494.
- 518 48. Bauer GR, Braimoh J, Scheim AI, Dharma C. Transgender-inclusive measures of
519 sex/gender for population surveys: Mixed-methods evaluation and recommendations. *Plos*
520 *One* 2017;12(5):1–28. doi: 10.1371/journal.pone.0178043
- 521 49. Sausa LA, Sevelius J, Keatley J, Iñiguez JR, Reyes M. Policy recommendations for
522 inclusive data collection of trans people in HIV prevention, care & services. Center of
523 Excellence for Transgender HIV Prevention: University of California, San Francisco,
524 2009.
- 525 50. Borsa JC, Damásio BF, Bandeira DR. Cross-cultural adaptation and validation of
526 psychological instruments: Some considerations. *Padeia* 2012;22(53):423–432.
527 doi:10.1590/1982-43272253201314
- 528 51. Fontanari AMV, Pase PF, Churchill S, Soll B, Schwarz K, Schneider MA, et al. Dealing
529 with gender-related and general stress: Substance use among Brazilian transgender youth.
530 *Addictive Behaviors Reports* 2019;9:100166.
- 531 52. Bowling A. Just one question: If one question works, why ask several? *Journal of*
532 *Epidemiology & Community Health* 2005;59:345–346.
- 533 53. Feldman BM, Grundland B, McCullough L, Wright V. Distinction of quality of life, health
534 related quality of life, and health status in children referred for rheumatologic care. *The*
535 *Journal of Rheumatology* 2000;27(1):226–233.
- 536 54. Norman SB, Hami Cissell S, Means-Christensen AJ, Stein MB. Development and
537 validation of an overall anxiety severity and impairment scale (OASIS). *Depression and*
538 *anxiety* 2006;23(4):245-249.
- 539 55. Dunn EC, Avenue H, Green JG. The Modified Depression Scale (MDS): A brief, no-cost
540 assessment tool to estimate the level of depressive symptoms in students and schools.
541 *School Mental Health* 2012;4(1):34–45. doi: 10.1007/s12310-011-9066-5

- 542 56. Lewis NJ, Batra P, Misiolak BA, Rockafellow S, Tupper C. Transgender/gender
543 nonconforming adults' worries and coping actions related to discrimination: Relevance to
544 pharmacist care. *American Journal of Health-System Pharmacy* 2019;76(8):512-520.
- 545 57. Müller A, Daskilewicz K. Are we doing alright? Realities of violence, mental health, and
546 access to healthcare related to sexual orientation and gender identity and expression in East
547 and Southern Africa: Research report based on a community-led study in nine countries,
548 2019.
- 549 58. World Health Organization. Depression and other common mental disorders: global health
550 estimates. 2017. Available from: <https://apps.who.int/iris/handle/10665/254610>
551 23/10/2021
- 552 59. Freese R, Ott MQ, Rood BA, Reisner SL, Pantalone DW. Distinct Coping Profiles Are
553 Associated With Mental Health Differences in Transgender and Gender Nonconforming
554 Adults. *J. Clin. Psychol.* 2018;74:136-146. doi: 10.1002/jclp.22490



Figure