Title

**Uncovering personalised glucose responses and circadian rhythms from
multiple wearable biosensors with Bayesian dynamical modelling**

Authors and affiliations

Nicholas E. Phillips [1,2], Tinh-Hai Collet [2,3]*, Felix Naef [1]*.

[1] Institute of Bioengineering, School of Life Sciences, Ecole Polytechnique Fédérale de
Lausanne, Lausanne, Switzerland.
[2] Nutrition Unit, Service of Endocrinology, Diabetology, Nutrition and Therapeutic Education,
Department of Medicine, Geneva University Hospitals, Geneva, Switzerland.
[3] Diabetes Centre, Faculty of Medicine, University of Geneva, Geneva, Switzerland.

* These authors contributed equally and act both as corresponding authors (tinh-
hai.collet@hcuge.ch; felix.naef@epfl.ch).

Motivation

Wearable biosensors measure physiological variables with high temporal resolution over multiple days and are increasingly employed in clinical settings, such as continuous glucose monitoring in diabetes care. Such datasets bring new opportunities and challenges, and patients, clinicians and researchers are today faced with a common challenge: how to best capture and summarise relevant information from multimodal wearable time series? Here, we aim to provide insights into individual blood glucose dynamics and their relationships with food and drink ingestion, time of day, and coupling with other physiological states such as physical and heart activity. To this end, we generate and analyse multiple wearable device data through the lens of a parsimonious mathematical model with interpretable components and parameters. A key innovation of our method is that the models are learnt on a personalised level for each participant within a Bayesian framework, which enables the characterisation of inter-individual heterogeneity in features such as the glucose response time following meals or underlying circadian rhythms. This framework may prove useful in other populations at risk of cardiometabolic diseases.

Summary

Wearable biosensors and smartphone applications can measure physiological variables over multiple days in free-living conditions. We measure food and drink ingestion, glucose dynamics, physical activity, heart rate (HR) and heart rate variability (HRV) in 25 healthy participants over 14 days. We develop a Bayesian framework to learn personal parameters that quantify circadian rhythms and physiological responses to external stressors. Modelling the effects of ingestion events on glucose levels reveals that slower glucose decay kinetics elicit larger postprandial glucose spikes, and we uncover a circadian baseline rhythm of glucose with high amplitudes in some individuals. Physical activity and circadian rhythms explain as much as 40-65% of the HR variance, whereas the variance explained for HRV is more heterogeneous across individuals (20-80%). A more complex model incorporating activity, HR and HRV explains up to 15% additional glucose variability, highlighting the relevance of integrating multiple biosensors to better predict glucose dynamics.

Keywords

Wearable devices, biosensors, glucose levels, circadian rhythms, mathematical modelling, Bayesian inference, time series

Introduction

Wearable biosensors and smartphone applications are increasingly used to measure multiple physiological variables, including glucose levels, food consumption, and physical and heart activity. In contrast to traditional lab measurements taken at a single time point, the high-resolution wearable time series data records dynamic changes of physiological variables in response to external perturbations and as a function of the time of day. While this wearable data has the potential to provide a dynamic view of health states[1], a major challenge in both clinical and research settings is how to extract physiologically meaningful information from wearable time series data, and in particular when multiple data modalities are combined.

Glucose regulation is a prime example of a dynamic and complex physiological system, as the body is confronted with irregular inputs (i.e. food intake, especially of carbohydrates) and controlled glucose uptake by organs (e.g. muscles, liver). As such, glycaemic regulation employs a range of homeostatic mechanisms including the glucose-insulin negative feedback loop, whereby insulin secretion by the pancreas is tightly regulated to avoid both low (hypoglycaemic) or high (hyperglycaemic) levels of glucose[2,3]. Understanding glucose regulation is an important health problem, as long-term chronic hyperglycaemia in diabetes can lead to micro- and macrovascular complications[4], and glucose levels show a non-linear association with vascular diseases even in populations without diabetes[5–7].

As glucose homeostasis is inherently dynamic and glucose levels fluctuate throughout the day, continuous glucose monitors (CGMs) have gained popularity due to the high temporal resolution. CGMs measure glucose in interstitial fluid continuously for up to 10-14 days with satisfactory clinical accuracy compared to reference capillary blood glucose values[8,9]. Standardised CGM-derived metrics such as the coefficient of variation (CV) and the time-in-range (the fraction of time spent within the desired range of 3.9-10.0 mmol/L, 70-180 mg/dL) have been adopted in clinical practice to assess glycaemic control in diabetes with insulin treatment[10–13]. At a more fine-grained level, CGMs have been combined with smartphone records of ingestion events to predict postprandial (post-meal) glycaemic responses (PPGR), where the PPGR is often defined as the area under the glucose curve for the two hours following a recorded ingestion event[14–16].

Nonetheless, neither the standardised CGM metrics nor the PPGR approach provide a complete picture of the entire glucose time series and its fluctuations over the 24-hour clock. Physiological processes in humans, including glucose metabolism, follow circadian rhythms[17–20], and responses to oral glucose tests are more pronounced in the evening than the morning[21]. A pre-breakfast rise in glucose levels, termed the "dawn phenomenon", has been observed since the early 1980s and is often linked with a concomitant early morning rise of cortisol[22,23], but the amplitude and phase of circadian rhythms in baseline glucose levels have thus far not been well described at

an individual level. Identifying the relative contribution of the circadian rhythms to the
glucose time series would be particularly helpful for the interpretation of the 24-hour
CGM report, which is often discussed with patients to identify patterns of low and
high glucose values, and to guide treatment[24].

In addition to glucose, other physiological responses are accessible with biosensors
such as heart rate (HR, beats per minute) and heart rate variability (HRV), where
HRV is typically quantified with metrics such as the root mean square of successive
differences (RMSSD) between heart beats[25]. Epidemiological data has linked low
HRV with high glucose levels[26,27] and a reduction in HRV has been shown to predict
the development of autonomic neuropathy before symptom onset amongst diabetic
patients[28]. The simultaneous measurement of HR and HRV can provide insights into
the autonomic nervous system activity[29], as HR receives inputs from both the
sympathetic nervous system (SNS, the "flight or fight" response) and
parasympathetic nervous system (PNS, the "rest and digest" response), while HRV-
derived RMSSD metric is dominated by the PNS via vagal nerve activity[30]. Both HR
and HRV are modulated by physical activity, which can now also be conveniently
measured with a triaxial accelerometer.

Regarding the analysis of wearable data, the methodological approaches depend on
the scientific questions and applications. A diverse range of glucose models have
been proposed over the last decades[31–33], ranging from minimal models[34] to more
detailed simulators with dozens of parameters[35] and neural networks[36–38]. Recent
efforts have also attempted to utilise additional multimodal wearable signals to either
improve glucose forecasting or provide more accessible proxies for glucose without
using CGMs[39–41]. Many of these methods are specialised towards short-range
forecasting, which is certainly useful in applications like the artificial pancreas[42]. In a
different context, researchers and clinicians need new wearable data analysis tools
to perform statistical comparisons between individuals and quantify changes in
glucose regulation across multiple time points and different disease states, but such
approaches to extract personalised summary metrics from the global recordings
remain comparatively unexplored.

In this study, we acquired multiple wearable biosensor data to monitor food and drink
ingestion, glucose excursions, physical activity, HR and HRV in individuals in free-
living conditions. Our aim was to quantify the role of external perturbations (such as
ingestion events and physical activity) and baseline circadian rhythms on a
personalised level. We develop models to analyse the wearable data with distinct
model components that capture the interactions between physiological variables, 24-
hour rhythms and random fluctuations. Individual-specific parameters are learnt
within a Bayesian framework, which provides parameter uncertainties and enables
statistical comparisons between participants. We subdivide the problem of analysing
the multiple signals by creating three successive mathematical models that include
different subsets of variables. Our three-tiered modelling reveals the high degree of

personalisation across a wide range of metrics even within a healthy population,
from glucose decay kinetics, circadian rhythms in baseline glucose levels,
dependence between HR and HRV and the boost in explained glucose variance
from adding physical and heart activity data. While our study assessed cross-
sectional differences between healthy individuals, future studies will be able to re-
use the framework to describe personalised longitudinal changes over time in
response to interventions and to cardiometabolic diseases.
136
137 **Results**
138
139 *Measuring multivariable physiological time series in free-living conditions*
140
To quantify the personalised dynamics of individuals in free-living conditions, we
measured ingestion events, glucose levels, physical activity, HR and HRV for 25
participants over a two-week period. Participants (16 males, 9 females) were young
(mean age 33.0 ± SD 11.0), had a normal weight (mean BMI 22.7 ± 2.8 kg/m^2, 1
person with overweight and 1 person with obesity) and a normal blood pressure
(systolic 117.6 ± 11.4 mmHg, diastolic 75.3 ± 7.9 mmHg). Participants were asked to
record all food and drink consumption and add a manual free text annotation of the
content with the smartphone application MyCircadianClock[43]. Each ingestion event
was automatically timestamped by the app. The adherence (defined as at least 2
meals separated by at least 5 hours in a given day[44] was above 83% for all
participants (Supplementary Table 1).
152
We measured glucose levels continuously using the Abbott FreeStyle Libre Pro
CGM device, which records interstitial glucose levels every 15 minutes over a two-
week period. As the device is blinded, participants were unable to access their
glucose data during the study period thus avoiding feedback on their eating
behaviour. Five participants wore two sensors (on different arms), with the aim of
validating that parameters estimated from the model were consistent between the
two sensors (noted ID .A and .B in the Figures). Physical activity, HR and HRV were
measured for each participant over the two-week study period using the CamNTech
ActiHeart version 5 device, and the physical and heart activity data was also blinded
to participants during the study.
163
164 *Multi-wearable time series data reveal complex dynamical responses as a function of*
165 *external inputs and time of day*
166
For initial data exploration, we superposed the recorded days of glucose data based
on time of day and found strong inter-individual heterogeneity in the mean 24-hour
pattern, with the highest mean glucose level occurring in the morning, afternoon or
evening depending on the individual (Figure 1A, all participants shown in
Supplementary Figure 1). These unique 24-hour trends could be caused by either
food or drink ingestion (i.e. external perturbations) and/or an underlying circadian

baseline rhythm in glucose. This motivated the inclusion of both ingestion events and
circadian rhythms in the model of glucose dynamics as separate components with
learnable parameters.
176
Further exploratory analysis of the multiple wearable signals showed rich interactions
between the five measured variables (i.e. ingestion, glucose, activity, HR and HRV).
As expected, glucose levels often rose following ingestion events, and for some of
the individuals, recorded meals seemed to lead to large, predictable peaks in
glucose (Figure 1B, ID 14 and 23), while others showed a more complex
relationship, with small postprandial glucose spikes that were barely larger than the
glucose fluctuations between meals (Figure 1B, ID 06). Based on these
observations, and compared to CGM analysis methods that focus exclusively on
PPGRs for 2-3 hours[14–16], our goal is now to dynamically model the entire glucose
time series over 2 weeks, including the fluctuating glucose levels occurring overnight
or during longer intervals between ingestion events.
188
Visual inspection of the physical and heart activity data showed that spikes in
physical activity typically coincided with an increased HR and HRV (as measured
with RMSSD^{-1}) (Figure 1C-D). By creating a joint dynamical model of the three
signals (physical activity, HR and HRV), we aimed to uncover the inter-individual
heterogeneity in the coupling between the multiple signals as well as the underlying
circadian rhythms.
195
Finally, we observed spikes in glucose levels following physical activity for some
individuals (Figure 1B-D, ID 23), which could be caused by the release of glucose
under the influence of adrenaline/epinephrine or glucagon. However, to establish
more firmly whether physical and heart activity signals can explain glucose variation,
we developed a dynamical model to mathematically assess the extent to which the
total glucose signal across the two-week study period is predictable by the combined
meal, physical, HR and HRV data.
203
204 *Slow glucose dynamics is associated with large postprandial glucose spikes*
205
The overall data modelling strategy is shown in Figure 2A-C, where we first focus on
ingestion events, glucose and circadian rhythms (Model 1, Figure 2A), then the
relationships between the physical and heart activity signals (Model 2, Figure 2B)
before finally adding interactions from the physical activity and heart signals to the
glucose levels (Model 3, Figure 2C). Based on the visual exploration (Figure 1) and
physiological knowledge, we first built a minimal dynamical model of glucose levels
(Model 1) that included the following four features: i) the ability to produce a
continuous postprandial glucose response following an ingestion event, ii) negative
feedback (due to the regulating action of insulin, represented as a feedback loop in
Figure 2A), iii) a random component that captures the glucose fluctuations between

ingestion events and overnight, iv) a circadian baseline rhythm (discussed in the next
section).
218
We modelled the glucose dynamics with a system of stochastic differential equations
(SDEs), including negative feedback regulation (see Methods), where ingestion
events act to perturb glucose to higher levels. In the deterministic part of the model,
after the ingestion causes a glucose increase, glucose levels return to their steady-
state values (i.e. reflecting homeostasis). The decay kinetics and precise shape of
the response will depend on the parameters of the model (Figure 2D), which are
learnt for each participant. Specifically, this individual-specific response to a meal
perturbation can be summarised with three parameters, namely a half-life reflecting
the time taken for glucose to return to baseline levels, the mean increase in glucose
levels caused by meal consumption (referred to as the average meal height) and a
damping coefficient specifying whether the response profile is akin to an
overdamped (a rapid glucose increase followed by a monotonous slower decay i.e.
non-dipping) or an underdamped (leading to a slower initial increase followed by
decay and overshoot, i.e. dipping) response. To account for noisy fluctuations in the
data, the glucose dynamics is also subjected to random perturbations via a process
noise term in the corresponding SDE, meaning that the glucose time series data can
show noisy deviations from the idealised meal response.
236
For each participant, the entire glucose time-series is probabilistically matched
(using exact likelihood calculations) to the model using a Gaussian state space
model (a.k.a. a Kalman filter), and we infer each of the model parameters using
Markov Chain Monte Carlo (MCMC) sampling within a Bayesian framework that
yields uncertainty estimates for each parameter (Methods and Supplementary
Information).
243
We first verified model performance by assessing the correlation coefficient between
the fitted meal response function and the data (Supplementary Figure 2). The
correlation coefficient generally ranged from 0.5 to 0.8 but was particularly low for
participant ID 04. Visual inspection of this participant's raw data showed large
glucose spikes following physical activity, which we address at a later stage of the
modelling. While all model parameters are shown in Supplementary Figure 2, here
we focus on the three summary metrics of the glucose dynamics.
251
Response half-lives ranged from 1h to 2.2h (Figure 3A) and average meal response
heights ranged from 0.5 mmol/L to 1.5 mmol/L (Figure 3B). The posterior parameter
distributions for each participant (Figure 3A-B) show the uncertainty associated with
the parameter estimates for each participant; in some cases the distributions were
overlapping between two individuals (and hence we lack sufficient resolution to claim
inter-individual differences) while in other cases the distributions were clearly
separated and would be significant within a Bayesian statistical testing framework
(e.g. half-life comparing ID 20 and 23, Figure 3A) [45]. Across participants we found a

positive relationship between response half-lives and average meal heights, with
slower glucose response half-lives associated with larger postprandial glucose
spikes (R = 0.44, p = 0.02, Figure 3C). This suggested that postprandial glucose
control (i.e. the height of glucose spikes following meals) depends on glucose
clearance time, which might be determined physiologically by insulin sensitivity or
beta cell function (Discussion).
266
The damping coefficients describing the shapes of glucose responses were
clustered around 0 across all participants (Supplementary Figure 2), where values of
0 represent "critical" damping at the border between overdamped (non-dipping
profiles, damping coefficient >0) and underdamped (profiles with a dip, damping
coefficient <0). Interestingly, glucose responses were proposed to be critically
damped in an early glucose model(Bolie, 1961), which would be consistent with our
finding that the inferred values are scattered around zero. However, subject-to-
subject variability is clearly found, with participant ID 14 showing a clear
underdamped glucose response compared to the critically damped response in ID
23 (Figure 3D-F). The inferred meal response and circadian time functions (orange)
are smoother than the glucose data (blue, Figure 3E-F), but the full model that also
adds random fluctuations produces glucose traces that closely resemble the glucose
data (Supplementary Figure 3).
280
The measured glucose coefficient of variation (CV), a metric of glycaemic control
used in clinical settings[10,11], showed significant associations with both the response
half-lives (p=0.03) and average meal heights (p=0.01, R^2 using both variables =
0.63). While the damping coefficient was not significantly associated with glucose
CV, the individual shapes of glucose responses might play a role in other aspects of
glucose dynamics such as overshooting and hunger[46]. Our results highlight that
glucose response half-lives play a role in glycaemic control and may be a relevant
metric for both fundamental research and clinical purposes.
289
290 *Circadian rhythms in baseline glucose levels show large inter-individual variability*
291
292 In addition to the input from ingestion events, the model also allows for an underlying
293 circadian rhythm in glucose levels described with three parameters: a baseline level
294 that specifies the glucose at the trough of the oscillation, an amplitude parameter
295 denoting the difference between the trough and peak of the oscillation, and the peak
296 time of the oscillation.
297
The amplitudes of underlying circadian glucose rhythms were highly variable
between participants (Figure 4A), being virtually null for some participants while
exceeding 1 mmol/L for others (Figure 4C, ID 03 and 07, respectively). Notably, the
parameter uncertainty was small enough that there was no overlap in the estimates
for participants ID 03 and 07 (Figure 4A). To identify subjects whose profiles do not
support a circadian baseline trend, we fitted an alternative model that lacked a

circadian baseline and compared the two models using the Bayesian information
criterion (BIC). For IDs 10, 13, 08, 25, 03 and 04 (which have the weakest amplitude
according to Figure 4A), the BIC indicated evidence for the model lacking the
circadian baseline, while the BIC favoured the model with an additional circadian
component for all remaining participants (Supplementary Figure 4). The combination
of the amplitude posterior estimates and heterogeneous model preference according
to BIC suggests that circadian baseline glucose oscillations are highly unique
physiological characteristics.
312
The peak times of the glucose circadian oscillations similarly showed large
differences between participants (Figure 4B, with the same participant order as in
Figure 4A). While the peak times of the circadian underlying trend typically fell
around the mid-afternoon, there were large phase differences between some
individuals. For example, participant ID 20 had a peak time at 10:00 while it occurred
much later for participant ID 17, falling at 20:00 (Figure 4D). The peak time
distributions showed tight confidence intervals for participants with large amplitudes
and wide intervals for participants with weaker amplitudes (Figure 4B). This
relationship is probably caused by a low signal-to-noise ratio and suggests that the
circadian peak time cannot be reliably inferred for participants with a low circadian
amplitude.
324
325 Overall, the underlying circadian glucose rhythm can explain >15% of glycaemic
326 variability in addition to the meal model for participants with large amplitudes (Figure
327 4E). While we have not tested whether it would be possible to modify either the peak
328 time or amplitude of this rhythm, these personalised parameters should prove to be
329 useful in applications such as personalised meal timing (Discussion).
330
331 *HR is well predicted by physical activity and circadian time but the predictability of*
332 *HRV is highly variable between individuals*
333
We next focused on the physical activity, HR and HRV data, where the aim was to
model the dependencies between the variables and quantify the ability of a subset of
the three signals to explain the variance of another, in addition to the contribution of
circadian rhythms. For this, we created a new model (Model 2, Figure 2B) that
incorporated the influence of physical activity on HR and HRV, and we used MCMC
to sample from model parameters and quantify differences between individuals (all
parameters shown in Supplementary Figure 5).
341
For HR the combination of circadian rhythms and physical activity as two inputs was
consistently predictive, explaining 40-65% of HR variance across all participants
(Figure 5A). Figure 5C-D shows an example of the predicted HR (orange) for two
different participants using both the underlying circadian rhythm (black) and
integrating the physical activity (green). While the circadian contribution to the
explained HR variance differs for these two participants (Figure 5A), the correlation

between the predicted and observed HR was ~0.8 for both participants,
demonstrating that time of day and activity state are necessary for optimal
personalised modelling of HR, which is consistent with previous studies[47].
351
The ability to predict HRV was, in contrast, much more heterogeneous between
participants with total variance explained between 20% and 80% (Figure 5B). This
notable difference in predictability is illustrated with two participants, showing a
favourable prediction for participant ID 25 (R = 0.89, Figure 2F) compared to ID 14
(R = 0.44, Figure 2E). In addition to inputs from physical activity and the underlying
circadian rhythm, we evaluated whether the correlations between HR and HRV could
be exploited by using HR to predict HRV (which is technically more difficult to
measure than HR). The dependence between HR and HRV showed marked inter-
individual differences, where for ID 25 the HR signal explained 40% of the variance
compared to using just activity and circadian rhythm, but for ID 14 the addition of HR
makes no difference to HRV prediction (Figure 5E-F). Given that HR and HRV
receive different inputs from the SNS and PNS[25], the strength of this dependence
may be a function of the autonomic nervous system. Of note, ID 14 was previously
diagnosed with diabetes currently treated only with lifestyle measures (and not
pharmacological treatment) and autonomic dysfunction is a known complication of
diabetes[48].
368
369   *Integrating physical and heart activity signals helps explain glycaemic dynamics for*
370   *some individuals*
371
As a final modelling step, we integrated the physical and heart activity signals with
the glucose-ingestion model to quantify how much of the glucose dynamics can be
accounted for with physical activity, HR and HRV (Model 3, Figure 2C). To simplify
the model inference problem, the parameters describing the physical and heart
activity model in isolation (Model 2, Figure 2B) were locked to their posterior mean
values, and we added three new parameters that describe the input of physical
activity, HR and HRV on glucose levels, respectively (Model 3, Figure 2C). These
influences were left unconstrained and could have a positive, negative or zero effect
on glucose levels.
381
Fitting the parameters to the data revealed that the effect of physical activity on
glucose (parameter $C_{5,1}$) was generally negative, the effect of HR (parameter $C_{5,2}$)
was generally positive and the effect of HRV (parameter $C_{5,3}$) was typically neutral
across all participants (Figure 6A-C). Given that the raw data showed glucose spikes
during some periods of exercise (Figure 1D), the negative influence of physical
activity on glucose ($C_{5,1}$) was not expected. To test the robustness of this prediction,
we therefore refitted the data using three simpler models, where there was only one
input at a time from the physical and heart activity signals (Supplementary Figure 6).
The influence of physical activity on glucose remained negative even when it was the
sole input from Model 2 into glucose levels, further suggesting that the overall effect

of physical activity is to deplete glucose levels amongst the participants of our study.
Meanwhile, as HR acts to increase glucose levels ($C_{5,2}$), increased HR during
intense exercise can still lead to a net increase in predicted glucose levels. Overall,
the amount of additional explained variance contributed by the physical and heart
activity signals was modest for most participants but contributed up to 15% in certain
individuals (Figure 6D).
398
Amongst the participants whose glucose dynamics benefit most from the physical
and heart activity signals, the mode of action also differed. For participant ID 04,
where the prediction of the glucose was the lowest with Model 1 (Figure 4E), the
inclusion of the physical and heart activity signals allowed for partial prediction of
exercise-induced glucose spikes that were otherwise absent from the glucose model
that contained meals only (Figure 6E). For participant ID 08, which also showed a
benefit of including physical and heart activity data, there were no notable isolated
glucose spikes that were predicted by the physical and heart activity data (Figure
6F). Instead, the linear combination of the physical and heart activity variables
(weighted according to the inferred coefficients $C_{5,1}$-$C_{5,3}$) appeared to track the
glucose levels (Figure 6F), and the cross-correlation profile showed a maximum
correlation between the two signals with no time delay (Figure 6G). This suggests
that for this mode of action, the physical and heart activity signals make a more
diffuse contribution to the prediction of glucose levels that is more related to baseline
trends spread over the time series.
414
415   **Discussion**
416
This study makes two major contributions to the quantitative analysis of glucose
dynamics in terms of study design and computational analysis. Firstly, we used
several wearable biosensors to measure multiple, interconnected data streams
simultaneously. Without such high-resolution time series detailed dynamic inferences
are impossible. On a practical level, one of the main advantages of health
measurement with wearable biosensors is the rapidity of the experimental pipeline:
in our study it took a team of two researchers two days to complete questionnaires
and set up devices to record data for all 25 participants simultaneously. Secondly,
we developed a data analysis method that combines stochastic dynamical modelling
with Bayesian inference to learn personal parameters along with their associated
uncertainty, and this parameter uncertainty was necessary to compare participants.
The main insight of this personalised modelling is the clear inter-individual
differences across a range of parameters and metrics, from the glucose half-lives
and circadian oscillations, the coupling between HR and HRV, to the benefit of
adding additional signals to predicting glucose.
432
The output of this modelling has wider implications for both understanding the
biological underpinnings of cardiometabolic dysfunction as well as consequences for
the use of wearables in a clinical setting. We first started by combining the food and

drink events with the glucose time series, where we found that slow glucose
dynamics are associated with large postprandial glucose spikes. Mechanistically,
slow glucose disposal could relate to the quantity of ingested carbohydrates, the rate
of gut absorption[49], its metabolism[50], suppression of endogenous glucose
production[51], insulin resistance or beta cell function[2]. Moreover, we also detected
highly personalised circadian rhythms in the underlying glucose levels. While diurnal
rhythms in beta cell function and insulin sensitivity have been shown at an average
level within healthy populations[52–54], it was unexpected to see such large differences
in circadian glucose amplitude and phase between individuals. Future studies will
determine whether this circadian glucose baseline trend is predictive of responses to
specific meal times, e.g. in time-restricted eating, an intervention which restricts
eating to a specific window within the 24-hour clock[55].
448
Our results also have practical implications for clinicians as the physical and heart
activity data explained up to 15% of glucose variability in our study, although this
was highly variable between participants and may only be useful for some
individuals. Never-the-less, from a clinical perspective, the incorporation of these
additional signals might help both patients and clinicians understand glucose
dynamics that seem otherwise disconnected from meal consumption (e.g. Figure
6E). As outlined in the Motivation paragraph, clinicians and patients with diabetes
can link glucose excursion with ingestion events and intensive physical activity, but
struggle to do this for the rest of the glucose dynamics measured over 24 hours.
458
There are multiple possible approaches to modelling our collected multi-modal data,
and the particular glucose modelling methodology has often been historically
dictated by the data available and by the stated goal[31–33]. Models based on
differential equations range from simple, minimal models[34] to more mechanistically
detailed descriptions that include more variables, more spatial compartments and
dozens of additional parameters[35]. Inspired by this work, there are many possible
extensions that could be added to our glucose model, such as a meal-specific
ingestion rates due to food content in carbohydrates, but also fat, fibre and protein
content which are known to slow down nutrient absorption[14], although the addition of
meal-specific response shapes would effectively double the number of meal-related
parameters.
470
An area that has seen a broad spectrum of time series models is in short-term
glucose forecasting, typically for applications in closed-loop insulin delivery
systems[42]. Gaussian state-space models that are conceptually similar to our own but
with more variables have been deployed in artificial pancreas devices[56,57], but a wide
range of methods for short-range forecasting have also been used (reviewed in
Woldaregay et al.[41]), including SVM and neural networks[36–38]. The advantage of
such methods is that more complex nonlinear dependencies and long-range memory
could possibly be captured. Here we traded some of this flexibility for explainability
by using a relatively simple dynamical model with interpretable components and

parameters, as our focus was on explaining the total time series rather than short-
term forecasting. In the future, we envisage several applications, such as larger
scale epidemiological studies (i.e. do inferred parameters track with health state?)
and clinical trials to see whether parameters change in response to an intervention.
Particularly for the second application, point estimation of parameters is not
adequate, and uncertainty estimates are required to perform statistical tests for a
given individual, which we achieve here through MCMC. In healthcare, there is
increasing interest in the use of digital twins[58,59] to integrate multiple clinical data
streams, devise personalised treatments and perform risk modelling. As our
approach contains interpretable parameters, it lends itself readily to exploring
hypothetical situations by altering parameters (such as circadian amplitude or
glucose response time) and simulating from the model. Overall, our method
transforms a multivariable wearable data input into a series of metrics that describe
the dependencies between physiological variables, including the relaxation
timescales after external perturbations and circadian properties, and this approach
provides a platform for probing physiological changes across circadian perturbations,
ageing and cardiometabolic disorders.
497
498 <u>Limitations of the study</u>
499
500 Our study population was young and in good health overall, and we lack additional,
501 more detailed health information or standard clinical metrics such as HbA1c. We
502 have therefore generally sought to identify differences between individuals without
503 attempting to associate them with either good or bad health outcomes.
504
With respect to the modelling, a potential limitation of our approach is the use of a
relatively simple linear model, which may not be able to capture more complex
phenomena such as eventual decreases in hepatic glucose production during
prolonged exercise[60]. A recent study based on deep learning found the addition of
wristband activity data improved the root mean square error of 60-minute glucose
forecasting by 2.25 mg/dL (0.1 mmol/L) from a baseline of 35.3 mg/dL (2.0 mmol/L),
and hence more substantial improvements in glucose predictions may prove to be a
difficult challenge even with more flexible models[61].
513
514 **Methods**
515
516 *Devices and experimental design*
517
The Multi-Sensor Study (MSS) was approved by the local ethics committee (CER-
VD, BASEC no. 2019-02245) and each participant signed a written informed
consent. Recruitment was performed via posters at the École Polytechnique
Fédérale de Lausanne (EPFL), Lausanne University Hospital (CHUV) and the
University of Lausanne (UNIL) and via presentations given in the EPFL School of
Life Sciences.

We included adults aged ≥ 18 years, with a smartphone compatible with the
MyCircadianClock app (iOS or Android systems[43]) and able to take pictures of
food/drinks, and who self-identified as disciplined enough and motivated to record all
data for two weeks. The exclusion criteria were major illness/fever, surgery over the
previous month, eating disorder, major mental illness, unable to give informed
consent, taking medicines including paracetamol, aspirin or vitamin C supplements,
enrolled in another interventional clinical trial (medication, medical device), shift work
or travel to a different time zone before and during the study.

At baseline, we collected data on demographics, medical history, physical activity
(short form of International Physical Activity Questionnaire, IPAQ-SF[62], chronotype
(The Munich ChronoType Questionnaire[63], sleeping habits (Pittsburgh Sleep Quality
Index[64] and eating timing (with a custom questionnaire on eating habits during work
and free days).

For each participant, we collected data for two weeks using the following devices: 1)
Timestamps of food/drinks and text annotations collected with the smartphone
application (app) MyCircadianClock[43]; 2) Continuous glucose monitoring (CGM)
using the Abbott FreeStyle Libre Pro device; 3) Physical activity, heart rate (HR) and
heart rate variability (HRV using RMSSD^{-1}) using the CamNTech ActiHeart device
version 5. Participants were instructed to take pictures of all consumed food and
drink with the research-dedicated myCircadianClock smartphone. Recorded entries
included a timestamped picture and a free-text annotation, and entries with the same
annotation were considered as the same meal type. Participants could annotate
photographs either immediately or in the following hours. Optionally, participants
could type text-only entries without any picture, e.g., if the smartphone ran out of
battery, or if it was not socially acceptable to take pictures in the current context.
Participants were also asked to optionally log physical exercise using the app. While
the CamNTech Actiheart device is waterproof, participants were permitted to briefly
remove the device during showers and baths. Information on device technical failure,
handling of missing data and data quality is included in the Supplementary
Information.

Pre-processing CGM data

We used nonparametric regression with Gaussian processes (GPs) to remove the
long-term trends observed in the data. After mean-centring the data, we fitted a GP
with a squared exponential kernel $K_{SE}(t, t') = (-|t - t'|^2 / 2l^2)$ and a length scale
l=48 hours using GPflow[65].

    Data analysis

See the Supplementary Information for detailed computational methods which are

summarised here. We use a linear Gaussian state space model (otherwise known as

a Kalman filter[66]) to analyse the time series generated by the wearable devices,

which was implemented using the 'LinearGaussianStateSpaceModel' distribution

within TensorFlow Probability[67]. We will first describe the general data analysis

framework before providing details on each of the three models used. For each

model we define a dynamic model that describes the time evolution of the underlying

physiological variables and a measurement model that incorporates measurement

noise. For the dynamic model, we use a system of stochastic differential equations

(SDEs)

580
$$\mathrm{d}\boldsymbol{x}(t) = W\boldsymbol{x}(t)\mathrm{d}t + \mathrm{d}\beta$$

where W is a matrix describing the interactions between the variables $\boldsymbol{x}(t)$, and β is

a brownian noise term with covariance matrix Q. The specific forms of W and Q are

unique for each model and will be described below. To keep the model exact while

benefiting from the generic framework of Gaussian state space models (a.k.a. as

Kalman filters), we then convert this system of continuous-time SDEs into a model

where time is discrete (see Supplementary Information for details).

590
$$\boldsymbol{x}(t_k) = F_{k-1}\boldsymbol{x}(t_{k-1}) + N(0, \Sigma_{k-1}),$$

591   where F_k is the state-transition model and Σ_k is the covariance of the process noise.

592   The measurement model describes the observation process and assumes that

593   variables are observed with normally distributed measurement noise

595
$$\boldsymbol{y}(t_k) = H_k\boldsymbol{x}(t_k) + N(\boldsymbol{m}_k, R_k),$$

597   where H_k is the observation matrix and $\boldsymbol{m_k}$ and R_k represents the mean and

598   covariance of the observation noise, respectively. The goal is to use the wearable

599   time series data $\boldsymbol{y}_{1:T}$ to estimate parameters (denoted by $\boldsymbol{\theta}$) for each participant.

600   Within a Bayesian inference framework, the parameters of the model can be

601   estimated from the data as follows

603
$$p(\boldsymbol{\theta} \mid \boldsymbol{y}_{1:T}) \propto p(\boldsymbol{\theta})p(\boldsymbol{y}_{1:T} \mid \boldsymbol{\theta}),$$

605   where $p(\boldsymbol{\theta})$ is the prior distribution of parameters and $p(\boldsymbol{y}_{1:T} \mid \boldsymbol{\theta})$ is the likelihood of

606   observing the temporal data $\boldsymbol{y}_{1:T}$ given the set of parameters $\boldsymbol{\theta}$. Considering the time

607   series sequence of data, the likelihood term for a given set of parameters $\boldsymbol{\theta}$ can be

608   expressed as

609
$$p(\boldsymbol{y}_{1:T} \mid \boldsymbol{\theta}) = p(\boldsymbol{y_1} \mid \boldsymbol{\theta}) \prod_{k=2}^{T} p(\boldsymbol{y}_k \mid \boldsymbol{y}_{1:k-1}, \boldsymbol{\theta}),$$

and the sequence of distributions $p(\mathbf{y_k} \mid \mathbf{y_{1:k-1}}, \boldsymbol{\theta})$ are calculated within a Kalman
filtering framework. Once the likelihood and priors are specified for each model, we
used the Hamiltonian Monte Carlo sampler provided within TensorFlow Probability to
sample model parameters from the posterior distribution using 4 different chains with
10,000 samples each. See Supplementary Information for further details on MCMC
scheme. We then estimate the explained variance using $1 - \text{Var}(\boldsymbol{y} - \widetilde{\boldsymbol{y}})/\text{Var}(y)$,
using the model predictions $\widetilde{\boldsymbol{y}}$ from the MCMC parameter samples.

Model 1: glucose model

We model glucose dynamics (Figure 2A) with a two-dimensional system of SDEs,
where the second variable x_{GLUC2} represents the glucose levels and the first variable
x_{GLUC1} represents an unobserved latent variable that allows negative feedback within
the system. In matrix form, the model is expressed as follows

$$d\mathbf{x}(t) = W\mathbf{x}(t)dt + d\boldsymbol{\beta}$$

$$\mathbf{x}(t) = \begin{bmatrix} x_{\text{GLUC1}}(t) \\ x_{\text{GLUC2}}(t) \end{bmatrix}, W = \begin{bmatrix} -A_{11} & -A_{12} \\ A_{21} & -A_{22} \end{bmatrix}, Q = \begin{bmatrix} 0 & 0 \\ 0 & B_{22} \end{bmatrix},$$

and where the coefficients A_{ij} are constrained to be positive. The covariance of the
brownian noise term β is given by Q. The 'damping coefficient' is determined by
whether the eigenvalues of the matrix W are real or complex. For the 2x2 matrix W,
this damping coefficient can be determined by $-\det(W - I\,\text{tr}(W)/2)/(\text{tr}(W)/2)^2$.

We incorporate meal events (that are recorded at time t_m) as producing a response
function $r_m(t)$ by perturbing the first variable x_{GLUC1} to higher values (see
Supplementary Information for precise functional form), and then the total meal
function is the sum over all M individual meal responses

$$r(t) = \sum_{}^{M} r_m(t, \boldsymbol{\theta})$$

We define the glucose half-life parameter as the model-predicted time to return to
0.5 mol/L after a standardised increase of 1mmol/L. We also add an underlying
circadian rhythm in glucose levels using a sinusoidal function

$$g(t) = A_{0,\text{GLUC}} + A_{1,\text{GLUC}}(1 + \cos(\omega t - \phi_{\text{GLUC}}))/2$$

where $A_{0,\text{GLUC}}$ is the baseline level, $A_{1,\text{GLUC}}$ is the amplitude, ω is the frequency (fixed
at $2\pi/24$), and ϕ_{GLUC} is the peak time of the maximum. The observation model for
the glucose model is then as follows

$$\mathbf{y}(t_k) = H_k\mathbf{x}(t_k) + N(\mathbf{m}_k, R_k)$$

$$H_k = [0, 1], \mathbf{m}_k = r(t) + g(t), R_k = \sigma_{glu}^2$$

We compared Model 1 with an alternative version without circadian oscillations using

the Bayesian Information Criterion (BIC) $\mathrm{BIC} = k\ln(n) - 2\ln(p(\mathbf{y}_{1:T} \mid \boldsymbol{\theta}))$, where k is

the number of parameters and n is the number of data points. We calculated the

difference in BIC score using Model 1 both with and without a circadian components

and used a cut-off of $\ln 10$ to indicate that the strength of evidence favoured a

particular model[68].

Model 2 : physical and heart activity model

We model glucose dynamics (Figure 2B) with a with a three-dimensional system of

SDEs, where the first variable x_{ACT} represents physical activity, the second variable

x_{HR} represents heart rate and the third variable x_{HRV} represents heart rate variability,

where we use the inverse of the root mean square of successive differences

between normal heartbeats (RMSSD^{-1}). We normalise all three variables by their

respective standard deviations before inferring parameters. In matrix form, the

model is expressed as follows

$$\mathrm{d}\mathbf{x}(t) = W\mathbf{x}(t)\mathrm{d}t + \mathrm{d}\boldsymbol{\beta},$$

$$\mathbf{x}(t) = \begin{bmatrix} x_{\mathrm{ACT}}(t) \\ x_{\mathrm{HR}}(t) \\ x_{\mathrm{HRV}}(t) \end{bmatrix} \quad W = \begin{bmatrix} -C_{11} & 0 & 0 \\ C_{21} & -C_{22} & 0 \\ C_{31} & 0 & -C_{33} \end{bmatrix} \quad Q = \begin{pmatrix} D_{11} & 0 & 0 \\ 0 & D_{22} & \rho\sqrt{D_{22}D_{33}} \\ 0 & \rho\sqrt{D_{22}D_{33}} & D_{33} \end{pmatrix},$$

666   and where the coefficients C_{ij} are constrained to be positive and the covariance of

667   the brownian noise term β is given by Q. The observation model is then given by

$$\mathbf{y}(t_k) = H_k\mathbf{x}(t_k) + N(\mathbf{m}_k, R_k),$$

$$\mathbf{m}_k = \begin{bmatrix} g_{\mathrm{ACT}}(t) \\ g_{\mathrm{HR}}(t) \\ g_{\mathrm{HRV}}(t) \end{bmatrix}, H_k = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, R_k = \begin{bmatrix} \sigma^2_{\mathrm{ACT}} & 0 & 0 \\ 0 & \sigma^2_{\mathrm{HR}} & 0 \\ 0 & 0 & \sigma^2_{\mathrm{HRV}} \end{bmatrix}$$

where $g_{\mathrm{ACT}}(t)$, $g_{\mathrm{HR}}(t)$ and $g_{\mathrm{HRV}}(t)$ are circadian oscillatory functions (Supplementary

Information).

Model 3: combined model

The final model (Figure 2C) connects the CamNTech ActiHeart signals with CGM

dynamics by stitching the previous glucose and physical and heart activity models

together. Both models are otherwise left unchanged, but there is an introduction of

three new parameters C_{51}, C_{52} and C_{53} that describe the effect of physical activity, HR
and HRV on glucose levels. These three parameters are left unconstrained and can
take either positive or negative values.

$$\mathrm{d}\mathbf{x}(t) = W\mathbf{x}(t)\mathrm{d}t + \mathrm{d}\boldsymbol{\beta}$$

$$\mathbf{x}(t) = \begin{bmatrix} x_{\mathrm{ACT}}(t) \\ x_{\mathrm{HR}}(t) \\ x_{\mathrm{HRV}}(t) \\ x_{\mathrm{GLUC1}}(t) \\ x_{\mathrm{GLUC2}}(t) \end{bmatrix} \quad W = \begin{bmatrix} -C_{11} & 0 & 0 & 0 & 0 \\ C_{21} & -C_{22} & 0 & 0 & 0 \\ C_{31} & 0 & -C_{33} & 0 & 0 \\ 0 & 0 & 0 & -A_{11} & -A_{12} \\ C_{51} & C_{52} & C_{53} & A_{21} & -A_{22} \end{bmatrix} \quad Q = \begin{bmatrix} D_{11} & 0 & 0 & 0 & 0 \\ 0 & D_{22} & \rho\sqrt{D_{22}D_{33}} & 0 & 0 \\ 0 & \rho\sqrt{D_{22}D_{33}} & D_{33} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & B_{22} \end{bmatrix}$$

682
683
684 The observation model is then given by

$$\mathbf{y}_k = H_k\mathbf{x}(t_k) + N(\mathbf{m}_k, R_k)$$

$$\mathbf{m}_k = \begin{bmatrix} g_{\mathrm{ACT}}(t) \\ g_{\mathrm{HR}}(t) \\ g_{\mathrm{HRV}}(t) \\ g_{\mathrm{GLUC}}(t) \end{bmatrix}, H_k = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}, R_k = \begin{bmatrix} \sigma_{\mathrm{ACT}}^2 & 0 & 0 & 0 \\ 0 & \sigma_{\mathrm{HR}}^2 & 0 & 0 \\ 0 & 0 & \sigma_{\mathrm{HRV}}^2 & 0 \\ 0 & 0 & 0 & \sigma_{\mathrm{gluc}}^2 \end{bmatrix}$$

685
686

Acknowledgments
The authors wish to thank all participants in the study and the team of Professor
Satchidananda Panda (Salk Institute) for the use of the *myCircadianClock*
smartphone application. This project was supported by the grant #2018-427 of the
Strategic Focal Area "Personalized Health and Related Technologies (PHRT)" of the
ETH Domain to N.E.P. Dr. T-H Collet's research is supported by grants from the
Swiss National Science Foundation (PZ00P3-167826, 32003B-212559), the
Leenaards Foundation, the Vontobel Foundation, the Nutrition 2000plus Foundation,
the SwissLife Jubiläumsstiftung Foundation, the Swiss Society of Endocrinology and
Diabetes, and the Swiss Multiple Sclerosis Society. The work in the Naef lab and
purchase of the wearable devices was supported by the EPFL.

Author contributions
Conceptualization: N.E.P., T.-H.C., F.N.; Methodology, N.E.P., T.-H.C., F.N.; Data
collection: N.E.P., T.-H.C.; Computation and data analysis: N.E.P.; Data
visualisation, N.E.P., T.H.C., F.N.; Writing: original draft preparation, N.E.P.; Writing:
review and editing, N.E.P., T.-H.C., F.N.; Project administration: N.E.P., T.-H.C.,
F.N.; Supervision: T.-H.C., F.N.; Funding acquisition: N.E.P., T.-H.C., F.N. All
authors have read and agreed to the published version of the manuscript.
706

707 **Competing interests**
708 The authors declare no competing interests.
709

 Data availability
 The data will be made available after publication in an anonymised form.

 Code availability
 The code for the method can be found at https://github.com/naef-lab/MultiSensor.

 Figure legends

Figure 1. Exploratory analysis of wearable signals: examples of 24-hour trends and responses to external stressors. (A) Continuous glucose monitoring (CGM) data: superposition of all recorded days of data shown on the same 24-hour scale for three different participants (see related Supplementary Figure 1 for all participants). Black: average over all days; coloured lines: data for individual days; time axis: wall clock time. (B) Selected day examples of CGM glucose levels alongside recorded ingestion events for three participants (same individuals shown in panels B-D). Blue: glucose levels; green shade: recorded activity events; time axis: wall clock time; vertical dashed lines: ingestion events. (C) Selected day examples of physical activity measured with the CamNTech Actiheart device. Green: physical activity; green shade: recorded activity events; time axis, clock time. (D) Selected day examples of HR and HRV measured with the CamNTech Actiheart device. Green shade: recorded activity events; purple: heart rate variability (HRV) (quantified with $RMSSD^{-1}$ in ms^{-1}); red: heart rate (HR) in beats per min. (bpm); time axis: wall clock time.

Figure 2. Schematic showing the three different models and parameter interpretation. (A) The glucose and ingestion events interaction (Model 1), (B) the physical and heart activity interaction model (Model 2), (C) the full model (Model 3). Solid arrows represent direct unidirectional influences, while dashed lines represent correlated fluctuations that are not specifically directional. (D) A meal or drink event causes a glucose increase to a specific meal height. The response half-life determines how quickly glucose returns to baseline. Underdamping (defined as a negative damping coefficient) leads to an overshoot below the baseline values.

Figure 3. Characterising participant-specific post-meal glycaemic responses. (A) The inferred glucose response half-life for each participant, defined as the model-predicted time it would take for glucose levels to fall to 0.5 mmol/L following a peak of 1 mmol/L. The boxplots represent the 25th, median (50th) and 75th percentiles of the posterior distribution and the whiskers represent the 5th and 95th percentiles. (B) The average meal glucose spike height calculated as the mean height over all meals consumed during the experiment. (C) The average meal height as a function of the glucose meal response half-life. Points represent the mean posterior values for each participant. (D) Average meal response profiles using the posterior mean parameter values. (E-F) Examples comparing the CGM data (blue) with the model prediction incorporating circadian dynamics (black) plus meal consumption (orange) for two

participants with overdamped and underdamped dynamics, respectively. The
timestamps of meals are shown as dashed lines. Participant order is the same in
Figures 3A-B and 4A-B.

Figure 4. Amplitudes and peak times of circadian baseline levels of glucose are
highly heterogeneous between participants. (A) The amplitude of the 24-hour
sinusoidal circadian rhythm in baseline glucose levels after model fitting to the CGM
data for all participants. The boxes represent the 25th, median (50th) and 75th
percentiles of the posterior distribution and the whiskers represent the 5th and 95th
percentiles. (B) The circadian phase of the glucose circadian rhythm across all
participants. (C) Example of participants with a high (ID 07) and low (ID 03)
amplitude glucose circadian rhythm. Blue, CGM data; black, fitted model of circadian
baseline (using mean posterior parameter values). (D) Examples showing two
participants with large phase difference in underlying glucose rhythm (ID 20 peak
phase: 10:00, ID 17 peak phase: 20:00). (E) The explained variance in glucose
levels using just the meal component of the model (light orange) compared with the
inclusion of the circadian rhythm (dark orange). Error bars represent the 5^{th} and 95^{th}
percentiles of the posterior distribution. Participant order is the same in Figures 3A-B
and 4A-B.

Figure 5. HRV predictions using multi-signal inputs and circadian rhythms are more
heterogeneous than for HR. (A) The amount of variance of the HR signal explained
by circadian rhythms (light orange) and a combined model with circadian rhythms
and physical activity (dark orange). Error bars represent the 5^{th} and 95^{th} percentiles
of the posterior distribution. (B) The amount of variance of the HRV ($RMSSD^{-1}$)
signal explained by circadian rhythms (light orange), a combined model with
circadian rhythms and physical activity (medium orange) and a combined model with
circadian rhythms, physical activity and HR (dark orange). (C,D) Examples
comparing HR data with model predictions for two participants. Red: HR data; black:
baseline circadian rhythm; green: physical activity (shown on normalised scale
where 1 represents the maximum value); orange: model prediction with circadian
rhythm and integrating activity. (E,F) Examples comparing HRV data with model
predictions for two participants. Red: HRV data; black: baseline circadian rhythm;
green: physical activity (shown on normalised scale where 1 represents the
maximum value); orange: model prediction using circadian rhythms, physical activity
and HR. Participant order is the same in panels A and B.

Figure 6. Adding physical activity, HR and HRV into the glucose dynamics model can
help explain glucose dynamics. (A-C) Posterior distributions (shown as boxplots) of
the model coefficients C_{51}, C_{52}, C_{53} across all participants, which correspond to the
influence on glucose of physical activity, HR and HRV, respectively. The boxplots
represent the 25th, median (50th) and 75th percentiles of the posterior distribution
and the whiskers represent the 5th and 95th percentiles. (D) A comparison of the
variance explained in the glucose signal using just the meal and circadian rhythm

model (light orange) compared to the prediction that also incorporates physical
activity, HR and HRV (dark orange). Error bars represent the 5[th] and 95[th] percentiles
of the posterior distribution. (E) Example from ID 04 shows that the physical and
heart activity data partially explain an exercise-induced glucose spike. Blue: glucose
data; light orange: prediction using meal and circadian model components; dark
orange: prediction including meal and circadian model components, physical activity,
HR and HRV; green shade: recorded activity events; brown: the weighted sum of the
physical activity, HR and HRV variables according to the inferred coefficients $C_{5,1}$,
$C_{5,2}$ and $C_{5,3}$; vertical dashed lines: ingestion events. (F) Example from ID 08
showing how glucose dynamics (blue) track with the weighted sum of the physical
activity, HR and HRV variables according to the inferred coefficients C_{51}, C_{52} and C_{53}
(dark brown: data only; light brown: using filtered estimations from Model 2 to fill
missing physical and heart activity data; vertical dashed lines: ingestion events). (G)
the cross-correlation of the glucose (blue in (F)) with the weighted sum of the
physical activity, HR and HRV variables (brown in (F)) using all recorded data.
Participant order is the same between panels A, B and C.
814
815   **References**
816

1. Trachana, K. *et al.* Taking systems medicine to heart. *Circulation Research* **122**,
1276–1289 (2018).
2. Kahn, S. E., Cooper, M. E. & Del Prato, S. Pathophysiology and treatment of type
2 diabetes: Perspectives on the past, present, and future. *The Lancet* **383**, 1068–
1083 (2014).
3. Tirone, T. A. & Brunicardi, F. C. Overview of glucose regulation. *World Journal of*
Surgery **25**, 461–467 (2001).
4. Forbes, J. M. & Fotheringham, A. K. Vascular complications in diabetes: old
messages, new thoughts. *Diabetologia* **60**, 2129–2138 (2017).
5. Coutinho, M., Gerstein, H. C., Wang, Y. & Yusuf, S. The relationship between
glucose and incident cardiovascular events. *Diabetes Care* **22**, 233–240 (1999).
6. Sarwar, N. *et al.* Diabetes mellitus, fasting blood glucose concentration, and risk of
vascular disease: A collaborative meta-analysis of 102 prospective studies. *The*
Lancet **375**, 2215–2222 (2010).
7. Selvin, E. *et al.* Glycated Hemoglobin, Diabetes, and Cardiovascular Risk in
Nondiabetic Adults. *New England Journal of Medicine* **362**, 800–811 (2010).
8. Agarwal, S. *et al.* Continuous Glucose Monitoring in the Intensive Care Unit During
the COVID-19 Pandemic. *Diabetes Care* **44**, 847–849 (2021).
9. Hansen, E. A. *et al.* Accuracy, satisfaction and usability of a flash glucose
monitoring system among children and adolescents with type 1 diabetes
attending a summer camp. *Pediatric Diabetes* **19**, 1276–1284 (2018).
10. Battelino, T. *et al.* Clinical targets for continuous glucose monitoring data
interpretation: Recommendations from the international consensus on time in
range. *Diabetes Care* **42**, 1593–1603 (2019).
11. Ceriello, A., Monnier, L. & Owens, D. Glycaemic variability in diabetes: clinical
and therapeutic implications. *The Lancet Diabetes and Endocrinology* **7**, 221–
230 (2019).

12. Danne, T. *et al.* International consensus on use of continuous glucose
monitoring. *Diabetes Care* **40**, 1631–1640 (2017).
13. Monnier, L. *et al.* Toward defining the threshold between low and high glucose
variability in diabetes. *Diabetes Care* **40**, 832–838 (2017).
14. Berry, S. E. *et al.* Human postprandial responses to food and potential for
precision nutrition. *Nature Medicine* **26**, 964–973 (2020).
15. Korem, T. *et al.* Bread Affects Clinical Parameters and Induces Gut Microbiome-
Associated Personal Glycemic Responses. *Cell Metabolism* **25**, 1243-1253.e5
(2017).
16. Zeevi, D. *et al.* Personalized Nutrition by Prediction of Glycemic Responses. *Cell*
163, 1079–1095 (2015).
17. Flanagan, A., Bechtold, D. A., Pot, G. K. & Johnston, J. D. Chrono-nutrition: From
molecular and neuronal mechanisms to human epidemiology and timed feeding
patterns. *Journal of Neurochemistry* **157**, 53–72 (2021).
18. Levi, F. & Schibler, U. Circadian Rhythms: Mechanisms and Therapeutic
Implications. *Annual Review of Pharmacology and Toxicology* **47**, 593–628
(2007).
19. Qian, J. & Scheer, F. A. J. L. Circadian System and Glucose Metabolism:
Implications for Physiology and Disease. *Trends in Endocrinology and*
Metabolism **27**, 282–293 (2016).
20. Van Cauter, E., Polonsky, K. S. & Scheen, A. J. Roles of circadian rhythmicity
and sleep in human glucose regulation. *Endocrine Reviews* **18**, 716–738 (1997).
21. Aparicio, N. J. *et al.* Circadian variation of the blood glucose, plasma insulin and
human growth hormone levels in response to an oral glucose load in normal
subjects. *Diabetes* **23**, 132–137 (1974).
22. Porcellati, F., Lucidi, P., Bolli, G. B. & Fanelli, C. G. Thirty Years of Research on
the Dawn Phenomenon: Lessons to Optimize Blood Glucose Control in Diabetes.
Diabetes Care **36**, 3860–3862 (2013).
23. Schmidt, M. I., Hadji-Georgopoulos, A., Rendell, M., Margolis, S. & Kowarski, A.
The dawn phenomenon, an early morning glucose rise: Implications for diabetic
intraday blood glucose variation. *Diabetes Care* **4**, 579–585 (1981).
24. Johnson, M. L. *et al.* Utilizing the Ambulatory Glucose Profile to Standardize and
Implement Continuous Glucose Monitoring in Clinical Practice. *Diabetes*
Technology and Therapeutics **21**, S2-17-S2-25 (2019).
25. Shaffer, F. & Ginsberg, J. P. An Overview of Heart Rate Variability Metrics and
Norms. *Frontiers in Public Health* **5**, 1–17 (2017).
26. Singh, J. P. *et al.* Association of hyperglycemia with reduced heart rate variability
(The Framingham Heart Study). *American Journal of Cardiology* **86**, 309–312
(2000).
27. Wu, J. S. *et al.* Epidemiological evidence of altered cardiac autonomic function in
subjects with impaired glucose tolerance but not isolated impaired fasting
glucose. *Journal of Clinical Endocrinology and Metabolism* **92**, 3885–3889
(2007).
28. Ewing, D. J., Campbell, I. W. & Clarke, B. F. Mortality in diabetic autonomic
neuropathy. *The Lancet* **307**, 601–603 (1976).
29. Wehrwein, E. A., Orer, H. S. & Barman, S. M. Overview of the Anatomy,
Physiology, and Pharmacology of the Autonomic Nervous System.
Comprehensive Physiology **6**, 1239–1278 (2016).

30. Shaffer, F., McCraty, R. & Zerr, C. L. A healthy heart is not a metronome: an
integrative review of the heart's anatomy and heart rate variability. *Frontiers in*
Psychology **5**, 1–19 (2014).
31. Cobelli, C. *et al.* Diabetes : Models , Signals , and Control. **2**, 54–96 (2009).
32. McGrath, T., Murphy, K. G. & Jones, N. S. Quantitative approaches to energy
and glucose homeostasis: Machine learning and modelling for precision
understanding and prediction. *Journal of the Royal Society Interface* **15**, 1–33
(2018).
33. Pattaranit, R. & Van Den Berg, H. A. Mathematical models of energy
homeostasis. *Journal of the Royal Society Interface* **5**, 1119–1135 (2008).
34. Dalla Man, C., Caumo, A. & Cobelli, C. The oral glucose minimal model:
Estimation of insulin sensitivity from a meal test. *IEEE Transactions on*
Biomedical Engineering **49**, 419–429 (2002).
35. Dalla Man, C., Rizza, R. A. & Cobelli, C. Meal simulation model of the glucose-
insulin system. *IEEE Transactions on Biomedical Engineering* **54**, 1740–1749
(2007).
36. Deng, Y. *et al.* Deep transfer learning and data augmentation improve glucose
levels prediction in type 2 diabetes patients. *npj Digital Medicine* **4**, (2021).
37. Li, K., Daniels, J., Liu, C., Herrero, P. & Georgiou, P. Convolutional Recurrent
Neural Networks for Glucose Prediction. *IEEE Journal of Biomedical and Health*
Informatics **24**, 603–613 (2020).
38. Rubin-Falcone, H., Fox, I. & Wiens, J. Deep residual time-series forecasting:
Application to blood glucose prediction. *CEUR Workshop Proceedings* **2675**,
105–109 (2020).
39. Bent, B. *et al.* Engineering digital biomarkers of interstitial glucose from
noninvasive smartwatches. *npj Digital Medicine* **4**, 1–11 (2021).
40. Rashtian, H., Torbaghan, S. S., Rahili, S., Snyder, M. & Aghaeepour, N. Heart
Rate and CGM Feature Representation Diabetes Detection from Heart Rate:
Learning Joint Features of Heart Rate and Continuous Glucose Monitors Yields
Better Representations. *IEEE Access* **9**, 83234–83240 (2021).
41. Woldaregay, A. Z. *et al.* Data-driven modeling and prediction of blood glucose
dynamics: Machine learning applications in type 1 diabetes. *Artificial Intelligence*
in Medicine **98**, 109–134 (2019).
42. Boughton, C. K. & Hovorka, R. New closed-loop insulin systems. *Diabetologia*
64, 1007–1015 (2021).
43. Gill, S. & Panda, S. A Smartphone App Reveals Erratic Diurnal Eating Patterns in
Humans that Can Be Modulated for Health Benefits. *Cell Metabolism* **22**, 789–
798 (2015).
44. Chow, L. S. *et al.* Time-Restricted Eating Effects on Body Composition and
Metabolic Measures in Humans who are Overweight: A Feasibility Study. *Obesity*
28, 860–869 (2020).
45. Kruschke, J. K. Bayesian estimation supersedes the t test. *Journal of*
Experimental Psychology: General **142**, 573–603 (2013).
46. Wyatt, P. *et al.* Energy Intake in Healthy Individuals. *Nature Metabolism* **3**,
(2021).
47. Bowman, C. *et al.* A method for characterizing daily physiology from widely used
wearables. *Cell Reports Methods* 100058 (2021)
doi:10.1016/j.crmeth.2021.100058.
48. Schroeder, E. B. *et al.* Diabetes, Glucose, Insulin, and Heart Rate Variability.
Diabetes Care **28**, 668–674 (2005).

49. Tambascia, M. A. ntonio, Malerbi, D. A. ugusto C. & Eliaschewitz, F. G. oldberg.
Influence of gastric emptying on the control of postprandial glycemia: physiology
and therapeutic implications. *Einstein (São Paulo, Brazil)* **12**, 251–253 (2014).
50. Moore, M. C., Coate, K. C., Winnick, J. J., An, Z. & Cherrington, A. D. Regulation
of hepatic glucose uptake and storage in vivo. *Advances in Nutrition* **3**, 286–294
(2012).
51. Bansal, P. & Wang, Q. Insulin as a physiological modulator of glucagon
secretion. *American Journal of Physiology - Endocrinology and Metabolism* **295**,
(2008).
52. Petrenko, V. *et al.* In pancreatic islets from type 2 diabetes patients, the
dampened circadian oscillators lead to reduced insulin and glucagon exocytosis.
Proc. Natl. Acad. Sci. U.S.A. **117**, 2484–2495 (2020).
53. Saad, A. *et al.* Diurnal pattern to insulin secretion and insulin action in healthy
individuals. *Diabetes* **61**, 2691–2700 (2012).
54. Sinturel, F. *et al.* Cellular circadian period length inversely correlates with HbA1c
levels in individuals with type 2 diabetes. *Diabetologia* **62**, 1453–1462 (2019).
55. Chaix, A., Manoogian, E. N. C., Melkani, G. C. & Panda, S. Time-Restricted
Eating to Prevent and Manage Chronic Metabolic Diseases. *Annual review of
nutrition* **39**, 291–315 (2019).
56. Soru, P. *et al.* MPC based Artificial Pancreas: Strategies for individualization and
meal compensation. *Annual Reviews in Control* **36**, 118–128 (2012).
57. Toffanin, C. *et al.* Artificial pancreas: Model predictive control design from clinical
experience. *Journal of Diabetes Science and Technology* **7**, 1470–1483 (2013).
58. Coorey, G., Figtree, G. A., Fletcher, D. F. & Redfern, J. The health digital twin:
advancing precision cardiovascular medicine. *Nature Reviews Cardiology* **18**,
803–804 (2021).
59. Kamel Boulos, M. N. & Zhang, P. Digital twins: From personalised medicine to
precision public health. *Journal of Personalized Medicine* **11**, (2021).
60. Roy, A. & Parker, R. S. Dynamic modeling of exercise effects on plasma glucose
and insulin levels. *Journal of Diabetes Science and Technology* **1**, 338–347
(2007).
61. Zhu, T. *et al.* Enhancing self-management in type 1 diabetes with wearables and
deep learning. *npj Digital Medicine 2022 5:1* **5**, 1–11 (2022).
62. Craig, C. L. *et al.* International physical activity questionnaire: 12-Country
reliability and validity. *Medicine and Science in Sports and Exercise* **35**, 1381–
1395 (2003).
63. Roenneberg, T., Wirz-Justice, A. & Merrow, M. Life between clocks: Daily
temporal patterns of human chronotypes. *Journal of Biological Rhythms* **18**, 80–
90 (2003).
64. Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R. & Kupfer, D. J. The
Pittsburgh sleep quality index: A new instrument for psychiatric practice and
research. *Psychiatry Research* **28**, 193–213 (1989).
65. De, A. G. *et al.* GPflow: A Gaussian Process Library using TensorFlow Mark van
der Wilk. *Journal of Machine Learning Research* **18**, 1–6 (2017).
66. Särkkä, Simo & Solin, Arno. Applied Stochastic Differential Equations.
(Cambridge University Press, 2019).
67. Martín Abadi, Ashish Agarwal, Paul Barham, E. B., Zhifeng Chen, Craig Citro,
Greg S. Corrado, A. D. & Jeffrey Dean, Matthieu Devin, S. G. TensorFlow: Large-
Scale Machine Learning on Heterogeneous Distributed Systems. (2015)

68. Kass, R. E. & Raftery, A. E. Bayes Factors. *Journal of the American Statistical*
Association **90**, 773–795 (1995).
993

Figure 1

Figure 2

A Model 1

B Model 2

C Model 3

Circadian clocks

Food + drinks

Heart rate variability

Glucose levels

Heart rate

Physical activity

D

Decreased half-life

Ingestion event (t = 0)

Response height

Response half-life

Underdamped

Overshoot

Glucose (mmol/L)

Time (hours)

Figure 3

Figure 4

Figure 6

A Activity → glucose

B HR → glucose

C HRV → glucose

D

E ID 04

F ID 08

G ID 08 – Cross-correlation of glucose vs. physical + heart activity weighted sum