Humoral and cellular response induced by a second booster of an inactivated

SARS-CoV-2 vaccine in adults.

Felipe Melo-González[1,2,3#], Constanza Méndez[1,2#], Hernán F Peñaloza[1,2#], Bárbara

M Schultz[1,2#], Alejandro Piña-Iturbe[1,2], Mariana Ríos[1,2], Daniela Moreno-Tapia[1,2],

Patricia Pereira-Sánchez[1,2], Diane Leighton[1,2], Claudia Orellana[1,2], Consuelo

Covarrubias[1,2], Nicolás MS Gálvez[1,2], Jorge A Soto[1,2,3], Luisa F Duarte[1,2], Daniela

Rivera-Pérez[1,2], Yaneisi Vázquez[1,2], Alex Cabrera[4], Sergio Bustos[4], Carolina

Iturriaga[5], Marcela Urzua[5], María S Navarrete[6], Álvaro Rojas[6], Rodrigo Fasce[7],

Jorge Fernández[7], Judith Mora[7], Eugenio Ramírez[7], Aracelly Gaete-Argel[1,8],

Mónica Acevedo[1,8], Fernando Valiente-Echeverría[1,8], Ricardo Soto-Rifo[1,8], Daniela

Weiskopf[9], Alba Grifoni[9], Alessandro Sette[9,10], Gang Zeng[11], Weining Meng[11],

CoronaVac03CL Study Group, José V González-Aramundiz[12], Pablo A

González[1,2], Katia Abarca[1,5], Susan M Bueno*[1,2,14], Alexis M Kalergis*[1,2,13,14].

[1]Millennium Institute on Immunology and Immunotherapy, Santiago, Chile.

[2]Departamento de Genética Molecular y Microbiología, Facultad de Ciencias

Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile.

[3]Departamento de Ciencias Biológicas, Facultad de Ciencias de la Vida,

Universidad Andrés Bello, Santiago, Chile. [4]Flow Cytometry Facility, Facultad de

Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile.

[5]Departamento de Enfermedades Infecciosas e Inmunología Pediátrica, División

de Pediatría, Escuela de Medicina, Pontificia Universidad Católica de Chile,

Santiago, Chile. [6]Departamento de Enfermedades Infecciosas del Adulto, División

de Medicina, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. [7]Departamento de Laboratorio Biomédico, Instituto de Salud Pública de Chile. [8]Laboratorio de Virología Molecular y Celular, Programa de Virología, Instituto de Ciencias Biomédicas, Facultad de Medicina, Universidad de Chile, Santiago de Chile. [9]Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology (LJI), La Jolla, CA 92037, USA. [10]Department of Medicine, Division of Infectious Diseases and Global Public Health, University of California, San Diego (UCSD), La Jolla, CA 92037, USA. [11]Sinovac Biotech, Beijing, China. [12]Departamento de Farmacia, Facultad de Química y de Farmacia, Pontificia Universidad Católica de Chile, Santiago, Chile. [13]Departamento de Endocrinología, Facultad de Medicina, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile. [14]Lead contacts.

*Corresponding author:

Alexis M Kalergis, Pontificia Universidad Católica de Chile. Av. Libertador Bernardo O'Higgins Nº 340, Santiago 8331010, Santiago, Chile. Phone 56-2-6862846. Email: akalergis@bio.puc.cl.

*Alternate corresponding author:

Susan M Bueno, Pontificia Universidad Católica de Chile. Av. Libertador Bernardo O'Higgins Nº 340, Santiago 8331010, Santiago, Chile. Phone 56-2-6862846. Email: sbueno@bio.puc.cl

[#]These authors contributed equally to this work.

Summary

The SARS-CoV-2 Omicron variant has challenged the control of the COVID-19

pandemic even in highly vaccinated countries. While a second booster of mRNA

vaccines improved the immunity against SARS-CoV-2, the humoral and cellular

responses induced by a second booster of an inactivated SARS-CoV-2 vaccine

have not been studied. In the context of a phase 3 clinical study, we report that a

second booster of CoronaVac$^®$ increased the neutralizing response against the

ancestral virus yet showed poor neutralization against the Omicron variant.

Additionally, isolated PBMCs displayed equivalent activation of specific CD4$^+$ T

lymphocytes and IFN-γ production when stimulated with a mega-pool of peptides

derived from the spike protein of the ancestral virus or the Omicron variant. In

conclusion, a second booster dose of CoronaVac$^®$ does not improve the

neutralizing response against the Omicron variant compared with the first booster

dose, yet it helps maintain a robust spike-specific CD4$^+$ T cell response.

Key words: CoronaVac$^®$, second booster dose, SARS-CoV-2, Omicron variant,

humoral immunity, cellular immunity.

Introduction.

The development of vaccines that grant long-lasting protection against SARS-CoV-2 is essential to control the current COVID-19 pandemic. Although several vaccines were developed in record time, three dynamic phenomena have prevented the global control of the COVID-19 pandemic. First, the continuous emergence of SARS-CoV-2 variants of concern (VOCs), such as Omicron (BA.1, BA.2), and its subvariants (BA.2.12.1, BA.4, BA.5) with high transmissibility and immune evasion profiles[1]. Second, the waning of neutralizing antibodies in fully vaccinated subjects[2]. Third, the difficulties to mass producing and globally distributing enough vaccines or implementing affordable and effective vaccination programs.

Several platforms have been used to develop vaccines against SARS-CoV-2[3]. Due to their novelty, mRNA vaccines (BNT162b2 and mRNA-1273) have been the most studied and are highly effective in protecting individuals from symptomatic infection, severe disease, and death[4]. Inactivated virus-based vaccine, a more traditional vaccine platform, has also been used to develop a SARS-CoV-2 vaccine. CoronaVac®, an inactivated vaccine developed by Sinovac Life Sciences Co., Ltd. (Beijing, China)[5], has been administered so far in 52 countries[6], showing a good safety profile in the population[7-9] and a robust immune protection against severe disease, hospitalization, and death[10].

During a phase 3 clinical trial in Chile, our group demonstrated that a two-dose vaccination schedule of CoronaVac® induced a strong neutralizing response and T cell activation against SARS-CoV-2 in adults[11]. Further studies from our laboratory determined that fully vaccinated subjects with CoronaVac® showed a strong

neutralizing response and IFN-γ production in stimulated peripheral blood

mononuclear cells (PBMCs), against different VOCs of SARS-CoV-2, such as

Alpha, Beta, Gamma, and Delta[12].

Different studies have reported a reduction of the neutralizing response against

SARS-CoV-2 in immunized subjects with BNT162b2 and mRNA1273 vaccines[13-15].

These studies also showed that a booster dose was required to keep an effective

neutralizing response against the ancestral SARS-CoV-2 (WT SARS-CoV-2) and

circulating variants at that time[13,14,16].

Consistently with the mentioned studies, our group reported a considerable

reduction of the neutralizing response five months after the administration of the

second dose of CoronaVac®, response that was recovered after the administration

of a booster dose of CoronaVac®[17]. Furthermore, the enhanced neutralizing

response initially detected against WT SARS-CoV-2 was effective against the

Delta variant, but it failed to neutralize the Omicron variant[17], which is currently the

most prevalent variant of SARS-CoV-2 worldwide.

Although previous studies have shown that a second booster with BNT162b2 and

mRNA1273 vaccines would prevent the decrease of neutralizing antibodies and

may offer protection against symptomatic disease, severe disease, hospitalization,

and death caused by the Omicron variant[4,18], the effect of a second booster dose

of CoronaVac® in the humoral and cellular response against SARS-CoV-2, with

special emphasis on the Omicron variant remains to be elucidated. In the present

report, we study the dynamics of the humoral and cellular immune responses in

individuals that received a second booster of CoronaVac® 6 months after the

administration of a first booster of the same vaccine. Our data shows that a second

booster of CoronaVac[®] induces a strong production of antibodies with neutralizing

capacities against WT SARS-CoV-2, although it has poor activity against the

Omicron variant. We also show that a second booster dose of CoronaVac[®] is

required to keep elevated levels of SARS-CoV-2-specific CD4[+] T lymphocytes in

circulation that are reactive against the WT SARS-CoV-2, the Delta, and the

Omicron variants.

Results.

1. Participants, sampling and experimental applications included in the study.

From a total of 2,302 individuals enrolled in the clinical trial CoronaVac03CL (clinicaltrials.gov #NCT04651790) in Chile (November 2020- to current date), 138 fully vaccinated subjects with CoronaVac[®] (0-28 schedule) that received two booster doses were initially considered for this study (Fig 1A). After the exclusion of 51 subjects due to SARS-CoV-2 infection during the trial or missing data, longitudinal analyses of the humoral response were performed in up to 87 subjects, whereas the cellular response was studied in a subgroup of 46 subjects (Fig 1A). Blood samples were collected before vaccination (T1), four-seven weeks after the second dose (T2), at least 9 weeks before the first booster dose (T3), three-six weeks after the first booster dose (T4), at least nine-cero weeks before the second booster dose (T5) and four-nine weeks after second booster dose (T6) (Fig 1B).

2. Humoral response against SARS-CoV-2 induced by a second booster dose of CoronaVac[®].

Neutralizing response of serum was evaluated by three different and complementary methodologies, surrogate virus neutralization test (sVNT), conventional virus neutralization test (cVNT), and pseudotype virus neutralization test (pVNT) (see method section). Consistently with previous studies[11,17], individuals vaccinated with two doses of CoronaVac[®] presented a significant increase in neutralizing antibodies against WT SARS-CoV-2 four weeks after the administration of the second dose when compared with the pre-immune serum (T1

vs. T2) (16.8 vs. 199.7 GMUs p<0.0001; 2.9 vs. 21.1 GMTs p<0.0001) (Fig. 2A-B,

supplementary tables 1 and 3). Then, a significant reduction in the neutralizing

antibody against SARS-CoV-2 was observed three to five months (15-20 weeks)

after the administration of the second dose when compared with four weeks after

the administration of the second dose (T2 vs. T3) (199.7 vs. 53.1 GMUs p<0.0001;

21.1 vs. 10.0 GMTs p<0.0001) (Fig. 2A-B supplementary table 1 and 3).

Consistently with other studies[13-15], the administration of a first booster dose

resulted in a rapid improvement of the neutralizing response against WT SARS-

CoV-2 after 4 weeks when compared with at least nine weeks before its

administration (T3 vs. T4) (53.1 vs. 586.0 GMUs p<0.0001; 10.0 vs. 95.1 GMTs

p<0.0001) (Fig. 2A-B supplementary table 1 and 3). Interestingly, the neutralizing

capacity against WT SARS-CoV-2 observed six months (24 weeks) after the

administration of the first boost was partially reduced in comparison to the

response detected 4 weeks after the administration of the first booster dose (T5 vs.

T4) (220.4 vs. 586.0 GMUs p<0.001; 54.9 vs. 95.1GMTs p>0.05) (Fig. 2A-B,

supplementary table 1 and 3). Finally, 4 to 9 weeks after the administration of a

second booster dose, an increase in the neutralizing response against WT SARS-

CoV-2 was observed when compared with the time before the administration of the

second booster dose (T5 vs. T6) (220.4 vs. 549.2 GMUs p<0.001; 54.9 vs. 149.3

GMTs p>0.05) (Fig. 2A-B, supplementary table 1 and 3). Importantly, the

neutralization response against WT SARS-CoV-2 found after the administration of

the second booster dose was not significantly higher than the response induced by

the first booster dose (T4 vs. T6) (586.0 vs. 549.2 GMUs p>0.05; 95.1 vs. 149.3

GMTs p>0.05) (Fig. 2A-B, supplementary table 1 and 3). This data indicates that a

second booster dose of CoronaVac® is required to keep high levels of neutralizing

antibodies against WT SARS-CoV-2. Moreover, given that the neutralization

observed after the first and the second booster dose was equivalent, we

hypothesize that the neutralizing response against SARS-CoV-2 induced by the

administration of the first booster of CoronaVac® has reached a peak of

neutralizing antibodies that are sustained by the administration of a second boost.

The administration of the second booster dose of CoronaVac® kept the

seropositivity rate higher than 96% for both sVNT and cVNT, with seroconversion

levels of 93.1% and 95.2% for sVNT and cVNT, respectively (supplementary table

3).

Next, we evaluated whether the neutralizing antibodies generated after the second

booster dose was effective against the Delta (B.1.617.2) and the Omicron

(B.1.1.529) variants. Although serum from vaccinated individuals collected four

weeks after the second booster dose presented a mildly reduced ability to

neutralize WT SARS-CoV-2 when compared with samples collected four weeks

after the first booster dose (Fig 2C), the neutralization response against WT SARS-

CoV-2 was significantly higher when compared with the Delta (B.1.617.2) and

especially with the Omicron (B.1.1.529) variants (33.0 vs. 3.7 GMTs p<0.001, 8.9-

fold reduction) (Fig 2C, supplementary table 2). Moreover, compared with the first

booster dose, the administration of the second booster dose of CoronaVac® did not

significantly impact the seropositivity rate (93.1% vs. 94.3%) and the

seroconversion (80.5% vs. 67.8%) against the Delta variant, although it slightly

increased the seropositivity rate (17.2% vs. 32.2%) and the seroconversion (4.6%

vs. 18.4%) against the Omicron variant (supplementary table 4).

Neutralization against the Omicron variant was later confirmed with a pseudotype-

based neutralization assay in sixty subjects (supplementary figure 1) and indicate

that a second booster dose of CoronaVac® maintains high neutralizing antibody

levels against WT SARS-CoV-2 that poorly neutralizes the Omicron variant.

3. Cellular response against SARS-CoV-2 induced by a second booster dose

of CoronaVac®.

We next evaluated the cellular response in a subgroup of fully vaccinated

volunteers (n=46) that received two booster doses of CoronaVac® (Fig 1A).

PBMCs were stimulated with mega-pools of peptides of SARS-CoV-2 theoretically

able to activate CD4$^+$ T lymphocytes (S+R) and CD8$^+$ T lymphocytes (CD8A+B)

and SARS-CoV-2-specific OX40$^+$CD137$^+$CD4$^+$T cells (AIM$^+$CD4$^+$ T cells), as well

as SARS-CoV-2 specific CD69$^+$CD137$^+$CD8$^+$ T cells (AIM$^+$CD8$^+$ T cells) were

quantified by flow cytometry.

Our data show that the activation of SARS-CoV-2-specific AIM$^+$CD4$^+$ T cells was

higher after the second dose of CoronaVac® in comparison with the pre-immune

sample (0.13% vs. 0.33% p=0.0241) (Fig. 3A; supplementary table 5). Although no

significant increase in the percentage of AIM$^+$CD4$^+$ T cells was observed after the

administration of the first or the second booster dose, the activation of AIM$^+$CD4$^+$ T

cells has remained stable over time, showing only a significant decrease 4-6

months after the first booster dose, that is successfully recovered after the

administration of the second booster dose (Fig 3A, supplementary table 5).

Further, we did not detect a significant activation of SARS-CoV-2-specific

AIM$^+$CD8$^+$ T cells in fully vaccinated volunteers with CoronaVac® after the

administration of the first or the second booster dose (Fig. 3B; supplementary

table 5).

Next, we evaluated the production of IFN-γ in stimulated-PBMCs by ELISPOT (Fig.

3C-D). The production of IFN-γ by stimulated PBMCs was consistent with the

activation of $CD4^+$ and $CD8^+$ lymphocytes in subjects that received a second

booster dose of CoronaVac[®]. A significant increase in IFN-γ[+] SFCs was observed

in PBMCs stimulated with MP-S+R after the second dose of CoronaVac[®] (4.1-fold

increase in PBMCs stimulated with MP-S+R p=0.0004) (Fig 3C). Moreover, the

increase in IFN-γ[+]SFCs observed after the second dose of CoronaVac[®] remained

stable over time and was also observed after the administration of the first and the

second booster dose (Fig 3C, supplementary table 5). In contrast, no significant

changes in IFN-γ[+]SFCs were observed in PBMCs stimulated with MP-CD8A+B

after the second dose of CoronaVac[®] (1.5-fold increase in PBMCs stimulated with

MP-CD8A+B p>0.05) (Fig. 3D) nor after the administration of a first or second

booster of CoronaVac[®] (Fig 3D, supplementary table 5).

Even though we did not detect a significant increase in the frequency of

AIM[+]CD4[+]T cells after the administration of the first or the second booster doses of

CoronaVac[®], we observed a progressive increment in volunteers with detectable

levels of AIM[+]CD4[+]T cells over time. Whereas 60.9% (28/46) of volunteers

presented detectable levels of SARS-CoV-2-specific AIM[+]CD4[+]T cells before

vaccination, two doses of CoronaVac[®] increased this frequency to 84.9% (39/46),

frequency that was further increased to 93.5% (43/46) due to the first booster of

CoronaVac[®] (Table 1). Consistently, the frequency of subjects whose PBMCs

produced IFN-γ after stimulation with MP-SR increased after the administration of

CoronaVac®, starting from 67.4% (31/46) before vaccination to 95.6% (44/46) after

two doses of CoronaVac® and to 100% (46/46) after the first booster (Table 1). The

second booster of CoronaVac® did not have a significant impact in the frequency of

volunteers that have specific AIM⁺CD4⁺T cells in circulation 91.3% (42/46)

compared with the first booster dose, although it maintained the production of IFN-

γ in the 100% (46/46) of studied volunteers (Table 1).

No major differences were found in the frequency of AIM⁺CD8⁺ T cells after two

doses of CoronaVac® (45.7%, 21/46) compared to before vaccination (43.5%,

20/46) (Table 1). Although, the administration of a first booster increased the

frequency of specific AIM⁺CD8⁺ T cells to 63% (29/46), frequency that drastically

decreased after the second booster dose (52.2%, 24/46) (Table 1). In contrast, the

frequency of IFN-γ-producing PBMCs stimulated with MP-CD8A+B increased over

time, starting at 60.9% (28/46) before vaccination, to 73.9% (34/46) after two doses

of CoronaVac® (Table 1). Frequency that was further increased after the first

booster dose (78.3%, 36/46) and maintained by the second booster dose (78.3%,

36/46) (Table 1).

Finally, we evaluated the activation of AIM⁺CD4⁺ T lymphocytes and IFNγ⁺

production by PBMCs in response to a mega-pool of peptides derived from the

Spike protein of the Delta and the Omicron variants by flow cytometry and

ELISPOT, respectively, four weeks after the administration of the first booster dose

and four weeks after the administration of the second booster dose. Our data show

that the activation AIM⁺CD4⁺ T cells and IFNγ⁺SFCs were equivalent when PBMCs

were stimulated with a mega-pool of peptides derived from the WT SARS-CoV-2,

the Delta, or the Omicron variants (Fig. 3E-F) and no major differences were

detected in the activation of AIM$^+$CD4$^+$ T cells and IFNγ$^+$SFCs in PBMCs, between

the first and the second booster dose against each of these variants (Fig. 3E-F).

Finally, we detected a small decrease in the percentage of subjects that presented

CD4$^+$AIM$^+$, CD8$^+$AIM$^+$ and IFNγ$^+$SFCs after the second booster compared with

after the first booster dose (supplementary table 6). These data show that

CoronaVac$^®$ induces a robust and stable CD4$^+$ T lymphocyte response to react

against the Delta and the Omicron variants after a second booster dose. Although

the humoral immunity against Delta and Omicron variants decreases, cellular

immunity remains stable across time and reacts against these variants.

Discussion.

In line with other reports[4,18], our data show that a second booster dose of

CoronaVac$^®$ restores the neutralizing response against the RBD of SARS-CoV-2 to

similar levels reached after the administration of the first booster dose.

Neutralizing antibodies induced by vaccines has been acknowledged as the first

line of defense against SARS-CoV-2 infection. However, follow-up studies have

shown that fully vaccinated individuals show a gradual decrease in their levels of

circulating neutralizing antibodies over time[13-15,17]. This response can be restored

with the administration of a booster dose[13,17], and some studies have even shown

that a second booster dose may grant better protection against severe disease,

hospitalization, and death due to COVID-19[4,19].

Our data show that despite a second booster dose induces the production of

neutralizing antibodies against WT SARS-CoV-2, these antibodies only weakly

neutralized the Omicron variant. These results are consistent with other studies

that show that the Omicron variant and its subvariants are incredibly efficient in

evading serum neutralization from individuals who have received one or two

booster doses[1,18,20]. Therefore, it is unlikely that nowadays, the increased

protection against severe disease, hospitalization, and death granted by a second

booster dose could be mainly mediated by the action of neutralizing antibodies.

Nonetheless, we detected a slight but non-significant increment of 1.6-fold change

in the neutralization of WT SARS-CoV-2 between four weeks after the first booster

dose and four weeks after the second booster dose by cVNT (Fig 1B). This modest

increase in WT SARS-CoV-2 neutralization granted by the second booster was not

observed through sVNT (Fig 1A, C). sVNT directly neutralizes the binding of the

RBD with ACE-2, neglecting the potential role of neutralizing antibodies against

other portions of the Spike protein and even against other viral proteins. Therefore,

it is possible that neutralizing antibodies against other domains of the Spike protein

and potentially against other proteins such as the membrane or the nucleocapsid

proteins actively not considered in the sVNT approach may participate in viral

neutralization.

The second layer of protection corresponds to the proliferation, activation, and

activity of T lymphocytes. Currently, the leading hypothesis regarding the protective

mechanism of SARS-CoV-2 vaccines against severe disease and death relies on

the induction of long-lasting T lymphocyte responses rather than on the availability

of circulating neutralizing antibodies[21,22].

In the present study, we report that the activation and proliferation of AIM^+CD4^+ T

lymphocytes in fully vaccinated individuals remain stable over time after

administering a first and a second booster of CoronaVac®. In addition, one study

has identified a robust activation of CD4$^+$OX40$^+$CD137$^+$ T lymphocytes,

CD4$^+$OX40$^+$sCD40L$^+$ T lymphocytes, and follicular CD4$^+$CXCR5$^+$OX40$^+$ T

lymphocytes in individuals that received two doses of mRNA-1273, BNT162b2 or

NVX-CoV2373 (Novavax), and one dose of Ad26.COV2.S (Janssen)[23]. Our data is

consistent with these findings, although the ability of CoronaVac$^®$ to induce a

robust memory T cell response remains to be elucidated.

On the other hand, it has also been reported a significant expansion and activation

of IFN-γ$^+$CD8$^+$ T lymphocytes in PBMCs of individuals vaccinated with two doses

of mRNA-1273, BNT162b2, and NVX-CoV2373, and one dose of Ad26.COV2.S[23].

However, in our experimental setting, we did not find a significant increase in

CD8$^+$AIM$^+$ T lymphocytes (Fig. 3B) nor a significant increase in IFN-γ production

(Fig 3D) in stimulated PBMCs from subjects immunized with CoronaVac$^®$ when

compared with the pre-immune sample. In contrast, previous studies have

described that two doses of CoronaVac$^®$ induce the production of IFN-γ by CD8$^+$ T

lymphocytes[24,25]. Therefore, more exhaustive studies aimed to understand how

CoronaVac$^®$ influences the expansion and activation of CD8$^+$ T lymphocytes,

memory T cells, and memory B cells are required to fully elucidate the protective

mechanisms driven by CoronaVac$^®$ in immunized adults, children, and the elderly.

Four interesting findings emerge from our study. First is the presence of

neutralizing antibodies (Fig 2A-B) in pre-immune samples of some individuals.

Most of the pre-immune neutralizing response was identified against the infective

virus. Asymptomatic SARS-CoV-2 infections[26] and antibodies generated against

non-spike proteins of seasonal coronaviruses[27] could explain this response,

although serologic studies are needed to confirm these hypotheses. Moreover, our

data show that whereas 28/46 and 20/46 of the analyzed volunteers presented

SARS-CoV-2-specific AIM^+CD4^+ or AIM^+CD8^+ T cells, respectively before the

administration of CoronaVac[®]; 4/46 and 22/46 volunteers did not present

AIM^+CD4^+ or AIM^+CD8^+ T cells, respectively after the second booster of

CoronaVac[®]. A previous study has shown a possible cross-reactive response of

seasonal coronaviruses-specific T cells against SARS-CoV-2[28], and it is likely the

reason why a high proportion of unvaccinated individuals presented SARS-CoV-2

specific T cells, although asymptomatic infections cannot be ruled out. Further

studies are needed to fully understand the factors involved in the generation and

maintain of an efficient T cell response and the effect on vaccine-induced

protection.

Controlling the COVID-19 pandemic requires multiple efforts to prevent severe

disease and death of infected patients and reduce viral infection and circulation in

the community.

The data provided in this report and other studies suggest that CoronaVac[®] and

other current vaccines effectively protect the population from severe disease,

hospitalization, and death. However, the immune response induced by these

vaccines poorly neutralizes the circulating SARS-CoV-2 variants and cannot

prevent viral infection.

Therefore, new strategies that include the design of new vaccines that target the

current variants, new types of vaccines that enforce immunity in the upper

respiratory tract, and global vaccine distribution programs are essential to control

and end the COVID-19 pandemic.

Methods.

Volunteers and sample collection

Blood samples were obtained from volunteers recruited in the clinical trial

CoronaVac03CL (clinicaltrials.gov #NCT04651790) in Chile (November 2020- to

current date).

Of the 2,302 individuals enrolled at baseline, 409 subjects entered the study and

received two doses with the homologous CoronaVac[®] 0-28 schedule, and 138

subjects ended up receiving two booster doses of CoronaVac[®]. In addition, blood

samples from 87 subjects were collected from before vaccination (T0) to up to four-

nine weeks after the second booster administration (T6). Volunteers who did not

have a blood sample at some of the planned times of the study or who had

previously exceeded the definitive time limit and volunteers who had developed

SARS-COV-2 during the study were excluded (Figure 1A).

Volunteers received two doses of CoronaVac® (3 µg or 600SU of inactivated

SARS-CoV-2 inactivated in the presence of alum adjuvant) in a four-week interval

(0-28 days), a booster dose five months after the second dose, and a second

booster dose 6 months after the first (Figure 1B). Blood samples collected 6 times

were analyzed: T1: Pre-immune, T2: 2[nd] dose + 4 weeks (+3 weeks), T3: before

the administration of the third dose (-9 weeks), T4: at 4 weeks (±2 weeks) after the

third dose, T5: before the administration of the fourth dose (-9 weeks), T6: and at 4

weeks (+5 weeks) after the fourth dose (T6), and vaccination schedule.

Experimental procedures

Neutralizing antibodies against RBD of WT SARS-CoV-2 were measured in the

serum of 87 volunteers recruited in the clinical trial CoronaVac03CL

(clinicaltrials.gov #NCT04651790). Briefly, blood samples were obtained at T1, T2,

T3, T4, T5, and T6. Neutralization of SARS-CoV-2 RBD by circulating antibodies at

each time point was evaluated by a surrogate virus neutralization test (sVNT)

(Genscript Cat#L00847-A)[11]. The antibody levels in international units per mL

(IU/mL) were estimated by interpolating the sVNT absorbance data in the standard

curve made with the WHO International Standard 20/136, using the 4-parameter

Logistic model. In addition, sVNT was used to evaluate the neutralizing capacity of

the tools against the Delta and Omicron variants, the RBDs for the S protein of

SARS-CoV-2 for Delta (Cat# Z03516) and Omicron B.1.1.529 (Cat#Z03730) in 87

volunteers.

Infective virus neutralization assays were performed as previously described[11,17].

Briefly, Vero E6 cells (4×10^4 cells/well) were plated in 96-well plates. 100 μL of

33782CL-SARS-CoV-2 (100 $TCID_{50}$) were incubated with serial dilutions of heat-

inactivated individual serum samples (dilutions of 1:4, 1:8, 1:16, 1:32, 1:64, 1:128,

1:256, and 1:512) for 1h at 37 °C. Then, the mix was added to the 96-well plates

with the Vero E6 cells, and the cytopathic effect was analyzed after 7 days. A

serum sample from uninfected patients (negative control) and a neutralizing

COVID-19 patient serum sample (positive control) was used for each test.

A pseudotyped virus neutralization test (pVNT) assay was performed to assess the

capacity of the antibodies against SARS-CoV-2 VOC in samples from forty-eight

volunteers as previously reported[29]. Briefly, a HIV-1 backbone expressing firefly

luciferase as a reporter gene and pseudotyped with the SARS-CoV-2 spike

glycoproteins (HIV-1-SΔ19) from lineage B.1 (D614G) and variant Omicron (A67V,

ΔH69-V70, T95I, Y145D, ΔG142 -V143- Y144, ΔN211, EPE 213-214, G339D,

S371L, S373P, S375F, K417N, N440K, G446S, S477N, T478K, E484A, Q493R,

G496S, Q498R, N501Y, T547K, D614G, H655Y, N679K, P681H, N764K, N865K,

Q954H, N969K, L981F) was prepared. Serum samples were diluted, and the

estimation of the ID50 was obtained using a 4-parameter nonlinear regression

curve fit measured as the percent of neutralization determined by the difference in

average relative light units (RLU) between test samples and pseudotyped virus

controls.

Seropositivity is considered when titers are increased compared to pre-immune

condition[30]. On the other hand, seroconversion was thought to be when the titer of

neutralizing results increased 4 times with respect to the pre-immune condition.

The expression of Activation-Induced Markers (AIM) by T cells and the number of

Spot Forming Cells (SFC) for IFN-γ were determined by ELISPOT and were

evaluated by flow cytometry in a subgroup of 46 volunteers as previously

described[11,12,17]. Briefly, PBMCs were stimulated with mega-pools (MPs) of

peptides derived from SARS-CoV-2: MP-S, MP-R, MP-CD8-A, MP-CD8-B, Delta,

and Omicron for 24h or 48h for flow cytometry and ELISPOT respectively. Then, 3-

5×10^5 cells per well were plated and stimulated with MP-S, MP-R, MP-CD8-A,

MP-CD8-B, DMSO (negative control), or Phorbol-12-myristate-13-acetate

(1.62mM) (Sigma #P8139)/Ionomycin (0.6mM) (Sigma #I0634). Antibodies used to

identify $AIM^+CD4^{+,}$ and AIM^+CD8^+ T lymphocytes by flow cytometry are detailed in

supplementary table 7. Samples were analyzed in a BD LSR-FORTESSA flow

cytometer located in the flow cytometry core at the Pontificia Universidad Católica

de Chile.

For ELISPOT, IFN-γ was measured using a Immunospot® (#hIFNgIL-4M-10)

following the manufacturer's instructions. Plates were read in an Immunospot S6

Micro Analyzer.

Ethical considerations.

The current study protocol was reviewed and approved by the Institutional

Scientific Ethical Committee of Health Sciences at the Pontificia Universidad

Católica de Chile (#200708006) and the trial was approved by the Chilean Public

Health Institute (#24204/20) and conducted according to the current Tripartite

Guidelines for Good Clinical Practices, the Declaration of Helsinki, and local

regulations. Informed consent was obtained from all volunteers upon enrollment.

Statistical analyses

To statistically compare the neutralizing response against WT SARS-CoV-2,

ANOVA with the Geisser-Greenhouse correction test followed by Sidak multiple

tests were carried out on the log transformed data. Percentage of AIM^+CD4^+ and

AIM^+CD8^+ T cells and IFN-γ production by stimulated PBMCs against WT SARS-

CoV-2 were compared with a non-parametric Friedman test followed by a Dunn's

test for multiple comparisons. To compare the neutralizing and the cellular

responses against WT, Delta and Omicron, a two-way ANOVA test followed by a

Dunn's test for multiple comparisons was used.

The significance level was set at 0.05 for all the analyses. All data were analyzed

with GraphPad Prism 9.3.1.

Funding

The CoronaVac03CL Study was funded by The Ministry of Health, Government of

Chile, the Confederation of Production and Commerce (CPC), Chile and SINOVAC

Biotech. NIH NIAID, under Contract 75N93021C00016, supports AS and Contract

75N9301900065 supports AS, AG and DW. The Millennium Institute on

Immunology and Immunotherapy, **Agencia Nacional de Investigación**

y Desarrollo (ANID) – Millennium Science Initiative Program – ICN09_016 /

ICN 2021_045: Millennium Institute on Immunology and Immunotherapy

(ICN09_016 / ICN 2021_045; former P09/016-F) supports SMB, KA, PAG and

AMK; The Innovation Fund for Competitiveness FIC-R 2017 (BIP Code: 30488811-

0) supports SMB, PAG and AMK.

Competing interests

GZ and WM are SINOVAC Biotech employees and contributed to the

conceptualization of the study (clinical protocol and eCRF design) and did not

participate in the analysis or interpretation of the data presented in the manuscript.

A.S. is a consultant for Gritstone Bio, Flow Pharma, ImmunoScape, Moderna,

AstraZeneca, Avalia, Fortress, Repertoire, Gilead, Gerson Lehrman Group,

RiverVest, MedaCorp, and Guggenheim. La Jolla Institute for Immunology (LJI)

has filed for patent protection for various aspects of T cell epitope and vaccine

design work. All other authors declare no conflict of interest.

Acknowledgments

We would like to thank the support of the Ministry of Health, Government of Chile; Ministry of Science, Technology, Knowledge, and Innovation, Government of Chile; The Ministry of Foreign Affairs, Government of Chile, and the Chilean Public Health Institute (ISP). We also would like to thank Rami Scharf, Jessica White, Jorge Flores and Miren Iturriza-Gomara from PATH for their active support in experimental design and scientific discussion. We also thank the Vice Presidency of Research (VRI), the Direction of Technology Transfer and Development (DTD) and the Legal Affairs Department (DAJ) of the Pontificia Universidad Católica de Chile. We are grateful to the Administrative Directions of the School of Biological Sciences and the School of Medicine of the Pontificia Universidad Católica de Chile for their administrative support. We would also like to thank to the members of the independent data safety monitoring committee (members in the SA) for their oversight, and finally to the subjects enrolled in the study for their participation and commitment with this trial. This project has been funded in whole or in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, under Contract No. 75N93021C00016 to A.S. and Contract No. 75N93019C00065 to A.S and D.W.

Author contributions.

Conceptualization: AMK, KA, SMB, PAG, JVG, GZ, WM, FM-G, CM, HFP, BMS.

Visualization: AMK, KA, SMB, PAG, JVG, GZ, WM.

Methodology and research: FM-G, CM, HFP, BMS, AP-I, MR, DM-T, PP-S, DL,

CO, CC, NMSG, JAS, LFD, DR-P, YV, AC, SB, CI, MU, MSN, AR, RF, JF, JM, ER,

AG-A, MA, FV-E, RS-R, DW, AG, AS, GZ, WM, JVG-A

Data analysis: FM-G, CM, HFP, BMS, AP-I, MR, DM-T, LFD, AG-A, MA, FV-E,

RS-R

Funding acquisition: AMK, SMB.

Project administration: AMK, KA, SMB, PAG.

Supervision: AMK, KA, SMB, PAG.

Writing – original draft: FM-G, CM, HFP, BMS.

Writing – review & editing: AMK, SMB, PAG, FM-G, CM, HFP, BMS.

Verifying underlying data: AMK, SMB, FM-G, CM, HFP, BMS.

Members of the CoronaVac03CL study group are listed in the supplementary

material

References.

1. Planas, D., Saunders, N., Maes, P., Guivel-Benhassine, F., Planchais, C., Buchrieser, J., Bolland, W.H., Porrot, F., Staropoli, I., Lemoine, F., et al. (2022). Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature *602*, 671-675. 10.1038/s41586-021-04389-z.

2. Levin, E.G., Lustig, Y., Cohen, C., Fluss, R., Indenbaum, V., Amit, S., Doolman, R., Asraf, K., Mendelson, E., Ziv, A., et al. (2021). Waning Immune Humoral Response to BNT162b2 Covid-19 Vaccine over 6 Months. N Engl J Med *385*, e84. 10.1056/NEJMoa2114583.

3. Nagy, A., and Alhatlani, B. (2021). An overview of current COVID-19 vaccine platforms. Comput Struct Biotechnol J *19*, 2508-2517. 10.1016/j.csbj.2021.04.061.

4. Magen, O., Waxman, J.G., Makov-Assif, M., Vered, R., Dicker, D., Hernan, M.A., Lipsitch, M., Reis, B.Y., Balicer, R.D., and Dagan, N. (2022). Fourth Dose of BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Setting. N Engl J Med *386*, 1603-1614. 10.1056/NEJMoa2201688.

5. Gao, Q., Bao, L., Mao, H., Wang, L., Xu, K., Yang, M., Li, Y., Zhu, L., Wang, N., Lv, Z., et al. (2020). Development of an inactivated vaccine candidate for SARS-CoV-2. Science *369*, 77-81. 10.1126/science.abc1932.

6. Chen, Z., Zheng, W., Wu, Q., Chen, X., Peng, C., Tian, Y., Sun, R., Dong, J., Wang, M., Zhou, X., et al. (2022). Global diversity of policy, coverage, and demand of COVID-19 vaccines: a descriptive study. BMC Med *20*, 130. 10.1186/s12916-022-02333-0.

7. Tanriover, M.D., Doganay, H.L., Akova, M., Guner, H.R., Azap, A., Akhan, S., Kose, S., Erdinc, F.S., Akalin, E.H., Tabak, O.F., et al. (2021). Efficacy and safety of an inactivated whole-virion SARS-CoV-2 vaccine (CoronaVac): interim results of a double-blind, randomised, placebo-controlled, phase 3 trial in Turkey. Lancet *398*, 213-222. 10.1016/S0140-6736(21)01429-X.

8. Han, B., Song, Y., Li, C., Yang, W., Ma, Q., Jiang, Z., Li, M., Lian, X., Jiao, W., Wang, L., et al. (2021). Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial. Lancet Infect Dis *21*, 1645-1653. 10.1016/S1473-3099(21)00319-4.

9. Seyahi, E., Bakhdiyarli, G., Oztas, M., Kuskucu, M.A., Tok, Y., Sut, N., Ozcifci, G., Ozcaglayan, A., Balkan, II, Saltoglu, N., et al. (2021). Antibody response to inactivated COVID-19 vaccine (CoronaVac) in immune-mediated diseases: a controlled study among hospital workers and elderly. Rheumatol Int *41*, 1429-1440. 10.1007/s00296-021-04910-7.

10. Jara, A., Undurraga, E.A., Gonzalez, C., Paredes, F., Fontecilla, T., Jara, G., Pizarro, A., Acevedo, J., Leo, K., Leon, F., et al. (2021). Effectiveness of an Inactivated SARS-CoV-2 Vaccine in Chile. N Engl J Med *385*, 875-884. 10.1056/NEJMoa2107715.

11. Bueno, S.M., Abarca, K., Gonzalez, P.A., Galvez, N.M.S., Soto, J.A., Duarte, L.F., Schultz, B.M., Pacheco, G.A., Gonzalez, L.A., Vazquez, Y., et al. (2021). Safety and Immunogenicity of an Inactivated SARS-CoV-2

Vaccine in a Subgroup of Healthy Adults in Chile. Clin Infect Dis.
10.1093/cid/ciab823.

12. Melo-Gonzalez, F., Soto, J.A., Gonzalez, L.A., Fernandez, J., Duarte, L.F.,
Schultz, B.M., Galvez, N.M.S., Pacheco, G.A., Rios, M., Vazquez, Y., et al.
(2021). Recognition of Variants of Concern by Antibodies and T Cells
Induced by a SARS-CoV-2 Inactivated Vaccine. Front Immunol *12*, 747830.
10.3389/fimmu.2021.747830.

13. Falsey, A.R., Frenck, R.W., Jr., Walsh, E.E., Kitchin, N., Absalon, J.,
Gurtman, A., Lockhart, S., Bailey, R., Swanson, K.A., Xu, X., et al. (2021).
SARS-CoV-2 Neutralization with BNT162b2 Vaccine Dose 3. N Engl J Med
385, 1627-1629. 10.1056/NEJMc2113468.

14. Choi, A., Koch, M., Wu, K., Chu, L., Ma, L., Hill, A., Nunna, N., Huang, W.,
Oestreicher, J., Colpitts, T., et al. (2021). Safety and immunogenicity of
SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim
analysis. Nat Med *27*, 2025-2031. 10.1038/s41591-021-01527-y.

15. Pegu, A., O'Connell, S.E., Schmidt, S.D., O'Dell, S., Talana, C.A., Lai, L.,
Albert, J., Anderson, E., Bennett, H., Corbett, K.S., et al. (2021). Durability
of mRNA-1273 vaccine-induced antibodies against SARS-CoV-2 variants.
Science *373*, 1372-1377. 10.1126/science.abj4176.

16. Moreira, E.D., Jr., Kitchin, N., Xu, X., Dychter, S.S., Lockhart, S., Gurtman,
A., Perez, J.L., Zerbini, C., Dever, M.E., Jennings, T.W., et al. (2022). Safety
and Efficacy of a Third Dose of BNT162b2 Covid-19 Vaccine. N Engl J Med
386, 1910-1921. 10.1056/NEJMoa2200674.

17. Schultz, B.M., Melo-Gonzalez, F., Duarte, L.F., Galvez, N.M.S., Pacheco,
G.A., Soto, J.A., Berrios-Rojas, R.V., Gonzalez, L.A., Moreno-Tapia, D.,
Rivera-Perez, D., et al. (2022). A Booster Dose of CoronaVac Increases
Neutralizing Antibodies and T Cells that Recognize Delta and Omicron
Variants of Concern. mBio, e0142322. 10.1128/mbio.01423-22.

18. Regev-Yochay, G., Gonen, T., Gilboa, M., Mandelboim, M., Indenbaum, V.,
Amit, S., Meltzer, L., Asraf, K., Cohen, C., Fluss, R., et al. (2022). Efficacy of
a Fourth Dose of Covid-19 mRNA Vaccine against Omicron. N Engl J Med
386, 1377-1380. 10.1056/NEJMc2202542.

19. Arbel, R., Sergienko, R., Friger, M., Peretz, A., Beckenstein, T., Yaron, S.,
Netzer, D., and Hammerman, A. (2022). Effectiveness of a second
BNT162b2 booster vaccine against hospitalization and death from COVID-
19 in adults aged over 60 years. Nat Med *28*, 1486-1490. 10.1038/s41591-
022-01832-0.

20. Tuekprakhon, A., Nutalai, R., Dijokaite-Guraliuc, A., Zhou, D., Ginn, H.M.,
Selvaraj, M., Liu, C., Mentzer, A.J., Supasa, P., Duyvesteyn, H.M.E., et al.
(2022). Antibody escape of SARS-CoV-2 Omicron BA.4 and BA.5 from
vaccine and BA.1 serum. Cell *185*, 2422-2433 e2413.
10.1016/j.cell.2022.06.005.

21. Hurme, A., Jalkanen, P., Heroum, J., Liedes, O., Vara, S., Melin, M.,
Terasjarvi, J., He, Q., Poysti, S., Hanninen, A., et al. (2022). Long-Lasting T
Cell Responses in BNT162b2 COVID-19 mRNA Vaccinees and COVID-19
Convalescent Patients. Front Immunol *13*, 869990.
10.3389/fimmu.2022.869990.

22. Tarke, A., Coelho, C.H., Zhang, Z., Dan, J.M., Yu, E.D., Methot, N., Bloom,
N.I., Goodwin, B., Phillips, E., Mallal, S., et al. (2022). SARS-CoV-2
vaccination induces immunological T cell memory able to cross-recognize
variants from Alpha to Omicron. Cell *185*, 847-859 e811.
10.1016/j.cell.2022.01.015.
23. Zhang, Z., Mateus, J., Coelho, C.H., Dan, J.M., Moderbacher, C.R., Galvez,
R.I., Cortes, F.H., Grifoni, A., Tarke, A., Chang, J., et al. (2022). Humoral
and cellular immune memory to four COVID-19 vaccines. Cell *185*, 2434-
2451 e2417. 10.1016/j.cell.2022.05.022.
24. Escobar, A., Reyes-Lopez, F.E., Acevedo, M.L., Alonso-Palomares, L.,
Valiente-Echeverria, F., Soto-Rifo, R., Portillo, H., Gatica, J., Flores, I.,
Nova-Lamperti, E., et al. (2021). Evaluation of the Immune Response
Induced by CoronaVac 28-Day Schedule Vaccination in a Healthy
Population Group. Front Immunol *12*, 766278. 10.3389/fimmu.2021.766278.
25. Mok, C.K.P., Cohen, C.A., Cheng, S.M.S., Chen, C., Kwok, K.O., Yiu, K.,
Chan, T.O., Bull, M., Ling, K.C., Dai, Z., et al. (2022). Comparison of the
immunogenicity of BNT162b2 and CoronaVac COVID-19 vaccines in Hong
Kong. Respirology *27*, 301-310. 10.1111/resp.14191.
26. Reynolds, C.J., Swadling, L., Gibbons, J.M., Pade, C., Jensen, M.P., Diniz,
M.O., Schmidt, N.M., Butler, D.K., Amin, O.E., Bailey, S.N.L., et al. (2020).
Discordant neutralizing antibody and T cell responses in asymptomatic and
mild SARS-CoV-2 infection. Sci Immunol *5*. 10.1126/sciimmunol.abf3698.
27. Galipeau, Y., Siragam, V., Laroche, G., Marion, E., Greig, M., McGuinty, M.,
Booth, R.A., Durocher, Y., Cuperlovic-Culf, M., Bennett, S.A.L., et al. (2021).
Relative Ratios of Human Seasonal Coronavirus Antibodies Predict the
Efficiency of Cross-Neutralization of SARS-CoV-2 Spike Binding to ACE2.
EBioMedicine *74*, 103700. 10.1016/j.ebiom.2021.103700.
28. Grifoni, A., Weiskopf, D., Ramirez, S.I., Mateus, J., Dan, J.M., Moderbacher,
C.R., Rawlings, S.A., Sutherland, A., Premkumar, L., Jadi, R.S., et al.
(2020). Targets of T Cell Responses to SARS-CoV-2 Coronavirus in
Humans with COVID-19 Disease and Unexposed Individuals. Cell *181*,
1489-1501 e1415. 10.1016/j.cell.2020.05.015.
29. Beltran-Pavez, C., Riquelme-Barrios, S., Oyarzun-Arrau, A., Gaete-Argel,
A., Gonzalez-Stegmaier, R., Cereceda-Solis, K., Aguirre, A., Travisany, D.,
Palma-Vejares, R., Barriga, G.P., et al. (2021). Insights into neutralizing
antibody responses in individuals exposed to SARS-CoV-2 in Chile. Sci Adv
7. 10.1126/sciadv.abe6855.
30. Zhang, Y., Zeng, G., Pan, H., Li, C., Hu, Y., Chu, K., Han, W., Chen, Z.,
Tang, R., Yin, W., et al. (2021). Safety, tolerability, and immunogenicity of
an inactivated SARS-CoV-2 vaccine in healthy adults aged 18-59 years: a
randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. Lancet
Infect Dis *21*, 181-192. 10.1016/S1473-3099(20)30843-4.
653
654

Figure legends.

Figure 1. Study profile, vaccination scheme and sampling, enrolled

volunteers, and cohort included in the study. (A) Schematic representation and

sample distribution of performed experiments. From a total of 138 individuals that

received two booster doses of CoronaVac®, the neutralizing antibodies were

analyzed in blood samples from 87 volunteers by surrogate virus neutralization test

(sVNT), 63 by conventional virus neutralization test (cVNT) and 60 by Pseudovirus-

based neutralization assay (pVNT). sVNT and pVNT were used to evaluate the

neutralizing response induced by a second booster dose against the Delta and/or

the Omicron variants. Cellular immunity was analyzed in blood samples from 46

volunteers at each time point. (B) Blood sampling times, before the first

vaccination/pre-immune (T1), at 4 weeks (+3 weeks) after the second dose (T2),

before the administration of the third dose (-9 weeks) (T3), at 4 weeks (-1/+2

weeks) after the third dose (T4), before the administration of the fourth dose (-9

weeks) (T5) and at 4 weeks (+5 weeks) after the fourth dose (T6), and vaccination

schedule.

Figure 2. Humoral response of volunteers who received a second booster

dose of CoronaVac®. The neutralizing capacity of circulating antibodies in adults

was evaluated in Blood samples collected before the first vaccination/pre-immune

(T1), at 4 weeks (+3 weeks) after the second dose (T2), before the administration

of the third dose (-9 weeks) (T3), at 4 weeks (-1/+2 weeks) after the third dose

(T4), before the administration of the fourth dose (-9 weeks) (T5) and at 4 weeks

(+5 weeks) after the fourth dose (T6), and vaccination schedule. (A) Neutralizing

capacity of circulating antibodies against SARS-CoV-2 in serum of 87 volunteers

was determined by a surrogate Viral Neutralization Test (sVNT) expressed as

IU/ml. (B) Reciprocal dilution of sera required to prevent in vitro infection obtained

sera from 63 adults required to prevent in vitro infection of Hela Cells. Numbers on

top of each data set represents the GMT, and horizontal lines represent the 95%

CI. (C) Geometric mean titter (GMTs) of neutralizing antibodies against the WT-

spike, Delta-spike and Omicron-spike proteins detected in the serum of 87

volunteers immunized with CoronaVac® through sVNT. Dashed line: limit of

detection. Red values under the significance line: indicate a decrease in the means

of the two compared time points; Blue values: indicate an increase in the means of

the two compared time points. (A-B) Data was analyzed with ANOVA with the

Geisser-Greenhouse correction test followed by a post-hoc Sidak multiple test.

*P<0.05; **P<0.01; ***P<0.001; ****P<0.0001. (C) Two-way ANOVA test followed

by a post-hoc Dunn's multiple test. *P<0.05; **P<0.01; ***P<0.001, ****P<0.0001.

Figure 3. Cellular response of volunteers that received a second booster

dose of CoronaVac®. Cellular response in adults was evaluated in PBMCs

collected before the first vaccination/pre-immune (T1), at 4 weeks (+3 weeks) after

the second dose (T2), before the administration of the third dose (-9 weeks) (T3),

at 4 weeks (-1/+2 weeks) after the third dose (T4), before the administration of the

fourth dose (-9 weeks) (T5) and at 4 weeks (+5 weeks) after the fourth dose (T6),

and vaccination schedule. (A) The percentage of AIM+ (OX40+CD137+) CD4+ T

cells and (B) AIM+ (CD69+CD137+) CD8+ T cells was determined in PBMCs of 46

adult volunteers by flow cytometry. PBMCs were stimulated for 24h with mega-

pools of peptides derived from proteins of WT SARS-CoV-2. The number of IFN-g

producing SFCs was determined by ELISPOT upon stimulation for 48h with mega-

pools of (C) S+R peptides or with (D) CD8A+B peptides. (E) The percentage of

AIM+ (OX40+CD137+) CD4+ T cells from PBMCs of 46 adult volunteers were

analyzed by flow cytometry after the stimulation for 24h with mega-pools of

peptides derived from the Spike protein of the WT SARS-CoV-2, the Delta and the

Omicron variants. (F) The number of IFN-g producing SFCs was determined by

ELISPOT assays PBMCs of 46 adult volunteers were analyzed by flow cytometry

after the stimulation for 48h with mega-pools of peptides derived from the Spike

protein of the WT SARS-CoV-2, the Delta and the Omicron variants. Horizontal

lines represent mean and 95% CI. Flow cytometry data was normalized against the

DMSO control. (A-D) Data was analyzed using a non-parametric Friedman test

followed by a post-hoc Dunn's test for multiple comparisons. $*P<0.05$; $**P<0.01$;

$***P<0.001$; $****P<0.0001$. (E-F) Two-way ANOVA test followed by a post-hoc

Dunn's multiple test. $*P<0.05$; $**P<0.01$; $***P<0.001$, $****P<0.0001$.

Table 1: Frequency of volunteers with detectable T cell response against SARS-CoV-2.

Methodology	Indicators Cellular positivity	T1	T2	T3	T4	T5	T6
Flow cytometry	CD4+AIM+(OX40$^+$CD13)	28/46	39/46	40/46	43/46	43/46	42/46
	%	60.9	84.9	86.9	93.5	93.5	91.3
	CD8+AIM+(CD69$^+$CD13)	20/46	21/46	24/46	29/46	25/46	24/46
	%	43.5	45.7	52.2	63.0	54.3	52.2
ELISPOT	IFN-γ$^+$ SFC MP-S+R	31/46	44/46	46/46	46/46	43/46	46/46
	%	67.4	95.6	100	100	93.5	100
	IFN-γ$^+$ SFC MP-CD8A+B	28/46	34/46	29/46	36/46	39/46	36/46
	%	60.9	73.9	63.0	78.3	84.8	78.3

AIM: Activation-Induced-Marker; SFC: Spot Forming Cells; MP-S: Spike mega-pool of peptides