Younger age of initiation of selling sex and depressive symptoms among female sex workers in Eswatini

Ashley Grosso[1*], Rebecca Fielding-Miller[2], Sindy Matse[3], Bhekie Sithole,[4] Stefan Baral[5]

[1]Institute for Health, Health Care Policy and Aging Research, Department of Urban-Global Public
Health, Rutgers School of Public Health, New Brunswick, NJ, United States

[2]Herbert Wertheim School of Public Health, University of California, San Diego, CA, United States

[3]Ministry of Health, Mbabane, Eswatini

[4]Health Communication Capacity Collaborative, Mbabane, Eswatini

[5]Key Populations Program, Center for Public Health and Human Rights, Department of
Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, United States

* Correspondence:
Ashley Grosso
agrosso@ifh.rutgers.edu

Younger age of initiation of selling sex and depressive symptoms among female sex workers in Eswatini

Abstract

Background: Minors who sell sex are likely to have complex mental health needs that may persist
into adulthood. This topic is understudied in Sub-Saharan Africa. This study hypothesized that adult
female sex workers in Eswatini who started selling sex as minors have a higher prevalence of
depression than those who started as adults. We also examined correlates of depression and underage
initiation of selling sex, including stigma and condom-related behaviors.

Methods: From October-December 2014, women aged 18 or older who sold sex in the past 12
months in Eswatini were recruited through venue-based sampling. Participants completed a survey
including the 9-item Patient Health Questionnaire (PHQ-9) and a question about the age at which
they first sold sex for money. T-tests, $\chi 2$ tests and multivariable logistic regression were used to
assess associations.

Results: Overall, 43.1% of participants (332/770) had probable depression, and 16.6% (128/770)
started selling sex as minors under the age of 18. Over half (55.5%, 71/128) of those who started
selling sex as minors had depression. This was significantly higher than the 40.7% (261/642)
prevalence of depression among participants who started selling sex as adults (p=0.002). After
adjusting for confounders, female sex workers who started selling sex as minors had higher odds of
depression than those who started as adults (adjusted odds ratio 1.70, 95% confidence interval 1.11-
2.60). Both depression and underage initiation of selling sex were associated with anticipating stigma
toward sex workers in healthcare settings and the number of times the participant reported a condom
slipped off or broke in the past month.

Conclusion: Results highlight the need for trauma-informed and adolescent-friendly mental health
services in settings free of stigma toward female sex workers in Eswatini.

Keywords: sex workers, Eswatini, depression, adolescent, barriers to care

1   Introduction

Intersectional stigma and depression both commonly affect cisgender female sex workers (FSW) globally.[1,2] The pooled prevalence of depression among FSW from low and middle income countries is 41.8%.[3] Adolescents under the age of 18 who sell sex are considered trafficked or commercially sexually exploited children and may be particularly vulnerable to immediate and long-term mental health problems, including depression.[4] Many FSW across Sub-Saharan Africa started selling sex while underage and experience more negative sexual health outcomes compared to FSW who started as adults.[5-8] However, evidence on the relationship between mental health and underage initiation of selling sex is mixed. In Malawi, FSW who started selling sex while underage were 2.3 times more likely to have probable depression (15%, 5/33 v. 6%, 10/167) compared to those who started as adults, but this was not statistically significant.[9] In Ethiopia, over one third (37.5%, 144/384) of FSW who started selling sex as minors had mild to severe depression, which was slightly higher compared to those who started as adults but not statistically significant (35.9%, 147/409).[10,11] In contrast, in South Africa FSW who started selling sex at younger ages had lower odds of depression or PTSD.[12] Research findings on these topics among FSW in other geographic regions are also conflicting. In the United States, FSW who started selling sex as minors were had higher odds of ever attempting suicide than those who started as adults.[13] However in Bangladesh, FSW with any mental disorder started selling sex at significantly older ages than those without a mental disorder.[14] The researchers surmised that younger FSW may be more resilient than older FSW.

In prior research, both depression and underage initiation of selling sex have been found to be associated with socioeconomic and demographic variables including income,[15,16] food insecurity,[17-19] and orphanhood.[20,21] Depression and selling sex while underage have both been linked to healthcare issues such as perceiving or experiencing sex work-related stigma in healthcare settings[17,22] and not being tested for HIV or aware of one's HIV status.[6,23] Behavioral risk factors, including not carrying condoms[5,24] and reporting condom use errors such as slipping and breaking[25,26] are related to both depression and underage initiation of selling sex. Both FSW with depression and FSW who started selling sex as minors report a longer length of time in the sex trade[8,27] and greater frequency of selling sex[28,29] compared to FSW who are not depressed and FSW who started selling sex as adults.

There remains limited research about mental health in Eswatini (formerly Swaziland), a country in southern Africa. One psychiatric hospital and one psychiatrist are available to serve the population of 1.1 million people.[30,31] The prevalence of depressive disorders in Eswatini is 4%.[32] Over one quarter of people living with HIV in Eswatini are estimated to have depressive symptoms.[33] In a nationally representative study in Eswatini, one in ten reported suicidal ideation, plans and/or attempts in the past 12 months.[34,35] The prevalence of suicidal ideation in the past 12 months is 17% among students aged 12-17[36] but 42% among sexually active female students in that age group.[37] Research has suggested that youth unemployment in Eswatini increases mental health problems and leads to selling sex.[38] Sex work is criminalized in Eswatini.[39] In a 2011 study, 59% of FSW in Eswatini ever had suicidal ideation, and 63% felt sad or had a depressed mood for more than two weeks at a time in the last three years.[40] HIV prevalence among FSW was 60%, compared to about 25% among the adult population overall.[41] About one quarter of FSW reported that they started selling sex as minors.[42] The present study examined the relationship between depression and underage initiation of selling sex among FSW in Eswatini.

2   Methods

2.1   Participants

FSW who exchanged sex for money, favors, or goods in the past 12 months and who were at least 18
years old, assigned female sex at birth, and provided written informed consent in English or SiSwati
were eligible to participate.[43,44] 781 FSW participants completed an in-person interviewer-
administered survey. For these analyses, the sample was restricted to 770 FSW with data on age of
initiation of selling sex and depressive symptoms.

2.2   Data collection

Data collection took place from October to December of 2014 in five towns in Eswatini: Mbabane,
Manzini, Piggs Peak, Lavumisa, and Nhlangano. Participants lived in the towns and surrounding
geographic areas. These sites were selected based on region, proximity to the border, population size,
presence of local partner organizations, and the existence of key populations programming, health
services, and other sex work "hotspots".

A modified version of the Priorities for Local AIDS Control Efforts (PLACE) methodology was
used.[45] FSW were recruited through sampling at "hotspot" venues that were identified through key
informant interviews. Study staff verified and visited the venues and approached potential
participants. Venue managers and community leaders assisted in identifying and introducing FSW to
the study staff. This venue-based sampling was supplemented through peer referral and snowball
sampling to reach FSW who do not frequent public venues and increase the representativeness of the
sample.

Upon completion of the survey, each participant was given a 50 Emalangeni [approximately 5 United
States dollars] reimbursement for transportation and offered condoms and condom-compatible
lubricants. The Eswatini Ministry of Health and Scientific Ethics and the Johns Hopkins School of
Public Health Institutional Review Board approved the study.

2.3   Measures

2.3.1 Depressive symptoms

Depressive symptoms in the past two weeks were measured using the nine-item Patient Health
Questionnaire (PHQ-9).[46] The PHQ-9 in this sample had a Cronbach alpha of 0.91. PHQ-9 scores of
ten or higher were considered indicative of probable depression. Depression scores were categorized
by severity as minimal (0-4), mild (5-9), moderate (10-14), moderately severe (15-19), and severe
(20-27). Participants were asked how difficult these symptoms made it for them to do their work,
take care of things at home, or get along with other people (not difficult at all, somewhat difficult,
very difficult, or extremely difficult). Participants who reported any depressive symptoms were asked
if they ever sought help or treatment from a healthcare provider. Participants who never sought help
or treatment from a healthcare provider were asked if the reason they did not was related at all to the
fact that they sell sex.

2.3.2 Underage initiation of selling sex

Age of initiation of selling sex for money was dichotomized to compare those who started before the
age of 18 to those who started at age 18 or older.

2.3.3 Covariates

Continuous variables included the participant's age at the time of the study, number of years selling
sex, days per month selling sex, and weekly income in emalangeni and the number of times the
participant reported a condom slipped off or broke in the past month.

Dichotomous variables included whether the participant reported that her parents died before she was
18 years old, started selling sex to feed herself/her family, and was ever afraid of or avoided seeking
healthcare due to worrying that someone would find out she sells sex.

Categorical variables included the participant's frequency of carrying condoms while selling sex
(never, almost never, sometimes, almost always/often, or always) and the participant's self-reported
HIV status (never tested, HIV-negative, or living with HIV).

2.4   Analyses

Analyses were conducted using Stata/MP17.0. PHQ-9 scores by younger or older age of initiation of
selling sex were compared using a t-test. $\chi 2$ tests were used to compare by age of initiation of selling
sex the prevalence of depression, depression severity, each depressive symptom (i.e., any response
other than "not at all"), seeking treatment, not seeking treatment because of selling sex, and
difficulties due to depressive symptoms.

The relationships between each covariate and probable depression and each covariate and underage
initiation of selling sex were assessed using $\chi 2$ tests for categorical variables and t-tests for
continuous variables. Covariates that were significantly related to both probable depression and
underage initiation of selling sex in bivariate models ($p<0.05$) were included in a multivariable
logistic regression model of factors associated with probable depression. Missing data ($<5\%$) were
handled using listwise deletion.

3   Results

3.1   Descriptive statistics

As Table 1 shows, the mean age of participants at the time of the study was 27.5. 7.6% of
participants (57/747) lost both of their parents and 34.7% (259/747) lost at least one parent before the
age of 18. Most participants (81.4%, 627/770) started selling sex to feed themselves or their families.
Participants had been selling sex on an average of 6.2 years and sold sex on an average of 14.8 days
per month. Their average weekly income was 908.5 emalangeni, or about $62 USD. Most
participants (87.4%, 673/770) always carried condoms when selling sex. The mean number of times
participants reported that a condom slipped off or broke was once in the past month. About one
quarter of participants (25.3%, 195/770) had ever been afraid of or avoided seeking healthcare
because of worrying someone would find out they sell sex. 12.9% of participants (97/750) had never
been tested for HIV, 54.0% (405/750) had tested negative, and 33.1% (248/750) had tested positive.
16.6% of participants (128/770) started selling sex as minors. 43.1% of participants (332/770) had
probable depression.

3.2   Relationship between selling sex while underage and depression symptoms

Over half (55.5%, 71/128) of FSW who started selling sex as minors had probable depression. This
was significantly higher than the 40.7% (261/642) prevalence of depression among FSW who started
as adults (p=0.002). Over one in five (21.4%, 71/332) FSW who were depressed started selling sex as
minors, compared to 13.0% (57/438) of FSW who were not depressed.

As Table 2 shows, the mean PHQ-9 score was 9.18. The mean PHQ-9 score was significantly higher
among FSW who started selling sex as minors compared to FSW who started as adults (10.45 v.
8.93, p=0.020).

As Figure 1 shows, a higher proportion of FSW who started selling sex as minors had moderate,
moderately severe, or severe depression symptoms, while a higher proportion of FSW who started as
adults had minimal or mild symptoms (p=0.032).

Table 3 shows the proportion of FSW who reported each depression symptom (i.e., any response
other than "not at all"). The most common symptom was feeling down, sad, or hopeless (74.6%,
574/770). The least common symptom was suicidal ideation. This was still highly prevalent; 42.7%
(329/770) of FSW had thoughts that they would be better off dead or hurting themselves in some
way. A significantly higher proportion of FSW who sold sex as minors compared to FSW who
started as adults reported experiencing each depression symptom except suicidal ideation and feeling
tired or having little energy.

Less than half of those with depression symptoms (46.3%, 304/656) had sought help or treatment
from a healthcare provider. This did not differ significantly based on age of initiation of selling sex
(p=0.559). A small proportion (14.4%, 50/348) did not seek treatment due to the fact that they sell
sex. This did not differ significantly based on age of initiation of selling sex (p=0.269).

Of those who reported any depressive symptoms, 19.1% (125/656) said these problems made it not
difficult at all to do their work, take care of things at home, or get along with other people. 52.6%
(345/656) said it was somewhat difficult, 21.3% (140/656) said it was very difficult, and 7.0%
(46/656) said it was extremely difficult (Figure 2). This did not significantly vary based on age of
initiation of selling sex (p=0.295).

3.3   Covariates associated with probable depression

As Table 1 shows, a higher proportion of FSW who were depressed started selling sex to feed
themselves or their families compared to FSW who were not depressed (87.1% v. 77.2%, p<0.001).
FSW who were depressed had been selling sex for longer (6.6 years v. 5.9 years, p=0.048) and sold
sex for more days per month (15.6 v. 14.3, p=0.017) than FSW who were not depressed. The average
income of FSW who were depressed was higher than that of FSW who were not depressed (1079.8 v.
774.4 emalangeni, p<0.001). Compared to FSW who were not depressed, FSW who were depressed
reported that a condom slipped off or broke more often in the past month (1.3 v. 0.8 times, p<0.001).
38.0% of FSW who were depressed had ever been afraid of or avoided seeking healthcare due to
selling sex, while only 15.8% of FSW who were not depressed reported this (p<0.001). One fifth
(20.3%) of FSW who were depressed had never been tested for HIV, while less than one in thirteen
(7.3%) FSW who were not depressed had never been tested (p<0.001). Depression was not
significantly associated with current age, loss of parents, or frequency of carrying condoms.

3.4   Covariates associated with underage initiation of selling sex

As Table 1 shows, FSW who started selling sex as minors were younger at the time of the study than
FSW who started as adults (24.8 v. 28.0, p<0.001). A higher proportion of FSW who started selling
sex as minors had lost at least one (53.7% v. 3.9%, p<0.001) or both parents (17.1% v. 5.8%,
p<0.001) before the age of 18 than FSW who started as adults. Three quarters (75.0%) of FSW who
started selling sex as minors had done so to feed themselves or their families, while 82.7% of FSW
who started as adults reported this reason (p=0.041). FSW who started selling sex as minors had been
selling sex for longer (8.0 years v. 5.8 years, p<0.001) and sold sex for more days per month (16.2 v.
14.5, p=0.026) than FSW who started as adults. Compared to FSW who started selling sex as adults,
a higher proportion of FSW who started as minors never carried condoms when selling sex (7.0% v.
2.0%, p=0.023). The average number of times a condom slipped off or broke in the past month was

higher among FSW who started selling sex as minors than among those who started as adults (1.3 v.
0.9, p=0.020). Over one third (34.4%) of FSW who started selling sex as minors were ever afraid of
or avoided seeking healthcare due to selling sex compared to less than one quarter (23.5%) of FSW
who started as adults (p=0.010). Underage initiation of selling sex was not significantly associated
with income or self-reported HIV status.

217　3.5　Multivariable results

The categorical covariates in the multivariable model that were significantly related to both underage
initiation of selling sex and depression were whether the participant started selling sex to feed herself
or her family and whether she was afraid of or avoided seeking healthcare due to selling sex. The
continuous covariates in the multivariate model were the participants' number of years selling sex,
days per month selling sex, and times condoms slipped or broke in the past month (Table 4). In the
multivariable model, FSW who started selling sex as minors had higher odds of having probable
depression (adjusted odds ratio=1.70, 95% confidence interval=1.11-2.60). Starting sex work to feed
oneself or one's family, frequency of condoms breaking or slipping, and fear or avoidance of
healthcare remained significantly associated with depression. The number of years and the number of
days per month that a participant sold sex were not significantly associated with probable depression
in the multivariable model.

229　4　Discussion

Over 40% of FSW in Eswatini in this study reported depression. These findings are consistent with
other studies of FSW in sub-Saharan Africa that have shown a high burden of depression.[17,23,47-49] In
this study, 16.6% of FSW in Eswatini started selling sex as minors, which is comparable to the
prevalence in other countries in the region such as South Africa (18%),[20], Lesotho (20%),[5] Zimbabwe
(10%),[50] and Malawi (17%).[9] Depression and underage initiation of selling sex in this study were
both associated with selling sex to feed oneself or one's family, perceived sex work stigma in
healthcare settings, and condoms breaking or slipping. Depression was also associated with income
and never testing for HIV. Underage initiation of selling sex was also associated with younger age at
the time of the study, being orphaned, and carrying condoms less frequently.

239　4.1　Underage initiation of selling sex and depression

This study's findings strengthen the evidence base on the relationship between underage initiation of
selling sex and depression in Africa. In Malawi, FSW who started selling sex as minors had over
twice the odds of depression (aOR 2.4), but the results were not statistically significant. In our study,
which had a larger sample size and higher prevalence of depression, a slightly smaller (aOR 1.70) but
statistically significant difference was found. Our findings suggest the need to address the mental
health needs of people selling sex in Eswatini, particularly those who are underage or started while
underage. Given the limited availability of psychiatric services and high prevalence of HIV in
Eswatini, a nurse counsellor-delivered brief psychological intervention that was piloted with people
living with HIV, tuberculosis, and depression in Eswatini could be implemented with FSW and
adolescents.[51] Trauma-informed approaches to mental health treatment are needed for women who
started selling sex as minors,[52] while being mindful of the fact that they may still be selling sex or
may not consider themselves victims of trafficking or child sexual exploitation. Preventing underage
initiation of selling sex and providing alternatives to minors already selling sex to exit the sex trade
through programs such as cash transfers and social protection[53] could reduce a broad range of harms,
including depression.

4.2   Socioeconomic implications

Consistent with research in South Africa, depression among FSW in Eswatini was associated with selling sex due to food insecurity.[17] Scaling up existing programs that address food insecurity may have the additional benefit of decreasing depression by reducing reliance on sex work among those who do not want to sell sex.[54] In contrast, recent depression among FSW in this study was associated with higher current income, consistent with research findings among FSW in China.[15] Once FSW have enough income to meet their basic needs such as food, they may become more aware of their mental health needs.

Similar to research in Mexico, entry into selling sex due to food insecurity was less common among FSW in Eswatini who started selling sex as minors.[18] Over half of FSW who started selling sex as minors had lost at least one of their parents before the age of 18, and 17.1% lost both of their parents before age 18. Those who start selling sex as minors, particularly those who were orphaned, may do so for reasons not captured in our survey, such as to pay school fees or purchase drugs or alcohol.[55] Increasing services for orphans in Eswatini may prevent underage initiation of selling sex and subsequent negative health outcomes.

4.3   Healthcare service implications

Over one third of FSW who were depressed and FSW who started selling sex as minors had ever been afraid of or avoided seeking healthcare due to fear of someone learning about their involvement in selling sex. This is consistent with research in South Africa showing that depression was related to perceived or experienced sex work-related stigma in healthcare settings.[17] Anticipating stigma toward FSW may contribute toward feelings of depression. Conversely, FSW who are depressed may have heightened fear of stigma in health settings. This fear of stigma may be related to the finding in our study that less than half of FSW with depressive symptoms sought mental health treatment, and one in eight FSW who did not seek treatment stated the reason was related to selling sex. Qualitative research with minors who sell sex in other settings has also identified stigma in healthcare as a barrier to care.[22,56,57] Interventions to reduce stigma toward sex workers in healthcare settings that have been found to be effective in other countries could be adapted and implemented in Eswatini. This may be particularly beneficial to improve the uptake of physical and mental health services among FSW who are depressed and those who started selling sex as minors.[58,59] The HIV Prevention 2.0 intervention in Senegal included training healthcare workers to increase their clinical and social competence for addressing the needs of FSW. A web-based referral system provided FSW with information on where to access nonstigmatizing health services. The intervention was found to reduce perceived healthcare stigma among FSW.

Many prior studies have shown that people living with HIV have higher rates of depression.[60] In contrast, our study showed that depression was most prevalent among those who never tested for HIV and least prevalent among those who were living with HIV. Our findings are consistent with research conducted with FSW in Uganda and Zambia showing that knowledge of one's HIV status was associated with a decrease in depressive symptoms and research in South Africa showing that FSW who were depressed had lower odds of being tested for STIs.[17,23] Depression could reduce the likelihood of HIV being diagnosed and treated. Addressing mental health could be a strategy to improve the HIV care continuum among FSW. Alternatively, uncertainty about their HIV status may contribute to symptoms of depression among FSW, and increasing uptake of testing may improve mental health. Integrating mental health services with HIV prevention, testing, and treatment services is warranted.

4.4 Behavioral program implications

FSW who were depressed and FSW who started selling sex as minors both reported that a condom had slipped or broken 1.3 times on average in the past month, potentially increasing their risk for pregnancy and HIV and STI acquisition and transmission. Access to services such as HIV and STI testing and treatment, pre- and post-exposure prophylaxis for HIV prevention, and contraception for those who do not want to become pregnant may be particularly important for these populations. This study did not include data on potential mediators of the relationship between condom breakage, depression, and underage initiation of selling sex such as substance use and condom application skills,[25] but additional education for FSW on these topics may be necessary. Treating depressive symptoms among FSW could reduce breakage and slippage and enhance the protective benefits of condoms. However, an alternative explanation for our findings may be that FSW who are depressed or begin selling sex while underage may be more likely to notice when a condom slips or breaks.

17.2% of FSW in this study who started selling sex as minors did not always carry condoms when selling sex. This is consistent with research from Lesotho showing that underage initiation of selling sex was associated with avoiding carrying condoms due to fear of trouble with police (e.g., condoms being used as evidence to arrest them for selling sex).[5] Further research is needed into specific barriers to carrying condoms among FSW in Eswatini. Providing condoms at sex work venues may particularly help FSW who started selling sex as minors.

4.5 Limitations

Data in this study were self-reported, which may lead to social desirability bias or inaccurate recall. The results from this sample of FSW recruited through venue-based sampling may not be generalizable to all FSW in Eswatini. Due to the cross-sectional nature of this study, it is not clear whether depression is the result of selling sex while underage or a result of other factors that led to underage initiation of selling sex. Longitudinal studies if ethically feasible could elucidate the timing of development of depressive symptoms among those who sell sex as minors.

4.6 Conclusion

This study demonstrates that depression is highly prevalent among FSW in Eswatini and associated with underage initiation of selling sex and other social and behavioral covariates. Taken together, these results indicate the need to comprehensively address intersecting risks and health conditions including early initiation of selling sex, mental health, stigma, food insecurity, and HIV risk among FSW in Eswatini.

342 Table 1. Characteristics of female sex workers in Eswatini by probable depression and age of entry into selling
343 sex, 2014

	Total	Not depressed	Depressed	P value	Started selling sex 18+	Started selling sex <18	P value
Depressed	43.1% (332/770)	-	-	-	40.7% (261/642)	55.5% (71/128)	0.002
Started selling sex <18	16.6% (128/770)	13.0% (57/438)	21.4% (71/332)	0.002	-	-	-
Mean current age	27.5	27.5	27.5	0.868	28.0	24.8	<0.001
Orphaned <18							
2 parents died	7.6% (57/747)	8.0% (34/427)	7.2% (23/320)	0.693	5.8% (36/624)	17.1% (21/123)	<0.001
1+ parents died	34.7% (259/747)	36.5% (156/427)	32.2% (103/320)	0.217	30.9% (193/624)	53.7% (66/123)	<0.001
Started selling sex to feed self/family	81.4% (627/770)	77.2% (338/438)	87.1% (289/332)	<0.001	82.7% (531/642)	75.0% (96/128)	0.041
Mean years selling sex	6.2 (n=757)	5.9 (n=432)	6.6 (n=325)	0.048	5.8 (n=630)	8.0 (n=127)	<0.001
Mean days per month selling sex	14.8 (n=767)	14.3 (n=436)	15.6 (n=331)	0.017	14.5 (n=639)	16.2 (n=128)	0.026
Mean weekly income in emalangeni	908.5 (n=747)	774.4 (n=419)	1079.8 (n=328)	<0.001	885.6 (n=622)	1022.0 (n=125)	0.110
Frequency of carrying condoms when selling sex							
Never	2.9% (22/770)	1.8% (8/438)	4.2% (14/332)	0.333	2.0% (13/642)	7.0% (9/128)	0.023
Almost never	0.3% (2/770)	0.2% (1/438)	0.3% (1/332)		0.2% (1/642)	0.8% (1/128)	
Sometimes	4.0% (31/770)	4.6% (20/438)	3.3% (11/332)		4.1% (26/642)	3.9% (5/128)	
Almost always/often	5.5% (42/770)	5.5% (24/438)	5.4% (18/332)		5.5% (35/642)	5.5% (7/128)	
Always	87.4% (673/770)	87.9% (385/438)	86.8% (288/332)		88.3% (567/642)	82.8% (106/128)	
Mean times condom slipped off or broke in the past month	1.0 (n=753(	0.8 (n=427)	1.3 (n=326)	<0.001	0.9 (n=628)	1.3 (n=125)	0.020
Ever afraid of/avoided seeking healthcare because of selling sex	25.3% (195/770)	15.8% (69/438)	38.0% (126/332)	<0.001	23.5% (151/642)	34.4% (44/128)	0.010
Self-reported HIV status							
Never tested	12.9% (97/750)	7.3% (31/425)	20.3% (66/325)	<0.001	12.6% (79/628)	14.8% (18/122)	0.695
Tested negative	54.0% (405/750)	55.8% (237/425)	51.7% (168/325)		53.8% (338/628)	54.9% (67/122)	
Tested positive	33.1% (248/750)	36.9% (157/425)	28.0% (91/325)		33.6% (211/628)	30.3% (37/122)	

344
345

346   Table 2. Average PHQ-9 score by age of entry into selling sex among female sex workers in
347   Eswatini, 2014

	PHQ-9 score	
	Median	Mean
Started selling sex <18	11	10.45
Started selling sex 18+	8.5	8.93
Total	9	9.18
N=770; p=0.020; PHQ-9=Patient Health Questionnaire		

369 Table 3. Differences in PHQ-9 items and seeking treatment by age of entry into selling sex among
370 female sex workers in Eswatini, 2014

	Started selling sex 18+	Started selling sex <18	Total	P value
Over the last 2 weeks, have you been bothered by any of the following problems?				
Little interest or pleasure in doing things	71.2% (457/642)	80.5% (103/128)	72.7% (560/770)	0.031
Feeling down, depressed (sad), or hopeless	72.9% (468/642)	82.8% (106/128)	74.6% (574/770)	0.019
Either trouble falling or staying asleep, or sleeping too much	67.6% (434/642)	77.3% (99/128)	69.2% (533/770)	0.029
Feeling tired or having little energy	69.5% (446/642)	77.3% (99/128)	70.8% (545/770)	0.074
Either poor appetite or over-eating	68.2% (438/642)	77.3% (99/128)	69.7% (537/770)	0.040
Feeling bad about yourself or that you are a failure or have let yourself or your family down	60.0% (385/642)	70.3% (90/128)	61.7% (475/770)	0.028
Trouble concentrating on things, such as reading the newspaper or watching TV	54.8% (352/642)	67.2% (86/128)	56.9% (438/770)	0.010
Either moving or speaking so slowly that other people could have noticed OR being so fidgety or restless that you have been moving around a lot more than usual	47.7% (306/642)	59.4% (76/128)	49.6% (382/770)	0.016
Thoughts that you would be better off dead, or hurting yourself in some way	41.9% (269/642)	46.9% (60/128)	42.7% (329/770)	0.299
Sought help or treatment from a healthcare provider	46.9% (254/542)	43.9% (50/114)	46.3% (304/656)	0.559
Did not seek health services because of selling sex	13.4% (38/284)	18.8% (12/64)	14.4% (50/348)	0.269
PHQ-9=patient health questionnaire				

380 Table 4. Multivariable logistic regression analysis of factors associated with probable depression
381 among female sex worker study participants in Eswatini, 2014

	Adjusted odds ratio (95% confidence interval)

Started selling sex <18	1.70 (1.11, 2.60)
Started selling sex to feed self/family	2.33 (1.51, 3.59)
Number of years selling sex	1.00 (0.97, 1.04)
Days per month selling sex	1.00 (0.98, 1.03)
Number of times a condom slipped off or broke in the last month	1.21 (1.09, 1.35)
Ever afraid of or avoided seeking healthcare due to fear of someone learning she sells sex	3.17 (2.21, 4.56)
N=737	

416 Figure 1 Depression severity by younger or older initiation of selling sex among female sex worker
417 study participants in Eswatini, 2014

447   Figure 2 If you are having any of the problems above, how difficult have these problems made it for
448   you to do your work, take care of things at home, or get along with other people?

450   %

References

1.   Millan-Alanis JM, Carranza-Navarro F, de León-Gutiérrez H, et al. Prevalence of suicidality,
depression, post-traumatic stress disorder, and anxiety among female sex workers: a systematic
review and meta-analysis. *Archives of Women's Mental Health* 2021; **24**(6): 867-79.

2.   Ma PHX, Chan ZCY, Loke AY. The socio-ecological model approach to understanding
barriers and facilitators to the accessing of health services by sex workers: a systematic review. *AIDS*
and Behavior 2017; **21**(8): 2412-38.

3.   Beattie TS, Smilenova B, Krishnaratne S, Mazzuca A. Mental health problems among female
sex workers in low- and middle-income countries: a systematic review and meta-analysis. *PLOS*
Medicine 2020; **17**(9): e1003297.

4.   Laird JJ, Klettke B, Hall K, Clancy E, Hallford D. Demographic and psychosocial factors
associated with child sexual exploitation: a systematic review and meta-analysis. *JAMA Network*
Open 2020; **3**(9): e2017682-e.

5.   Grosso A, Busch S, Mothopeng T, et al. HIV risks and needs related to the Sustainable
Development Goals among female sex workers who were commercially sexually exploited as
children in Lesotho. *Journal of the International AIDS Society* 2018; **21**: e25042.

6.   Grosso A, Ketende S, Dam K, et al. Structural determinants of health among women who
started selling sex as minors in Burkina Faso. *Journal of Acquired Immune Deficiency Syndromes*
2015; **68**: S162-S70.

7.   Parcesepe AM, L'Engle KL, Martin SL, Green S, Suchindran C, Mwarogo P. Early sex work
initiation and violence against female sex workers in Mombasa, Kenya. *Journal of Urban Health*
2016; **93**(6): 1010-26.

8.   Parcesepe AM, L'Engle KL, Martin SL, Green S, Suchindran C, Mwarogo P. Early sex work
initiation and condom use among alcohol-using female sex workers in Mombasa, Kenya: a cross-
sectional analysis. *Sexually Transmitted Infections* 2016; **92**(8): 593-8.

9.   MacLean SA, Lancaster KE, Lungu T, et al. Prevalence and correlates of probable depression
and post-traumatic stress disorder among female sex workers in Lilongwe, Malawi. *International*
Journal of Mental Health and Addiction 2018; **16**(1): 150-63.

10.   Erulkar A, Medhin G. Factors associated with depression among young female migrants and
commercial sex workers in Ethiopia. 2021.

11.   Erulkar A, Medhin G, Negeri L. The journey of out-of-school girls in Ethiopia: examining
migration, livelihoods, and HIV. 2017.

12.   Coetzee J, Buckley J, Otwombe K, Milovanovic M, Gray GE, Jewkes R. Depression and Post
Traumatic Stress amongst female sex workers in Soweto, South Africa: a cross sectional, respondent
driven sample. *PLOS ONE* 2018; **13**(7): e0196759.

13.   Clarke RJ, Clarke EA, Roe-Sepowitz D, Fey R. Age at entry into prostitution: relationship to
drug use, race, suicide, education level, childhood zbuse, and family experiences. *Journal of Human*
Behavior in the Social Environment 2012; **22**(3): 270-89.

14.   Hengartner MP, Islam MN, Haker H, Rössler W. Mental health and functioning of female sex
workers in Chittagong, Bangladesh. *Frontiers in Psychiatry* 2015; **6**: 176.

15.   Shen H, Zou H, Huang S, et al. Depression and HIV risk behaviors among female sex
workers in Guangdong, China: a multicenter cross-sectional study. *BioMed Research International*
2016; **2016**: 6986173.

16.   Goldenberg SM, Rangel G, Vera A, et al. Exploring the impact of underage sex work among
female sex workers in two Mexico–US border cities. *AIDS and Behavior* 2012; **16**(4): 969-81.

17.   Rossouw J, Schwartz S, Rao A, et al. Exploring the association between depression and social
and biobehavioral HIV risk factors among female sex workers in Nelson Mandela Bay Municipality,
South Africa. *AIDS Research and Human Retroviruses* 2021.

18.   Loza O, Strathdee SA, Lozada R, et al. Correlates of early versus later initiation into sex work
in two Mexico–US border cities. *Journal of Adolescent Health* 2010; **46**(1): 37-44.

19.   Footer KH, White RH, Park JN, Decker MR, Lutnick A, Sherman SG. Entry to sex trade and
long-term vulnerabilities of female sex workers who enter the sex trade before the age of eighteen.
Journal of Urban Health 2020; **97**(3): 406-17.

20.   Grosso A, Pitche V, Kochelani D, et al. Experiences of loss of a parent before age 18 among
female sex workers across seven western and southern African countries. International AIDS
Conference. Amsterdam, Netherlands; 2018.

21.   Sahad SM, Mohamad Z, Shukri M. Differences of mental health among orphan and non-
orphan adolescents. *International Journal of Academic Research in Psychology* 2018; **5**(1): 556-65.

22.   Ijadi-Maghsoodi R, Bath E, Cook M, Textor L, Barnert E. Commercially sexually exploited
youths' health care experiences, barriers, and recommendations: a qualitative analysis. *Child Abuse*
and Neglect 2018; **76**: 334-41.

23.   Ortblad KF, Musoke DK, Chanda MM, et al. Knowledge of HIV status is associated with a
decrease in the severity of depressive symptoms among female sex workers in Uganda and Zambia.
Journal of Acquired Immune Deficiency Syndromes 2020; **83**(1): 37.

24.   Andrew BJ. A test of the theory of planned behaviour and an implementation intentions
intervention for condom use behaviours among men who have sex with men: University of Sydney;
2015.

25.   Bradley J, Rajaram SP, Moses S, et al. Why do condoms break? A study of female sex
workers in Bangalore, south India. *Sexually Transmitted Infections* 2012; **88**(3): 163-70.

26.   Lang DL, Salazar LF, DiClemente RJ, Markosyan K, Darbinyan N. Predictors of condom
errors among sex workers in Armenia. *International Journal of STD & AIDS* 2011; **22**(3): 126-30.

27.   Wondie Y, Yigzaw T, Koester L. Post-traumatic stress disorder, depression and substance
abuse among female street-based sex workers in Addis Ababa. *Ethiopian Renaissance Journal of*
Social Sciences and the Humanities 2019; **6**(1).

28.   Jewkes R, Milovanovic M, Otwombe K, et al. Intersections of sex work, mental ill-health,
IPV and other violence experienced by female sex workers: findings from a cross-sectional
community-centric national study in South Africa. *International Journal of Environmental Research*
and Public Health 2021; **18**(22): 11971.

29.   Martin L, Hearst MO, Widome R. Meaningful differences: comparison of adult women who
first traded sex as a juvenile versus as an adult. *Violence Against Women* 2010; **16**(11): 1252-69.

30.   Dlamini SB, Shongwe MC. Exploring mental health nurses' perceptions on factors
contributing to psychiatric readmissions in Eswatini: A qualitative study. *International Journal of*
Africa Nursing Sciences 2019; **11**: 100157.

31.   Dlamini LP, Mahanya S, Dlamini SD, Shongwe MC. Prevalence and factors associated with
postpartum depression at a primary healthcare facility in Eswatini. *South African Journal of*
Psychiatry 2019; **25**(1): 1-7.

32.   World Health Organization. Depression and other common mental disorders: global health
estimates: World Health Organization, 2017.

33.   Lofgren S, Bond D, Nakasujja N, Boulware D. Burden of depression in outpatient HIV-
infected adults in sub-Saharan Africa; Systematic review and meta-analysis. *AIDS and Behavior*
2019: 1-13.

34.   Pengpid S, Peltzer K. The prevalence and correlates of suicidal ideation, plans and suicide
attempts among 15-to 69-year-old persons in Eswatini. *Behavioral Sciences* 2020; **10**(11): 172.

35.   Motsa MPS, Chiou H-Y, Chen Y-H. Association of chronic diseases and lifestyle factors with
suicidal ideation among adults aged 18–69 years in Eswatini: evidence from a population-based
survey. *BMC Public Health* 2021; **21**(1): 2245.

36.   Almansour AM, Siziya S. Suicidal ideation and associated factors among school going
adolescents in Swaziland. *African Health Sciences* 2017; **17**(4): 1172-7.

37.   Quarshie EN-B, Atorkey P, García KPV, Lomotey SA, Navelle PL. Suicidal behaviors in a
nationally representative sample of school-going adolescents aged 12–17 years in Eswatini. *Trends in*
Psychology 2021: 1-30.

38.   Dlamini Z, Hlope TG, Mohammed M. Exploring the relationship between unemployment and
mental illness among the youth in selected communities in Eswatini. 2019.

39.   Criminal Law and Procedure ACT 6/1889.

40.   Berger BO, Grosso A, Adams D, et al. The prevalence and correlates of physical and sexual
violence affecting female sex workers in Swaziland. *Journal of Interpersonal Violence* 2018; **33**(17):
2745-66.

41.   Baral S, Ketende S, Green JL, et al. Reconceptualizing the HIV epidemiology and prevention
needs of female sex workers (FSW) in Swaziland. *PLOS ONE* 2014; **9**(12).

42.   Grosso A, Ketende S, Dam K, et al. Prevention and treatment needs of women who started
selling sex as minors. International AIDS Conference. Melbourne, Australia; 2014.

43.   Rao A, Stahlman S, Hargreaves J, et al. Sampling key populations for HIV surveillance:
results from eight cross-sectional studies using respondent-driven sampling and venue-based
snowball sampling. *JMIR Public Health and Surveillance* 2017; **3**(4): e8116.

44.   Health Communication Capacity Collaborative. Characterizing the HIV prevention and
treatment needs among key populations, including men who have sex with men and female sex
workers in Swaziland: from evidence to action., 2015.

45.   Weir SS, Pailman C, Mahlalela X, Coetzee N, Meidany F, Boerma JT. From people to places:
focusing AIDS prevention efforts where it matters most. *AIDS* 2003; **17**(6): 895-903.

46.   Kroenke K, Spitzer RL, Williams JB. The PHQ□9: validity of a brief depression severity
measure. *Journal of General Internal Medicine* 2001; **16**(9): 606-13.

47.   Bhardwaj A, Comins C, Guddera V, et al. Prevalence of depression and key associated factors
among female sex workers living with HIV in Durban, South Africa. *Journal of the International*
AIDS Society 2020; **23**: 141-2.

48.   Stockton MA, Pence BW, Mbote D, et al. Associations among experienced and internalized
stigma, social support, and depression among male and female sex workers in Kenya. *International*
Journal of Public Health 2020; **65**(6): 791-9.

49.   Leis M, McDermott M, Koziarz A, et al. Intimate partner and client perpetrated violence are
associated with reduced HIV pre exposure prophylaxis (PrEP) uptake, depression and generalized
anxiety in a cross sectional study of female sex workers from Nairobi, Kenya. *Journal of the*
International AIDS Society 2021; **24**: e25711.

50.   Cowan FM, Davey C, Fearon E, et al. Targeted combination prevention to support female sex
workers in Zimbabwe accessing and adhering to antiretrovirals for treatment and prevention of HIV
(SAPPH-IRe): a cluster-randomised trial. *The Lancet HIV* 2018; **5**(8): e417-e26.

51.   Putnis N, Riches N, Nyamayaro A, et al. Depression and HIV/TB In southern Africa: a mixed
method evaluation of an integrated primary care mental health programme In Eswatini. 2021.

52.   Laser-Maira JA, Peach D, Hounmenou CE. Moving towards self-actualization: a trauma-
informed and needs-focused approach to the mental health needs of survivors of commercial child
sexual exploitation. *International Journal of Social Work* 2019; **6**(2): 27.

53.   Cluver LD, Orkin FM, Yakubovich AR, Sherr L. Combination social protection for reducing
HIV-risk behavior among adolescents in South Africa. *Journal of Acquired Immune Deficiency*
Syndromes 2016; **72**(1): 96-104.

54.   Fielding-Miller R, Mnisi Z, Adams D, Baral S, Kennedy C. "There is hunger in my
community": a qualitative study of food security as a cyclical force in sex work in Swaziland. *BMC*
Public Health 2014; **14**(1): 79.

55.   Pufall EL, Eaton JW, Robertson L, Mushati P, Nyamukapa C, Gregson S. Education,
substance use, and HIV risk among orphaned adolescents in Eastern Zimbabwe. *Vulnerable Children*
and Youth Studies 2017; **12**(4): 360-74.

56.   Prior A, Shilo G, Peled E. Help-seeking and help-related experiences of commercially
sexually exploited youth: a qualitative meta-synthesis. *Trauma, Violence, and Abuse* 2022:
15248380221074333.

57.   Thorburn N. Surviving shame. *New Zealand Journal of Counselling* 2015; **35**(2): 14-26.

58.   Lyons CE, Olawore O, Turpin G, et al. Intersectional stigmas and HIV-related outcomes
among a cohort of key populations enrolled in stigma mitigation interventions in Senegal. *AIDS*
2020; **34 Suppl 1**(Suppl 1): S63-s71.

59.   Lyons CE, Ketende S, Diouf D, et al. Potential impact of integrated stigma mitigation
interventions in improving HIV/AIDS service delivery and uptake for key populations in Senegal.
Journal of Acquired Immune Deficiency Syndromes 2017; **74**(Suppl 1): S52.

60.   Ciesla JA, Roberts JE. Meta-analysis of the relationship between HIV infection and risk for
depressive disorders. *American Journal of Psychiatry* 2001; **158**(5): 725-30.