<u>Phenome-wide Mendelian randomisation analysis identifies causal factors for age-related macular</u>

<u>degeneration.</u>

Thomas H Julian[1,2], Johnathan Cooper-Knock[3], Stuart MacGregor[4], Hui Guo[5], Tariq Aslam[2,6], Eleanor

Sanderson[7], Graeme C Black[1,8], Panagiotis I Sergouniotis[1,2,8,9]

[1] Division of Evolution, Infection and Genomics, School of Biological Sciences, Faculty of Biology,

Medicine and Health, University of Manchester, Manchester, UK.

[2] Manchester Royal Eye Hospital, Manchester University NHS Foundation Trust, Manchester, UK.

[3] Department of Neuroscience, Sheffield Institute for Translational Neuroscience (SITraN), University of

Sheffield, Sheffield, UK.

[4] Statistical Genetics, QIMR Berghofer Medical Research Institute, Brisbane, Australia.

[5] Centre for Biostatistics, Division of Population Health, Health Services Research and Primary Care,

School of Health Sciences, Faculty of Biology, Medicine and Health, University of Manchester,

Manchester, UK.

[6] Division of Pharmacy and Optometry, Faculty of Biology, Medicine and Health, School of Health

Sciences, University of Manchester, Manchester, UK.

[7] MRC Integrative Epidemiology Unit, University of Bristol, Bristol, United Kingdom.

[8] Manchester Centre for Genomic Medicine, Saint Mary's Hospital, Manchester University NHS

Foundation Trust, Manchester, UK.

[9] European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI), Wellcome

Genome Campus, Cambridge, UK.

Correspondence to: Panagiotis Sergouniotis (panagiotis.sergouniotis@manchester.ac.uk) and Graeme

Black (graeme.black@manchester.ac.uk)

<u>**IMPACT STATEMENT:**</u>

A phenome-wide Mendelian randomisation analysis revealed a causal link between age-related macular

degeneration and a number of lipid, complement, immune cell, and serum protein traits, highlighting

potential treatment targets.

<u>**ABSTRACT:**</u>

<u>**Background:**</u> Age-related macular degeneration (AMD) is a leading cause of blindness in the

industrialised world and is projected to affect >280 million people worldwide by 2040. Aiming to identify

causal factors and potential therapeutic targets for this common condition, we designed and undertook a

phenome-wide Mendelian randomisation (MR) study.

<u>**Methods:**</u> We evaluated the effect of 4,591 exposure traits on early AMD using univariable MR.

Statistically significant results were explored further using: validation in an advanced AMD cohort; MR

Bayesian model averaging (MR-BMA); and multivariable MR.

<u>**Results:**</u> Overall, 44 traits were found to be putatively causal for early AMD in univariable analysis.

Serum proteins that were found to have significant relationships with AMD included S100-A5 (odds ratio

[OR]=1.07, p-value=6.80E-06), cathepsin F (OR=1.10, p-value=7.16E-05) and serine palmitoyltransferase

2 (OR=0.86, p-value=1.00E-03). Univariable MR analysis also supported roles for complement and

immune cell traits. Although numerous lipid traits were found to be significantly related to AMD, MR-BMA

suggested a driving causal role for serum sphingomyelin (marginal inclusion probability [MIP]=0.76;

model-averaged causal effect [MACE]=0.29).

<u>**Conclusions:**</u> The results of this MR study support several putative causal factors for AMD and highlight

avenues for future translational research.

<u>**Funding:**</u> This project was funded by the Wellcome Trust (224643/Z/21/Z); the University of Manchester's

Wellcome Institutional Strategic Support Fund (Wellcome ISSF) grant (204796/Z/16/Z); the UK National

Institute for Health Research (NIHR) Academic Clinical Fellow and Clinical Lecturer Programmes; Retina

INTRODUCTION

Age-related macular degeneration (AMD) is a common retinal condition that affects individuals who are ≥50 years old. It is caused by the complex interplay of multiple genetic and environmental risk factors and genome-wide association studies (GWAS) have identified AMD-implicated variants in at least 69 loci. These include important risk alleles in the 1q32 and 10q26 genomic regions (corresponding to the *CFH* [complement factor H] and *ARMS2/HTRA1* locus respectively) (Fritsche et al., 2016; Winkler et al., 2020). Other key risk factors include age, smoking, alcohol consumption and low dietary intake of antioxidants (carotenoids, zinc) (Chakravarthy et al., 2010).

AMD can be categorised according to severity (early, intermediate, or advanced) or based on the presence of neovascularization (neovascular or non-neovascular). Advanced AMD results in loss of central vision, often leading to severe visual impairment (Fleckenstein et al., 2021). Notably, AMD is a major cause of blindness in the elderly population and represents a substantial global burden that is expected to continue to grow into the future as an ageing population expands worldwide (Chakravarthy et al., 2010).

Mendelian randomization (MR) is a statistical approach that uses genetic variation to look for causal relationships between *exposures* (such as smoking) and *outcomes* (such as risk of a specific disease) (Julian et al., 2021). MR is increasingly being utilised as it can, to a degree, address a major limitation of observational studies: unmeasured confounding (Sanderson et al., 2022). To minimise issues with certain types of confounding and to support causal inference statements, MR uses genetic variation as an *instrument* (*i.e.* as a variable that is associated with the exposure but is independent of confounders and is not associated with the outcome, other than through the exposure). The principles of MR are based on Mendel's laws of segregation and independent assortment, which state that offspring inherit alleles randomly from their parents and randomly with respect to other locations in the genome. A key concept is the use of genetic variants that are related to an exposure of interest to proxy the part of the exposure that is independent of possible confounding influences (e.g. from the environment or from other traits). It is noted that analogies have been drawn between MR and randomised controlled trials with these two approaches considered proximal in terms of hierarchy of evidence (Julian et al., 2021). To date, the use

of MR approaches in the context of AMD has been limited although these methods have been

successfully implemented to explore the relationship between AMD and a small number of traits including

lipids, thyroid function, CRP and complement factors (Cipriani et al., 2021; Han et al., 2021, 2020b; Li et

al., 2022; Zuber et al., 2020).

In this study, we developed a systematic, broad ('phenome-wide') MR-based analytical approach and

used it to investigate the relationship between early AMD and several thousand exposure variables. A set

of traits that are robustly associated with genetic liability to AMD were identified.

METHODS

Data sources

Outcome data

Two AMD phenotypes were used as outcome measures in this study. The first one was early AMD. The GWAS summary statistics for this phenotype were taken from a meta-analysis by Winkler and colleagues (Winkler et al., 2020). This meta-analysis focused on populations of European ancestries and used data from the ARIC, AugUR, CHS, GHS, IAMDGC, KORA S4, LIFE-Adult NICOLA, UKBB and WHI studies (14,034 early AMD cases and 91,214 controls overall). A full description of how these studies classified participants as "early AMD" can be found in the relevant publication (Winkler et al., 2020); briefly, a number of approaches considering drusen size/area and the presence or absence of pigmentary abnormalities were utilised including the Three Continent Consortium (3CC) severity scale (Klein et al., 2014), the Rotterdam Eye Study classification (Korb et al. 2014), the Beckman clinical classification (Ferris et al. 2013), and the AREDS-9 step classification scheme (Davis et al. 2005). All relevant studies used colour fundus photography for grading purposes. The second phenotype that we studied was advanced AMD. For this trait, we drew on GWAS summary statistics from a multiple trait analysis of GWAS (MTAG) study by Han and colleagues (Han et al., 2020a). This meta-analysis also focused on individuals with European ancestries and derived data from the IAMDGC 2013 (17,181 cases and 60,074 controls) (Fritsche et al., 2013) and IAMDGC 2016 (16,144 advanced AMD cases and 17,832 controls) (Fritsche et al., 2016) studies as well as the GERA study (4,017 cases and 14,984 controls) (Kvale et al., 2015). The relevant summary statistics are primarily reflective of advanced AMD, but the GERA cohort included both advanced and intermediate AMD cases. Advanced AMD was broadly defined by the presence of geographic atrophy or choroidal neovascularisation, although there was a degree of variability in the criteria used in the included studies. Notably, the MTAG approach can leverage the high genetic correlation between the input phenotypes to detect genetic associations relevant only to advanced AMD.

Exposure data

A phenome-wide screen was performed to make causal inferences on the role of a wide range of traits in

early and advanced AMD. To achieve this, both published and unpublished GWAS data from the IEU

open GWAS database were used; these were accessible via the TwoSampleMR programme in R

(Hemani et al., 2018). All European GWAS within this database were included with the exception of

imaging phenotypes and expression quantitative trait locus (eQTL) related data which were removed. The

restriction to European datasets limits the generalisability of the results to other populations but is

necessary to produce reliable findings. In the early AMD analysis, studies from the "ukb" and "met-d"

batches were excluded as data for these studies were entirely from the UK Biobank resource and, as a

result, there was extensive population overlap with the early AMD GWAS (Sudlow et al., 2015). In the

advanced AMD analysis, the "ukb" and "met-d" batches were included. The early AMD analysis was

conducted on 30/12/2021 and a total of 10,979 traits were considered for analysis. The advanced AMD

analysis was conducted on 08/01/2022. On 26/01/2022 we added the newly published "finn-b" (n=2803)

traits to the analysis in place of the outdated "finn-a" traits (n=1489). It was impractical to manually inspect

the degree of population overlap for all traits prior to conducting the analysis; instead, the degree of

overlap for all significant traits was inspected after the analysis.

Instrument selection

A statistically driven approach to instrumental variable selection was used. Typically, an arbitrary p-value

threshold is set for the identification of appropriate single-nucleotide variants (SNVs); these are

subsequently used as instrumental variables (referred to thereafter as *instruments*). A conventional p-

value threshold for the selection of instruments is >5E-08. This approach however can, in some cases, be

problematic. For example, when the number of instruments exceeding this threshold is small, the analysis

can be underpowered or, in certain cases of unbiased screens, the results can be inflated (Boddy et al.,

2022). With this in mind, the p-value for instrument selection for each trait was set to the level where >5

instruments were available for each analysis. More specifically, for each trait, the analysis would first be

conducted with a p-value threshold for inclusion of 5E-8 before sequentially increasing the threshold by a

factor of 10 each time until >5 eligible instruments are identified. A predefined maximum p-value of 5E-05

was used and the final range of p-values for inclusion was 5E-06 to 5E-08.

Proxies

Where an exposure instrument was not present in the outcome dataset, a suitable proxy was identified

(Hartwig et al., 2016). In the early AMD analysis, this was achieved by using the TwoSampleMR software

with a linkage disequilibrium R^2 value of ≥0.9 (Purcell et al., 2007). For the advanced AMD phenotype,

data that was not derived from the TwoSampleMR resource were used and therefore the Ensembl server

was utilised to identify proxies (Cunningham et al., 2022; Hemani et al., 2018).

Clumping

SNVs were clumped using a linkage disequilibrium R^2 value of 0.001 and a genetic distance cut-off of

10,000 kilo-bases. A European reference panel was used for clumping.

Harmonisation

The effects of instruments on outcomes and exposures were harmonised to ensure that the beta values

(*i.e.* the regression analysis estimates of effect size) were expressed per additional copy of the same

allele (Hartwig et al., 2016). Palindromic alleles (*i.e.* alleles that are the same on the forward as on the

reverse strand) with a minor allele frequency (MAF) >0.42 were omitted from the analysis in order to

reduce the risk of errors.

Removal of pleiotropic genetic variants and outliers

Pleiotropic instruments and outliers were removed from the analysis by using a statistical approach that

removes instruments which are found to be more significant for the outcome than for the exposure

(Hemani et al., 2017). Radial MR, a simulation-based approach that detects outlying instruments, was

also utilised (Bowden et al., 2018).

Causal inference

MR relies on three key assumptions with regards to the instrumental variable: (1) the instrumental SNV

should be associated with the exposure; (2) the SNV should not be associated with confounders; (3) the

SNV should influence the outcome only through the exposure (Julian et al., 2021).

MR estimation was primarily performed using a multiplicative random effects (MRE) inverse variance

weighted (IVW) method. MRE IVW was selected over a fixed effects (FE) approach as it allows inclusion

of heterogeneous instruments (this was certain to occur within the breadth of this screen) (Burgess et al.,

2019). A range of "robust measures" were used to increase the accuracy of the results and to account for

violations of the above key MR assumptions (Burgess et al., 2019); these measures included weighted

median (Bowden et al., 2016a), Egger (Burgess and Thompson, 2017a), weighted mode (Hartwig et al.,

2017) and radial MR with modified second order weights (Bowden et al., 2018).

Further quality control

The instrument strength was determined using the F-statistic (which tests the association between the

instruments and the exposure) (Burgess et al., 2011). F-statistics were calculated against the final set of

instruments that were included. A mean F-statistic >10 was considered sufficiently strong.

The Cochran's Q test was performed for each analysis. Cochran's Q is a measure of heterogeneity

among causal estimates and serves as an indicator of the presence of horizontal pleiotropy (which occurs

when an instrument exhibits effects on the outcome through pathways other than the exposure) (Bowden

and Holmes, 2019). It is noted that a heterogeneous instrument is not necessarily invalid, but rather calls

for a primary assessment with an MRE IVW rather than an FE approach; this has been conducted as

standard throughout our analysis.

The MR-Egger intercept test was used to detect horizontal pleiotropy. When this occurs, the Egger

regression is robust to horizontal pleiotropy (under the assumption that that pleiotropy is uncorrelated with

the association between the SNV and the exposure) (Burgess and Thompson, 2017b). Unless otherwise

indicated by the Egger intercept, the assumption that no demonstrable horizontal pleiotropy is present

was made and Egger regression was not utilised to determine causal effects (given the low power of this

approach in the context of a small number of SNVs) (Bowden et al., 2015).

The I^2 statistic was calculated as a measure of heterogeneity between variant-specific causal estimates.

An I^2 <0.9 indicates that Egger is more likely to be biased towards the null through violation of the `NO

Measurement Error' (NOME) assumption (Bowden et al., 2016b).

Leave-one-out (LOO) cross-validation was performed for every analysis to determine if any particular

SNV was driving the significance of the causal estimates.

Management of duplicate traits/GWAS

As the GWAS database that was used contained multiple different GWAS for certain traits, some

exposures were analysed on multiple occasions. Where this occurred, the largest sample size study was

considered to be the primary analysis. Where there were duplicate studies in the same population, the

study with the largest F-statistic was used.

Identification of significant results

Before considering an MR result to be significant, the results of a range of causal inference and quality

control tests should be taken into account. Notably, it is not necessary for a study to find significance in all

measures to determine a true causal relationship. MR is a low power study type and, as such, an overly

conservative approach to multiple testing can be excessive (Burgess et al., 2019). However, in the

context of the present study the results of the early AMD phenome-wide screen were considered

significant only if they remained: significant after false discovery rate correction (FDR) in the MRE IVW;

nominally significant in weighted mode and weighted median; and nominally significant throughout the

leave-one-out analysis (IVW-MRE) (Benjamini and Hochberg, 1995). This conservative approach was

selected as a large number of phenotypes was studied and because we wanted to focus on high

confidence signals.

Where causal traits for early AMD were identified, the relationship between these traits and advanced

AMD was studied. These two AMD classifications are phenotypically distinct but are generally part of the

same disease spectrum. When traits failed to replicate as causal factors in the advanced AMD dataset it

could not be inferred that these traits are not truly causal for early AMD. However, significance in both

AMD phenotypes provided support for the detected causal links and evidence that a factor plays a role

across the disease spectrum.

Multivariable mendelian randomisation

Multivariable MR was performed in circumstances where it was important to estimate the effect of >1

closely related (and/or potentially confounding) exposure traits (Sanderson et al., 2019). P-values for the

inclusion of instruments for the exposures of interest were optimised to obtain sufficiently high (>10)

conditional F-statistics for reliable analysis (Sanderson et al., 2021). With this in mind, selection for

exposures began at a p-value threshold of > 5E-08. Where a trait's instruments had a conditional F-

statistic <10, the p-value for selection was reduced in an automated manner by factor of 10 until an F-

statistic >10 was obtained. The same clumping procedure as in the univariable MR analysis was used.

Adjusted Cochran's Q-statistics were calculated, with a p-value of <0.05 indicating significant

heterogeneity. Where the Cochran's Q-statistic indicated heterogeneity, a Q-statistic minimisation

procedure was used to evaluate the causal relationship; testing assumed both high (0.9) and low (0.1)

levels of phenotypic correlation (Sanderson et al., 2021).

Two-sample multivariable Mendelian randomisation approach based on Bayesian model
averaging (MR-BMA)

Multivariable MR can be used to obtain effect estimates for a few (potentially related) traits. However, it

cannot be directly applied when many traits need to be considered. In contrast, MR-BMA, a Bayesian

approach first described by Zuber and colleagues (Zuber et al., 2020), can search over large sets of

potential risk factors to determine which are most likely to be causal.

Notably, Zuber and colleagues previously performed an in-depth analysis which considered the role of

lipids against an older AMD GWAS. The relevant study served as proof-of-concept for the MR-BMA

method and demonstrated that several lipid traits have causal roles in AMD. However, the analysis had

two potential limitations. First, it downweighed fatty acid traits through limiting composite traits for SNV

identification to HDL, LDL, and triglycerides. Second, numerous lipid traits with a potential role in AMD

were not included in the analysis. For these reasons, we chose to conduct a more comprehensive

analysis with a slightly altered approach. The following study design modifications were made compared

to the study by Zuber and colleagues (Zuber et al., 2020):

(1) Fatty acids were included as a composite trait (utilising GWAS data for serum fatty acids derived from

the Nightingale Health 2020 resource as listed in TwoSampleMR package (Hemani et al., 2018)).

(2) All lipid and fatty acid measures included in a GWAS by Kettunen and colleagues (Kettunen et al.,

2016) were considered as potential causal traits (n=102 traits).

(3) A more recent AMD GWAS was used (Winkler et al., 2020).

(4) In general, multivariable MR (of any sort) cannot produce reliable results where the studied traits are

≥0.99 correlated with respect to the included instruments. For this reason, where two traits were highly

correlated, one was removed at random rather than by manually selecting traits in a manner which risks

selection bias.

MR-BMA for immune cell and complement phenotypes was additionally performed. In this analysis,

instruments were obtained at genome wide significance (p-value>5E-08) for every included exposure in

the model (given that composite traits were not available/applicable). For the immune cell MR-BMA, all

immune traits that were studied in a GWAS by Orru and colleagues (Orrù et al., 2020) and were present

in the TwoSampleMR package were used as exposures. In the complement analysis, all complement

traits available in relevant studies by Sun and colleagues (Sun et al., 2018) and Suhre and colleagues

(Suhre et al., 2017) were used, with the exception of complement subfractions.

For the MR-BMA analysis, the prior probability was set to 0.1 and the prior variance was set to 0.25. A

stochastic search with 10,000 iterations was undertaken and empirical p-values with 100,000

permutations were calculated.

Presentation of effect sizes:

Effect sizes are presented to enable appraisal of the impact of putative risk factors. For the univariable

MR analysis, these are presented as OR per 1 standard deviation (SD) of change in the exposure for

continuous traits, and as beta values for binary exposure traits. This approach was selected because

ORs are uninformative for binary exposure traits (Burgess and Labrecque, 2018). Furthermore, beta

values are not presented by all MR authorities where an exposure variable is binary, and it is often

highlighted that these values are only indicative. Multivariable MR effect sizes are presented as beta

estimates irrespective of the nature of the exposure variable (given that the role of multivariable MR within

this study was purely to identify confounders).

MR-BMA effect sizes are presented in the form of a model-averaged causal estimate (MACE). The MACE

is a conservative estimate of the direct causal effect of an exposure on an outcome averaged across

models. It is noted that the primary function of MR-BMA IS to highlight the probable causal trait among a

number of candidate causal risk factors. Although the MR-BMA findings can be used to interpret the

direction of effect, they should not be necessarily viewed as an absolute guide to magnitude (Zuber et al.,

2020).

<u>Software</u>

R version 4.1.0.

TwoSampleMR version 0.5.6.

RadialMR version 1.0.

MVMR version 0.3

MR-BMA code was sourced from https://github.com/verena-zuber/demo_AMD.

RESULTS

Overview

Focusing on early AMD, univariable MR analysis was applied to a broad range of traits. Following quality

control, significant results were replicated in an advanced AMD dataset and further analyses were

conducted using multivariable MR and MR-BMA (Zuber et al., 2020).

Overall, 4,591 traits were eligible for analysis. Among these, 44 were found to be putatively causal for

early AMD **(Table 1 and Table 1 - Source Data 1 & 2)**. Most of these causal traits were serum lipoprotein

concentration and compositional measures (n=29). Other significant traits identified included immune cell

phenotypes (n=5), serum proteins (n=6) and disease phenotypes (n=4).

Table 1. Significant results detected in a phenome-wide univariable Mendelian randomisation (MR) analysis of early age-related macular degeneration (AMD). Only traits passing the stringent quality control described in the methods are listed.

Trait name	Odds ratio	FDR-adjusted IVW p-value	MRE IVW beta
Rheumatoid arthritis	NA	1.51E-06	0.08
Unswitched memory B cell % B cell	1.11	5.40E-05	0.10
CD62L- Dendritic Cell % dendritic cell	0.95	3.32E-02	-0.05
Effector Memory CD8+ T cell absolute count	1.08	1.79E-04	0.07
CD25 on IgD+ CD38- naive B cell	0.93	1.25E-04	-0.07
CD80 on plasmacytoid dendritic cell	0.96	1.63E-02	-0.04
Total cholesterol in IDL*	0.80	6.48E-08	-0.22
Free cholesterol in IDL*	0.80	5.51E-08	-0.22
Total lipids in IDL*	0.79	3.35E-09	-0.23
Concentration of IDL particles*	0.80	9.83E-09	-0.23
Phospholipids in IDL*	0.79	1.86E-09	-0.23
Triglycerides in IDL*	0.84	2.40E-07	-0.18
Total cholesterol in large LDL*	0.83	9.85E-08	-0.19
Cholesterol esters in large VLDL*	0.83	1.41E-07	-0.19
Free cholesterol in large LDL*	0.83	7.07E-07	-0.18
Total lipids in large LDL*	0.83	1.04E-07	-0.19
Concentration of large LDL particles*	0.83	2.40E-07	-0.19
Phospholipids in large LDL*	0.83	1.33E-06	-0.18

Cholesterol esters in large VLDL*	0.82	6.55E-03	-0.20
18:2 linoleic acid (LA)*	0.80	2.93E-07	-0.23
Total cholesterol in LDL*	0.82	4.46E-08	-0.19
Total cholesterol in medium LDL*	0.82	3.64E-09	-0.20
Cholesterol esters in medium LDL*	0.82	5.93E-09	-0.20
Total lipids in medium LDL*	0.81	3.35E-09	-0.21
Concentration of medium LDL particles*	0.81	3.64E-09	-0.21
Phospholipids in medium LDL*	0.81	1.86E-09	-0.21
Total cholesterol in small LDL*	0.81	9.85E-08	-0.21
Total lipids in small LDL*	0.81	2.65E-07	-0.21
Total cholesterol in small VLDL*	0.85	2.54E-02	-0.16
Serum total cholesterol*	0.77	3.67E-08	-0.26
Total phosphoglycerides*	0.81	3.93E-03	-0.21
Triglycerides in very large HDL*	0.87	4.37E-04	-0.14
Total lipids in very small VLDL*	0.84	1.97E-03	-0.18
Concentration of very small VLDL particles*	0.84	9.04E-04	-0.18
Phospholipids in very small VLDL*	0.83	3.63E-03	-0.19
Interferon alpha-10	1.14	1.83E-02	0.13
Protein S100-A5	1.07	6.94E-04	0.07
Serine palmitoyltransferase 2	0.86	3.17E-02	-0.15
CD59 glycoprotein	1.10	1.77E-03	0.09
Complement factor H-related protein 5	1.09	9.30E-05	0.09
Cathepsin F	1.10	4.44E-03	0.10
Benign neoplasm: Skin, unspecified	NA	8.23E-03	0.07
Psychiatric diseases	NA	8.54E-05	-0.43
Myotonic disorders	NA	2.32E-02	0.00

* these traits (trait ID 'met-c' in **Table 1 - Source Data 1**) had 1.9% sample overlap with the early AMD dataset, a minor degree of overlap which is unlikely to bias results; the remaining causal traits had no sample overlap with early AMD.

FDR, false discovery rate; IVW, inverse variance weighted; MRE, multiplicative random effects; %, as a proportion of; NA, not applicable (as exposure traits of binary nature do not produce accurate odds ratios; beta values can be used instead to infer direction of effect but not necessarily magnitude).

Lipoprotein metabolism is linked to AMD risk

Univariable MR demonstrated significant causal relationships for 28 serum lipoprotein measures and 1

serum fatty acid concentration (18:2 linoleic acid) in early AMD (**Table 1)**. These relationships were also

strongly supported by the results of our advanced AMD analysis **(Table 1 - Source Data 3).**

Serum metabolites are highly correlated traits, and the instruments for serum lipoprotein and fatty acid
measures in the present study were demonstrably correlated **(Figure 1 - Figure supplement 1)**. MR-
BMA was used in order to discern which traits were driving the causal relationships. In our initial analysis,
two genetic variants (rs11065987 [*BRAP*] and rs10455872 [*LPAL2*]) were found to be outliers in terms of
Q-statistic (Q-statistic 13.84-15.71 and 6.79-10.46 respectively across models) **(Figure 1)**. These SNVs
were therefore omitted and the analysis was re-run. In subsequent analyses, no outlier SNVs were
identified. The top 10 models in terms of posterior probability are presented in **Table 1 - Source Data 4**;
the top 10 individual risk factors in terms of marginal inclusion probability (MIP; defined as the sum of the
posterior probabilities over all the models where the risk factor is present) are shown in **Table 2**. The
MIPs of all the traits included this analysis are plotted in **Figure 2.** The top four traits with respect to their
MIP were serum sphingomyelins (MIP=0.76, MACE=0.29), triglycerides in IDL (MIP=0.32, MACE=-0.16),
free cholesterol (MIP=0.20, MACE=0.07) and phospholipids in very small VLDL (MIP=0.63, MACE= -
0.31). It is noted that whilst MR-BMA is designed to select the likely causal risk factor among a set of
candidate causal traits, it is often not possible to achieve this with certainty. As such, a definitive
statement of causality for individual lipid traits cannot be obtained within the utilised MR causal inference
framework.
352
Serine palmitoyltransferase 2, an enzyme which catalyses the first committed step in sphingolipid
biosynthesis, was robustly associated with early AMD. The detected effect size highlighted that genetic
liability to increasing serum enzyme levels is protective of AMD (**Table 1, Table 1 - Source Data 1**; OR
0.86, p-value=1.00E-03). It is of interest that increasing serum levels of the related enzyme serine
palmitoyltransferase 1 also appeared to be protective of early AMD in most measures (**Table 1 - Source
Data 2**; OR 0.94, IVW p-value = 2.00E-03, FDR-adjusted IVW p-value = 0.05, weighted median p-value =
0.05; significant throughout leave-one-out analysis). Serine palmitoyltransferase enzymes were not
significantly related to the risk of advanced AMD **(Table 1 - Source Data 3)**.
361
362
363
364

Table 2. Lead causal traits identified by MR-BMA of lipid-related phenotypes in early age-related macular degeneration (AMD) ranked according to their marginal inclusion probability (MIP).

Rank	Risk factor (Trait ID)	Marginal inclusion probability (MIP)	Average effect	Nominal p-value	FDR-adjusted p-value
1	Sphingomyelins (met-c-935)	0.76	0.30	2.40E-04	5.76E-03
2	Phospholipids in very small VLDL (met-c-955)	0.63	-0.31	1.00E-05	7.20E-04
3	Triglycerides in IDL (met-c-872)	0.32	-0.16	2.10E-04	5.76E-03
4	Free cholesterol (met-c-858)	0.20	0.07	2.83E-03	5.09E-02
5	Omega-3 fatty acids (met-c-855)	0.07	0.01	3.70E-02	2.34E-01
6	Free cholesterol in very large HDL (met-c-944)	0.07	0.01	2.74E-02	2.34E-01
7	Total lipids in very small VLDL (met-c-953)	0.06	-0.02	2.08E-02	2.34E-01
8	Cholesterol esters in medium VLDL (met-c-910)	0.05	0.01	3.66E-02	2.34E-01
9	Cholesterol esters in very large HDL (met-c-943)	0.05	0.01	4.87E-02	2.34E-01
10	Ratio of bisallylic groups to double bonds (met-c-844)	0.05	0.01	4.41E-02	2.34E-01

Reinforcing a causal role for complement

Increasing serum levels of the complement proteins CD59 glycoprotein (OR 1.10, IVW p-value = 2.04E-05) and Complement factor H-related protein 5 (OR 1.10, IVW p-value = 7.70E-07) were found to be associated with genetic liability to early AMD and passed multiple testing correction. Whilst Complement C4, Complement factor B and Complement factor I were also related to early AMD risk at nominal level of significance, they did not exceed the conservative criteria for causal inference imposed in this study with respect to robust measures **(Table 1 - Source Data 2)**. In subsequent advanced AMD analyses (**Table 1 -Source Data 3**), CD59 glycoprotein remained related to disease risk (OR 1.07, IVW p-value = 0.04, weighted median p-value = 9.00E-04, MR-Egger p-value = 2.00E-03, weighted mode p-value = 7.00E-03). Complement factor I reached a nominal level of significance in the IVW measure (OR 0.95, p-value

= 0.05) but did not remain significant in robust tests. No other complement measures that were nominally

significant for early AMD were found to be significantly related to advanced AMD.

Complement traits are correlated with one another, and as such MR-BMA analysis was performed. In the

initial analysis, the relationship between complement proteins was strongly suggested to be driven by

Complement factor H (posterior probability=0.83, MIP=1.00, MACE=-0.68). However, rs2274700

(*CFHR3*, Cook's distance 26.26-55.24 and Cochran's Q 0.87-2.12 across models) and rs10824796

(*MBL2*, Cook's distance 0.13-9.93 and Cochran's Q 0.87-22.12 across models) were identified as outlier

instruments and therefore warranted removal from subsequent analyses. Rs2274700, a genetic variant

known to be associated with AMD risk (Liao et al., 2016), represented the only instrument strongly

associated with Complement factor H in the MR-BMA analysis and, as such, its removal precluded an

informative high-throughput analysis of the complement cascade. Whilst it is clear that there is a causal

relationship between complement and AMD, it is not possible to comment with precision about the

specific complement-related molecule driving this.

Other immune traits have mixed protective and causal effects in AMD

Our screen identified other serum immune traits with potential causal roles in early AMD, with several of

these related to dendritic cell populations. These traits were: Unswitched memory B cell as a proportion of

B cells (OR=1.11, p-value=4E-07); CD62L- dendritic cell as a proportion of dendritic cells (OR=0.95, p-

value=1.22E-03); Effector Memory CD8+ T cell absolute count (OR=0.95, p-value=1.56E-06); CD80 on

plasmacytoid dendritic cell (OR=0.96, p-value=4.20E-04); and Interferon alpha 10 (OR=1.14, p-

value=5.38E-04). Notably, numerous immune cell traits were also found to be significantly associated with

advanced AMD (**Table 1 - Source Data 3)**. As with lipids, these traits are highly correlated and it is not

possible to confidently pinpoint which immune cell trait is truly causal in a univariable analysis due to

potential overlap in terms of genetic instruments. MR-BMA (using data from a GWAS by Orru and

colleagues (Orrù et al., 2020)) for the above immune traits was not possible due to the presence of too

few genetic variants strongly instrumenting across the exposure variables which, in turn, led to a failure to

construct a meaningful model. For this reason, it was not possible to determine which of these specific

immune traits are driving the casual relationship between immune cell traits and AMD (though dendritic

cell traits dominate the univariable analysis).

Disease phenotypes are related to AMD risk

There are four disease phenotypes/groups which were found to be significant at a level which satisfy the

criteria for causal inference defined in this study. These traits are rheumatoid arthritis (beta= 0.08, IVW p-

value=9.86E-09), psychiatric diseases (beta=-0.43, IVW p-value=6.88E-07), 'benign neoplasm: skin,

unspecified' (beta=0.07, IVW p-value= 1.76E-04) and myotonic disorders (beta=-0.003, IVW p-value=

7.49E-04). None of these disorders were significant in advanced AMD. The very broad and non-specific

definitions encapsulated in the GWAS of benign skin neoplasms, psychiatric disease and myotonic

disorder make it challenging to apply further analyses within the scope of this study, but these may be

interesting phenotypes for exploration in future studies.

In univariable MR, rheumatoid arthritis was identified to be causally related to AMD as detailed above.

Multivariable MR with effector memory CD8+ T cell absolute count as a second exposure variable

(utilising a Q-statistic minimisation procedure) demonstrated that the causal effect of rheumatoid arthritis

was either mediated by immune cells or underpinned by correlated pleiotropy (rheumatoid arthritis effect

size: 0.03, 95% CI: -0.007 to 0.055; effector memory CD8+ T cell absolute count effect size 0.07, 95% CI:

0.001 to 0.19). This finding persists in models assuming both very high (0.9) and low (0.1) phenotypic

correlation.

With respect to the relationship between AMD and psychiatric disease, no reverse causation was

identified (p=0.31). More specific psychiatric traits which could be analysed were not significantly related

to genetic liability to early AMD **(Table 1 -Source Data 2)**. An extensive range of psychiatric disorders

were also explored in the advanced AMD dataset and but no significant relationships were detected

(Table 1 - Source Data 3). Due to the broad nature of this trait, and the negative results for all disorders

explored, it was not possible to discern the driving signal behind the significant result for psychiatric

disease. It appears probable that psychiatric disease is a false positive result given that it lacks: the

supporting evidence of biological plausibility; validation in an advanced AMD dataset; or supportive

evidence through significance of other similar traits in the early AMD analysis.

Serum cathepsin F and S100 proteins have a causal role in AMD

In addition to the four previously mentioned molecules (**Table 1**, Interferon alpha-10, CD59 glycoprotein,

Complement factor H-related protein and serine palmitoyltransferase 2) a further two serum proteins were

identified to have likely causal effects. These are S100-A5 (OR 1.07, IVW p-value = 6.80E-06) and

Cathepsin F (OR= 1.10, IVW p-value = 7.16E-05).

Serum cathepsin F was shown to be causally related to early AMD. However, no other molecules from

the serum cathepsin group produced significant results for early AMD and no cathepsin group protein was

robustly significant for advanced AMD (although cathepsin S and G have significant IVW p-values of

1.71E-05 and 0.001 respectively, **Table 1 - Source Data 3**).

Serum protein S100-A5 was demonstrated to be causal for AMD (OR 1.07, IVW p-value = 6.80E-06).

Additionally, protein S100-A13 was significantly related to early AMD risk in both IVW (OR=1.07, p-

value=1.22E-05) and robust measures (weighted median p-value=3.31E-05, weighted mode p-

value=9.00E-04 , Egger p-value= 1.00E-03) but did not remain significant throughout the leave-one-out

analysis, suggesting that the relationship for this particular protein could be driven by a small number of

influential variants. Whilst protein S100-A2 was nominally significant in the IVW analysis (p=0.04), it did

not pass any robust measures. No S100 group proteins were significantly related to advanced AMD risk

(Table 1 - Source Data 3).

DISCUSSION

In this study we used MR to advance our understanding of AMD pathogenesis. Our findings indicate protective effects for serum VLDL and IDL compositional and concentration traits in AMD. Causal effects for serum free cholesterol and sphingolipid metabolism were also highlighted. Notably, our results suggest that increasing serum levels of sphingomyelins are causally linked to AMD and we report protective relationships for specific enzymes implicated in sphingolipid biosynthesis. It is known that sphingolipids play key roles in retinal physiology, and there is emerging evidence that some sphingolipids may play a role in AMD pathogenesis (Simon et al., 2021). Further, our findings are in keeping with those of a meta-analysis of metabolomic studies in which pathway enrichment analysis suggested a role for sphingolipid metabolism in AMD (Hou et al., 2020). Whilst sphingomyelin itself has not previously been described as a causative factor for AMD, other substances involved in sphingolipid metabolism (including ceramide, which can serve as both a substrate for sphingomyelin synthesis or a product of sphingomyelin metabolism) have been suggested as causal factors through regulation of retinal cell death, inflammation and neovascularization (Simon et al., 2021). It is known that the balance of sphingolipids has crucial roles in determining cell fates, and, therefore, the contrasting effects of serine palmitoyltransferase and sphingomyelin observed in this study are noteworthy (Taniguchi and Okazaki, 2020). Our study therefore adds weight to the assertion that sphingolipids play a key role in AMD pathophysiology. Further study of sphingolipid metabolism in individuals with AMD is expected to provide important insights and to highlight potential treatment targets.

Our MR findings support the well-documented assertion that complement plays a key role in AMD pathogenesis (Clark and Bishop, 2018; Jha et al., 2007; Sivaprasad and Chong, 2006). Whilst MR-BMA was not feasible for this set of traits, univariable analysis supported roles for Complement factor H-related protein 5 and CD59 glycoprotein. A causal signal was obtained for Complement factor H-related protein 5, a finding in keeping with recent studies highlighting the role of this and other factor-H-related proteins (FHR-1 to FHR4; all involved in the regulation of complement factor C3b turnover) in AMD (Cipriani et al., 2021). An intriguing observation was the apparently causal role of increasing serum levels of the complement-related CD59 glycoprotein (an inhibitor of membrane attack complex formation) in both early

and advanced AMD. This finding appears to be at odds with previous studies that investigated the impact

of high levels of CD59 in a model of AMD and found a potential therapeutic benefit (Cashman et al.,

2011). The relationship between systemic and local complement activation is poorly understood and

studies have produced conflicting results with respect to the roles of each complement molecule in AMD

(Clark and Bishop, 2018; Jha et al., 2007). It can therefore be speculated that the superficially

inconsistent results between our study and previous reports on CD59 reflect the complicated relationship

between serum and retinal CD59 expression.

Notably, a connection between complement and lipid accumulation has been proposed in AMD. This is

due to the observation that dysregulation of the complement system results in lipid deposition both

systemically and in the retina. There is also metabolomic evidence that decreased VLDL and increased

HDL levels are associated with increased complement activation independent of AMD status (Armento et

al., 2021). Given our results, the relationship between VLDL and complement is an area warranting

further study.

Cathepsins are a very diverse family of lysosomal proteases which play an important part in cellular

homeostasis by participating in antigen processing and by degrading chemokines and proteases (Yadati

et al., 2020). Cathepsins are susceptible to age-related alterations and have been linked to modulation of

pro-inflammatory signalling pathways. Additionally, cathepsins D and S play a direct role in the retina as

they help with the degradation of photoreceptor outer segments. It has also been shown that

homozygosity for variant B cystatin C (an inhibitor of cysteine proteases) causes reduced secretion of

mature cystatin C and is associated with increased susceptibility to neovascular AMD (Turk et al., 2012;

Zurdel et al., 2002). For these reasons, it has been suggested that dysregulation of cathepsin activity may

be a factor in AMD pathophysiology. Here, we show data suggesting that serum cathepsin F, a

ubiquitously expressed cysteine cathepsin, has a causal relationship with early AMD. The roles of specific

cathepsins in AMD is an area that has not yet been mechanistically explored.

S100 proteins act as intracellular regulators and extracellular signalling proteins. Intracellularly, S100

proteins have a wide range of roles including regulating proliferation, differentiation, apoptosis, calcium

homeostasis, metabolism, and inflammation (Donato et al., 2013). Extracellular S100 proteins have

important roles in immunity and inflammation. Of the S100 protein family, serum S100-A5 is the only

protein to be robustly identified as a causal factor for AMD in our study. Whilst S100 proteins have

received little study in AMD thus far, the binding of S100B to RAGE (Receptor for Advanced Glycation

End products) and the subsequent increase in VEGF have previously been shown to be linked to the

development of AMD (Ma et al., 2007; Xia et al., 2017). S100-A5 is a protein which has received little

study and, as such, its biological functions are mostly uncharacterised, (although is known to be involved

in inflammation via the activation of RAGE) (Wheeler and Harms, 2017). Whilst it is outside the scope of

this study to assert a mechanistic hypothesis for this finding, it seems likely that S100-A5 could be linked

to inflammatory processes.

To a large extent, the present study has captured known AMD risk factors. A notable omission however is

smoking, an accepted AMD contributor. In this study, cigarettes smoked per day was associated with

early AMD at a nominal level of significance (IVW p-value = 0.009) but was not significant with FDR

adjustment or in other measures. This finding is most probably reflective of the underpowered nature of

MR. Given the hypothesis-free design of this study and the low power of MR, it would be inappropriate to

conclude that non-significant results provide evidence of no relationship between certain environmental

exposures and AMD.

In summary, this study has identified previously undescribed causal factors for AMD in addition to

reinforcing several previously suggested AMD risk factors. Future research will: explore how the

interaction between causal traits affects AMD risk; investigate how the casual and protective factors

described here alter the progression of established disease; and follow-up these causal factors from a

mechanistic perspective in order to open up avenues for therapeutic interventions.

Ethical approval

This study utilised publicly available data and no additional ethical approval is required.

ACKNOWLEDGEMENTS

We acknowledge funding from the Wellcome Trust (224643/Z/21/Z); the University of Manchester's

Wellcome Institutional Strategic Support Fund (Wellcome ISSF) grant (204796/Z/16/Z); the UK National

Institute for Health Research (NIHR) Academic Clinical Fellow and Clinical Lecturer Programmes; Retina

UK and Fight for Sight (GR586); the Australian National Health and Medical Research Council (NHMRC)

(1150144).

COMPETING INTERESTS

TA is involved on advisory boards and has received grants and speaker fees from Allergan, Novartis,

Bayer, Roche, Bausch and Lomb, Heidelberg, Topcon and Canon. The remaining authors declare no

competing interests.

DATA AVAILABILITY

All data associated with this study are provided within the supplementary data.

REFERENCES

Armento A, Ueffing M, Clark SJ. 2021. The complement system in age-related macular degeneration. *Cell*
Mol Life Sci **78**:4487–4505.
Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful approach to
multiple testing. *J R Stat Soc* **57**:289–300.
Boddy S, Islam M, Moll T, Kurz J, Burrows D, McGown A, Bhargava A, Julian TH, Harvey C, Marshall
JNG, Hall BPC, Allen SP, Kenna KP, Sanderson E, Zhang S, Ramesh T, Snyder MP, Shaw PJ,
McDermott C, Cooper-Knock J. 2022. Unbiased metabolome screen leads to personalized medicine
strategy for amyotrophic lateral sclerosis. *Brain Commun* **4**. doi:10.1093/braincomms/fcac069
Bowden J, Davey Smith G, Burgess S. 2015. Mendelian randomization with invalid instruments: effect
estimation and bias detection through Egger regression. *Int J Epidemiol* **44**:512–525.
Bowden J, Davey Smith G, Haycock PC, Burgess S. 2016a. Consistent Estimation in Mendelian
Randomization with Some Invalid Instruments Using a Weighted Median Estimator. *Genet Epidemiol*
40:304–314.
Bowden J, Del Greco M. F, Minelli C, Smith GD, Sheehan NA, Thompson JR. 2016b. Assessing the
suitability of summary data for two-sample Mendelian randomization analyses using MR-Egger
regression: the role of the I2 statistic. *International Journal of Epidemiology*. doi:10.1093/ije/dyw220
Bowden J, Holmes MV. 2019. Meta-analysis and Mendelian randomization: A review. *Res Synth Methods*
10:486–496.
Bowden J, Spiller W, Del Greco M F, Sheehan N, Thompson J, Minelli C, Davey Smith G. 2018.
Improving the visualization, interpretation and analysis of two-sample summary data Mendelian
randomization via the Radial plot and Radial regression. *Int J Epidemiol* **47**:1264–1278.
Burgess S, Labrecque JA. 2018. Mendelian randomization with a binary exposure variable: interpretation
and presentation of causal estimates. *European Journal of Epidemiology*. doi:10.1007/s10654-018-

0424-6
Burgess S, Smith GD, Davies NM, Dudbridge F, Gill D, Maria Glymour M, Hartwig FP, Holmes MV,
Minelli C, Relton CL, Theodoratou E. 2019. Guidelines for performing Mendelian randomization
investigations. *Wellcome Open Research* **4**. doi:10.12688/wellcomeopenres.15555.2
Burgess S, Thompson SG. 2017a. Interpreting findings from Mendelian randomization using the MR-
Egger method. *Eur J Epidemiol* **32**:377–389.
Burgess S, Thompson SG. 2017b. Erratum to: Interpreting findings from Mendelian randomization using
the MR-Egger method. *Eur J Epidemiol* **32**:391–392.
Burgess S, Thompson SG, CRP CHD Genetics Collaboration. 2011. Avoiding bias from weak instruments
in Mendelian randomization studies. *Int J Epidemiol* **40**:755–764.
Cashman SM, Ramo K, Kumar-Singh R. 2011. A Non Membrane-Targeted Human Soluble CD59
Attenuates Choroidal Neovascularization in a Model of Age Related Macular Degeneration. *PLoS*
ONE. doi:10.1371/journal.pone.0019078
Chakravarthy U, Wong TY, Fletcher A, Piault E, Evans C, Zlateva G, Buggage R, Pleil A, Mitchell P.
2010. Clinical risk factors for age-related macular degeneration: a systematic review and meta-
analysis. *BMC Ophthalmology*. doi:10.1186/1471-2415-10-31
Cipriani V, Tierney A, Griffiths JR, Zuber V, Sergouniotis PI, Yates JRW, Moore AT, Bishop PN, Clark SJ,
Unwin RD. 2021. Beyond factor H: The impact of genetic-risk variants for age-related macular
degeneration on circulating factor-H-like 1 and factor-H-related protein concentrations. *Am J Hum*
Genet **108**:1385–1400.
Clark SJ, Bishop PN. 2018. The eye as a complement dysregulation hotspot. *Semin Immunopathol*
40:65–74.
Cunningham F, Allen JE, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Austine-Orimoloye O, Azov
AG, Barnes I, Bennett R, Berry A, Bhai J, Bignell A, Billis K, Boddu S, Brooks L, Charkhchi M,
Cummins C, Da Rin Fioretto L, Davidson C, Dodiya K, Donaldson S, El Houdaigui B, El Naboulsi T,
Fatima R, Giron CG, Genez T, Martinez JG, Guijarro-Clarke C, Gymer A, Hardy M, Hollis Z, Hourlier
T, Hunt T, Juettemann T, Kaikala V, Kay M, Lavidas I, Le T, Lemos D, Marugán JC, Mohanan S,
Mushtaq A, Naven M, Ogeh DN, Parker A, Parton A, Perry M, Piližota I, Prosovetskaia I, Sakthivel
MP, Salam AIA, Schmitt BM, Schuilenburg H, Sheppard D, Pérez-Silva JG, Stark W, Steed E,
Sutinen K, Sukumaran R, Sumathipala D, Suner M-M, Szpak M, Thormann A, Tricomi FF, Urbina-
Gómez D, Veidenberg A, Walsh TA, Walts B, Willhoft N, Winterbottom A, Wass E, Chakiachvili M,
Flint B, Frankish A, Giorgetti S, Haggerty L, Hunt SE, IIsley GR, Loveland JE, Martin FJ, Moore B,
Mudge JM, Muffato M, Perry E, Ruffier M, Tate J, Thybert D, Trevanion SJ, Dyer S, Harrison PW,
Howe KL, Yates AD, Zerbino DR, Flicek P. 2022. Ensembl 2022. *Nucleic Acids Res* **50**:D988–D995.
Donato R, Cannon BR, Sorci G, Riuzzi F, Hsu K, Weber DJ, Geczy CL. 2013. Functions of S100 proteins.
Curr Mol Med **13**:24–57.
Fleckenstein M, Keenan TDL, Guymer RH, Chakravarthy U, Schmitz-Valckenberg S, Klaver CC, Wong
WT, Chew EY. 2021. Age-related macular degeneration. *Nat Rev Dis Primers* **7**:31.
Fritsche LG, Chen W, Schu M, Yaspan BL, Yu Y, Thorleifsson G, Zack DJ, Arakawa S, Cipriani V, Ripke
S, Igo RP Jr, Buitendijk GHS, Sim X, Weeks DE, Guymer RH, Merriam JE, Francis PJ, Hannum G,
Agarwal A, Armbrecht AM, Audo I, Aung T, Barile GR, Benchaboune M, Bird AC, Bishop PN,
Branham KE, Brooks M, Brucker AJ, Cade WH, Cain MS, Campochiaro PA, Chan C-C, Cheng C-Y,
Chew EY, Chin KA, Chowers I, Clayton DG, Cojocaru R, Conley YP, Cornes BK, Daly MJ, Dhillon B,
Edwards AO, Evangelou E, Fagerness J, Ferreyra HA, Friedman JS, Geirsdottir A, George RJ,
Gieger C, Gupta N, Hagstrom SA, Harding SP, Haritoglou C, Heckenlively JR, Holz FG, Hughes G,
Ioannidis JPA, Ishibashi T, Joseph P, Jun G, Kamatani Y, Katsanis N, N Keilhauer C, Khan JC, Kim
IK, Kiyohara Y, Klein BEK, Klein R, Kovach JL, Kozak I, Lee CJ, Lee KE, Lichtner P, Lotery AJ,
Meitinger T, Mitchell P, Mohand-Saïd S, Moore AT, Morgan DJ, Morrison MA, Myers CE, Naj AC,
Nakamura Y, Okada Y, Orlin A, Ortube MC, Othman MI, Pappas C, Park KH, Pauer GJT, Peachey
NS, Poch O, Priya RR, Reynolds R, Richardson AJ, Ripp R, Rudolph G, Ryu E, Sahel J-A,
Schaumberg DA, Scholl HPN, Schwartz SG, Scott WK, Shahid H, Sigurdsson H, Silvestri G,
Sivakumaran TA, Smith RT, Sobrin L, Souied EH, Stambolian DE, Stefansson H, Sturgill-Short GM,
Takahashi A, Tosakulwong N, Truitt BJ, Tsironi EE, Uitterlinden AG, van Duijn CM, Vijaya L,
Vingerling JR, Vithana EN, Webster AR, Wichmann H-E, Winkler TW, Wong TY, Wright AF, Zelenika
D, Zhang M, Zhao L, Zhang K, Klein ML, Hageman GS, Lathrop GM, Stefansson K, Allikmets R,
Baird PN, Gorin MB, Wang JJ, Klaver CCW, Seddon JM, Pericak-Vance MA, Iyengar SK, Yates

JRW, Swaroop A, Weber BHF, Kubo M, Deangelis MM, Léveillard T, Thorsteinsdottir U, Haines JL,
Farrer LA, Heid IM, Abecasis GR, AMD Gene Consortium. 2013. Seven new loci associated with
age-related macular degeneration. *Nat Genet* **45**:433–9, 439e1–2.

Fritsche LG, Igl W, Bailey JNC, Grassmann F, Sengupta S, Bragg-Gresham JL, Burdon KP, Hebbring SJ,
Wen C, Gorski M, Kim IK, Cho D, Zack D, Souied E, Scholl HPN, Bala E, Lee KE, Hunter DJ, Sardell
RJ, Mitchell P, Merriam JE, Cipriani V, Hoffman JD, Schick T, Lechanteur YTE, Guymer RH,
Johnson MP, Jiang Y, Stanton CM, Buitendijk GHS, Zhan X, Kwong AM, Boleda A, Brooks M, Gieser
L, Ratnapriya R, Branham KE, Foerster JR, Heckenlively JR, Othman MI, Vote BJ, Liang HH,
Souzeau E, McAllister IL, Isaacs T, Hall J, Lake S, Mackey DA, Constable IJ, Craig JE, Kitchner TE,
Yang Z, Su Z, Luo H, Chen D, Ouyang H, Flagg K, Lin D, Mao G, Ferreyra H, Stark K, von Strachwitz
CN, Wolf A, Brandl C, Rudolph G, Olden M, Morrison MA, Morgan DJ, Schu M, Ahn J, Silvestri G,
Tsironi EE, Park KH, Farrer LA, Orlin A, Brucker A, Li M, Curcio CA, Mohand-Saïd S, Sahel J-A,
Audo I, Benchaboune M, Cree AJ, Rennie CA, Goverdhan SV, Grunin M, Hagbi-Levi S,
Campochiaro P, Katsanis N, Holz FG, Blond F, Blanché H, Deleuze J-F, Igo RP Jr, Truitt B, Peachey
NS, Meuer SM, Myers CE, Moore EL, Klein R, Hauser MA, Postel EA, Courtenay MD, Schwartz SG,
Kovach JL, Scott WK, Liew G, Tan AG, Gopinath B, Merriam JC, Smith RT, Khan JC, Shahid H,
Moore AT, McGrath JA, Laux R, Brantley MA Jr, Agarwal A, Ersoy L, Caramoy A, Langmann T,
Saksens NTM, de Jong EK, Hoyng CB, Cain MS, Richardson AJ, Martin TM, Blangero J, Weeks DE,
Dhillon B, van Duijn CM, Doheny KF, Romm J, Klaver CCW, Hayward C, Gorin MB, Klein ML, Baird
PN, den Hollander AI, Fauser S, Yates JRW, Allikmets R, Wang JJ, Schaumberg DA, Klein BEK,
Hagstrom SA, Chowers I, Lotery AJ, Léveillard T, Zhang K, Brilliant MH, Hewitt AW, Swaroop A,
Chew EY, Pericak-Vance MA, DeAngelis M, Stambolian D, Haines JL, Iyengar SK, Weber BHF,
Abecasis GR, Heid IM. 2016. A large genome-wide association study of age-related macular
degeneration highlights contributions of rare and common variants. *Nat Genet* **48**:134–143.

Han X, Gharahkhani P, Mitchell P, Liew G, Hewitt AW, MacGregor S. 2020a. Genome-wide meta-
analysis identifies novel loci associated with age-related macular degeneration. *J Hum Genet*
65:657–665.

Han X, Ong J-S, An J, Hewitt AW, Gharahkhani P, MacGregor S. 2020b. Using Mendelian randomization
to evaluate the causal relationship between serum C-reactive protein levels and age-related macular
degeneration. *Eur J Epidemiol* **35**:139–146.

Han X, Ong J-S, Hewitt AW, Gharahkhani P, MacGregor S. 2021. The effects of eight serum lipid
biomarkers on age-related macular degeneration risk: a Mendelian randomization study. *Int J*
Epidemiol **50**:325–336.

Hartwig FP, Davey Smith G, Bowden J. 2017. Robust inference in summary data Mendelian
randomization via the zero modal pleiotropy assumption. *Int J Epidemiol* **46**:1985–1998.

Hartwig FP, Davies NM, Hemani G, Davey Smith G. 2016. Two-sample Mendelian randomization:
avoiding the downsides of a powerful, widely applicable but potentially fallible technique. *Int J*
Epidemiol **45**:1717–1726.

Hemani G, Tilling K, Davey Smith G. 2017. Orienting the causal relationship between imprecisely
measured traits using GWAS summary data. *PLoS Genet* **13**:e1007081.

Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, Laurin C, Burgess S, Bowden J,
Langdon R, Tan VY, Yarmolinsky J, Shihab HA, Timpson NJ, Evans DM, Relton C, Martin RM,
Davey Smith G, Gaunt TR, Haycock PC. 2018. The MR-Base platform supports systematic causal
inference across the human phenome. *Elife* **7**. doi:10.7554/eLife.34408

Hou X-W, Wang Y, Pan C-W. 2020. Metabolomics in Age-Related Macular Degeneration: A Systematic
Review. *Invest Ophthalmol Vis Sci* **61**:13.

Jha P, Bora PS, Bora NS. 2007. The role of complement system in ocular diseases including uveitis and
macular degeneration. *Mol Immunol* **44**:3901–3908.

Julian TH, Boddy S, Islam M, Kurz J, Whittaker KJ, Moll T, Harvey C, Zhang S, Snyder MP, McDermott C,
Cooper-Knock J, Shaw PJ. 2021. A review of Mendelian randomization in amyotrophic lateral
sclerosis. *Brain*. doi:10.1093/brain/awab420

Kettunen J, Demirkan A, Würtz P, Draisma HHM, Haller T, Rawal R, Vaarhorst A, Kangas AJ, Lyytikäinen
L-P, Pirinen M, Pool R, Sarin A-P, Soininen P, Tukiainen T, Wang Q, Tiainen M, Tynkkynen T, Amin
N, Zeller T, Beekman M, Deelen J, van Dijk KW, Esko T, Hottenga J-J, van Leeuwen EM, Lehtimäki
T, Mihailov E, Rose RJ, de Craen AJM, Gieger C, Kähönen M, Perola M, Blankenberg S, Savolainen
MJ, Verhoeven A, Viikari J, Willemsen G, Boomsma DI, van Duijn CM, Eriksson J, Jula A, Järvelin M-

R, Kaprio J, Metspalu A, Raitakari O, Salomaa V, Slagboom PE, Waldenberger M, Ripatti S, Ala-
Korpela M. 2016. Genome-wide study for circulating metabolites identifies 62 loci and reveals novel
systemic effects of LPA. *Nat Commun* **7**:11122.

Klein R, Meuer SM, Myers CE, Buitendijk GHS, Rochtchina E, Choudhury F, de Jong PTVM, McKean-
Cowdin R, Iyengar SK, Gao X, Lee KE, Vingerling JR, Mitchell P, Klaver CCW, Wang JJ, Klein BEK.
2014. Harmonizing the classification of age-related macular degeneration in the three-continent AMD
consortium. *Ophthalmic Epidemiol* **21**:14–23.

Kvale MN, Hesselson S, Hoffmann TJ, Cao Y, Chan D, Connell S, Croen LA, Dispensa BP, Eshragh J,
Finn A, Gollub J, Iribarren C, Jorgenson E, Kushi LH, Lao R, Lu Y, Ludwig D, Mathauda GK, McGuire
WB, Mei G, Miles S, Mittman M, Patil M, Quesenberry CP Jr, Ranatunga D, Rowell S, Sadler M,
Sakoda LC, Shapero M, Shen L, Shenoy T, Smethurst D, Somkin CP, Van Den Eeden SK, Walter L,
Wan E, Webster T, Whitmer RA, Wong S, Zau C, Zhan Y, Schaefer C, Kwok P-Y, Risch N. 2015.
Genotyping Informatics and Quality Control for 100,000 Subjects in the Genetic Epidemiology
Research on Adult Health and Aging (GERA) Cohort. *Genetics* **200**:1051–1060.

Liao X, Lan C-J, Cheuk I-W-Y, Tan Q-Q. 2016. Four complement factor H gene polymorphisms in
association with AMD: A meta-analysis. *Arch Gerontol Geriatr* **64**:123–129.

Li X, Li H, Cheng J, Wang M, Zhong Y, Shi G, Yu A-Y. 2022. Causal Associations of Thyroid Function
and Age-Related Macular Degeneration: A Two-Sample Mendelian Randomization Study. *Am J
Ophthalmol* **239**:108–114.

Ma W, Lee SE, Guo J, Qu W, Hudson BI, Schmidt AM, Barile GR. 2007. RAGE ligand upregulation of
VEGF secretion in ARPE-19 cells. *Invest Ophthalmol Vis Sci* **48**:1355–1361.

Orrù V, Steri M, Sidore C, Marongiu M, Serra V, Olla S, Sole G, Lai S, Dei M, Mulas A, Virdis F, Piras
MG, Lobina M, Marongiu M, Pitzalis M, Deidda F, Loizedda A, Onano S, Zoledziewska M, Sawcer S,
Devoto M, Gorospe M, Abecasis GR, Floris M, Pala M, Schlessinger D, Fiorillo E, Cucca F. 2020.
Complex genetic signatures in immune cells underlie autoimmunity and inform therapy. *Nat Genet*
52:1036–1045.

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW,
Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association and population-based
linkage analyses. *Am J Hum Genet* **81**:559–575.

Sanderson E, Davey Smith G, Windmeijer F, Bowden J. 2019. An examination of multivariable Mendelian
randomization in the single-sample and two-sample summary data settings. *Int J Epidemiol* **48**:713–
727.

Sanderson E, Glymour MM, Holmes MV, Kang H, Morrison J, Munafò MR, Palmer T, Schooling CM,
Wallace C, Zhao Q, Davey Smith G. 2022. Mendelian randomization. *Nat Rev Methods Primers* **2**.
doi:10.1038/s43586-021-00092-5

Sanderson E, Spiller W, Bowden J. 2021. Testing and correcting for weak and pleiotropic instruments in
two-sample multivariable Mendelian randomization. *Stat Med* **40**:5434–5452.

Simon MV, Basu SK, Qaladize B, Grambergs R, Rotstein NP, Mandal N. 2021. Sphingolipids as critical
players in retinal physiology and pathology. *J Lipid Res* **62**:100037.

Sivaprasad S, Chong NV. 2006. The complement system and age-related macular degeneration. *Eye*
20:867–872.

Sudlow C, Gallacher J, Allen N, Beral V, Burton P, Danesh J, Downey P, Elliott P, Green J, Landray M,
Liu B, Matthews P, Ong G, Pell J, Silman A, Young A, Sprosen T, Peakman T, Collins R. 2015. UK
biobank: an open access resource for identifying the causes of a wide range of complex diseases of
middle and old age. *PLoS Med* **12**:e1001779.

Suhre K, Arnold M, Bhagwat AM, Cotton RJ, Engelke R, Raffler J, Sarwath H, Thareja G, Wahl A, DeLisle
RK, Gold L, Pezer M, Lauc G, El-Din Selim MA, Mook-Kanamori DO, Al-Dous EK, Mohamoud YA,
Malek J, Strauch K, Grallert H, Peters A, Kastenmüller G, Gieger C, Graumann J. 2017. Connecting
genetic risk to disease end points through the human blood plasma proteome. *Nat Commun*
8:14357.

Sun BB, Maranville JC, Peters JE, Stacey D, Staley JR, Blackshaw J, Burgess S, Jiang T, Paige E,
Surendran P, Oliver-Williams C, Kamat MA, Prins BP, Wilcox SK, Zimmerman ES, Chi A, Bansal N,
Spain SL, Wood AM, Morrell NW, Bradley JR, Janjic N, Roberts DJ, Ouwehand WH, Todd JA,
Soranzo N, Suhre K, Paul DS, Fox CS, Plenge RM, Danesh J, Runz H, Butterworth AS. 2018.
Genomic atlas of the human plasma proteome. *Nature* **558**:73–79.

Taniguchi M, Okazaki T. 2020. Ceramide/sphingomyelin rheostat regulated by sphingomyelin synthases

and chronic diseases in Murine models. *J Lipid Atheroscler* **9**:380–405.

Turk V, Stoka V, Vasiljeva O, Renko M, Sun T, Turk B, Turk D. 2012. Cysteine cathepsins: from structure,
function and regulation to new frontiers. *Biochim Biophys Acta* **1824**:68–88.

Wheeler LC, Harms MJ. 2017. Human S100A5 binds Ca2+ and Cu2+ independently. *BMC Biophys* **10**:8.

Winkler TW, Grassmann F, Brandl C, Kiel C, Günther F, Strunz T, Weidner L, Zimmermann ME, Korb CA,
Poplawski A, Schuster AK, Müller-Nurasyid M, Peters A, Rauscher FG, Elze T, Horn K, Scholz M,
Cañadas-Garre M, McKnight AJ, Quinn N, Hogg RE, Küchenhoff H, Heid IM, Stark KJ, Weber BHF.
2020. Genome-wide association meta-analysis for early age-related macular degeneration highlights
novel loci and insights for advanced disease. *BMC Med Genomics* **13**:120.

Xia C, Braunstein Z, Toomey AC, Zhong J, Rao X. 2017. S100 Proteins As an Important Regulator of
Macrophage Inflammation. *Front Immunol* **8**:1908.

Yadati T, Houben T, Bitorina A, Shiri-Sverdlov R. 2020. The Ins and Outs of Cathepsins: Physiological
Function and Role in Disease Management. *Cells* **9**. doi:10.3390/cells9071679

Zuber V, Colijn JM, Klaver C, Burgess S. 2020. Selecting likely causal risk factors from high-throughput
experiments using multivariable Mendelian randomization. *Nat Commun* **11**:29.

Zurdel J, Finckh U, Menzer G, Nitsch RM, Richard G. 2002. CST3 genotype associated with exudative
age related macular degeneration. *Br J Ophthalmol* **86**:214–219.

FIGURE LEGENDS

Figure 1: Plots outlining the top-ranking models with respect to their posterior probability in the first run of Mendelian randomisation Bayesian model averaging (MR-BMA) of lipid-related traits in early age-related macular degeneration (AMD). Plots 1A and 1C present Cook's distance while plots 1B and 1D present Cochran's Q. Outlier instruments are annotated.

The Cochran's Q is a measure which serves to identify outlier variants with respect to the fit of the linear model. The Q statistic is used to identify heterogeneity in a meta-analysis, and to pinpoint specific variants as outliers. The contribution of variants to the overall Q statistic is measured (defined as the weighted squared difference between the observed and predicted association with the outcome) in order to identify outliers. Cook's distance on the other hand is utilised to identify influential observations (*i.e.* those variants which have a strong association with the outcome). Such variants are removed from the analysis because they may have an undue influence over variable selection, leading to models which fit that variant well but others poorly.

Figure 2: Graph detailing the results of a Mendelian randomisation Bayesian model averaging (MR-BMA) analysis that aims to identify causal lipid-related risk factors for early age-related macular degeneration (AMD). The studied phenotypes are ranked according to their marginal inclusion probability (MIP); four likely causal traits are highlighted.

Figure 1 - Supplementary Figure 1: Plot illustrating the correlations between the beta values for metabolites considered in our Mendelian randomisation Bayesian model averaging (MR-BMA) analysis for early age-related macular degeneration (AMD). This plot visually represents the correlation matrix between the genetic associations of the exposure variables with respect to their instruments. The traits are labelled according to their 'Trait ID'; further information can be found in the source data.

Model 1: Free cholesterol and phospholipids in very small VLDL

Model 2: Sphingomyelins and phospholipids in very small VLDL