

Title Page:

Execution by lethal injection: Autopsy findings of pulmonary edema

Joel B. Zivot, MD, FRCP(C), MA (Corresponding Author)

Associate Professor of Anesthesiology and Surgery

Emory University School of Medicine

2015 Uppergate Drive

Emory University, Atlanta, GA 30307

Mark A. Edgar, MD

Department of Pathology

Mayo Clinic

4500 San Pablo Rd South

Jacksonville, FL 32224

David A. Lubarsky MD

CEO, UC Davis Health and Vice Chancellor for Human Health Sciences

UC Davis Health

4619 X Street, Suite 3101

Sacramento, CA 95817

August 17, 2022

Execution by lethal injection: Autopsy findings of pulmonary edema

Joel B. Zivot, MD, FRCP(C), MA

Mark A. Edgar, MD

David A. Lubarsky MD

Authors' contributions:

All authors contributed equally to the creation of this manuscript.

Conflict of interest statements:

The authors have no conflicts of interests to disclose.

Role of funding source:

No funding source to disclose.

Ethics committee approval:

N/A (excluded from review according to applicable HIPPA rules)

Abstract:

Lethal injection is the common method of capital punishment in the United States (US), but problematic executions feed controversy centered on the 8th amendment of the US constitution that proscribes cruel punishment.¹ However, the definition of cruelty is contentious, and lethal injection displays few signs of the hidden course of a potentially cruel death. Few survive and those who die cannot give witness. In this study, we reviewed autopsy reports of prisoners executed by intravenous midazolam (usually followed by a neuromuscular blocking agent and potassium chloride) or pentobarbital as a single drug. Autopsy reports for 43 inmates obtained from eight states were reviewed to identify any abnormal findings. Pulmonary edema

(often fulminant with froth filling airways) emerged as a consistent and unexpected abnormality in 33/43 autopsies (76.74%), 23/28 midazolam executions (82.14%), and 10/15 (66.66%) pentobarbital executions. Injecting heavy overdoses of either acidic or basic solutions into the bloodstream may be directly toxic to pulmonary capillary endothelial cells. The injury manifests by the sudden escape of edema fluid into normally air-filled lungs, resulting in pulmonary edema. These findings explain witnessed respiratory distress during executions and increase concern about an inability of an execution witness to observe such a finding when a paralytic is added.

Introduction:

In the United States (US), lethal injection is currently the most common method of state-sponsored judicial execution. Forensic pathologist, Dr. Jay Chapman, originally proposed a three-drug protocol² involving a drug to render the inmate unconscious followed by a neuromuscular blocking agent to paralyze the muscles of respiration and potassium chloride to stop electrical activity in the heart as a more humane alternative to death in the electric chair. Lethal injection protocols have varied over time in the US, but a three-drug protocol initiated with a large intravenous dose midazolam (usually 500 mg) has been used in seven states and a protocol employing pentobarbital (usually as a single agent in a dose of 2500 mg or 5000 mg) has been employed in 14 states. Protocols involving midazolam have generated controversy with eyewitness reports of movement following administration of the drug and observations indicating respiratory distress (grunting, straining, stridor, gurgling, coughing).

Lethal injection, as a lawful form of capital punishment in the United States, meets its purpose when the method of execution succeeds in killing the inmate. While the pharmacokinetic

and pharmacodynamics of the drugs have been either directly studied or postulated^{3 4}, structural effects of drugs employed in the specific context of lethal injection have not been investigated until now. Death certificates provide information on what contributes to the cause of death but in lethal injection, the cause of death is always listed as ‘homicide’⁵, a legal term for a human death caused by a human, and can be legal or not. The purpose of this study was to review autopsy reports of 43 inmates executed by lethal injection using two separate protocols, one that includes midazolam and the other utilizing pentobarbital as the single agent, in order to investigate the anatomic effects of execution by lethal injection.

Materials and Methods:

Autopsy reports were obtained from eight states (Alabama, Arkansas, Arizona, Florida, Georgia, Ohio, Oklahoma, and Virginia) for inmates executed by lethal injection protocols that included either pentobarbital alone (15 cases) or midazolam accompanied by one or more additional drugs that include Potassium Chloride/Acetate and a paralytic such as Vecuronium Bromide (28 cases). One of the autopsies (case 24) was performed in Ohio by one of the authors (ME). Autopsies are in the public domain and are not considered part of PHI and are therefore not constrained for release by HIPPA. The final data set under review in this series does not include the names of any individuals. Midazolam was either administered as the first of three drugs (followed next by a neuromuscular blocking agent such as vecuronium bromide and then potassium chloride/acetate in 27 cases) or in combination with hydromorphone (one case). Gross descriptions of organs described in each report were reviewed for evidence of any abnormality. Microscopic descriptions (available in 11 midazolam executions) were also reviewed. Detailed timelines released by the State were available for 22 midazolam executions,

some of which were accompanied by witness observations made at the time of execution by attorneys or representatives of the media. Postmortem interval was calculated using time of death (either obtained from state records or media reports) and time of autopsy.

Results:

Inmates ranged in age from 35 to 83 years in age; 42 were male and 1 female. Where known, postmortem intervals ranged from 4 hours and 22 minutes to 2 days; postmortem intervals were not known for pentobarbital executions. The autopsy was conducted the day following execution in 4 cases, but postmortem interval could not be calculated from available data.

Gross descriptions varied in level of detail. For example, cardiac valve circumferences were measured in some cases and not others. There were most often two intravenous lines in place (one in each antecubital fossa), but additional puncture wounds and/or vascular access catheters were described in some cases, with a maximum of five intravenous catheters in place (right and left antecubital, right hand, left wrist and right groin).

Among the expected findings for this population were varying degrees of coronary atherosclerosis (with up to 75% stenosis of left anterior descending coronary artery), hepatomegaly, and an incidental thyroid mass, but no consistent changes were found in organ systems other than respiratory. A cerebellar tumor was noted in case 10, but histologic findings were not reported. Blood was described as “present from the mouth” in one midazolam execution, and in three others there was either watery or bloody fluid in the mouth and nose. An unexpected macroscopic finding in 33 of 43 autopsies (24 midazolam and 10 pentobarbital

executions) was the presence of fulminant pulmonary edema as evidenced by the presence of froth, frothy fluid or blood-tinged froth located in the in the tracheobronchial tree (8 midazolam and 6 pentobarbital executions) or fluid in lung parenchyma or small airways (15 midazolam and 4 pentobarbital executions). The presence of pulmonary edema was confirmed microscopically in 8 midazolam autopsies, including one case in which there was no macroscopic evidence of pulmonary edema. Altogether, 34 of 43 cases showed some evidence of edema in lung parenchyma or airways. Lung weights ranged from 430 to 1150 grams for the right (mean 765 grams) and from 460 to 980 grams for the left (mean 681 grams).

Microscopic examination of brain tissue was performed in 4 cases; 3 cases (cases 8, 22, and 24) reported histologic evidence of hypoxic-ischemic neuronal injury and one showed no histologic abnormality in the brain. (See table 1)

Discussion:

In this study of autopsy findings of 43 inmates executed by a protocol involving Midazolam or single drug Pentobarbital, we unexpectantly found evidence of lung edema in 34 of 43 cases. Lethal injection is not a medical act and protocol revisions have been driven by drug availability and not as an act of technical refinement or process improvement. Lethal injection has not been subject to the oversight of medical science because of a prohibition of involvement according to medical ethical opinion.⁶

In *Baze v. Rees*, the United States Supreme Court asserts that since capital punishment is constitutional, “it necessarily follows that there must be a means of carrying it out”.⁷ Lethal injection gained public approval based on a perception that death by this method was not cruel. When lethal injection is set next to the electric chair, lethal gas, or the firing squad, it seemed the

clear winner as compared to these other methods, witnessing death by lethal injection generally reveals little. Even in that case though, documented “botched” executions reportedly occur at a higher rate than other methods⁸. It is conceivable the courts have also struggled to consider the possibility of an 8th amendment violation in an action that outwardly seemed so bloodless. Although adopted as a non-cruel alternative to other methods of execution, lethal injection has increasingly resulted in problematic executions with serious complications. Though the court seems determined to defend lethal injection as a constitutional method of execution, death row prisoners regard this quite differently. Recent execution jurisprudence has seen a rise in claims by prisoners who now seek to be executed by lethal gas, the firing squad, and the electric chair. In *Bucklew v. Presythe*⁹, Bucklew argued death by lethal gas would be a less cruel method of execution as he suffered from cavernous hemangiomas within his airway. In the majority opinion, the court did not find in favor of this claim. The State of Tennessee employs the three-drug method beginning with Midazolam. Death-row prisoners in Tennessee chose to die by electric chair and not lethal injection as a result of fear about cruelty in the Tennessee lethal injection method (personal communication with Zivot, February 7, 2019). In *Glossip v. Gross*, the Supreme Court referred to 11 executions involving 500 mg of intravenous midazolam that “appear to have been conducted with no significant problems”¹⁰. Nine of those cases were part of this study, and all showed acute pulmonary edema; there is no autopsy data available for the other two. This statement requires reassessment considering new evidence that adds pulmonary edema to the list of unanticipated complications of lethal injection using midazolam and pentobarbital. The additional likelihood of awareness during execution by the 3-drug midazolam method with attendant sensations of air hunger and drowning raises grave concerns about the

suitability of midazolam (a drug which cannot induce general anesthesia) as the first drug in three-drug lethal injection protocols.

According to the 8th amendment of the US constitution, punishment must not be cruel. The evolving definition of cruelty tracks with the evolution of civil society. Witnesses to executions are necessary in order to engage their empathy to judge the cruelty of the act but lethal injection execution witnesses see a state-curated event. According to Austin Sarat¹¹, in a review of 1,054 lethal injection executions, witnesses report that the execution appeared botched in 75 cases (7.12%). Before now, physical evidence for cruelty has been lacking and these autopsy findings suggest that the rate of cruelty in lethal injection executions may be significantly higher than what can be seen by traditional visual observation alone.

The autopsy has existed in some form for centuries, and the discipline of pathology as a medical subspecialty has its roots in the modern autopsy as practiced in Central Europe during the 19th century. Autopsy is from the Greek *autopsia* and translates to “see for oneself”.¹² The alternate term necropsy, means “to look at the dead”.¹³ The purpose of the autopsy has varied considerably over time, but most autopsies performed today are carried out to pinpoint a cause (and sometimes manner) of death and to thoroughly document the causes and effects of disease on the body. An autopsy may be considered the only complete physical examination ever made of a body, and just as antemortem physical examinations vary in thoroughness and specific focus depending on the indication for the procedure, so too do autopsies differ in the level of detail in dissection and description depending on the context in which they are performed. Not all states require autopsies following execution, but some jurisdictions perform autopsies for all homicides. Other purposes for autopsy in this setting include documentation of lethal injection, confirmation of correct intravenous catheter placement, exclusion of any other contributing cause

of death, and to search for unexpected complications of the procedure. Autopsies on executed prisoners also have focused on determination of the presence and quantity of execution drugs in post-mortem blood samples. Lethality claims about death by injected pentobarbital drawn from literature suggesting fatality with a blood concentration of 30 mg/L.¹⁴ Rapid and accurate methods to detect the presence of barbiturates have been available for a number of years but uncertainty remains about what blood level of a barbiturate is necessary to produce death.¹⁵ Midazolam is a benzodiazepine with low associated toxicity and death by the exclusive use of midazolam is, to our knowledge, unreported.

Our intent in this study was to review a set of autopsy reports (written by pathologists from seven different states) following lethal injection to further our understanding of the process by which death is brought about in this situation. Significant pathology in the lungs emerged as a consistent and unexpected finding. Pulmonary edema and congestion are common findings in hospital autopsies where death is usually a gradual process, often as a result of progressive cardiac failure or overwhelming infection, but it is not seen in cases of sudden death¹⁶. Likewise, froth in the airways (while commonly seen in acute cardiogenic pulmonary edema) suggests physical or chemical injury as seen in death by drowning, electrocution, inhalation of toxic gas, or heroin overdose. It requires breathing to create froth, so the timing of pulmonary edema must, but definition, occur prior to the administration of a paralyzing agent or heart stopping medication (after which breathing does not occur). The cause of pulmonary edema in heroin overdose has been debated with proposed etiologies including central respiratory depression, hypoxia-induced increase in capillary permeability, primary toxic effect on alveolar capillaries, and myocardial depression¹⁷.

The finding of “vascular congestion or parenchymal edema” is an alarming one for the question of cruelty in this method of execution. The subjective experience of acute pulmonary edema has been listed for patients partly in these words: “A feeling of suffocating or drowning that worsens when lying down”¹⁸

If it is assumed that agents used in the 3-drug midazolam protocol function as they do in clinical settings, then midazolam would cause sedation, the paralytic would quickly induce apnea, and potassium chloride would quickly stop the heart. In this sequence of events there is no reason to expect pulmonary congestion or edema; in fact, lung weights close to those seen in very sudden death (i.e. normal weights) would be anticipated. Normal adult lung weights vary according to gender with the right lung generally weighing more than the left. In one paper, the average lung weight was reported as 234 grams (s +/- 39)¹⁹. In another paper describing the weight of lungs in men, authors report the weight of the right lung to range between 155-720 grams and the left lung to range between 112-675 grams²⁰. The lung weights observed in this study were all above 400 grams (average weights for right and left lungs, respectively, in midazolam executions were 787 grams and 707 grams and for pentobarbital executions 723 grams and 631 grams), indicating some combination of vascular congestion and parenchymal edema.

As pentobarbital is not administered to patients in a clinical setting in the massive doses used for execution (e.g. 5 grams), its effects on the body at high concentration are unknown. Pentobarbital is a highly alkaline solution (pH between 9.8-11)²¹. Pentobarbital is a Barbiturate and these drugs have long been known to cause vascular injury if improperly administered, with inadvertent intra-arterial injection sometimes causing gangrene²². In contrast, midazolam preparations for injection are adjusted to a more acidic pH of 3 to 3.6²³. It is possible that the

massive quantities of acidic or basic solution entering the bloodstream during lethal injection may be directly toxic to pulmonary capillary endothelial cells and that the earliest manifestation of this injury is the escape of edema fluid into the normally air-filled lungs. The statement by inmate 25, “My body is on fire” raises the possibility that intravascular acid-sensing ion channel activation by the large quantity of highly acidic midazolam. Pain at the injection site is reported in cases when the injected solution has a pH less than 4 or greater than 11²⁴.

It is intuitive that pulmonary edema in the three-drug midazolam protocol is caused by midazolam itself for several reasons. First, witnesses report respiratory distress early in the procedure (sometimes almost immediately after announcement that the execution has begun), before the paralytic agent or potassium chloride could have taken effect. Second, the production of froth in the setting of pulmonary edema requires the active mixing of fluid, proteins, and surfactant, and this process ceases after the paralytic takes effect; non-aerated edematous lung tissue does not produce froth²⁵. Third, intravenous acid is known to cause diffuse lung injury in animal models²⁶, a process that begins with exudation of fluid and protein into the alveolar space. Pulmonary edema in this setting is appropriately described as “flash pulmonary edema” in contrast with that accompanying death from natural causes (agonal pulmonary edema). Furthermore, even in massive overdoses, benzodiazepines very rarely cause death and would not be expected to cause pulmonary edema as an agonal affect associated with gradual multi-organ and cardiac failure.

We speculate that pentobarbital produces pulmonary edema via a similar mechanism, specifically the caustic action of alkaline solution on pulmonary alveolar capillaries, but because of the different mechanism of action of pentobarbital which includes a direct respiratory depressant effect, it is difficult to be confident of this. As indicated above, heavy, congested,

edematous lungs are common following death from a wide variety of causes, but fulminant pulmonary edema with froth filling the respiratory tree is a noteworthy finding more commonly associated with lung injury than pure respiratory depression.

There were a number of limitations inherent in this study. First, the authors did not “see for themselves” firsthand the anatomic findings detailed above (except in case 24), but rather relied on written autopsy reports. This has the advantage of limiting biased observations as pathologists from eight different states recorded consistent findings of pulmonary edema, but also makes it possible that minor degrees of pulmonary edema may have been overlooked if not specifically sought. Similarly, histologic examination (which would have afforded a second opportunity to detect mild pulmonary edema) was performed in a minority of cases. It is also true that detailed medical histories seeking conditions which might predispose to pulmonary edema were not collected on these inmates, but it would be extraordinary if the majority of subjects in this series suffered from a medical predisposition to acute pulmonary edema.

Negative pressure pulmonary edema (caused by the development of a very high negative intrathoracic pressure) can develop as the result of airway obstruction in the context of sedation, but this typically manifests as a struggle for breath against upper airway obstruction followed by a deep inspiration. In the setting of negative pressure pulmonary edema, the observed lung fluid has a low protein concentration suggesting a hydrostatic, as opposed to a direct toxic effect.²⁷ It is not clear if the pulmonary edema fluid observed in our review had high or low levels of protein. Witnesses to executions, including one of the authors (JZ), do not report this pattern of breathing, making it unlikely to have played a major role in these cases.

Conclusion:

In our review of the autopsy findings of 43 inmates executed by a protocol involving Midazolam or single drug Pentobarbital, we found evidence of lung edema in 34 of 43 cases (79%) The occurrence of pulmonary edema in the setting of lethal injection execution may not be easily observed by witnesses to execution. Pulmonary edema is an extremely uncomfortable experience²⁸, and these findings raise the concern that lethal injection execution is far more uncomfortable an experience than previously believed. The mechanism of the development of pulmonary edema in these case remains uncertain, but it is possibly the result of the injection of large quantities of unphysiological pH solutions. The protein content in the observed pulmonary edema fluid was not measured and further analysis of this fluid would be warranted in order to determine if the mechanism of pulmonary edema was hydrostatic as opposed to cytotoxic.

¹ US const. amend. VIII

² <https://www.latimes.com/opinion/op-ed/la-oe-alper-3-drug-cocktail-20170420-story.html#:~:text=The%20three%2Ddrug%20protocol%20has,chloride%20to%20stop%20his%20heart.>

³ Koniaris, LG, Zimmers TA, Lubarsky DA, Sheldon, JP “Inadequate anaesthesia in lethal injection for execution” *Lancet* 2005 Apr 16; 365(9468): 1412-4

⁴ Zimmers TA, Sheldon J, Lubarsky DA, Lopez-Munoz F, Waterman L, Weisman R, Koniaris L G “Lethal injection for execution:chemical asphyxiation?” *PLoS Medicine* 2007 Apr 4 (4): e156

⁵ Zivot, J, “For All Those Executed, the Cause of Death is Homicide” KevinMD.com, 25 Mar.2014, www.kevinmd.com/blog/2014/04/executed-death-homicied.html.

⁶ *AMA J Ethics.* 2017;19(9):911-912. doi: 10.1001/journalofethics.2017.19.9.coet1-1709.

⁷ *Baze v. Rees*, 553 U.S. 35 (2008)

⁸ Sarat, Austin, “Gruesome Spectacles: Botched Executions and America’s Death Penalty” (Stanford University Press 2014)

⁹ *Bucklew v. Precythe*, 587 U.S. (2019)

¹⁰ Glossip v. Gross, 576 U.S. 28 (2015)

¹¹ Sarat, Austin, “Gruesome Spectacles: Botched Executions and America’s Death Penalty” (Stanford University Press 2014)

¹² Burton JL. The history of the autopsy. In: Burton JL, Ruty GN, eds. *The Hospital Autopsy*. London: Arnold, 2001, pp. 1–6.

¹³ Wood, MJ, Guha AK, “Declining clinical autopsy rates versus increasing medicolegal autopsy rates in Halifax, Nova Scotia, Why the difference? A Historical perspective, *Archives of Pathology & Laboratory Medicine*: July 2001, Vol 125, No. &, pp. 924-930

¹⁴ Druda DF, Gone S, Graudins A, “Deilberate self-poisoning with a lethal dose of pentobarbital with confirmatory serum drug concentrations: survival after cardiac arrest with supportive care”, *Journal of Medical Toxicology*, 2018 Aug 2; (15): 45-48

¹⁵ Gillett R, Warburton F G, “Barbiturate blood levels found at necropsy in proven cases of acute barbiturate poisoning” *Journal of Clinical Pathology*, 1970 Jul; 23 (5): 435-9

¹⁶ Molina DK, DiMaio VJ. Normal organ weights in men: part II – the brain, lungs, liver, spleen, and kidneys. *Am J Forensic Med Pathol*. 2012, 33(4):368-72.

¹⁷ Dinis-Oliveira RJ, Santos A, Magalhaes T. “Foam cone” exuding from the mouth and nostrils following heroin overdose. *Toxicol Mech Methods*. 2012, 22(2):159-160.

¹⁸ Mayo Clinic; Pulmonary edema; Sudden (acute) pulmonary edema signs and symptoms. Retrieved 2019, 15 Apr. <https://www.mayoclinic.org/diseases-conditions/pulmonary-edema/symptoms-causes/syc-20377009?p=1>

¹⁹ Joachim H, Riede U N, Mittermayer C, “The weight of human lungs as a diagnostic criterium (distinction of normal lungs from shock lungs by histologic, morphometric and biochemical investigations)” *Pathol Res Pract* 1978 May;162(1):24-40

²⁰ Milina D K, DiMaio V J M, “Normal organ weights in men: part II-the brain, lungs, liver, spleen, and kidneys” *Am J Forensic Med Pathol* 2012 Dec;33(4):368-72

²¹ Abdollahi, Baghaei A, “Encyclopedia of Toxicology” (Third Edition), 2014

²² Stone H H, Donnelly C C, “The accidental intraarterial injection of thiopental” *Anesthesiology* 1961 Nov-Dec;22:995-1006

²³ *Midazolam - Food and Drug Administration*. (n.d.). Retrieved July 21, 2022, from https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/208878Orig1s000lbl.pdf

²⁴ Klement W, Arndt J O, “Pain on i.v. injection of some anaesthetic agents is evoked by the unphysiological osmolality or pH of their formulations” *Br J Anaesth* 1991 Feb;66(2):189-95

²⁵ Said SI et al. Pulmonary surface activity in induced pulmonary edema. *J Clin Invest*. 1965, 44(3): 458-464.

²⁶ Pedoto A et al. Acidosis stimulates Nitric Oxide Production and lung damage in rats. *Am J Respir Crit Care Med*. 1999, 159: 397-402.

²⁷ Bhattacharya M, Kallet RH, Ware MB, Matthay MA, “Negative pressure pulmonary edema” Chest 2016 Oct; 150 (4) : 927-933

²⁸ Mayo Clinic <https://www.mayoclinic.org/diseases-conditions/pulmonary-edema/.../syc-20377009>).

medRxiv preprint doi: <https://doi.org/10.1101/2022.08.24.22279183>; this version posted August 27, 2022. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Inmate	State	R lung	L lung	Micro	Drug(s)	Observations	PMI
1	AL	598 g; parenchyma oozing moderate amounts of yellow-tinged frothy fluid	470 g	None noted	3 drug MZ	Shallow breathing for 10 min after administration of MZ	7 h 45 min
2	FL	555 g; congestion and edema	680 g	None noted	3 drug MZ	None available	10 h 45 min
3	AL	1020g; major bronchi contain froth; lung oozing large amounts of yellow-tinged frothy fluid	870 g	None noted	3 drug MZ	Rapid breathing; chest heaved up and down (for at least 3 minutes)	13 h 22 min
4	FL	900g; frothy red fluid oozes from surface on manual compression	880 g	Acute pulmonary congestion and edema	3 drug MZ	Feet twitched, yawned	14 h 58 min
5	FL	610 g; nonspecific congestion	565 g	None noted	3 drug MZ	Body convulsed for 10 seconds; eyes fluttered; mouth opened	13 h 24 min
6	FL	1120 g; ooze froth with slight manual compression; marked congestion and edema	965 g	None noted	3 drug MZ	Chest heaved up and down for more than 5 minutes; at one point breath grew shallow	14 h 17 min
7	AL	1150 g; upper airways contain blood-tinged frothy edema fluid; intra-pulmonary airways show blood-tinged edema fluid	910 g	None noted	3 drug MZ	Heavy breathing	12 h 32 min
8	VA	886 g; upper airways foamy liquid	715 g	Lung: edema and RBC; brain: HIE	3 drug MZ	Labored breathing for 5 minutes; snoring loudly; taking a giant breath; making crying sounds; having stomach movements	11 h 33 min
9	FL	985 g; bloody froth expressed on slight manual compression	825 g	None noted	3 drug MZ	Eye and head movements	16 h 24 min
10	FL	750 g; ooze bloody froth with slight manual compression	595 g	None noted	3 drug MZ	Visible breathing for 5 minutes	15 h 24 min
11	FL	735 g; acute edema and congestion	720 g	None noted	3 drug MZ	Breathing and movement slowed	13 h 17 min
12	FL	815 g; sectioned surfaces ooze blood and frothy fluid	775 g	None noted	3 drug MZ	Breathing ceased after 5 minutes	15 h 14 min
13	FL	760 g; pink tinged frothy fluid tracheobronchial tree; frothy fluid oozes from lung surface on compression	545 g	Acute pulmonary edema and congestion	3 drug MZ	Left arm and shoulder twitching for 15-30 sec	14 h 28 min
14	AR	835 g; mild diffuse edematous change	735 g	None noted	3 drug MZ	Chest moving up and down for at least 5 minutes; cheeks moving for 7 minutes	1 day
15	FL	860 g; slight congestion	590 g	Mild alveolar edema	3 drug MZ	None available	2 days
16	FL	840 g; ooze frothy red fluid; acute congestion and edema	620 g	None noted	3 drug MZ	Chest heaved and face turned purple	19 h 49 min
17	AR	755 g; white frothy fluid expressed on compression	660 g	None noted	3 drug MZ	Appeared to swallow multiple times	1 day
18	OK	740 g; edema	580 g	Pulmonary edema	3 drug MZ	Variety of movements including breathing heavily, writhing, mumbling, grimacing, grunting, and straining against restraints	1 d 13 h 19 min
19	AL	430 g; major bronchi contain froth	600 g	None noted	3 drug MZ	Breathing quickened; other extremity movements including raising arm	15 h 12 min
20	AL	680 g; mild vascular congestion	540 g	None noted	3 drug MZ	Labored breathing for 7 minutes before breathing slowed	10 h 48 min
21	AL	780 g; congested	980 g	Lungs show mild interstitial anthracosis	3 drug MZ	Chest moved during early part of execution	11 h 58 min
22	VA	800 g; moderate amount of froth upper airway	628 g	Lung: within normal limits; scattered hypoxic-ischemic	3 drug MZ	Deep breathing; diaphragm contracted sharply several times	12 h 15 min

				neurons hippocampus			
23	AL	850 g; congested	920 g	Pigmented macrophages, mucus plugs in bronchioles	3 drug MZ	Difficulty breathing, struggling for breath, heaving, asthmatic-sounding barking coughs and gasping respiration; coughed regularly for 15 minutes	13 h 55 min
24	OH	709 g; sections exude serosanguineous frothy fluid	665 g	Pulmonary edema; HIE	3 drug MZ	Chest was rapidly rising and falling; labored breathing; puffing lips out as he exhaled; gasping and wheezing loudly	23 h 46 min
25	OK	550 g; edematous; exudes moderate amount of clear frothy fluid	550 g	Congestion and edema	MZ, PB, KCl	Stated: "My body is on fire"	16 h 34 min
26	AR	590 g; serosanguineous fluid nares, oral cavity, upper and lower airways	555 g	None noted	3 drug MZ	Coughing, struggling for air, made noises like gasping and straining for breath; heaving and choking with chest pumping up and down; labored breathing	1 day
27	AR	750 g; abundant watery secretion nares, oral cavity, upper, lower airways	720 g	None noted	3 drug MZ	Deep and heavy breathing; sucked in air and repeatedly arched his back over 5-6 minutes	1 day
28	GA	610 g; exude marked frothy fluid	640 g	Intra-alveolar hemorrhage and edema fluid bilaterally	MZ, HM	Gasped more than 600 times over 1 hour and 40 minutes; heavy then shallow breathing; struggling efforts to breathe; gulped "like a fish on land"; snoring and sucking noises	16 h 14 min
29	GA	860 g	820 g	None noted	PB	N/A	N/A
30	GA	720 g; frothy fluid in tracheobronchial tree	640 g	None noted	PB	N/A	N/A
31	GA	860 g; congestion	720 g	None noted	PB	N/A	N/A
32	GA	740 g; moderate congestion	580 g	None noted	PB	N/A	N/A
33	GA	820 g; mild degree frothy fluid lower airways	720 g	None noted	PB	N/A	N/A
34	GA	760 g; cut surface exudes minimal frothy fluid	700 g	None noted	PB	N/A	N/A
35	GA	620 g; moderate amount of white foam upper airway	520 g	None noted	PB	N/A	N/A
36	GA	760 g; congestion and edema	640 g	None noted	PB	N/A	N/A
37	GA	800 g; frothy material main bronchi and small airways	640 g	None noted	PB	N/A	N/A
38	GA	720 g; congestion	600 g	None noted	PB	N/A	N/A
39	GA	580 g; edema	580 g	None noted	PB	N/A	N/A
40	GA	580 g; edema; edematous fluid major airways	460 g	None noted	PB	N/A	N/A
41	GA	520 g; white frothy fluid tracheobronchial tree	540 g	None noted	PB	N/A	N/A
42	GA	760 g	620 g	None noted	PB	N/A	N/A
43	GA	780 g; upper airways moderate amounts of white foam	680 g	None noted	PB	N/A	N/A

Table 1: Pathologic findings at autopsy and pertinent witness observations for 43 inmates executed with lethal injection protocols using midazolam. Abbreviations: HIE: hypoxic-ischemic encephalopathy; HM: hydromorphone; KCL: potassium chloride; Micro: microscopic findings; MZ: midazolam; N/A: not available; PB: pentobarbital; PMI: postmortem interval; RBC: red blood cells.