T cell responses to SARS-CoV-2 vaccination in people with multiple sclerosis differ between disease-modifying therapies

Asia-Sophia Wolf*[1], Anthony Ravussin*[1], Marton König*[2], Mathias H. Øverås[2], Guri Solum[1], Ingrid Fadum Kjønstad[1], Adity Chopra[3,4], Trygve Holmøy[3,5], Hanne F. Harbo[2,3], Silje Watterdal Syversen[6], Kristin Kaasen Jørgensen[7], Einar Høgestøl[2,3], Jon Torgils Vaage[3,4], Elisabeth G. Celius[2,3], Fridtjof Lund-Johansen[3,4], Ludvig A. Munthe[3,4,8], Gro Owren Nygaard[†2], Siri Mjaaland[†1]

* These authors contributed jointly to this publication

[†] Joint last authors

[1] Division of Infection Control, Section for Immunology, Norwegian Institute of Public Health, Oslo, Norway

[2] Department of Neurology, Oslo University Hospital, Oslo, Norway

[3] Institute of Clinical Medicine, University of Oslo, Oslo, Norway

[4] Department of Immunology, Oslo University Hospital, Oslo, Norway

[5] Department of Neurology, Akershus University Hospital, Lørenskog, Norway

[6] Center for Treatment of Rheumatic and Musculoskeletal Diseases (REMEDY), Diakonhjemmet Hospital, Oslo, Norway

[7] Department of Gastroenterology, Akershus University Hospital, Lørenskog, Norway

[8] KG Jebsen Centre for B cell malignancy, University of Oslo, Oslo, Norway

Corresponding author: Marton König, MD, PhD, Department of Neurology, Oslo University Hospital, PO Box 4956, Nydalen, N-0424 Oslo, Norway (makoni@ous-hf.no).

Conflicts of interest

MK reported receiving speaker honoraria from Novartis, Biogen, Merck and Sanofi outside

the submitted work.

TH has received speaker honoraria from Biogen, Merck, Novartis, Sanofi, Bristol Myers

Squibb and Roche, and participated in clinical trials organized by Roche, Merck and Biogen

EGC has participated in advisory boards and received speaker honoraria from Biogen,

Janssen, Merck, Novartis, Teva, Roche, Sanofi and Bristol Myers Squibb.

KKJ has received speaker bureaus from Bristol Myers Squibb and Roche.

Running title: Vaccine responses in multiple sclerosis

Key words: multiple sclerosis, vaccination, SARS-CoV-2, T cells, immunocompromised

Abstract

Immune responses in people with multiple sclerosis (pwMS) on disease-modifying therapies

(DMTs) have been of significant interest throughout the COVID-19 pandemic. Lymphocyte-

targeting immunotherapies including anti-CD20 treatments and sphingosine-1-phosphate

receptor (S1PR) modulators attenuate antibody responses after vaccination. Evaluation of

cellular responses after vaccination is therefore of particular importance in these populations.

In this study, we analysed CD4 and CD8 T cell functional responses to SARS-CoV-2 spike

peptides in healthy controls and pwMS on five different DMTs by flow cytometry. Although

pwMS on anti-CD20 and S1PR therapies had low antibody responses after both 2 and 3

vaccine doses, T cell responses in pwMS on anti-CD20 therapies were preserved after third

vaccination, even when additional anti-CD20 treatment was administered between vaccine

doses 2 and 3. PwMS taking S1PR modulators had low detectable T cell responses in

peripheral blood. CD4 and CD8 T cell responses to SARS-CoV-2 variants of concern Delta

and Omicron were lower than to the ancestral Wuhan-Hu-1 variant. Our results indicate the

importance of assessing both cellular and humoral responses after vaccination and suggest

that even in the absence of robust antibody responses vaccination can generate immune

responses in pwMS.

Introduction

Immune responses to SARS-CoV-2 in immunocompromised individuals have been of intense

interest throughout the COVID-19 pandemic. In the absence of vaccination, immunologically

vulnerable groups are especially susceptible to severe COVID-19 disease and hospitalisation

(reviewed in (1)); after vaccination, reduced responses to SARS-CoV-2 vaccines and

potential vaccine failure have been of particular concern.

Multiple sclerosis (MS) is an immune-mediated disease characterised by inflammation and

demyelination in the central nervous system. Current treatment involves modulation of the

immune system to alleviate inflammation. However, some disease-modifying therapies

(DMTs) can also impede an effective response to infectious diseases and vaccination

(reviewed in (2)). It is unclear whether people with MS (pwMS) are more susceptible to

severe COVID-19 disease in the absence of vaccination (3-6); current evidence suggests

that this varies depending on DMT usage, where treatment with anti-CD20 drugs presents an

increased risk factor (7, 8), as well as neurological disability, comorbidities, and age (6, 8). It

is therefore important to establish vaccine effectiveness in pwMS and whether vaccination

protects against COVID-19 disease to the same extent as in the general population. Certain

DMTs are known to be associated with increased risk of other infections: anti-CD20 drugs

such as rituximab, ocrelizumab, and ofatumumab are associated with a range of serious

infections, including respiratory tract infections (9, 10); sphingosine-1-phosphate receptor

(S1PR) modulators, including fingolimod, ozanimod, and siponimod, which sequester

lymphocytes in lymph nodes, are associated with increased risk of herpesvirus infections or

reactivations (11); and natalizumab, an anti-alpha-4 integrin monoclonal antibody (mAb), with

a risk of progressive multifocal leukoencephalopathy (9).

The primary focus of many vaccine efficacy studies to date has been the humoral immune

response (12, 13). Many DMTs, particularly anti-CD20 drugs, target B cells and people on

such treatments have reduced or non-existent antibody responses after vaccination (14-18);

pwMS on fingolimod have been found to have significantly reduced antibody responses (14,

19), By contrast, pwMS taking other DMTs including natalizumab (20, 21), cladribine (an

adenosine mimic which triggers lymphocyte apoptosis) (22), and alemtuzumab (an anti-

CD52 mAb that depletes T and B cells) (23) appear to have antibody responses comparable

to untreated control groups.

Nevertheless, it is unclear if vaccine-specific T cell responses are impaired. Several studies

have looked at cellular responses to SARS-CoV-2 spike peptides in pwMS after two doses of

SARS-CoV-2 vaccine (16, 21, 24-26) and found that IFN-γ+ T cell responses were

detectable in many, though not all, patients on a variety of DMTs. One exception was pwMS

treated with fingolimod (26), where T cell responses were significantly attenuated. Although

not all individuals on anti-CD20 therapies developed T cell responses, it further appears that

T cell responses and antibody titres are not well correlated, and so a lack of antibody

response is not in itself indicative of a failed response to vaccination (27). Additionally, data

on the effect of a third vaccine dose on both antibody levels and T cell responses are mixed;

some studies suggest no effect of additional vaccination on either humoral or cellular

immune responses (28), whereas others find boosted responses (29, 30).

The time period between receiving a dose of DMT and vaccination varies between DMTs.

Fingolimod, for example, is taken daily, whereas anti-CD20 treatments are administered at

six-month intervals, with a clear impact of this interval on the humoral response. An

increased gap between administration of anti-CD20 therapies and vaccination is associated

with stronger antibody responses (15, 27, 31, 32), which may be beneficial during

vaccination but can also lead to interruptions in ongoing treatment of MS or undesirable

delay in vaccine schedules. It is therefore of interest to establish what effect ongoing DMT

treatment has on vaccine responsiveness during both the primary vaccine course and for

subsequent boosters.

Recent register studies indicate that pwMS treated with high efficacy DMTs, including

alemtuzumab, natalizumab, cladribine, S1PR modulators, and anti-CD20 therapies, have the

best long-term outcomes for reduced worsening of disability and relapse outcomes (33, 34).

Although for safety reasons alemtuzumab is rarely given to newly diagnosed patients, many

people have been treated with this induction therapy during the last decade and comprise an

important subset of pwMS. This study therefore focused on the cellular response to these

five therapies that are among the most likely to be the treatments of choice for future pwMS.

The aim of this study was to investigate IgG antibody binding to the receptor binding domain

(RBD) on the spike protein of SARS-CoV-2 as well as functional spike-specific CD4 and CD8

T cell responses from pwMS on five different DMTs and a healthy control group after two

doses of SARS-CoV-2 vaccine. We also investigated whether a third vaccine dose improved

the humoral and/or cellular responses in individuals treated with rituximab or fingolimod who

had impaired IgG anti-spike RBD antibody responses after two vaccine doses.

Results

A cohort of pwMS living in Oslo or Akershus, Norway, who were treated with DMTs prior to

the beginning of the COVID-19 pandemic were recruited as part of an ongoing population-

based study of vaccine responses in pwMS in Norway (NevroVax) (15, 29). Cellular samples

were collected from a subset of individuals on different DMTs (fingolimod, rituximab,

cladribine, natalizumab and alemtuzumab) both before and after the primary course of 2

vaccine doses in April-July 2021 (Supplementary Figure 1). Healthy controls were recruited

from among healthcare workers at Diakonhjemmet Hospital and Akershus University

Hospital. Rituximab- and fingolimod-treated individuals who had low antibody responses after

vaccination were offered a 3rd vaccine dose in the summer of 2021 (EudraCT Number: 2021-

003618-37, see Methods), before recommendations for booster vaccines in Norway were

changed in September of that year to recommend a 3rd dose for all immunocompromised

individuals. The characteristics of this cohort are described in Table 1 according to DMT,

including age, sex, time since last drug administration, and vaccine type (primarily the mRNA

vaccines BNT162b2 (Comirnaty, Pfizer-BioNTech) and mRNA-1273 (Spikevax, Moderna);

see Methods for further details). Breakthrough COVID-19 infections >14 days after

vaccination are shown for each group (n=29 across all DMTs and healthy controls).

Infections were predominantly contracted between November 2021-February 2022,

representing a mixture of Delta and Omicron VOC infections. None of the individuals were

hospitalised or died.

DMTs vary based on mechanism of action and cellular target. We therefore assessed the

effect of each DMT on the lymphocyte, CD4 and CD8 T cell frequency and function by flow

cytometry (Figure 1). The full flow cytometry gating strategy is shown in Supplementary

Figure 2. Compared to healthy controls, pwMS on DMTs did not show altered frequencies of

CD3+ lymphocytes except for fingolimod-treated individuals, who had significant reductions

in the CD3+ cell populations (Figure 1A) (proportion of CD3+ live lymphocytes in fingolimod-

treated: median, 46.9%; IQR, 44.9%; healthy controls: median, 66.8%; IQR, 11%).

Additionally, fingolimod-treated individuals had sharply reduced CD4+ T cell populations

(fingolimod-treated: median, 19.2%; IQR, 26.0%; healthy controls: median, 63.1%; IQR,

14.1%) and a concomitant increase in CD8+ T cells (Figure 1B and Supplementary Figure 3).

The graphs show frequencies from the post-vaccination time point; however, the cell

population frequencies for individuals were consistent before and after vaccination. The

CD4:CD8 T cell ratio did not correlate with antibody responses in fingolimod-treated

individuals, and people with less distorted ratios of T cells did not have improved antibody

titres, which were low throughout the group (data not shown).

Antibody and T cell responses were measured to assess the immune response to two doses

of vaccine. Samples taken 3 weeks (median 20.5 days) after the 2nd vaccine dose were

assayed for antibody binding activity (Figure 1C). IgG anti-spike RBD responses were

classified as negative (<5 BAU/ml), very weak positive (5-20 BAU/ml), weak positive (20-200

BAU/ml), and positive (>200 BAU/ml) and are indicated on the graph for reference. All

healthy controls and individuals treated with alemtuzumab, cladribine and natalizumab had

strong antibody titres, predominantly in the 'positive' range (median per group: healthy

control, 1166 BAU/ml; alemtuzumab, 5591 BAU/ml; cladribine, 3081 BAU/ml; natalizumab,

3625 BAU/ml). However, individuals treated with fingolimod or rituximab had poor antibody

responses after two vaccine doses (median: fingolimod, 2.5 BAU/ml; rituximab, 0.5 BAU/ml

(below the level of detection for this assay)).

T cell responses were assessed using activation-induced marker (AIM) assays and

measured by flow cytometry (see Supplementary Figure. 2 for gating). Peripheral blood

mononuclear cells (PBMCs) were stimulated with SARS-CoV-2 spike peptides and CD4 T

cell activation was measured by CD40L and TNF-α coexpression (Figure 1D) before (V0,

baseline) and 2 weeks after (V2) vaccination. Samples were taken from the same individuals

at both time points wherever possible, indicated by paired dots. There was a significant

increase in the spike-specific CD4 T cell response after vaccination in the healthy controls

and alemtuzumab-treated patients. This suggested that most of the alemtuzumab-treated

pwMS had reconstituted their immune system within the time since last treatment, which was

more than three years for most patients. Responses were highly heterogeneous and did not

reach statistical significance in the other DMT groups. More than half (10/18) of the

fingolimod-treated group had too few CD4 T cells in our assay to accurately measure

activation responses and were excluded from the analysis as we had too few CD4+ events to

calculate the percent response. CD8 T cell responses (Figure 1E) producing IFN-γ and TNF-

α varied between individuals. Of interest, IFN-γ+ and TNF-α+ spike-specific CD8+ T cell

responses from fingolimod-treated individuals negatively correlated with higher proportions of

CD8+ T cells (i.e. individuals with more skewed CD4/CD8 ratios also had fewer cytokine

producing CD8 T cells), although this did not reach statistical significance (Supplementary

Figure 3I).

As the fingolimod- and rituximab-treated individuals had poor antibody responses, these

patients received a third dose of vaccine (see Methods) (Figure 2). Individuals treated with

rituximab did not show significant improvements in IgG anti-spike RBD after a third vaccine

dose (Figure 2A), although the overall responses and number of responders increased

(median and IQR at V2, 1.3 and 18.3 BAU/ml; at V3, 2.0 and 665.5 BAU/ml; 13/43 (30.2%)

individuals had >5 BAU/ml titres after 2 vaccine doses, increasing to 27/61 (44.3%) after 3

vaccine doses), suggesting that some, though not all, individuals improved their antibody

responses after repeated vaccination. However, the number of patients that were positive

(>200 BAU/ml) was significantly increased from 5/43 after 2 doses to 19/61 after 3 doses

(two-tailed p=0.0320, Fisher's exact test). Moreover, a significant proportion of individuals

had detectable spike-specific CD4 and CD8 T cell responses (Figure 2B-C), demonstrating

that rituximab treatment does not inhibit T cell responses to the same extent as antibody

responses.

The same effect was not seen for people treated with fingolimod. After a third vaccine dose,

fingolimod-treated patients showed no significant increase in IgG anti-spike RBD (Figure 2D),

and generally had even lower antibody responses than the rituximab-treated group, with no

patients reaching the 'positive' response classification of >200 BAU/ml (median and IQR at

V2, 2.1 and 5.35 BAU/ml; at V3, 8.0 and 33.5 BAU/ml). Despite an increase in weak

responders (20-200 BAU/ml), this was not statistically significant (1/13 individuals had weak

responses after V2 compared to 7/21 after V3; two-tailed p=0.1164, Fisher's exact test).

Spike-specific CD4 and CD8 responses also showed no significant response, suggesting

that fingolimod has a major impact on measurable T cell responses in blood as well as

antibody levels. We did see a small but statistically significant CD8 T cell response

(p=0.0429) to influenza (flu) peptides compared to the unstimulated control, suggesting that

existing T cell responses, possibly generated prior to beginning fingolimod treatment, are

maintained over time.

61.4% (51/83) of fingolimod- or rituximab-treated patients received an influenza vaccine

between September 2020 and February 2021. However, influenza-specific T cell responses

did not significantly differ between individuals who had received a seasonal influenza

vaccination during the previous winter (2020-21) and those who had not. This suggested that

T cell responses generated via previous vaccination or influenza infections prior to the

COVID-19 pandemic were still detectable in these patients. There was a weak but significant

positive correlation between CD4 responses to SARS-CoV-2 spike and CD4 responses to

influenza peptides in rituximab-treated patients, suggesting individual differences to vaccine

antigens in general (Supplementary Figure 4A). T cell responses to EBV and CMV peptides

were higher than responses to the vaccine peptides, which represents the difference

between vaccination and latent viral infection. In rituximab-treated individuals we saw strong

CMV-specific CD4 and CD8 T cell responses (p=0.0005 and p<0.0001 respectively,

Wilcoxon tests) (Figure 2B) and EBV-specific CD8 T cell responses (p<0.0001) (Figure 2C).

However, there was no correlation between CD4 T cell responses to spike and CMV in

rituximab-treated patients, (Supplementary Figure 4B), CD4 and CD8 T cell responses to

spike (Supplementary Figure 4C), or CD4 responses and antibody responses

(Supplementary Figure 4D), consistent with other studies showing low concordance between

these measures of immune responsiveness (27).

The administration interval of DMTs varies by drug, as described in Table 1. In the course of

this study, patients taking rituximab received treatment according to their individual

schedules. All patients received rituximab prior to the baseline (V0) sample and completed

the initial two-dose vaccine course without further rituximab infusions. Between the second

and third vaccine doses, approximately half (30/62) of the patients received another dose of

rituximab (median time 8.43 weeks before V3, range 1.86-19.7 weeks). We hypothesised

that this rituximab dosage impaired the ability to respond to vaccination. Antibody and T cell

responses in these two groups were therefore compared (Figure 3). There was no significant

difference in IgG anti-spike RBD between these two groups after the second vaccine dose

(V2), but after a third vaccine dose (V3) individuals who had recently received rituximab had

significantly lower antibody activity (Figure 3A) than those who had not (p=0.023, unpaired t

test). However, there was no such difference between the T cell responses of the two groups

(Figure 3B-C). Comparing the spike-specific responses between groups showed no

difference in CD4 (p=0.998, unpaired t test) or CD8 T cell activation (p=0.545), suggesting

that T cell responses are not affected by re-administration of rituximab after the primary

vaccine course.

Finally, the question of whether vaccination confers protection against SARS-CoV-2 variants

of concern (VOC) has been of particular concern since the initial emergence of the Alpha

(B.1.1.7) variant and subsequent Delta (B.1.617.2) and Omicron (B.1.1.529/BA.1-5) variants.

Mutations in the spike region of these variants are thought to reduce the ability of vaccine-

generated antibodies to recognise these variants and potentially to reduce protection against

them. To measure how T cell responses from vaccination were affected, we assessed CD4

and CD8 T cell responses to the mutated peptides of these three variants (Figure 4). PBMCs

from triple-vaccinated rituximab-treated patients were stimulated as before with only the

mutated peptide regions from the Alpha, Delta, or Omicron variants, as well as the

homologous peptides for each variant from the original Wuhan-Hu-1 sequence). The location

and number of mutated peptides (34, 32 and 83 peptides for Alpha, Delta, and Omicron

respectively) are shown in Figure 4A. There were no significant differences in CD4 T cell

responses to the Alpha variant compared to the homologous Wuhan-Hu-1 sequence, but

significantly reduced responsiveness to the mutated peptides of both the Delta (p=0.047,

Wilcoxon t test) and Omicron variants (p=0.0028) (Figure 4B). Although CD8 T cell

responses were reduced, particularly for the Delta VOC, these differences did not reach

significance (Figure 4C).

Discussion

Older and immunocompromised individuals are particularly at risk of severe COVID-19

disease. Vaccine efficacy in immunocompromised individuals is therefore important to

understand, particularly as many countries including Norway (35) have achieved high

vaccine coverage and have since lifted many or all infection-limiting measures such as social

distancing. However, SARS-CoV-2 variants continue to circulate, and vulnerable groups may

still be at risk of severe disease. These data show that pwMS treated with alemtuzumab,

cladribine, and natalizumab have robust humoral and CD4 and CD8 T cell responses after

two vaccine doses, in agreement with other studies (3, 14, 20, 22, 25). However, individuals

treated with fingolimod and rituximab have strongly reduced antibody responses compared

with both healthy controls and pwMS taking other DMTs. Upon receipt of a third vaccine

dose, both treatment groups showed small increases in IgG anti-spike levels and a

significantly increased percentage of patients developed high responses (>200 BAU/ml) in

the rituximab treated group, demonstrating that some individuals were capable of increasing

B cell responses. This finding was also found in a larger study where improved IgG anti-spike

responses were found after third vaccination (29). Moreover, triple-vaccinated rituximab-

treated individuals demonstrated both CD4 and CD8 T cell responses against SARS-CoV-2

spike peptides. These T cell responses were not reduced even when individuals received

rituximab between their second and third vaccine doses, suggesting that although re-

administration of anti-CD20 drugs does impair humoral responses, cellular responses are

preserved.

Additionally, we observed strong CD4 and CD8 T cell responses to the herpesviruses CMV

and EBV in pwMS, suggesting that specific T cell responses against antigens from long-term

latent infections are present. T cell activation against other vaccine antigens such as

influenza were comparable to the SARS-CoV-2 spike-specific responses, and rituximab-

treated individuals showed a positive correlation between spike-specific and flu-specific

responses. This suggests that vaccine responsiveness varies by individual but is not

necessarily associated with T cell responses to other infections such as CMV. Immune

responses may also differ based on whether individuals were already using DMTs at the time

of antigen exposure, which may affect the magnitude of the immune response, as well as the

availability and duration of antigen seen during vaccination or acute infection compared to

chronic infections.

In pwMS treated with fingolimod, a third vaccine dose did not appear to improve either the

antibody or T cell responses. As fingolimod is taken daily, the fluctuations in B cell counts

seen in individuals taking anti-CD20 drugs are not seen (36). The reduction in peripheral

lymphocyte and CD4 T cell counts we observed here was consistent in individuals at

different sampling points, suggesting that the administration of fingolimod causes lymphocyte

sequestration to different extents for each individual. Although antibody titres were strongly

reduced for all fingolimod-treated patients, we observed that individuals with less skewed

CD4:CD8 T cell ratios had stronger spike-specific CD8 T cell responses, suggesting that

people with higher circulating CD4 T cell frequencies are more likely to generate measurable

and potentially protective cellular responses.

Nevertheless, pwMS receiving fingolimod do not appear to be at higher risk of severe

COVID-19 or hospitalisation than the general population prior to vaccination (7, 37).

Fingolimod has been found to reduce proinflammatory cytokine release from dendritic cells

and monocytes (38) which may reduce detrimental uncontrolled inflammation associated with

severe COVID-19 disease (39). Additionally, as lymphocytes are sequestered rather than

destroyed by S1PR modulators (40), failure to detect T cell responses in peripheral blood

may not fully reflect the extent of the total T cell response, and non-circulating cellular

responses induced by vaccination may be present in the lymph nodes or other secondary

lymphoid organs.

Several large-scale studies have found that pwMS on fingolimod or ocrelizumab are at higher

risk of SARS-CoV-2 infection after vaccination than the general population or pwMS on other

DMTs (41-43), possibly reflecting the role of circulating antibodies in preventing infection.

However, the severity of these infections is still unclear, and where cases could be followed

on an individual level, there were no deaths from COVID-19 (42, 43). Further research has

found that, even after vaccination, pwMS on anti-CD20 drugs were at higher risk of

hospitalisation but not death; this risk was not seen with other DMTs including S1PR

modulators (44). In our study, 29 individuals across all DMTs contracted SARS-CoV-2 after

vaccination and none of these were hospitalised or died. Further large-scale studies are

required to determine vaccine protection against severe disease and death in pwMS and

particularly patients treated with anti-CD20 drugs and S1PR modulators.

The emergence of SARS-CoV-2 VOC has further complicated the question of vaccine

efficacy and protection. Neutralising antibodies against the Delta and Omicron VOCs have

been found to be sharply reduced compared to the original Wuhan-Hu-1 strain (45), while T

cell responses are more heterogeneous and show wide cross-reactivity to other human

coronaviruses as well as between variants (46-48). In triple-vaccinated rituximab-treated

individuals we found that CD4+ T cell responses to both the Delta and Omicron VOCs were

reduced compared to WT, suggesting that although T cells are responsive to the mutated

VOC regions, vaccine-generated T cell-mediated protection may be reduced. However,

these mutated regions cover only a fraction of the spike peptide sequences and further work

is needed to determine how these mutations affect T cell vaccine responses.

We found no correlation between any combinations of antibody titre, CD4 T cell responses,

or CD8 T cell responses, and therefore using only one of these parameters as an indication

of immune responsiveness cannot give a full picture of vaccine efficacy. Although the

correlates of protection against SARS-CoV-2 infection and severe COVID-19 disease are still

unclear and the relative roles of antibody-mediated virus neutralisation and T cell-dependent

protection are still being extensively studied (49-51), analysis of cellular responses in

addition to antibody titres can give a better understanding of whether immunosuppressed

individuals are likely to require additional protective measures. Further follow up studies are

required to determine whether T cell responses in the absence of antibody titres, such as

seen in our rituximab-treated population, are protective against severe disease, but the

current evidence supports the contention that T cell immunity is sufficient.

One limitation of this work is a lack of longitudinal sampling to measure changes in CD4 and

CD8 T cell responses between the second and third vaccine for the rituximab- and

fingolimod-treated patients. The question of whether repeated vaccination with antigens from

the Wuhan-Hu-1 variant can prevent disease from successive VOC remains to be seen.

In summary, we found that pwMS on DMTs that inhibit antibody responses are still capable

of mounting T cell responses comparable with healthy controls, and furthermore that

continued administration of the widely used anti-CD20 drug rituximab between the primary

vaccine course and subsequent vaccine doses does not impede cellular responses. Further

analyses of the efficacy and durability of cellular responses, and well as the impact of

additional vaccination, are needed to better understand how vaccines protect against severe

disease in immunocompromised individuals.

Methods

Participant recruitment and ethical approvals

All patients from the Norwegian MS registry (n=12000) in 2021 were invited to participate in

the humoral arm of the NevroVax study. A subgroup of patients from Oslo University Hospital

on the DMTs alemtuzumab, cladribine, natalizumab, fingolimod and rituximab (c. n=10 per

DMT) were recruited to provide PBMC samples, along with all patients who lacked antibody

responses after 2 vaccine doses (considered at the time to be <70 arbitrary units (AU)/ml by

ELISA). Individuals from Oslo University Hospital, Akershus University Hospital and

Haukeland University hospital with low humoral responses subsequently received a third

vaccine dose and those treated at Oslo University Hospital or Akershus University Hospital

comprised the fingolimod- and rituximab-treated individuals at V3. Healthy controls were

recruited among healthcare workers from Diakonhjemmet Hospital and Akershus University

Hospital and samples were stored in the Oslo University Hospital biobank.

<u>Vaccination and inclusion in vaccination trial</u>

PwMS were vaccinated as per guidelines of the Norwegian Corona Vaccination Program

where immunocompromised individuals and healthcare workers (who participated here as

healthy controls) were high priority. Vaccines were administered according to the

manufacturers' recommendations and health administration advice at the time, ranging from

three weeks between first and second doses for mRNA-1273 and 6-10 weeks for BNT162b2.

Some individuals received first doses of ChAdOx1-S (Vaxzevria, AstraZeneca), the

distribution of which was subsequently discontinued in Norway in March 2021, and received

second doses of BNT162b2. Individuals who had a COVID-19 infection before or during the

course of vaccination were excluded from further analyses. Individuals who failed to

seroconvert to IgG anti-spike (RBD) after the standard two doses were invited to participate

in a vaccination trial to receive a third dose of mRNA-1273 or BNT162b2 outside the

framework of the Norwegian Corona Vaccination Program (EudraCT Number: 2021-003618-

37). Further patients included in this study after 1st September 2021 received third dose

vaccines following revised guidelines in the Norwegian Corona Vaccination Program (where

all immunocompromised adults were advised to receive a third dose).

<u>Sample collection</u>

Venous blood for PBMC isolation was collected at Oslo University Hospital into BD

Vacutainer CPT tubes with sodium citrate. Tubes were centrifuged for 20 minutes at 1600xg

to isolate PBMCs, which were then pipetted into fresh tubes, washed twice with RPMI, and

frozen in 90% foetal calf serum (FCS) (Gibco, Thermo Fisher) with 10% dimethyl sulfoxide

(DMSO, Sigma Aldrich) in liquid nitrogen for future use.

<u>T cell stimulation and flow cytometry</u>

Cryopreserved PBMCs were thawed in RPMI, washed thrice to remove residual DMSO, and

counted. Cells were plated into 96-well U-bottomed plates at 200,000 cells per well and

stimulated for 24 hours in RPMI culture media containing 10% FCS, 1mM sodium pyruvate

(Gibco, Thermo Fisher), 1x MEM NEAA (Gibco), 50nM 1-thioglycerol and 12ug/ml

gensumycin. GolgiPlug (BD Biosciences) containing brefeldin A was added after 2 hours of

stimulation until the end of the incubation. Cells were stimulated with peptide pools:

PepTivator SARS-CoV-2 Prot_S covering the immunodominant sequence domains of the

spike glycoprotein from the SARS-CoV-2 (Wuhan-Hu-1 variant), EBV Consensus, and CMV

pp65 pool (used according to the manufacturer's recommendations at 0.75nmol/ml, all

Miltenyi Biotec) and pooled pan-influenza peptides for HLA class I and II (final concentration

1µg/ml) (GenScript). Peptide pools for mutated SARS CoV-2 Spike are outlined in the next

paragraph. Cytostim (Miltenyi Biotec) was used as a positive control according to the

manufacturer's recommendation.

After 24 hours, cells were centrifuged at 500xg for 5 minutes, the supernatant discarded, and

cells resuspended in FACS buffer (1% FCS in PBS). Cells were centrifuged again and the

supernatant removed. Cells were incubated with 10µl surface antibody cocktail (anti-human

CD3-BV605 (clone SK7) (BD Biosciences), CD4-eFluor 450 (OKT-4), CD8-AF488 (OKT-8),

and Fixable Live/Dead Near-IR (1:1000 dilution) (all ThermoFisher)) for 30 minutes at 4°C,

washed in FACS buffer, then fixed in Fix/Perm (BD Biosciences) for 20 minutes at room

temperature. Cells were then washed twice with PermWash (BD Biosciences) and incubated

with 10µl intracellular antibody cocktail (anti-human IFN-γ-BV711 (clone 4S.B3), CD40L-

BV510 (24-31) (both BioLegend), TNF-α-PE (Mab11), CD69-APC (FN50) (both BD

Biosciences)) for 30 minutes at room temperature. Cells were finally washed with PermWash

and resuspended in 200ul FACS buffer for analysis by flow cytometry within 24 hours.

Cells were acquired on a BioRad ZE5 flow cytometer and analysed with FlowJo™ v.10.7

Software (BD Life Sciences).

<u>Variants of Concern and mutated peptide sequences</u>

Three PepTivator SARS-CoV-2 VOC spike protein Mutation Pools and the three

corresponding spike protein WT Reference Pools (all Miltenyi Biotec) were used at a final

concentration of 0.75 nmol/ml per the manufacturer's recommendation. Prot_S B.1.1.7

Mutation Pool (cat. no. 130-127-844) included 34 peptides from 10 mutations: deletion 69,

deletion 70, deletion 144, N501Y, A570D, D614G, P681H, T716I, S982A, D1118H. The

corresponding non-mutated peptide pool control was Prot_S B.1.1.7 WT Reference Pool

(cat. no. 130-127-841).

Prot_S B.1.617.2 Mutation Pool (cat. no. 130-128-763) included 32 peptides from 10

mutations: T19R, G142D, E156G, deletion 157, deletion 158, L452R, T478K, D614G,

P681R, and D950N. This subvariant lacks the E484Q mutation. The non-mutated peptide

pool control was Prot_S B.1.617.2 WT Reference Pool (cat. no. 130-128-761).

Prot_S B.1.1.529/BA.1 Mutation Pool (cat. no. 130-129-928) included 83 peptides from 37

mutations: A67V, H69 deletion, V70 deletion, T95I, G142D, V143 deletion, Y144 deletion,

Y145 deletion, N211 deletion, L212I, insertion 214EPE, G339D, S371L, S373P, S375F,

K417N, N440K, G446S, S477N, T478K, E484A, Q493R, G496S, Q498R, N501Y, Y505H,

T547K, D614G, H655Y, N679K, P681H, N764K, D796Y, N856K, Q954H, N969K, L981F.

The non-mutated peptide pool control was Prot_S B.1.1.529/BA.1 WT Reference Pool (cat.

no. 130-129-927).

<u>Antibody quantification</u>

Semiquantitative measurement of antibodies to full-length spike protein (Spike-FL) and the

receptor-binding domain (RBD) from SARS-CoV-2 was performed using a multiplexed bead-

based assay as described in (52). Polymer beads with fluorescent barcodes were coupled to

successively to neutravidin (ThermoFisher) and biotinylated viral antigens to generate bead-

based protein arrays. Sera were diluted 1:100 in assay buffer (PBS, 1% Tween-20, 10ug/ml

D-biotin, 10 µg/ml neutravidin, 0.1% sodium azide). Diluted sera were incubated with bead-

based arrays in 384 well plates for 30 minutes at 22°C at constant agitation, washed three

times in PBS/1% Tween-20 (PBT) and labelled with R-Phycoerythrin (R-PE)-conjugated

goat-anti-human IgG (Jackson Immunoresearch). For measurement of neutralizing

antibodies, the beads were pelleted after incubation with serum and labelled successively

with digoxigenin-conjugated human ACE2 and mouse monoclonal anti-dixogigenin (Jackson

Immunoresearch), which was conjugated in-house to R-PE. The beads were analyzed with

an AttuneNxT flow cytometer (ThermoFisher), and raw data (fcs.3.1) were analyzed in

WinList 3D (Verity Softwarehouse). The median R-PE fluorescence intensity (MFI) of each

bead subset was exported to Excel. The MFI of beads coupled with viral antigens was

divided by that measured on beads coupled with neutravidin only (relative MFI, rMFI). A total

of 979 pre-pandemic sera and 810 sera from COVID-19 convalescents were analyzed to

establish cutoffs for seropositivity. A double cutoff of rMFI >5 for anti-RBD and anti-Spike FL

yielded a specificity of 99.7% and a sensitivity of 95% (53). Serum from an individual who

had received three doses of the Pfizer/BioNTech anti-COVID-19 vaccine was used as

standard to convert signals to binding antibody units per milliliter (BAU/ml).

<u>Statistics and analysis</u>

Statistical analyses were performed using GraphPad Prism v.9 for Windows, GraphPad

Software. Two-tailed p values are shown. For analysis of functional markers (CD40L+ TNF-

α+ CD4 T cells and IFN-γ+ TNF-α+ CD8 T cells), data from FACS plots with fewer than 1000

CD4 or CD8 T cells were excluded.

<u>Study approval</u>

The study was approved by the Norwegian South-Eastern Regional Ethical Committee

(Reference numbers 200631, 235424, 135924, and 204104), and the Norwegian Medicines

Agency (EudraCT Number: 2021- 003618-37). All participants gave written, informed

consent prior to inclusion in this study.

Author contributions

LAM and SM conceived and designed the study. A-SW drafted the paper. AR, MK, LAM, TH,

GON, and SM contributed to drafting the paper. A-SW, AR, GS, IFK, and AC performed

experiments. A-SW, AR, MK, GS, IFK, FL-J, GON, LAM, and SM analyzed and interpreted

data. MHØ, AC, TH, HFH, EH, JTV, and EGC contributed to data analysis and interpretation.

MK, GON, MHØ, SWS, KKJ, JTV, FL-J, and LAM organized the collection of samples and

information from the cohorts. All authors contributed to and approved the final manuscript.

Acknowledgements

The authors would like to thank Ingrid Egner, Katrine Persgård Lund, Viktoriia Chaban, and

Julie Røkke Osen at the Department of Immunology, Oslo University Hospital for biobanking

and technical help. We would also like to thank the participants in the NevroVax study and

staff at Diakonhjemmet Hospital and Akershus University Hospital for donating blood

samples, as well as the staff at Oslo University Hospital for taking the samples. This study

was supported by the Norwegian Institute of Public Health, the Norwegian Ministry of Health

through a program for corona vaccination surveillance, the Odd Fellows Foundation, the

Sanofi Research Fund for MS, the Norwegian MS Society, and the Research Council of

Norway (RCN) Covid (312693), LAM, SM; a KG Jebsen Foundation (grant 19), LAM; the

Coalition for Epidemic Preparedness Innovations (CEPI), LAM, SM, JTV, FL-J; and the

University of Oslo and Oslo University Hospital, LAM, FL-J, JTV.

Bibliography

1.   Brodin P. Immune determinants of COVID-19 disease presentation and severity.

Nature Medicine. 2021;27(1):28-33.

2.   Kelly H, Sokola B, Abboud H. Safety and efficacy of COVID-19 vaccines in multiple

sclerosis patients. Journal of Neuroimmunology. 2021;356:577599.

3.   Hada M, Mosholder AD, Leishear K, Perez-Vilar S. Systematic review of risk of

SARS-CoV-2 infection and severity of COVID-19 with therapies approved to treat multiple

sclerosis. Neurol Sci. 2022:1-11.

4.   Sharifian-Dorche M, Sahraian MA, Fadda G, Osherov M, Sharifian-Dorche A,

Karaminia M, et al. COVID-19 and disease-modifying therapies in patients with

demyelinating diseases of the central nervous system: A systematic review. Multiple

Sclerosis and Related Disorders. 2021;50:102800.

5.   Centonze D, Rocca MA, Gasperini C, Kappos L, Hartung H-P, Magyari M, et al.

Disease-modifying therapies and SARS-CoV-2 vaccination in multiple sclerosis: an expert

consensus. Journal of Neurology. 2021;268(11):3961-8.

6.   Sormani MP, De Rossi N, Schiavetti I, Carmisciano L, Cordioli C, Moiola L, et al.

Disease-Modifying Therapies and Coronavirus Disease 2019 Severity in Multiple Sclerosis.

Annals of Neurology. 2021;89(4):780-9.

7.   Sormani MP, Salvetti M, Labauge P, Schiavetti I, Zephir H, Carmisciano L, et al.

DMTs and Covid-19 severity in MS: a pooled analysis from Italy and France. Annals of

Clinical and Translational Neurology. 2021;8(8):1738-44.

8.   Salter A, Fox RJ, Newsome SD, Halper J, Li DKB, Kanellis P, et al. Outcomes and

Risk Factors Associated With SARS-CoV-2 Infection in a North American Registry of

Patients With Multiple Sclerosis. JAMA Neurology. 2021;78(6):699-708.

9.   Luna G, Alping P, Burman J, Fink K, Fogdell-Hahn A, Gunnarsson M, et al. Infection

Risks Among Patients With Multiple Sclerosis Treated With Fingolimod, Natalizumab,

Rituximab, and Injectable Therapies. JAMA Neurology. 2020;77(2):184-91.

10.   Salzer J, Svenningsson R, Alping P, Novakova L, Björck A, Fink K, et al. Rituximab in

multiple sclerosis. A retrospective observational study on safety and efficacy.

2016;87(20):2074-81.

11.   Zhao Z, Ma C-L, Gu Z-C, Dong Y, Lv Y, Zhong M-K. Incidence and Risk of Infection

Associated With Fingolimod in Patients With Multiple Sclerosis: A Systematic Review and

Meta-Analysis of 8,448 Patients From 12 Randomized Controlled Trials. Frontiers in

Immunology. 2021;12.

12.   Ghadiri F, Sahraian MA, Azimi A, Moghadasi AN. The study of COVID-19 infection

following vaccination in patients with multiple sclerosis. Mult Scler Relat Disord.

2022;57:103363.

13.   Dreyer-Alster S, Menascu S, Mandel M, Shirbint E, Magalashvili D, Dolev M, et al.

COVID-19 vaccination in patients with multiple sclerosis: Safety and humoral efficacy of the

third booster dose. J Neurol Sci. 2022;434:120155.

14.   Tallantyre EC, Vickaryous N, Anderson V, Asardag AN, Baker D, Bestwick J, et al.

COVID-19 Vaccine Response in People with Multiple Sclerosis. Ann Neurol. 2022;91(1):89-

100.

15.   König M, Lorentzen ÅR, Torgauten HM, Tran TT, Schikora-Rustad S, Vaage EB, et

al. Humoral immunity to SARS-CoV-2 mRNA vaccination in multiple sclerosis: the relevance

of time since last rituximab infusion and first experience from sporadic revaccinations.

Journal of Neurology, Neurosurgery & Psychiatry. 2021:jnnp-2021-327612.

16.   Katz JD, Bouley AJ, Jungquist RM, Douglas EA, O'Shea IL, Lathi ES. Humoral and T-

cell responses to SARS-CoV-2 vaccination in multiple sclerosis patients treated with

ocrelizumab. Mult Scler Relat Disord. 2022;57:103382.

17.   Apostolidis SA, Kakara M, Painter MM, Goel RR, Mathew D, Lenzi K, et al. Cellular

and humoral immune responses following SARS-CoV-2 mRNA vaccination in patients with

multiple sclerosis on anti-CD20 therapy. Nature Medicine. 2021;27(11):1990-2001.

18.   Bar-Or A, Calkwood JC, Chognot C, Evershed J, Fox EJ, Herman A, et al. Effect of

ocrelizumab on vaccine responses in patients with multiple sclerosis. The VELOCE study.

2020;95(14):e1999-e2008.

19.   Türkoğlu R, Baliç N, Kızılay T, Erol R, Akbayır E, Yılmaz V, et al. Fingolimod impairs

inactivated vaccine (CoronaVac)-induced antibody response to SARS-CoV-2 spike protein in

persons with multiple sclerosis. Mult Scler Relat Disord. 2022;58:103524.

20.   Altieri M, Capuano R, Conte M, Donnarumma G, Grimaldi E, Coppola N, et al. Six-

month humoral response to BNT162b2 mRNA COVID-19 vaccine in people with multiple

sclerosis treated with natalizumab. Neurol Sci. 2022.

21.   Iannetta M, Landi D, Cola G, Campogiani L, Malagnino V, Teti E, et al. B- and T-Cell

Responses After SARS-CoV-2 Vaccination in Patients With Multiple Sclerosis Receiving

Disease Modifying Therapies: Immunological Patterns and Clinical Implications. Front

Immunol. 2021;12:796482.

22.   Brill L, Rechtman A, Zveik O, Haham N, Levin N, Shifrin A, et al. Effect of cladribine

on COVID-19 serology responses following two doses of the BNT162b2 mRNA vaccine in

patients with multiple sclerosis. Mult Scler Relat Disord. 2022;57:103343.

23.   Ciampi E, Uribe-San-Martin R, Soler B, García L, Guzman J, Pelayo C, et al. Safety

and humoral response rate of inactivated and mRNA vaccines against SARS-CoV-2 in

patients with Multiple Sclerosis. Multiple Sclerosis and Related Disorders. 2022;59:103690.

24.   Habek M, Željko C, Savić Mlakar A, Bendelja K, Rogić D, Adamec I, et al. Humoral

and cellular immunity in convalescent and vaccinated COVID-19 people with multiple

sclerosis: Effects of disease modifying therapies. Mult Scler Relat Disord. 2022;59:103682.

25.   Sabatino JJ, Jr., Mittl K, Rowles WM, McPolin K, Rajan JV, Laurie MT, et al. Multiple

sclerosis therapies differentially impact SARS-CoV-2 vaccine-induced antibody and T cell

immunity and function. JCI Insight. 2022.

26.   Tortorella C, Aiello A, Gasperini C, Agrati C, Castilletti C, Ruggieri S, et al. Humoral-

and T-Cell–Specific Immune Responses to SARS-CoV-2 mRNA Vaccination in Patients With

MS Using Different Disease-Modifying Therapies. Neurology. 2022;98(5):e541-e54.

27.   Gadani SP, Reyes-Mantilla M, Jank L, Harris S, Douglas M, Smith MD, et al.

Discordant humoral and T cell immune responses to SARS-CoV-2 vaccination in people with

multiple sclerosis on anti-CD20 therapy. EBioMedicine. 2021;73:103636.

28.   Bajwa HM, Novak F, Nilsson AC, Nielsen C, Holm DK, Østergaard K, et al.

Persistently reduced humoral and sustained cellular immune response from first to third

SARS-CoV-2 mRNA vaccination in anti-CD20-treated multiple sclerosis patients. Mult Scler

Relat Disord. 2022;60:103729.

29.   König M, Torgauten HM, Tran TT, Holmøy T, Vaage JT, Lund-Johansen F, et al.

Immunogenicity and Safety of a Third SARS-CoV-2 Vaccine Dose in Patients With Multiple

Sclerosis and Weak Immune Response After COVID-19 Vaccination. JAMA Neurol. 2022.

30.   Brill L, Raposo C, Rechtman A, Zveik O, Levin N, Oiknine-Djian E, et al. Severe

Acute Respiratory Syndrome Coronavirus 2 Third Vaccine Immune Response in Multiple

Sclerosis Patients Treated with Ocrelizumab. Annals of Neurology. 2022;91(6):796-800.

31.   Disanto G, Sacco R, Bernasconi E, Martinetti G, Keller F, Gobbi C, et al. Association

of disease-modifying treatment and anti-CD20 infusion timing with humoral response to 2

SARS-CoV-2 vaccines in patients with multiple sclerosis. JAMA neurology.

2021;78(12):1529-31.

32.   Brill L, Rechtman A, Zveik O, Haham N, Oiknine-Djian E, Wolf DG, et al. Humoral and

T-Cell Response to SARS-CoV-2 Vaccination in Patients With Multiple Sclerosis Treated

With Ocrelizumab. JAMA Neurology. 2021;78(12):1510-4.

33.   Brown JWL, Coles A, Horakova D, Havrdova E, Izquierdo G, Prat A, et al. Association

of Initial Disease-Modifying Therapy With Later Conversion to Secondary Progressive

Multiple Sclerosis. JAMA. 2019;321(2):175-87.

34.   Spelman T, Magyari M, Piehl F, Svenningsson A, Rasmussen PV, Kant M, et al.

Treatment Escalation vs Immediate Initiation of Highly Effective Treatment for Patients With

Relapsing-Remitting Multiple Sclerosis: Data From 2 Different National Strategies. JAMA

Neurology. 2021;78(10):1197-204.

35.   Norwegian Institute of Public Health. Coronavirus vaccination - statistics. 2022

[updated 01/08/2022. Available from: https://www.fhi.no/en/id/vaccines/coronavirus-

immunisation-programme/coronavirus-vaccination---statistic/.

36. Francis G, Kappos L, O'Connor P, Collins W, Tang D, Mercier F, et al. Temporal

profile of lymphocyte counts and relationship with infections with fingolimod therapy. Multiple

Sclerosis Journal. 2014;20(4):471-80.

37. Sullivan R, Kilaru A, Hemmer B, Campbell Cree BA, Greenberg BM, Kundu U, et al.

COVID-19 Infection in Fingolimod- or Siponimod-Treated Patients. Case Series.

2022;9(1):e1092.

38. Thomas K, Sehr T, Proschmann U, Rodriguez-Leal FA, Haase R, Ziemssen T.

Fingolimod additionally acts as immunomodulator focused on the innate immune system

beyond its prominent effects on lymphocyte recirculation. J Neuroinflammation.

2017;14(1):41.

39. Leon J, Michelson DA, Olejnik J, Chowdhary K, Oh HS, Hume AJ, et al. A virus-

specific monocyte inflammatory phenotype is induced by SARS-CoV-2 at the

immune–epithelial interface. Proceedings of the National Academy of Sciences.

2022;119(1):e2116853118.

40. McGinley MP, Cohen JA. Sphingosine 1-phosphate receptor modulators in multiple

sclerosis and other conditions. The Lancet. 2021;398(10306):1184-94.

41. Garjani A, Patel S, Bharkhada D, Rashid W, Coles A, Law GR, et al. Impact of mass

vaccination on SARS-CoV-2 infections among multiple sclerosis patients taking

immunomodulatory disease-modifying therapies in England. Multiple Sclerosis and Related

Disorders. 2022;57:103458.

42. Schiavetti I, Cordioli C, Stromillo ML, Teresa Ferrò M, Laroni A, Cocco E, et al.

Breakthrough SARS-CoV-2 infections in MS patients on disease-modifying therapies.

Multiple Sclerosis Journal. 2022;0(0):13524585221102918.

43. Sormani MP, Schiavetti I, Inglese M, Carmisciano L, Laroni A, Lapucci C, et al.

Breakthrough SARS-CoV-2 infections after COVID-19 mRNA vaccination in MS patients on

disease modifying therapies during the Delta and the Omicron waves in Italy. eBioMedicine.

2022;80:104042.

44.   Bsteh G, Gradl C, Heschl B, Hegen H, Di Pauli F, Assar H, et al. Impact of

vaccination on COVID-19 outcome in multiple sclerosis. European Journal of Neurology.

2022;n/a(n/a).

45.   Servellita V, Syed AM, Morris MK, Brazer N, Saldhi P, Garcia-Knight M, et al.

Neutralizing immunity in vaccine breakthrough infections from the SARS-CoV-2 Omicron and

Delta variants. Cell. 2022;185(9):1539-48.e5.

46.   Yu ED, Wang E, Garrigan E, Goodwin B, Sutherland A, Tarke A, et al. Development

of a T cell-based immunodiagnostic system to effectively distinguish SARS-CoV-2 infection

and COVID-19 vaccination status. Cell Host & Microbe. 2022;30(3):388-99.e3.

47.   Liu J, Chandrashekar A, Sellers D, Barrett J, Jacob-Dolan C, Lifton M, et al. Vaccines

elicit highly conserved cellular immunity to SARS-CoV-2 Omicron. Nature.

2022;603(7901):493-6.

48.   Swadling L, Diniz MO, Schmidt NM, Amin OE, Chandran A, Shaw E, et al. Pre-

existing polymerase-specific T cells expand in abortive seronegative SARS-CoV-2. Nature.

2022;601(7891):110-7.

49.   Earle KA, Ambrosino DM, Fiore-Gartland A, Goldblatt D, Gilbert PB, Siber GR, et al.

Evidence for antibody as a protective correlate for COVID-19 vaccines. Vaccine.

2021;39(32):4423-8.

50.   Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, et al.

Neutralizing antibody levels are highly predictive of immune protection from symptomatic

SARS-CoV-2 infection. Nature Medicine. 2021;27(7):1205-11.

51.   McMahan K, Yu J, Mercado NB, Loos C, Tostanoski LH, Chandrashekar A, et al.

Correlates of protection against SARS-CoV-2 in rhesus macaques. Nature.

2021;590(7847):630-4.

52.   Tran TT, Vaage E, Mehta A, Chopra A, Kolderup A, Anthi A, et al. Multiplexed

measurement of binding-and neutralizing antibodies to SARS- CoV-2 variants in 12.000 post-

vaccine sera2022.

53.   Jyssum I, Kared H, Tran TT, Tveter AT, Provan SA, Sexton J, et al. Humoral and

cellular immune responses to two and three doses of SARS-CoV-2 vaccines in rituximab-

treated patients with rheumatoid arthritis: a prospective, cohort study. Lancet Rheumatol.

2022;4(3):e177-e87.

54.   Huang Y, Yang C, Xu X-f, Xu W, Liu S-w. Structural and functional properties of

SARS-CoV-2 spike protein: potential antivirus drug development for COVID-19. Acta

Pharmacologica Sinica. 2020;41(9):1141-9.

Figure legends

Table 1: Summary of participant characteristics. Participants are grouped by DMT. V0

indicates pre-vaccination baseline samples, V2 indicates samples taken after receiving 2

doses of vaccine, and V3 indicates samples taken after 3 doses of vaccine. Numbers of

individuals per DMT group, age, sex, time since last drug treatment, vaccine types, time

between sampling and the corresponding vaccine dose, and the number of participants in

each group who were subsequently infected with COVID-19 >2 weeks post-vaccination.

Figure 1: Lymphocyte proportions in peripheral blood and spike-specific vaccination

responses in pwMS on DMTs. Individuals are grouped by DMT (healthy controls (HC),

alemtuzumab (ALEM)-, cladribine (CLAD)-, natalizumab (NTZ)-, fingolimod (FIN)- and

rituximab (RTX)-treated MS patients). (A) CD3+ lymphocyte proportions and (B) the ratio of

CD4:CD8 T cells in different DMT groups showed reduced frequencies of CD3+ lymphocytes

and CD4+ T cells in fingolimod-treated patients. Violin plots show individuals as separate

points, lines indicate median, IQR, and min and max. Mann-Whitney test comparing drug-

treated groups with healthy controls, two-tailed p values were calculated, **** $p<0.0001$. (C)

Binding antibody units after 2 doses of vaccine. Responses below the lower limit of detection

are shown as 0.5 BAU/ml; titres <5 BAU/ml are considered negative, 5-20 BAU/ml as very

weak positives, 20-200 BAU/ml as weak positives and >200 BAU/ml as positives. (D) CD4 T

cell (CD40L+ TNF-α+) and (E) CD8 T cell responses (IFN-γ+ and/or TNF-α+) to spike

peptides before (V0) and after (V2) 2 doses of vaccine. Responses with 0 events are plotted

at 0.001% to indicate non-responses. Samples from the same individual before and after

vaccination are paired with a line. Patient numbers for each group are indicated along the x-

axis. Individuals with <1000 CD4 or CD8 T cells acquired by FACS were excluded from

further analysis. Statistical comparisons by Wilcoxon two-tailed paired t-tests, * p<0.05.

Figure 2: T cell and antibody responses in rituximab- and fingolimod-treated patients after 3[rd]

vaccine dose. Antibody responses (BAU/ml), CD4+ and CD8+ T cell responses after 3[rd]

vaccine dose in rituximab-treated (A-C) and fingolimod-treated patients (D-F) (A, n=43-61);

B, n=21-56; C, n=21-54; D, n=13-21; E, n=6-13; F, n=6-17). Individuals with <1000 CD4+ T

cells acquired by FACS were excluded from this analysis. Dotted lines in (A) and (D) indicate

classification of antibody responses as negative or positive as described previously. For B-C

and E-F, lines on scatter plots indicate the median. Statistical analyses by Wilcoxon paired t-

tests, two-tailed p values are shown, * p<0.05, *** p<0.001

Figure 3: Re-administration of rituximab between vaccine doses affects antibody but not T

cell responses. Rituximab-treated individuals were grouped by whether or not they received

a dose of RTX between vaccine doses 2 and 3. (A) Antibody titres (BAU/ml) after 2[nd] (V2)

and 3[rd] (V3) vaccine doses for patients who did not receive RTX between vaccines (empty

boxes) (n= 20-32) and patients who did receive RTX between vaccines (grey boxes) (n=23-

30). (B) CD4 T cell and (C) CD8 T cell responses without stimulation (unstim) or to SARS-

CoV-2 spike or CMV peptides after 3[rd] vaccine dose for patients without (n=28) or with RTX

administration (n=28) between vaccine doses, as previously. Statistical analyses for paired

responses by Wilcoxon t test, unpaired responses by Mann-Whitney, two-tailed p values are

shown, * p<0.05, ** p<0.01.

Figure 4: T cell responses to the Delta and Omicron SARS-CoV-2 variants after three

vaccine doses. Schematic of mutated regions in the Alpha, Delta and Omicron regions

stimulated by peptides (A). The SARS-CoV-2 spike protein is 1273 amino acids (aa) long,

consisting of the signal peptide, S1 and S2 subunits; the receptor binding domain (RBD) in

S1 is indicated in red (54). Regions covered by the SARS-CoV-2 Prot_S (WT) peptide used

for AIM assays are shown for reference. The control and mutant peptides for each variant

cover the same loci but with the mutated (mutant) or Wuhan-Hu-1 variant (control). Amino

acid mutations are listed in Methods. PBMCs from rituximab-treated patients after a 3[rd]

vaccine dose were stimulated with spike peptide pools from the mutated regions (blue

circles) of the Alpha (n=29), Delta (n=41) and Omicron VOC (n=21) and the same regions of

the WT sequence (empty circles) and the CD4+ (B) and CD8+ (C) T cell responses were

compared. Statistical differences were calculated by Wilcoxon paired t tests. * p<0.05, **

p<0.01

		Alemtuzumab		Cladribine		Natalizumab		Fingolimod			Rituximab			Healthy controls	
		V0	V2	V0	V2	V0	V2	V0	V2	V3	V0	V2	V3	V0	V2
n		11	11	8	8	10	8	12	18	20	5	13	63	15	15
Previous COVID-19 infection (%)		0 (0)		0 (0)		0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	4 (80.0)	5 (38.5)	3 (4.8)	0	
Age (years)	Median	37.4		42.4		37.0	37.0	41.8	45.3	47.7	29.6	39.1	43.3	46	
	Mean	35.3		41.5		38.1	37.8	41.1	45.1	47.0	31.1	41.1	46.0	46.0	
	Range	22 - 47.5		34.0 - 47.3		26.3 - 51.6	26.3 - 51.6	33.1 - 49.0	33.1 - 60.4	30.5 - 66.0	18.3 - 47.3	18.3 - 76.9	22.1 - 76.9	25.0 - 63.0	
Sex	Female (%)	8 (72.7)		8 (100)		7 (70.0)	6 (75.0)	6 (50.0)	8 (44.4)	11 (55.0)	3 (60.0)	9 (69.2)	53 (84.1)	12 (80.0)	
Time since last drug treatment (weeks)	Median	164.1	165.9	31.3	32.5	0	0.93				5.86	14.5	24.7/8.43 [b]		
	Mean	156.2	157.9	36.2	38.9	-1.7 [a]	1.89				9.37	18.5	28.1/8.31 [b]		
	Range	70.3 - 252.1	71.7 - 252.4	3.0 - 78.4	5.14 - 83.4	-4.57 - 5.14 [a]	0.29 - 6.0		Taken daily		-0.7 - 31.9 [a]	8.43 - 46.0	17.0 - 63.0/1.86 - 19.7 [b]	n/a	
Vaccine type - first dose (n)	BNT162b2		9		6		6		12	14		7	52 [d]		0
	mRNA-1273		2		2		2		6	6		3	8		9
	ChAdOx1-S		0		0		0		0	0		0	3		6
	COVID-19 infection		0		0		0		0	0	4	3 [c]	2 [d]		0
Vaccine type - second dose (n)	BNT162b2		9		6		6		12	14		10	55		6
	mRNA-1273		2		2		2		6	6		3	8		9
Vaccine type - third dose (n)	BNT162b2									6			12		
	mRNA-1273	n/a		n/a		n/a		n/a		14	n/a		51	n/a	
Time between sample and last vaccine dose (days)	Median		11		12		12		13	28.5		38	23		10
	Mean		11.2		19.4		13.9		22.6	29.3		33.5	25.6		10.3
	Range	n/a	8 - 15	n/a	7 - 73	n/a	9 - 26	n/a	7 - 49	20 - 43	n/a	8 - 71	20 - 42	n/a	8 - 16
Covid infection >14 days post-full vaccination (%)		n/a	2 (18.2)	n/a	3 (37.5)	n/a	3 (37.5)	n/a	n/a	2 (10.0)	n/a	n/a	14 (22.2)	n/a	5 (33.3)

[a] Negative time values indicate drug was administered after sample was taken

[b] Individuals are split into groups who last received RTX prior to V0 and those who received RTX between V2 and V3 (see Fig. 3)

[c] Some individuals were infected with COVID-19 at baseline, which was treated as analogous to a first vaccine dose

[d] Some individuals with prior COVID-19 infections still received a 'first' vaccine dose of BNT162b2 and are represented twice