

1 **Substance Use and other Correlates of HIV infection among**
2 **transwomen and men who have sex with men in Perú: Implications**
3 **for Targeted HIV Prevention Strategies for Transwomen**

4
5 Elena Cyrus, PhD, MPH^{1,2}, Javier R. Lama, MD, MPH³, Jorge Sanchez, MD, MPH⁴, Segundo
6 Leon, MPH⁵, Manuel V. Villaran, MD³, MPH, MSc³, Panagiotis Vagenas, PhD, MPH⁶, Daniell
7 S. Sullivan¹, David Vu¹, Makella Coudray PhD¹, Frederick L. Altice, MD, MA²

8
9 ¹University of Central Florida: Population Health Sciences, College of Medicine, Orlando, FL,
10 USA

11 ²Yale School of Public Health, Department of Global Health, New Haven, CT, USA

12 ³Asociación Civil Impacta Salud y Educación (IMPACTA), Lima, Perú

13 ⁴Centro de Investigaciones Tecnológicas, Biomédicas y Medioambientales; Universidad
14 Nacional Mayor de San Marcos, Lima, Peru

15 ⁵San Juan Bautista Privada, Lima, Perú

16 ⁶Berkeley Research Development Office, University of California Berkeley, Berkeley, CA

17 ⁷Univeristy of Central Florida: College of Medicine, Orlando, FL, USA

18
19 [§]Elena Cyrus, PhD, MPH

20 6850 Lake Nona Blvd.

21 Orlando, FL 32827 USA

22 407-266-1506

23 elena.cyrus@ucf.edu

24
25 EC: elena.cyrus@ucf.edu

26 JRL: jrlama@impactaperu.org

27 JS: jsanchez@citbm.pe

28 SL: segundo.leon@upsjb.edu.pe

29 MVV: mvillar2@fredhutch.org

30 PV: pvagenas@berkeley.edu

31 DSS: dannysullivan@knights.ucf.edu

32 DV: dvu140@knights.ucf.edu

33 MC: makella.coudray@ucf.edu

34 FLA: frederick.altice@yale.edu

35

36 Keywords: transwomen, MSM, HIV, risk behavior, substance use, Perú

37

38 Abstract Word Count: 290/350

39 Manuscript Word Count: 2452/3500

40 **Abstract**

41 **Introduction:** Characterization of HIV risk factors among transwomen and men who have sex
42 with men (MSM) should be assessed separately and independently. However, due to several
43 constraints, these populations continue to be conflated in clinical research and data. There are
44 limited datasets globally powered to make such comparisons. The study aimed to use one of the
45 largest surveys of transwomen and MSM in Latin America to determine differences in HIV risk
46 and related correlates between the two populations.

47
48 **Methods:** Secondary data analysis was completed using a cross-sectional biobehavioral survey
49 of 4413 MSM and 714 transwomen living in Perú. Chi Square analysis of selected HIV
50 correlates was conducted to examine differences between transwomen and MSM. Additionally,
51 stratified binary logistic regression was used to split data for further comparative analyses of
52 correlates associated with transwomen and MSM separately.

53
54 **Results:** HIV prevalence among transwomen was two-fold greater than among MSM (14.9% vs.
55 7.0%, $p < 0.001$). Transwomen had a higher prevalence of most HIV risk factors assessed,
56 including presence of alcohol dependence (16.4% vs. 19.0%; $p < .001$) and drug use in the past 3
57 months (17.0% vs. 14.9%). MSM were more likely to use marijuana (68.0% vs. 50.0%, $p < .001$),
58 and transwomen were more likely to engage in inhaled cocaine use (70.0% vs. 51.1%, $p < .001$).
59 The regression exposed differences in correlates driving sub-epidemics in transwomen vs. MSM,
60 with a trend of substance use increasing HIV risk for transwomen only.

61
62 **Conclusions:** Transwomen were more likely to be HIV-infected and had different risk factors
63 from MSM. Targeted prevention strategies are needed for transwomen that are at highest risk.
64 Additionally, further research is needed to determine if these observations in Perú regarding
65 substance use patterns and the role of substance use in HIV risk relate to other trans populations
66 globally.

67

68 **Introduction**

69 Men who have sex with men (MSM) and transwomen in Perú bear a disproportionate
70 burden of human immunodeficiency virus (HIV) infection.^{1,2} In 2016, the HIV seroprevalence
71 among Peruvian MSM and transwomen in Lima was >10% and >20%, respectively, compared to
72 the general Peruvian population prevalence of <1%.³ Similar to other Latin American countries,
73 MSM and transwomen in Perú are socially and medically vulnerable communities, subject to
74 discrimination, social exclusion and insufficient access to preventative and long-term care, partly
75 because of cultural and sexual biases that are pervasive against same-sex relations in the wider
76 Peruvian society.⁴ Additionally, MSM and transwomen are prone to engage in high-risk sexual
77 behaviors, such as transactional or survival sex work and substance use.²

78 While acknowledging these common characteristics among MSM and transwomen, the
79 scope of the problem and risk are not equal for both groups, and transwomen are at greater risk
80 for HIV transmission.^{2,3,5} A combination of social, biological, and structural factors contribute to
81 increased HIV vulnerability in transwomen. Deeply rooted stigma and discrimination in gender
82 expression and gender identity can result in physical and psychological harm towards trans
83 populations that can drive these individuals towards unhealthy coping mechanisms such as
84 substance use.⁶ The lack of a legal identity status for transwomen also limits economic
85 opportunities for these individuals, forcing them to pursue illegal and socially stigmatizing
86 economic activities such as sex work, which could lead to increased risky sexual behaviors and
87 subsequent HIV exposure.⁷ Furthermore, transwomen have been historically neglected in HIV
88 research studies compared to MSM, contributing to the lack of preventative and treatment
89 resources in these communities.⁸ Overall, these evidenced factors and more create increased HIV
90 and substance use rates in transwomen populations compared to MSM and the general
91 population.

92 Evidence of an enhanced disparity between MSM and transwomen can help guide policy
93 or interventional studies that address the specific needs of transwomen. Due to limited resources
94 and competing research priorities, or conflation for convenience, research endeavors often
95 address these groups simultaneously.⁹⁻¹¹ The purpose of this study was to further examine
96 differences in these two groups from Perú's largest biobehavioral surveillance study on
97 correlates and sexual risk behavior among MSM and transwomen and to compare characteristics

98 between them to justify the separation of these groups for singular and independent research
99 efforts.

100

101 **Methods**

102 **Study Design and Data Collection**

103 Between May and October 2011, 5,575 Peruvian MSM and transwomen were recruited
104 using modified snowball methods and peer-educator outreach in previously mapped venues
105 frequented by MSM and transwomen in Lima and four other cities: Ica, Piura, Iquitos, and
106 Pucallpa. Extensive details about recruitment and study procedures have been previously
107 described.^{1,2} Eligibility included individuals 18 years of age or older, being genetically male, and
108 self-reporting at least one male sexual partner in the previous 12 months. The two-step question
109 was used to classify individuals as either transgender or MSM.¹² Of the 5,755 MSM and
110 transwomen recruited, 5,148 completed all study procedures and were eligible for analysis. Of
111 these, 21 did not complete the sexual identity questions regarding gender identification and were
112 thus excluded from the analysis, resulting in a final analytic sample of 5,127.

113 The research project was approved by the Institutional Review Boards of IMPACTA Perú
114 and Yale University. After establishing eligibility, participants underwent informed consent
115 procedures. Activities sequentially included pre-test HIV counseling, phlebotomy and urine
116 testing, a computer self-assisted self-interview (CASI) for data collection, and post-test
117 counseling and review of test results. Those testing positive for HIV or sexually transmitted
118 infections (STIs) were treated according to Peruvian national guidelines.

119

120 **Variables**

121 For analysis, participants' ages were dichotomized as 25 years and younger or older than
122 25, based on median age. Residence was stratified as living within or outside of Lima. High-risk
123 sexual behaviors associated with HIV infection were decided a priori using the Alaska sexual
124 risk criteria¹² for behaviors during the previous 6 months: 1) having an STI; 2) self-identification
125 as a commercial sex worker; 3) no condom use during last anal intercourse; 4) anal intercourse
126 with more than 5 partners; and 5) being a sexual partner of an HIV-infected male.

127 Sexual identity was based on whether participants self-identified as being MSM or a
128 transwoman. Identification of transwomen was confirmed through a two-step question asking

129 about gender assigned at birth and current self-identified gender.¹³ Commercial or transactional
130 sex work (CSW) was defined as having exchanged sex for money, drugs or rent in the past 6
131 months. Standard Determine® (Third Generation) rapid enzyme-linked immunosorbent assay
132 (ELISA) HIV testing provided preliminary positive HIV test results that were confirmed with
133 Western Blot testing. Rapid plasma reagin (RPR) was used to detect the presence of syphilis and
134 confirmed through microhemagglutination assay for *Treponema pallidum* antibodies (MHA-TP).
135 Acute syphilis was based on having an RPR titer $\geq 1:16$; this correlates with >90% likelihood of
136 having an active infection.⁴ In addition to these two biomarkers to test for presence of HIV and
137 syphilis, participants also reported any other STI diagnosis within the past 6 months.
138 Additionally, we asked about the diagnosis of syphilis or any other STI in the past 12 months.

139 Alcohol use disorders (AUD) were assessed using the alcohol use disorders identification
140 test (AUDIT), dichotomized by 8 or higher for hazardous use and 20 or higher for alcohol
141 dependency.¹⁴ The AUDIT has adequate reliability and validity, and its sensitivity and
142 specificity are comparable to those of other screening measures.¹⁵ Drug use was self-reported in
143 the past 3 months and included any use of marijuana, ingested cocaine (often referred to as
144 “pasta” in Perú) or inhaled/powdered cocaine, amphetamines, poppers, or ecstasy.

146 **Analysis**

147 Statistical analysis compared HIV/STI prevalence, self-reported sexual risk behavior, and
148 demographic characteristics between MSM and transwomen. All variables were binary or
149 categorical, and categorical variables were ultimately dichotomized. Variables were first
150 analyzed using χ^2 , and p-values < 0.05 were considered significant. Multiple logistic regression
151 was conducted separately comparing the correlates of HIV infection among MSM and
152 transwomen. Parsimonious models where significant correlates on bivariate analysis at $p < 0.05$
153 were included in the final model. Backward and forward elimination models were also
154 conducted. Goodness-of-fit was determined using the Akaike Information Criterion, which was
155 used to select the best fit model. Statistical Package for the Social Sciences (SPSS) 22 software
156 was used for all analyses.¹⁶


158 **Results**

159 Of the 5,127 participants, 714 (13.9%) self-reported as transwomen. Figure 1 shows
160 comparisons of different characteristics and HIV-associated risk factors between MSM and
161 transwomen, and Figure 2 illustrates the breakdown of illicit or illegal drug use by type of drug.
162 HIV prevalence was significantly two-fold higher among transwomen compared to MSM (14.7%
163 vs. 7.0%, $p < .001$) and for syphilis and acute syphilis (27.0% vs. 12.3%, $p < .001$; 10.8% vs. 4.2%,
164 $p < .001$). In addition to the biomarkers, transwomen also had a higher self-report of syphilis or an
165 STI in the past 12 months (13.0% vs. 6.3%, $p < .001$; 29.3% vs. 24.4%, $p = .04$).

166 Compared to MSM, transwomen were significantly more likely to have more than five
167 sexual partners (61.0% vs. 40.9%, $p < .001$) and engage in commercial sex work (57.6% vs.
168 29.7%, $p < .001$). However, more MSM reported engaging in unprotected sex in the last sexual
169 encounter (47.9% vs. 36.9%, $p < .001$) and having an HIV infected sexual partner (10.2% vs.
170 5.1%, $p < .001$).

171 Compared to MSM, more transwomen met criteria for alcohol dependence (19.0% vs.
172 16.4%, $p = .06$). Differences in overall illicit/illegal drug use between transwomen and MSM were
173 not significant (17.0% vs. 14.9%, $p = .15$; data not presented); however, in further analysis by
174 drug type, there were significant differences in the two dominant drugs – inhaled cocaine and
175 marijuana. MSM were more likely to use marijuana (68.0% vs. 50.0%, $p < .001$), and transwomen
176 were more likely to engage in inhaled cocaine use (70.0% vs. 51.1%, $p < .001$).

177 For associations between HIV correlates and a positive HIV status, with the exception of
178 alcohol dependence, the selected risk factors were all found to be significantly associated with
179 MSM as depicted in Table 1. For transwomen, none of the correlates were statistically
180 significantly associated at $\alpha \leq .05$. However, at $\alpha \leq .10$, living in Lima (OR=3.27,
181 $p < .001$), having acute syphilis (aOR=1.81, $p = .10$), engaging in illicit/illegal drug use in the past
182 three months (aOR=1.65, $p = .08$), and being alcohol dependent (aOR=1.69, $p = .07$) increased the
183 likelihood of a positive HIV status for the study participant. Finally, being over the age of 25 was
184 not significantly associated with HIV infection for transwomen, whereas in the MSM cohort,
185 being over the age of 25 was associated with an increased risk of HIV infection.




186

187 **Figure 1: Demographics, risk factors, and STI/HIV prevalence by sexual identity:**

188 **Transwomen versus Men Who Have Sex With Men**

189



190

191 **Figure 2: Illegal drug use by Transwomen versus Men who have sex with men (n=5,127)**

192

Variable	Men who have sex with men (n=4413)			Transwomen (n=714)		
	AOR	95% CI	p-value	AOR	95%CI	p-value
Over age 25	1.25	.97 - 1.61	0.08	1.13	.70 - 1.80	0.62
Lima residence	2.86	2.16 - 3.78	<.001	3.27	1.81 - 5.91	<.001
Acute syphilis	2.6	1.63 - 4.15	<.001	1.81	.88 - 3.73	0.10
Syphilis positive	2.3	1.60 - 3.31	<.001	1.4	.78 - 2.52	0.26
Any STI past 12 mths	2.02	1.50 - 2.71	<.001	0.96	.51 - 1.82	0.90
Drugs in past 3 mths	0.49	.32 - .76	<.001	1.65	.94 - 2.87	0.08
Alcohol Dependent	1.21	.84 - 1.76	0.30	1.69	.94 - 3.03	0.07

193 **Table 1: Independent Risks for HIV among Transwomen and Men Who Have Sex with**
 194 **Men (N=5,127)**

195
 196 **Discussion**

197 Study findings demonstrate transwomen living in Perú have a higher HIV/STI prevalence
 198 and engaged in more high-risk behavior than MSM. HIV prevalence among transwomen was
 199 more than double that of MSM, which is consistent with other research that identifies
 200 transwomen as a higher risk group.¹⁷ Transwomen were more likely to engage in sex work, to
 201 have a higher mean number of sexual partners, and to have an alcohol use disorder.

202 In this study, drug use in the past three months was statistically associated with HIV in
 203 both MSM and transwomen; however, it was protective for MSM, although statistically non-
 204 significant at the designated alpha level. Drug use also had a trend of increasing HIV risk for
 205 transwomen.

206 In-depth analysis was performed for drug use, and there were differences in the drug type
 207 usage – MSM had more marijuana use, while transwomen had higher rates for inhaled or snorted
 208 cocaine (Figure 2). Upon further exploration of the literature, marijuana, which in this study is
 209 more frequently used among MSM compared to transwomen, might be commonly used in a
 210 recreational capacity in MSM populations. However, cocaine use among transwomen may be

211 used more often as a coping mechanism compared to MSM during transactional sexual
212 encounters.^{18,19} Theoretically, both scenarios of recreational marijuana use or cocaine during
213 survival sex can increase sexual risk behavior and HIV risk.^{12,18,19} However, in this study,
214 perhaps drug use, more specifically cocaine use, increases risk among transwomen because of
215 greater exposure to and frequency of illicit/illegal poly substance drug use.

216 MSM were more likely to report having sex with an HIV-infected partner in the past
217 three months than transwomen; however, transwomen also reported more commercial sex work
218 and engaging in sexual activities with more than five partners. If transwomen engage in more
219 commercial sex work and the HIV status of their sexual partner is not necessarily known, sexual
220 acts with an HIV positive partner may be underreported among transwomen. In this population,
221 little is known about the demography and health status of sex worker clients'; however, among
222 those who engage in sex work, unprotected sexual acts with regular or non-paying clients are
223 commonly reported²⁰⁻²³, and these unprotected sexual acts with regular clients can contribute to
224 the spread of HIV/STIs.

225 HIV infection being associated with syphilis and older MSM is consistent with a previous
226 similar study among Peruvian MSM.²⁴ However, there are two differences between this study
227 and the other study. First, age or being older than 25 does not seem to be related to HIV status
228 for transwomen in the present study. Additionally, in this study, testing positive for syphilis was
229 not associated with HIV infection among transwomen, despite syphilis prevalence being more
230 than double among transwomen than MSM. Only acute syphilis or early infection, which was
231 also more prevalent among transwomen, was associated with an HIV seropositive status.
232 However, the lack of association between a non-reactive syphilis screening test and HIV
233 seropositivity may be related to lack of power in the smaller sample size of the transwomen
234 cohort.

235 The impact of having syphilis or co-infection with syphilis and HIV is problematic
236 because syphilis can lead to a higher likelihood of HIV transmission, while being co-infected can
237 accelerate progression of HIV illness.²⁵ Therefore, this area deserves more investigation into
238 possible confounders or moderators, such as access to care or other structural issues²⁶, that
239 impact the relationship between these two variables. A more stable statistical model should be
240 used to confirm this relationship, especially now when there has been a further increase of
241 syphilis rates in Perú and globally²⁷.

242 Alcohol use and sex under the influence, which are known correlates for HIV
243 transmission¹², are more prevalent among transwomen than MSM living in Perú²⁸. This statistic
244 is consistent in this study. Furthermore, alcohol dependence is an area of concern for
245 transwomen, as more transwomen were identified as being alcohol dependent in this study, and
246 alcohol dependence was trending as being linked with positive HIV infection, which was not the
247 case for MSM in the sample.

248 There were several limitations to this study. The method of recruitment, modified
249 snowball sampling, may have introduced some bias; however, it is considered the most useful
250 method to sample hard-to-reach populations such as MSM and transwomen.²⁹ It should be noted
251 that the sample may not be representative of the target population as compared to a typical
252 random sample. The study was cross-sectional, which did not allow for a temporal assessment of
253 causality. Recall bias may have also been an issue for some of the self-reported variables. With
254 transwomen having one sixth of the MSM sample size and having a less than 15% HIV
255 prevalence, the power may not have been adequate in the logistic regression analysis to clearly
256 identify correlates associated with HIV-infection among transwomen. Although there were
257 biomarkers for HIV and syphilis, self-report was used to capture data on other STIs and alcohol
258 use, which could have led to some level of underreporting and underestimations.

259

260 **Conclusion**

261 By using one of the largest samples of surveillance data among Peruvian MSM and
262 transwomen, this study is one of few existing studies that examines differences in risk factors
263 between MSM and transwomen. Our study provides evidence for the need of expanded and
264 unique research efforts targeted towards transwomen in the Andean region and all South
265 America. These results may also be replicated on a global scale. Treating these two groups as the
266 same demographic may ultimately result in disappointing outcomes for primary prevention
267 programs aimed at curbing HIV incidence among transwomen and ultimately a misapplication of
268 valuable public health resources.

269 Both MSM and transwomen living in Perú engage in high-risk sexual behaviors, placing
270 them at increased risk for HIV/STI transmission. However, transwomen have a formidable
271 disadvantage between the two groups. Compared to MSM, transwomen engage in more high-risk
272 behaviors such as substance and alcohol use, commercial or transactional sex-work, sex with

273 multiple partners, and unprotected sex. Emphasis should be placed on this group as a most at-risk
274 population (MARP)^{28,30} in Perú and globally.

275 Additionally, substance use, in particular alcohol dependence and inhaled cocaine use,
276 may have an association with high-risk sexual behavior among transwomen, and this relationship
277 should be explored further. Emerging research^{5,6} continues to reinforce the importance of these
278 types of studies, as they are meaningful for developing a comprehensive body of literature on
279 this subject and providing tools for HIV researchers working with transwomen.

280 **Conflict of Interest Statement:** There are no conflicts of interest to disclose.

281

282 **Author Contributions:**

283 Elena Cyrus, PhD, MPH^{1,2}: Principal Investigator of the study. I developed the research
284 question, completed analysis and was the primary writer for the paper.

285 Javier R. Lama, MD, MPH³: Co-Investigator of parent study where the data was sourced in
286 Perú. Contributed to review and edits of final version of paper

287 Jorge Sanchez, MD, MPH⁴: Co-Principal Investigator of parent study from where the data was
288 sourced in Perú. Contributed to review and edits of final version of paper

289 Segundo Leon, MPH⁵: Contributed to interpretation of results and manuscript writing

290 Manuel V. Villaran, MD, MPH, MSc³: Contributed to interpretation of results and manuscript
291 writing

292 Panagiotis Vagenas, PhD, MPH⁶: Contributed to interpretation of results and manuscript writing

293 Daniell S. Sullivan⁷: Contributed to manuscript writing and editing

294 David Vu⁷: Contributed to manuscript writing and editing

295 Makella Coudray PhD¹: Contributed to manuscript writing and editing

296 Frederick L. Altice, MD, MA⁸: Senior mentor/author and Co-Principal Investigator of parent
297 study from where the data was sources

298

299 **Acknowledgements:** This project was supported by the National Institute on Drug Abuse

300 (NIDA) at the National Institutes of Health and Award K99/R00DA046311; The Loan

301 Repayment Program at the National Institute on Alcohol Abuse and Alcoholism (NIAAA); and

302 the NIH Research Training Grant # R25 TW009338 funded by the Fogarty International Center,

303 the National Heart, Lung and Blood Institute, the NIH Office of the Director: Office of Research

304 on Women's Health and the NIH Office of the Director Office of AIDS Research.

305

306 **References**

- 307 1. Joint United Nations Programme on HIV/AIDS (UNAIDS). Regional fact sheet 2020: Latin
308 America and the Caribbean. https://www.unaids.org/sites/default/files/media_asset/2020_aids-
309 [data-book_en.pdf](https://www.unaids.org/sites/default/files/media_asset/2020_aids-data-book_en.pdf).
- 310 2. Silva-Santisteban A, Eng S, de la Iglesia G, Falistocco C, Mazin R. HIV prevention among
311 transgender women in Latin America: implementation, gaps and challenges. *J Int AIDS Soc.*
312 2016 Jul 17;19(3 Suppl 2):20799.
- 313 3. Chow JY, Konda KA, Calvo GM, Klausner JD, Caceres CF. Demographics, Behaviors, and
314 Sexual Health Characteristics of High Risk Men Who Have Sex With Men and Transgender
315 Women Who Use Social Media to Meet Sex Partners in Lima, Peru. *Sex Transm Dis* 2017 44(3):
316 143-148.
- 317 4. Long JE, Montañó M, Cabello R, Sanchez H, Lama JR, Duerr A. Brief Report: Comparing
318 Sexual Risk Behavior in a High-Risk Group of Men Who Have Sex With Men and Transgender
319 Women in Lima, Peru. *J Acquir Immune Defic Syndr.* 2019;80(5):522-526.
320 doi:10.1097/QAI.0000000000001966.
- 321 5. Salazar X, Núñez-Curto A, Villayzán J, Castillo R, Benites C, Caballero P, Cáceres CF. How
322 Peru introduced a plan for comprehensive HIV prevention and care for transwomen. *J Int AIDS*
323 *Soc.* 2016 Jul 17;19(3 Suppl 2):20790.
- 324 6. Poteat T, Wirtz AL, Radix A, Borquez A, Silva-Santisteban A, Deutsch MB, et al. HIV risk
325 and preventive interventions in transgender women sex workers. *The Lancet.*
326 2015;385(9964):274-286.
- 327 7. Salazar X, Villayzán J. *La situación de la población Trans el Perú en el contexto del acceso*
328 *universal a tratamiento, atención y apoyo en VIH/SIDA.* IESSDEH, ONUSIDA (in Spanish);
329 Lima: 2010.
- 330 8. Operario D, Burton J, Underhill K, Sevelius J. Men who have sex with transgender women:
331 challenges to category-based HIV prevention. *AIDS Behav.* 2008;12:18–26

- 332 9. Castillo R, Konda KA, Leon SR, Silva-Santisteban A, Salazar X, Klausner JD, Coates TJ,
333 Cáceres CF. HIV and Sexually Transmitted Infection Incidence and Associated Risk Factors
334 Among High-Risk MSM and Male-to-Female Transgender Women in Lima, Peru. *J Acquir*
335 *Immune Defic Syndr.* 2015 Aug 15;69(5):567-75.
- 336 10. Chariyalertsak S, Kosachunhanan N, Saokhieo P, Songsupa R, Wongthanee A,
337 Chariyalertsak C, et al. HIV incidence, risk factors, and motivation for biomedical intervention
338 among gay, bisexual men, and transgender persons in northern Thailand. *PloS one.*
339 2011;6(9):e24295.
- 340 11. Herrera MC, Konda KA, Leon SR, Deiss R, Brown B, Calvo GM, Salvatierra HJ,
341 Cáceres CF, Klausner JD. Impact of alcohol use on sexual behavior among men who have sex
342 with men and transgender women in Lima, Peru. *Drug Alcohol Depend.* 2016 Apr 1;161:147-54.
- 343 12. Ludford KT, Vagenas P, Lama JR, Peinado J, Gonzales P, Leiva R, et al. Screening for drug
344 and alcohol use disorders and their association with HIV-related sexual risk behaviors among
345 men who have sex with men in Peru. *PLoS One.* 2013;8(8):e69966. Published 2013 Aug 6.
- 346 13. Reisner SL¹, Biello K, Rosenberger JG, Austin SB, Haneuse S, Perez-Brumer A, Novak DS,
347 Mimiaga MJ. Using a two-step method to measure transgender identity in Latin America/the
348 Caribbean, Portugal, and Spain. 2014 Nov;43(8):1503-14. doi: 10.1007/s10508-014-0314-2.
- 349 14. Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro M. AUDIT: The alcohol use disorders
350 identification test: Guidelines for use in Primary Care, 2nd edition. 2001.
- 351 15. Allen JP, Litten RZ, Fertig JB, Babor T. A review of research on the alcohol use disorders
352 identification test (AUDIT). *Alcohol Clin Exp Res.* 1997;21(4):613-619.
- 353 16. IBM Corp. IBM SPSS statistics for windows, version 21.0. 2012.
- 354 17. Centers for Disease Control and Prevention (CDC). HIV among transgender people.
355 <http://www.cdc.gov/hiv/risk/transgender/>. Updated 2013. Accessed July, 2014.
- 356 18. Santos G, Rapues J, Wilson EC, Macias O, Packer T, Colfax G, et al. Alcohol and substance
357 use among transgender women in San Francisco: Prevalence and association with human
358 immunodeficiency virus infection. *Drug Alcohol Rev.* 2014;33(3):287-295.

- 359 19. Baum MK, Rafie C, Lai S, Sales S, Page B, Campa A. Crack-cocaine use accelerates HIV
360 disease progression in a cohort of HIV-positive drug users. *J Acquir Immune Defic Syndr*.
361 2009;50(1):93-99.
- 362 20. Kerrigan D, Wirtz A, Baral S, Decker M, Murray L, Poteat T, et al. *The global HIV*
363 *epidemics among sex workers*. World Bank Publications; 2012.
- 364 21. Wang C, Hawes SE, Gaye A, Sow PS, Ndoye I, Manhart LE, et al. HIV prevalence, previous
365 HIV testing, and condom use with clients and regular partners among Senegalese commercial
366 sex workers. *Sex Transm Infect*. 2007;83(7):534-540.
- 367 22. Voeten H, Egesah O, Varkevisser C, Habbema J. Female sex workers and unsafe sex in
368 urban and rural Nyanza, Kenya: Regular partners may contribute more to HIV transmission than
369 clients. *Tropical Medicine & International Health*. 2007;12(2):174-182.
- 370 23. Platt L, Jolley E, Rhodes T, Hope V, Latypov A, Reynolds L, et al. Factors mediating HIV
371 risk among female sex workers in Europe: A systematic review and ecological analysis. *BMJ*
372 *open*. 2013;3(7).
- 373 24. Sanchez J, Lama JR, Peinado J, Paredes A, Lucchetti A, Russell K, et al. High HIV and
374 ulcerative sexually transmitted infection incidence estimates among men who have sex with men
375 in Perú: Awaiting for an effective preventive intervention. *J Acquir Immune Defic Syndr*.
376 2009;51 Suppl 1:S47-51.
- 377 25. Buchacz K, Patel P, Taylor M, Kerndt PR, Byers RH, Holmberg SD, et al. Syphilis increases
378 HIV viral load and decreases CD4 cell counts in HIV-infected patients with new syphilis
379 infections. *AIDS*. 2004;18(15):2075-2079.
- 380 26. Poteat T, Wirtz AL, Radix A, Borquez A, Silva-Santisteban A, Deutsch MB, et al. HIV risk
381 and preventive interventions in transgender women sex workers. *The Lancet*. 2014.
- 382 27. Hung P, Osias E, Konda KA, Calvo GM, Reyes-Diaz ME, Vargas SK, et al. High Lifetime
383 Prevalence of Syphilis in Men Who Have Sex With Men and Transgender Women Versus Low

384 Lifetime Prevalence in Female Sex Workers in Lima, Peru. *Sex Transm Dis.* 2020;47(8):549-
385 555.

386 28. Baral SD, Poteat T, Strömdahl S, Wirtz AL, Guadamuz TE, Beyrer C. Worldwide burden of
387 HIV in transgender women: A systematic review and meta-analysis. *The Lancet infectious*
388 *diseases.* 2013;13(3):214-222.

389 29. Heckathorn DD. Respondent-driven sampling II: Deriving valid population estimates from
390 chain-referral samples of hidden populations. *Social Problems.* 2002;49(1):11.

391 30. Beyrer C, Baral S, Kerrigan D, El-Bassel N, Bekker LG, Celentano DD. Expanding the
392 space: Inclusion of most-at-risk populations in HIV prevention, treatment, and care services. *J*
393 *Acquir Immune Defic Syndr.* 2011;57 Suppl 2:S96-9.

394

395