# The time-varying association between previous antibiotic use and antibiotic resistance

Avi Baraz[1,2], Michal Chowers[3,4], Daniel Nevo[5], Uri Obolski[1,2,*]

[1]School of Public Health, Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

[2]Porter School of the Environment and Earth Sciences, Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel

[3]Meir Medical Center, Kfar Saba, Israel

[4]Sackler Faculty of Medicine, Tel Aviv University, Israel

[5]Department of Statistics and Operations Research, Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel

[*]Corresponding author

E-mail: uriobols@tauex.tau.ac.il

# Abstract

**Objectives**

The objective of the study was to estimate how the time elapsed from previous antibiotic use is associated with antibiotic resistance.

**Methods**

Data comprised electronic medical records of all patients in an Israeli hospital who had a positive bacterial culture between 2016-19. These included susceptibility testing results, and clinical and demographic data. Mixed-effects time-varying logistic models were fitted to estimate the association between time elapsed since last use of aminoglycosides and gentamicin resistance ($n = 13,095$), cephalosporins and ceftazidime resistance ($n = 13,051$), and fluoroquinolones and ciprofloxacin resistance ($n = 15,364$), while adjusting for multiple covariates.

**Results**

For all examined antibiotics, previous antibiotic use had a statistically significant association with resistance ($p < 0.001$). These associations exhibited a clear decreasing pattern over time, which we present as a flexible function of time. Nonetheless, previous antibiotic use remained a significant risk factor for resistance for at least 180 days, with adjusted odds ratios of 1.94 (95% CI 1.40-2.69), 1.33 (95% CI 1.10-1.61), and 2.25 (95% CI 1.49-3.41), for gentamicin, ceftazidime and ciprofloxacin, respectively.

**Conclusions**

The association between prior antibiotic use and resistance decreases over time. Commonly used cut-offs for prior antibiotic use can either misclassify patients still at higher risk, when too

41 recent; or provide a diluted estimate of the effects of antibiotic use on future resistance, when

42 too distant. Hence, prior antibiotic use should be considered as a time-dependent risk factor for

43 resistance, both in epidemiological research and clinical practice.

# Introduction

Empiric antibiotic therapy is a common and crucial component of treating hospitalized patients.

However, increased antibiotic resistance frequencies often lead to inappropriate empiric therapy

[1]. Risk factors for antibiotic resistant infections are hence constantly investigated, for both

research and practical, clinical purposes [2]. A biologically plausible and empirically evident

cause for resistance is previous use of antibiotics [3]. Despite its importance, little is known

about the association between the time elapsed since antibiotic use on the risk for resistance.

Both prospective and retrospective studies ubiquitously report previous antibiotic use as a

binary variable, indicating prior antibiotic use in certain, often arbitrary time-periods, with

common cut-offs of 30, 90, or 365 days [2,4-11]. Such discretization of the data can disregard

patients with existing risk if the cut-off is too short, or dilute the effects of the risk if the cut-off is

too long. Here, we employed patients' electronic medical records (EMRs) to estimate the risk of

antibiotic resistance as a function of the time elapsed since their last antibiotic use. We applied

mixed-effects time-varying logistic regression and obtained the adjusted risk estimates as

flexible functions of time elapsed since last use of antibiotics for gentamicin, ceftazidime and

ciprofloxacin.

# Materials and methods

## Data

The dataset included the EMRs of all patients who had a positive bacterial culture between

2016-19, in Meir medical center, Israel. The data contained, patients' antibiotic susceptibility test

results from all positive cultures, as well as demographic and clinical data, including antibiotic

use within the hospital in the year preceding the susceptibility tests (see Supplementary

information, Tables S1-S3). Cultures were taken upon suspected infection, but were not verified

against clinical symptoms.

The main variable of interest was the time elapsed (in days) since a patients' last antibiotic use

in the hospital until the drawing of their bacterial cultures. Antibiotic use was considered only

when it commenced $\geq 5$ days and $\leq 365$ days before a culture was drawn. The first constraint

was set as to avoid date misclassifications that might lead to antibiotics prescribed after culture

results were obtained, and since it is a short period for resistance acquisition. The second

constraint was set due to data availability. To attain a sufficient sample size, antibiotic use was

grouped into cephalosporins (of all generations), fluoroquinolones and aminoglycosides,

administered either orally or intravenously. Resistance to three antibiotics, selected as markers

for resistance to the classes of antibiotics considered above, was modeled: gentamicin,

ceftazidime, and ciprofloxacin.

## Statistical analysis

Relationships between patients' clinical and demographic covariates, and antibiotic resistance

were modeled by mixed-effects time-varying logistic models [12]. These models extend

standard logistic regression by allowing the effect of previous antibiotic use to vary as flexible

functions of time, using splines (see Supplementary information for additional information). The

models addressed multiple cultures from the same patients by assigning a random intercept per

patient, although this addition did not substantially change the results. The models were

adjusted for multiple prominent clinical and demographic covariates, chosen based on prior

knowledge [2], and underwent sensitivity and subgroup analyses (see Supplementary

information, Tables S1-S3).

Additionally, we examined adjustments for prior use of the other classes of antibiotics modeled,

as well as to non-cephalosporin beta-lactam antibiotics, to account for potential cross-resistance

[13]. Each of these covariates was modeled as a dummy variable (used/not), with a linear time

trend (in the log-odds scale). These variables had little influence on the results, and hence only

prior use of non-cephalosporin beta-lactams was incorporated into the ceftazidime resistance

model (see Supplementary information, Figures S1-S3).

# Ethics declaration

95　The study was approved by the Institutional Review Board Committee of the Meir Medical

96　Center (MMC-0143-19).

# Results

We examined the association between previous aminoglycoside use and resistance to

gentamicin ($n = 13,095$) cultures, resistance frequency 15.58%); previous cephalosporin use

and resistance to ceftazidime ($n = 13,051$ cultures, resistance frequency 24.27%); and previous

fluoroquinolone use and resistance to ciprofloxacin ($n = 15,364$) cultures, resistance frequency

26.66%). For all examined antibiotics, the time-varying coefficient of previous antibiotic use in

the hospital had a statistically significant association with resistance ($p < 0.001$) (see

Supplementary information, Additional Information, Tables S7-S9). As expected, the spline-

based adjusted odds ratios (aORs) decayed as time passed from last antibiotic use (Figure 1).

**Fig. 1. Estimates of risk for antibiotic resistance from generalized additive mixed-effects models:** Spline-based estimates (curves) and their 95% CI (shaded) of the aORs for resistance, against days elapsed since last use of relevant antibiotics. (a) gentamicin resistance against last use of aminoglycosides; (b) ceftazidime resistance against last use of cephalosporins; and (c) ciprofloxacin resistance against last use of fluoroquinolones.

Importantly, the estimates across the three antibiotics examined maintained significant and positive associations between previous use and resistance for over 180 days. Summary of the aORs and their CIs at 30, 90, 180 and 365 days are provided in Table 1. For gentamicin and ceftazidime, the decrease of the aOR can be very well approximated by exponential decay of the forms        and       , respectively, where    is months since last antibiotic use (see Supplementary material for details). The decrease in the aOR of ciprofloxacin resistance did not follow such a form, and the estimate varied substantially as time elapsed from the last use increased, possibly due to limited sample size for distant periods (Figure 1C, shaded region).

|  | aOR at 30 days (95% CI) | aOR at 90 days (95% CI) | aOR at 180 days (95% CI) | aOR at 365 days (95% CI) |
|---|---|---|---|---|
| Resistance to gentamicin (aminoglycoside use) | 2.92 (2.24, 3.81) | 2.48 (1.94, 3.16) | 1.94 (1.40, 2.69) | 1.17 (0.59, 2.31) |
| Resistance to ceftazidime (cephalosporin use) | 1.71 (1.45, 2.00) | 1.54 (1.32, 1.80) | 1.33 (1.10, 1.61) | 0.98 (0.70, 1.37) |
| Resistance to ciprofloxacin, (fluoroquinolone use) | 4.54 (3.49, 5.90) | 3.36 (2.45, 4.59) | 2.25 (1.49, 3.41) | 2.11 (0.86, 5.14) |

**Table 1. Adjusted odds-ratios (aOR) for resistance to different antibiotics by time since**
**last antibiotic use.** The aOR were adjusted for patients' clinical and demographic
characteristics, as well as to bacterial species and sample source (see Supplementary
information, Tables S7-S9).

# Discussion

In this study, we provided time-varying estimates of the association of previous antibiotic use
with antibiotic resistance. We have characterized these estimates for gentamicin, ceftazidime
and ciprofloxacin, serving as proxies for resistance to aminoglycosides, cephalosporins and
fluoroquinolones, respectively. Whereas the estimates differed between the antibiotics
examined, they all exhibited a marked decrease over time. These differences might be
explained by the variation of the link between exposure and resistance acquisition between

antibiotic classes [8]. Higher aORs are expected for fluoroquinolones relative to
aminoglycosides [8]. However, the relatively low aORs obtained for cephalosporins might be
due to the aggregation of different cephalosporins administered to patients. 1st and 2nd
generation cephalosporins are more widely used in the studied hospital, and are expected to
confer lower resistance to ceftazidime, compared to 3rd generation cephalosporins [8,14].

Nonetheless, for all examined antibiotics, previous antibiotic use remained a statistically
significant risk factor for resistance for at least 180 days.

Our results have important implications for both research and clinical practice. Studies analyzing
risk factors for antibiotic resistance model previous antibiotic use as binary indicators, choosing
cut-offs ranging from several days to a year prior to the examined bacterial sample. Our results
demonstrated that this might lead to inaccuracies. Applying antibiotic use cut-offs close to the
outcome (e.g. 30 or 90 days), might lead to substantial underestimation of the risk of patients
using antibiotics before the cut-offs. Conversely, applying distant cut-offs (e.g. one year) might
lead to underestimation of the association between previous use and resistance for patients
who used antibiotics recently. One solution considers the time elapsed since antibiotic use,
either with splines as demonstrated here, or approximated by linear interactions with time in a
standard logistic regression model. Furthermore, clinicians should bear in mind that patients
who previously used antibiotics are more prone to have resistant infections, and that this risk
decays over time but persists for at least 180 days.

Our study has several limitations. Despite known biological mechanisms linking antibiotic use to
resistance, the results presented here do not claim to be causal estimates. Residual
confounding could still affect our estimates. For example, we did not possess patient data of
antibiotic use outside the hospital. This could have contributed to the high variance in the
fluoroquinolone estimates or affected the cephalosporin estimates, as fluoroquinolones and

older generation cephalosporins may be prescribed to outpatients. However, since our models

adjust for many other variables that could account for these unobserved differences between

patients, we hope that these effects are reasonably mitigated. Furthermore, the sample

collection dates in our data reflect clinical infection onset, but not necessarily the patient's

microbiota. Ideally, a study prospectively sampling patients could more accurately inform on the

formation of antibiotic resistant bacterial populations. Finally, this is a single center study, and

the estimates presented here could vary in settings with different antibiotic resistance

frequencies, patient populations, or other conditions affecting bacterial infections. Hence,

although the general time-varying patterns observed here may persist, they are best estimated

locally, for more accurate risk stratification of patients. Nonetheless, the implications of our

estimates are relevant for studies attempting to estimate risk factors, or predictors, for

resistance, regardless of causal interpretation. This is true both for epidemiological studies

using these variables, or when clinicians utilize previous antibiotic use to estimate the risk of

antibiotic resistant infections for hospitalized patients.

In conclusion, this study provides time-varying estimates of the association of antibiotic use with

future antibiotic resistance. These estimates are relevant both for epidemiological literature and

clinical practice. Commonly used cut-offs for previous use can either misclassify patients still at

higher risk, or provide diluted estimates of the association of antibiotic use with future

resistance. We therefore implore to consider prior antibiotic use as a time-dependent risk factor

for resistance, and to continue researching this phenomenon for various antibiotics and settings.

# Transparency declaration

There are no potential conflicts of interest for any authors.

## Funding

This work was supported by The Israel Science Foundation (ISF 1286/21).

## Author contributions

UO and DN conceived the study and supervised all analysis; AB performed the analyses; MC provided clinical insights; All authors interpreted the results; AB, DN, and UO wrote the initial draft; all authors critically revised and approved the paper.

## References

1. Carrara E, Pfeffer I, Zusman O, Leibovici L, Paul M. Determinants of inappropriate empirical antibiotic treatment: systematic review and meta-analysis. Int J Antimicrob Agents. 2018 Apr;51(4):548–53. DOI: https://doi.org/10.1016/j.ijantimicag.2017.12.013

2. Chatterjee A, Modarai M, Naylor NR, Boyd SE, Atun R, Barlow J, et al. Quantifying drivers of antibiotic resistance in humans: a systematic review. Lancet Infect Dis. 2018 Dec;18(12):e368–78. DOI: https://doi.org/10.1016/S1473-3099(18)30296-2

3. Holmes AH, Moore LSP, Sundsfjord A, Steinbakk M, Regmi S, Karkey A, et al. Understanding the mechanisms and drivers of antimicrobial resistance. The Lancet. 2016 Jan;387(10014):176–87. DOI: https://doi.org/10.1016/S0140-6736(15)00473-0

4. Bakhit M, Hoffmann T, Scott AM, Beller E, Rathbone J, Del Mar C. Resistance decay in individuals after antibiotic exposure in primary care: a systematic review and meta-analysis. BMC Med. 2018 Dec;16(1):126. DOI: https://doi.org/10.1186/s12916-018-1109-4

5. Bassetti M, Righi E, Vena A, Graziano E, Russo A, Peghin M. Risk stratification and treatment of ICU-acquired pneumonia caused by multidrug- resistant/extensively drug-resistant/pandrug-resistant bacteria: Curr Opin Crit Care. 2018 Oct;24(5):385–93. DOI: 10.1097/MCC.0000000000000534

6. Raman G, Avendano EE, Chan J, Merchant S, Puzniak L. Risk factors for hospitalized patients with resistant or multidrug-resistant Pseudomonas aeruginosa infections: a systematic

review and meta-analysis. Antimicrob Resist Infect Control. 2018 Dec;7(1):79. DOI:

https://doi.org/10.1186/s13756-018-0370-9

7. Prina E, Ranzani OT, Polverino E, Cillóniz C, Ferrer M, Fernandez L, et al. Risk Factors

Associated with Potentially Antibiotic-Resistant Pathogens in Community-Acquired Pneumonia.

Ann Am Thorac Soc. 2015 Feb;12(2):153–60. DOI: https://doi.org/10.1513/AnnalsATS.201407-

305OC

8. Sulis G, Sayood S, Katukoori S, Bollam N, George I, Yaeger LH, et al. Exposure to World

Health Organization's AWaRe antibiotics and isolation of multidrug resistant bacteria: a

systematic review and meta-analysis. Clin Microbiol Infect. 2022 Sep;28(9):1193–202. DOI:

https://doi.org/10.1016/j.cmi.2022.03.014

9. Minh NNQ, Toi PV, Qui LM, Tinh LBB, Ngoc NT, Kim LTN, et al. Antibiotic use and

prescription and its effects on Enterobacteriaceae in the gut in children with mild respiratory

infections in Ho Chi Minh City, Vietnam. A prospective observational outpatient study. Foster J,

editor. PLOS ONE. 2020 Nov 4;15(11):e0241760. DOI:

https://doi.org/10.1371/journal.pone.0241760

10. Dan S, Shah A, Justo JA, Bookstaver PB, Kohn J, Albrecht H, et al. Prediction of

Fluoroquinolone Resistance in Gram-Negative Bacteria Causing Bloodstream Infections.

Antimicrob Agents Chemother. 2016 Apr;60(4):2265–72. DOI:

https://doi.org/10.1128/AAC.02728-15

11. Shah A, Justo JA, Bookstaver PB, Kohn J, Albrecht H, Al-Hasan MN. Application of

Fluoroquinolone Resistance Score in Management of Complicated Urinary Tract Infections.

Antimicrob Agents Chemother. 2017 May;61(5):e02313-16. DOI:

https://doi.org/10.1128/AAC.02313-16

12. Hastie T, Tibshirani R. Varying-Coefficient Models. J R Stat Soc Ser B Methodol. 1993

Sep;55(4):757–79. DOI: https://doi.org/10.1111/j.2517-6161.1993.tb01939.x

13. Cherny SS, Nevo D, Baraz A, Baruch S, Lewin-Epstein O, Stein GY, et al. Revealing

antibiotic cross-resistance patterns in hospitalized patients through Bayesian network modelling.

:10. DOI: https://doi.org/10.1093/jac/dkaa408

14. Chowers M, Zehavi T, Gottesman BS, Baraz A, Nevo D, Obolski U. Estimating the impact of

cefuroxime versus cefazolin and amoxicillin/clavulanate use on future collateral resistance: a

retrospective comparison. J Antimicrob Chemother. 2022 Apr 15;dkac130. DOI:

https://doi.org/10.1093/jac/dkac130