Assessment of Knowledge on Human Mpox Virus among General Population in Bangladesh

Sudipta Deb Nath[1#], A.M. Khairul Islam[2*#], Koushik Majumder[3], Fahmida Hoque Rimti[3], Jyoti Das[4], Mustari Nailah Tabassum[3], Arefin Naher Oishee[5], Tarannum Mahmood[3], Monisha Paul[6], Muntasrina Akhter[7], Alok Bijoy Bhadra[8], Fariha Hoque Rimu[9], Snahasish Chakraborty[10], Preetom Shom[11], Sirajum Monira Nosaibah[12], Md Ashikur Rahman[4], Ahmed Safwan Khan[13], Anika Anjum[5], Sushmita Khan[14], M. Mahbub Hossain[15], Mohammad Delwer Hossain Hawlader[4]

1. (i) Department of Genetic Engineering and Biotechnology, University of Dhaka, Dhaka-1000, Bangladesh, (ii)Child Health Research Foundation (CHRF), Dhaka-1216, Bangladesh

2. International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,b), Dhaka-1212, Bangladesh

3. Chittagong Medical College, Chattogram-4203, Bangladesh

4. Department of Public Health, North South University, Dhaka-1229, Bangladesh

5. M Abdur Rahim Medical College, Dinajpur-5200, Bangladesh

6. Khulna Medical College, Khulna-9000, Bangladesh

7. Public Health Professional Development Society (PPDS), Dhaka-1215, Bangladesh

8. Sheikh Hasina Medical College, Habiganj-3300, Bangladesh

9. University of Rochester Medical Center, New York-14642, USA

10. Rangamati Medical College-4500, Bangladesh

11. Hajee Mohammad Danesh Science & Technology University, Dinajpur-5200, Bangladesh

12. Sylhet MAG Osmani Medical College, Sylhet-3100, Bangladesh

13. Sher -E- Bangla Medical College, Barishal-8200, Bangladesh

14. Rajshahi Medical College, Rajshahi-6100, Bangladesh

15. Department of Decision and Information Sciences, C.T. Bauer College of Business & Department of Health Systems and Population Health Sciences, Tilman J. Fertitta Family College of Medicine, University of Houston, TX 77204, USA

Conjoint first author

34   * **Corresponding Author:**

35   A.M.Khairul Islam

36   International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,b)

37   Dhaka-1212, Bangladesh

38   Email: ikhairul963@gmail.com

Abstract

Introduction: Mpox (Monkeypox) is a zoonotic Orthopox virus of the Poxviridae family. The general public in Bangladesh should be informed about Mpox to reduce the burden of a possible epidemic in the community.

Objective: The study aimed to determine the level of public awareness and knowledge about Mpox among the general community to provide information regarding future preventive measures.

Materials and Method: From May to June 2022, this nationwide cross-sectional study was conducted in eight administrative divisions in Bangladesh. We determined the sample size using Cochran's formula. Through a semi-structured questionnaire, data regarding sociodemographic characteristics and knowledge about Mpox were collected via face-to-face interviews. IBM SPSS v.25 was used to analyze the data.

Result: Among the total of 1,711 respondents to the questionnaire, almost two-thirds (N=1139) of the respondents had heard about Mpox prior to the study. We observed poor knowledge scores (mean $\pm$ sd) about the transmission pathways (0.71 ± 0.73), vaccination (0.09 ± 0.27), and the signs and symptoms of Mpox (1.91 ± 1.50). Most participants were also unaware of the treatment options of Mpox (0.22 ± 0.59). Educational status and occupation were found to affect the knowledge significantly (p value<0.001). This study showed that the higher the education level, the higher the knowledge level.

Conclusion: The general community has a minimal understanding of the spread of Mpox and its prevention. This virus requires additional research on its epidemiology, ecology, and biology in endemic regions to be comprehended and prevented.

Keywords: Awareness, Bangladesh, Endemic, Knowledge, Mpox, Viral

1. Introduction

Mpox (Monkeypox) is a zoonotic disease caused by infection with the Mpox virus, a member of the Orthopox virus genus (1,2). The virus was first detected in 1958 in the Cynomolgus monkey colony (1). Afterward, in 1971, the first Mpox case was identified in the Democratic Republic of Congo [DRC], and following this, the disease became endemic in central and West Africa (2–4). However, the first incidence outside Africa occurred in the United States in 2003, caused by a rodent imported from Ghana (1,2,4). The detection of cases in the UK and Israel in 2018 and in Singapore in 2019 was documented accordingly (5,6). More than 86,000 cases have been reported in 113 nations, prompting the World Health Organization (WHO) to declare this a public health emergency of international significance (7). This new emergence has more variations compared to its previous presentations such as younger patients, higher prevalence in men, higher rates of sexual transmission, anogenital lesions, an incubation period of up to 21 days, and infection without traveling to an endemic area (8–10). Moreover, this disease has similar presentations to smallpox including fever accompanied by headache, rash, back pain, malaise, and fatigue, except for the specific symptom of lymphadenopathy (1,2).

The virus is transmitted through bushmeat handling, animal hosts, such as squirrels, rodents, and prairie dogs, infected oropharyngeal secretion, and direct contact (1,2,11). Several risk factors, such as the interaction between humans and infected animals, termination of smallpox vaccination, and increased globalization, make the disease a future global public health concern (1,12). Also, the rising occurrence among the younger unprotected population raises concerns about the imminent spread of the human Mpox virus (13).

Although the reason for re-emergence is still unknown, its broad range of animal hosts with high adaptation power and ability to circulate among humans is already making the virus alarming for the future (14). A report by WHO revealed that a lack of information about the Mpox was one of the obstacles in preventing its re-emergence (15). Studies done on general practitioners revealed that only 10% (Indonesia), 18.6% (Saudi Arabia), and 27% (Italy) of the participants had sufficient knowledge about Mpox (16–18). Another study in Saudi Arabia showed that 48% of the general population had sufficient knowledge about Mpox (19). To limit the potential for the Mpox virus spreading across the world, the general public must be educated about the disease and know how to protect themselves. Additionally, due to the dense population, disease transmission is facilitated in Bangladesh (20). Thus, this study

aims to measure the level of awareness and knowledge about Mpox among the general population of Bangladesh, in order to guide future prevention and treatment planning.

2. Methodology

2.1. Study design, settings, and ethics

This was an observational type of cross-sectional study and comprised participants from all the eight divisions (Barishal, Chattogram, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur, and Sylhet) of Bangladesh to provide a nationwide representation. The survey was conducted from May to August 2022, immediately after the confirmation of the ongoing global outbreak of the human Mpox virus in 2022. We followed the simple random sampling method to collect our data. Informed consent was collected from every participant. The anonymity and confidentiality were ensured by excluding the name of the participants. Ethical clearance for the study was taken from the Institutional Review Board (IRB) of North South University.

2.2. Criteria of inclusion and exclusion

The criteria of inclusion for participants were 1) being a Bangladeshi resident, 2) being an adult ($\geq$ 18 years old), and 3) providing consent. The exclusion criteria included- 1) having any psychiatric illnesses, and 2) not providing consent.

2.3. Sample size

Considering an unknown prevalence (50%), 5% margin of error, 95% confidence interval, and 165,158,616 population size a minimum sample size was calculated of 383 (21). However, to increase the diversity and representativeness of the study sample, efforts were made to enrol the maximum number of participants, therefore a final sample size of 1743 was achieved.

2.4. Development and validation of the survey questionnaire

A 20-item semi-structured questionnaire was first developed following Center for Disease Control and Prevention (CDC) guidelines and a study by Harapan H. *et al.* (15,16,22). Then, a pilot study was undertaken on 85 participants to validate the questionnaire. The validation method of the questionnaire included conducting two interviews with the same 85 participants. The second interview was conducted one week after the first one to assess

whether the participants were able to comprehend the questions consistently across both sessions. Finally, the questionnaire was revised, upgraded, and divided into two sections to collect data (Supplementary File 1). The first section consisted of sociodemographic variables, such as gender, age, education, and profession. The second section consisted of 12 questions regarding knowledge of the Mpox outbreak, its signs, symptoms, prevention, and treatment. Moreover, we translated the questionnaire into Bengali for easy understanding.

2.5. Data Collection

All the data was collected via face-to-face interviews. Ten data collectors collected data after one week of training regarding the data collection tools and measures for ensuring optimal participation. It took approximately 10 and 12 minutes to complete the whole questionnaire. The objective of this study was described to all the participants in their native language (Bengali). In addition, including the pilot-study participants, we collected 101 data from Barishal, 309 from Chattogram, 286 from Dhaka, 279 from Khulna, 213 from Mymensingh, 176 from Rajshahi, 219 from Rangpur, and 213 from Sylhet.

2.6. Study variables

In our study, we observed the percentage of persons knowing/hearing the name of Mpox, which is termed as 'awareness'. The known details about the disease is termed as 'knowledge' (23). Knowledge-based questions were for those who at least knew/heard of the name 'Mpox'. For the questions regarding knowledge of Mpox, participants were given 1 for each correct answer and 0 for each incorrect/ do not know ones. For the questions with more than one correct answer, each correct option was given 1 point. However, only for question 12 in Table 3, the scoring was -2 to +2. The highest knowledge score was 26. The mean score per question is demonstrated in Table 3, and the mean percent knowledge score (total knowledge score * 100 / 26) difference concerning sociodemographic variable are in Table 4. We also collected the age (18-35 years/ 36-50 years/ above 50 years), gender (male/ female), education (no formal education/ primary/ secondary/ higher secondary/ graduate/ postgraduate), occupation (unemployed or retired/ student/ homemaker/ daily worker/ service holder/ business/ health professionals), and residence (rural/ urban) as sociodemographic variables.

2.7. Statistical analysis

All the data analysis was done using IBM SPSS v.25. The illustration was prepared by
GraphPad Prism (version 8.0). Descriptive analysis of the sociodemographic variables is
shown in Table 1. Simple and multiple binary logistic regression was used to examine the
odds of demographic variables on being aware of Mpox. Omnibus Tests of Model
Coefficients and Hosmer and Lemeshow Goodness of Fit Test were performed to see the
validity of the regression model. Results were presented as crude and adjusted odds ratios.
Independent-Samples T-test and One-way ANOVA test were conducted to observe the mean
knowledge-score difference among the categories of the sociodemographic information. P
values less than 0.05 were considered significant.

3. Results

3.1. Sociodemographic Information

In total, 1711 respondents completed the questionnaire. Male (N=979/1711; 57.2%) made up
most of the cohort. More than one-third of the respondents (N=1102/1711; 64.4 %) were
between 18 and 35, and more than one-third indicated a graduate degree as their highest
educational attainment (N=664/1711; 38.8 %). Three-fourths of the participants reported
living in an urban area (N=1284/1711; 75.0%), as shown in Table 1.

Table 1: Sociodemographic information of the participants

Sociodemographic information	Category	Number (%)
Age (in years) (N = 1711)	18-35	1102 (64.4%)
	36-50	349 (20.4%)
	Above 50	260 (15.2%)
Gender (N = 1711)	Male	979 (57.2%)
	Female	732 (42.8%)
Educational Status (N = 1711)	No formal education	142 (8.3%)
	Primary	107 (6.3%)
	Secondary	199 (11.6%)
	Higher secondary	346 (20.2%)

	Graduate	664 (38.8%)
	Postgraduate	253 (14.8%)
Occupation (N = 1711)	Unemployed/retired	97 (5.7%)
	Student	593 (34.7%)
	Homemaker	197 (11.5%)
	Daily worker	187 (10.9%)
	Service holder	307 (17.9%)
	Business	212 (12.4%)
	Health professionals	118 (6.9%)
Residence (N = 1711)	Rural	427 (25.0%)
	Urban	1284 (75.0%)

3.2. Sources of knowledge about Mpox

About 61.7% of the respondents (N=1055/1711) were aware of Mpox, while 38.3%
(N=656/1711) were unaware of such virus (Figure 1A). The respondents who were aware of
Mpox also reported the source from which they obtained Human Mpox virus-related
information. Most of the participants acquired their information from social media (N =
508/1055; 48.2%) and the internet (N= 355; 33.6%) followed by Television (340/1055;
32.2%), Newspaper (179/1055; 17.0%), friends/neighbours (148/1055; 14.0%), and family
(60/1055; 5.7%) (Figure 1B). On the contrary, healthcare personnel (N=16/1055; 1.5%) were
the least influential source of information for the participants (Figure 1B).

3.3. Awareness about Mpox

Gender, occupation, and place of residence had statistically significant effects on respondents'
awareness of Mpox (Table 2). Gender of the participants was associated with their awareness
of monkeypox, where males were more aware than females (AOR: 1.4, 95% CI [1-1.8],
p<0.001). Respondents having secondary level (AOR: 4, 95% CI [2.1- 7.4], p<0.001), higher
secondary level (AOR: 8.1, 95% CI [4.2-15.4], p<0.001), graduate level (AOR: 11.2, 95% CI

[5.9-21.4], p<0.001) and post-graduate level (AOR: 9.8, 95% CI [5.0-19.2], p<0.001)

education was significantly associated with awareness and the odds of respondents being

aware of this virus increased with their higher education level. Respondents from urban areas

were more likely to be aware of Mpox (AOR: 1.4, 95% CI [1.7-2.3], p<0.001). (Table 2)

Table 2: Factors affecting the awareness of the respondents about Mpox

Variables	Crude odds ratio (COR)	Range of COR (95% CI)	P-value	Adjusted odds ratio (AOR)	Range of AOR (95% CI)	P-value
Age (in years)						
18-35	2.7	2.0-3.5	<0.001*	1.3	0.9-1.9	0.118
36-50	1.2	0.9-1.7	0.193	1.3	0.9-1.9	0.207
Above 50 (Ref)	1.0.	-	-	1.0	-	-
Gender						
Male	1.0	0.8-1.2	0.972	1.4	1-1.8	0.021*
Female (Ref)	1.0	-	-	1.0	-	-
Educational Status						
No formal education (Ref)	1.0	-	-	1.0	-	-
Primary	3.0	1.6-5.8	0.001*	2.6	1.3-5.1	0.005*
Secondary	5.7	3.2-10.1	<0.001*	4.0	2.1-7.4	<0.001*
Higher secondary	15.4	8.8-26.7	<0.001*	8.1	4.2-15.4	<0.001*
Graduate	23.4	13.7-40.0	<0.001*	11.2	5.9-21.4	<0.001*
Post-graduate	18.5	10.4-32.9	<0.001*	9.8	5.0-19.2	<0.001*
Occupation						
Unemployed/retired (Ref)	1.0	-	-	1.0	-	-
Student	2.1	1.3-3.2	0.002*	1.2	0.7-2	0.533

Homemaker	0.7	0.4-1.1	0.090	0.9	0.5-1.7	0.772
Daily worker	0.2	0.1-0.3	<0.001*	0.6	0.3-1.1	0.094
Service holder	1.6	1.0-2.6	0.040*	1.1	0.6-1.8	0.829
Business	0.8	0.5-1.3	0.371	0.9	0.5-1.6	0.852
Health professionals	2.5	1.3-4.5	0.003*	1.4	0.7-2.6	0.337
Residence						
Rural (Ref)	1.0	-	-	1.0	-	-
Urban	2.8	2.2-3.5	<0.001*	1.7	1.4-2.3	<0.001*

194 *P-value significant; CI: Confidence Interval.

3.4 Knowledge about Mpox

196 A 12-item questionnaire was used to assess knowledge about Mpox among those who were
197 aware of the virus. The distribution of each knowledge item about Mpox and the mean scores
198 against each question are presented in Table 3

199 **Table 3:** Knowledge score assessment of Mpox by individual questions

Knowledge assessment	Scoring (mean ± sd)
1. Knowledge of Mpox prevalence in Africa. (0-1)	0.66 ± 0.47
2. Knowledge regarding the outbreak of human Mpox in Europe, the USA, UK. (0-1)	0.61 ± 0.49
3. Knowledge of Human Mpox cases in Bangladesh. (0-1)	0.57 ± 0.50
4. Knowledge of Mpox as a viral disease. (0-1)	0.68 ± 0.47
5. Knowledge regarding carrier of Mpox virus (squirrels, rat/mice, monkey). (0-3)	0.71 ± 0.73
6. Knowledge of signs and symptoms of Mpox. (0-5)	1.91 ± 1.50
7. Knowledge of similar signs and symptoms of Mpox and smallpox. (0-1)	0.48 ± 0.50
8. Knowledge on the transmission of Mpox. (0-5)	0.86 ± 1.12
9. Knowledge on international travellers as a source of imported cases of Mpox. (0-1)	0.81 ± 0.39

10. Knowledge of prevention of Mpox virus infection. (0-4)	2.00 ± 1.53
11. Knowledge on vaccination against the Mpox virus. (0-1)	0.09 ± 0.27
12. Knowledge regarding the treatment of Mpox virus infection. (-2 - +2)	0.22 ± 0.59

Most of the respondents answered correctly (0.66 ± 0.47) identifying Africa as the most
prevalent region for Mpox. Similarly, the mean score was also 0.61 ± 0.49 about the recent
outbreaks of Mpox in Europe, the USA, and the UK. In regard to any cases of Mpox in
Bangladesh, knowledge of the respondents about this item was 0.57 ± 0.50. The majority of
the participants could identify Mpox as a viral disease, thus the scoring was 0.68 ± 0.47. In
response to the question about the source of Mpox, participants were able to choose one or
more options from squirrels, rat/mice, and monkeys. As all these animals may be responsible
for spreading Mpox, the mean scoring of this item (0.71 ± 0.73) indicates that respondents are
largely unaware of the sources of Mpox (Table 3).

About signs and symptoms of Mpox, participants could select one or more of the following:
fever and headache, rash, muscle discomfort, swollen lymph node, and skin lesion. The mean
score (1.91 ± 1.50) was lower in this issue (Table 3). Most of the participants were unable to
correctly identify that Mpox and smallpox do indeed have similar symptoms (0.48 ± 0.50).
About the mode of transmission, participants could select one or more alternatives from an
animal bite or scratch, wild animal meat preparation/processing, significant respiratory,
indirect contact with lesion material, and mother to fetus. We found that the knowledge level
on this matter was very little (0.86 ± 1.12). The scoring about international travellers as a
source of imported cases of Mpox was 0.81 (± 0.39), indicating that most of the respondents
were aware on this regard (Table 3).

Regarding how they would prevent Mpox, participants were given the option of avoiding
contact with sick animals that could harbor the virus, isolating infected people from others,
avoiding direct contact with infected materials, or practicing good hand hygiene after contact
with infected animals or humans. As all of these are prevention methods of Mpox, the mean
scoring of this item (2.00 ± 1.53) indicates that knowledge level about the transmission
pathways was lower. Almost all respondents were unaware of a vaccine (0.09 ± 0.27).
Additionally, most of the participants were unable to correctly identify the treatments for
Mpox (0.22 ± 0.59) (Table 3).

3.5. Factors that influence knowledge scores

In Table 4, the factors affecting the knowledge score were demonstrated. According to the
findings, individuals with a higher degree of education have considerably higher mean
knowledge scores about Mpox, and the higher the education level, the higher the knowledge
level was.

Table 4: Factors affecting knowledge score regarding Mpox.

Variables	Knowledge percent score	P-value
Age (in years)		
18-35	37.22 ± 19.69	
36-50	36.66 ± 20.45	0.676
Above 50	35.53 ± 20.13	
Gender		
Male	36.57 ± 20.14	
Female	37.43 ± 19.49	0.485
Educational Status		
No formal education	16.74 ± 18.14	
Primary	25.56 ± 17.44	
Secondary	28.91 ± 19.78	
Higher secondary	35.72 ± 19.96	<0.001
Graduate	38.79 ± 19.39	
Post-graduate	41.05 ± 18.80	
Occupation		
Unemployed/retired	36.98 ± 19.43	
Student	35.55 ± 19.63	
Homemaker	29.23 ± 17.40	<0.001
Daily worker	24.86 ± 19.04	

Service holder	41.52 ± 18.40	
Business	32.68 ± 19.88	
Health professionals	51.46 ± 17.05	
Residence		
Rural	36.37 ± 20.85	0.683
Urban	37.06 ± 19.65	

P-value significant

Respondents with a primary level of education had a higher mean percent knowledge score
(25.56 ± 17.44) than those without formal education (16.74 ± 18.14). The mean knowledge
score of respondents with a higher secondary degree (35.72 ± 19.96) was greater than that of
respondents with a secondary degree (28.91 ± 19.78) (Table 4). A higher mean knowledge
score (41.05 ± 18.80) was also obtained by respondents with post-graduate degrees compared
to those with graduate degrees (38.79 ± 19.39). Education and occupational factors
significantly influenced the mean knowledge score (p<0.001). Among all the occupations,
healthcare professionals had the highest mean knowledge score (51.46 ± 17.05).

4. Discussion

The results of the study indicated that the general people lacked essential awareness and
understanding about Mpox. About two-thirds of the respondents heard about Mpox, while
38.3% had never heard of the name- 'Mpox' before completing the survey. In a Malaysian
study, researchers indicated that almost 95% of respondents were aware of dengue fever,
another viral illness (which is also more common), during its epidemic, despite their lack of
understanding of transmission and management (24). A pilot study in Karachi, Pakistan, was
conducted among people that visited hospitals. The findings indicated that 90% of the
participants were aware of the viral disease, among whom only 38.5% possessed adequate
knowledge (25). Similarly, another survey from Pakistan indicated that only 35% of the
adults from low and high socioeconomics groups possessed sufficient knowledge of dengue
during the outbreak (26). Given that dengue is more prevalent than monkeypox, it is not
surprising for our participants to be unfamiliar with the name Mpox. Further investigation is
needed in order to comprehend how to better inform the general population about a newly
discovered, potentially developing virus (27,28).

The news of potential Human Mpox virus cases in Bangladesh spread like wildfire on multiple social media channels. On the other hand, the Bangabandhu Sheikh Mujib Medical University (BSMMU) refuted these assertions, stating that no case of Mpox had yet been identified (29). Given the global discovery of the virus, there is an urgent need to understand the source better, the dynamics of transmission, and to give people the knowledge and assistance they require to safeguard themselves and others in various situations (30). To the best of our knowledge, this is the first nationwide survey in Bangladesh to look into the awareness of general public and understanding of the fundamentals, spread, transmission, symptoms, prevention, and treatment of Human Mpox virus infection.

Most research participants were generally ignorant of the origins of the virus. Moreover, the degree of understanding regarding the transmission channels is also insufficient with only 0.86 mean score according to our findings, while the maximum score for the question was five. A WHO study indicated that one of the difficulties in avoiding the re-emergence of Mpox was a lack of information about Mpox, which is supported by our findings (30). Although the majority of participants in the survey could not identify that Mpox and smallpox have comparable symptoms, almost all of the individuals identified that Mpox is a viral illness that affects them. Vaccination is one of the best ways to prevent infectious illnesses. As there are a few effective antiviral medications for Mpox, in some high-risk circumstances, vaccines to prevent the human smallpox infection (offering 95% Mpox protection), might be utilized (31,32). As per our findings, most of the respondents were unaware about the vaccine and unable to correctly name the treatments as well. Nevertheless, owing to a lack of information, the general population of Bangladesh is mostly in the dark about the identity and diversity of viruses prevalent worldwide.

We also demonstrated that those who use the internet or social media as a source of information are more knowledgeable than their peers about Mpox. This might be because social media and the internet make it simple for most individuals to access information that has been updated. This emphasizes the usefulness of the internet for health promotion, especially in the case of pandemics (33). Compared to other resources, online media have emerged as one of the primary and expedient sources for information access (34). These findings are similar to other studies performed in Egypt and Ethiopia that mentioned the internet and social media as the main information sources (35,36). However, the advent of social media has also accelerated the widespread dissemination of inaccurate data which can

be challenging to detect. On top of that its ability to impact the cognition and convictions of
individuals is evident (37). Newspapers, local leaders, and healthcare professionals appear to
be a less common source of knowledge regarding the ailment when compared to other media.
As a result, the Bangladeshi health system may increase the involvement of community
leaders and health extension workers to disseminate information to the populace efficiently.

In line with the majority of previous research done in developing countries (38), we observed
that men had a higher awareness of this viral illness than females in our study. In developing
countries like Bangladesh, rural males usually have more outside interactions than females.
This might have affected the overall gender-based knowledge differences in this study.
Consistent with previous research (39,40), we found that as the education level of the
participants grew, so did their awareness about the spread of this virus. Furthermore, the
knowledge of people from the urban area was 1.4% higher than that of their rural
counterparts. The outcome of a Chinese study (41) is consistent with this observation, where
the knowledge scores for urban and rural residents were 25.58 ± 3.73 and 24.35 ± 3.95,
respectively. The greater availability of mass media might explain this in urban areas, such as
television, radio, and online social networks (42). While health professionals exhibited the
highest knowledge score in this study, there is still room for further improvement through
targeted training. In situations of endemics or pandemics, healthcare workers consistently
receive guidance to effectively map the outbreak and implement essential interventions aimed
at limiting its transmission (43). The knowledge of general public in this field may also be
influenced by the availability of suitable facilities to support health educators, including
medical personnel and government officials.

The findings of this study may have several implications for public health policymaking and
practice. First, the lack of accurate knowledge of Mpox in the Bangladeshi population amidst
a global public health crisis requires multipronged interventions that promote education and
preventive health behavior. Community-based, social media and mass media interventions
have been effective in health promotion among the Bangladeshi population in the past, which
informs the potential of these media to promote Human Mpox virus knowledge and
prevention (44,45). Second, healthcare providers and social workers need to work with
multisectoral organizations to mobilize organizational resources that ensure optimal
preparedness for potential Human Mpox virus outbreaks and associated health hazards in
Bangladesh. Third, community stakeholders should be engaged in participatory community

prevention of Mpox and other infections that may affect individuals and families living in
diverse Bangladeshi communities. In this process, exploring and utilizing context-specific
knowledge would be critical for successful infection prevention. Fourth, situational analyses
at the community, Upazila, district, and divisional level should be conducted in the context of
Mpox and how a potential outbreak may impact the current state of healthcare and social
services. These insights, alongside other data sources, should be used to create a national-
level data dashboard that informs real-time decision-making for the prevention of Human
Mpox virus outbreaks and other infection outbreaks. News agencies and other public media
can use such a database to generate public knowledge and disseminate the same. Lastly,
policy-level efforts should be made proactively to mitigate the potential crises that may arise
from an outbreak and lessons learned from COVID-19, and Human Mpox virus outbreaks in
similar contexts should be analyzed to inform national and local actions preventing adverse
health outcomes in the future.

Conclusion

This study suggests that the general population of Bangladesh lacks sufficient knowledge
about the source of infection and the dynamics of Mpox transmission. Therefore, considering
the ongoing virus emergence, it is vital to offer the necessary knowledge and support to help
people protect themselves and others. In light of the current global public health crisis,
healthcare professionals, public health workers, and healthcare policymakers should
collaborate to establish a robust disease surveillance system and implement appropriate
policies and interventions that enable effective health education and preventive health
behaviors to promote public awareness, increase the precautionary behaviors and mitigate the
potential crisis.

Strengths and Limitations

This is one of the earliest analyses regarding the knowledge of the general people of
Bangladesh on Human Mpox virus. As it is a preliminary study and due to limited logistic
support, we have been unable to reach out to all the areas in Bangladesh within a short span.
However, a larger sample size would provide more extensive and unbiased (narrower
confidence intervals with more precision in regression models) insight into the current
situation in Bangladesh. Moreover, our data collection plan from each division included
considering the corresponding population density of the divisions. However, this aspect could
not be pursued due to both time and resource limitations. In future research regarding Mpox,
more precise data collection method might be followed.

Author Contribution:

Sudipta Deb Nath: Conceptualization, Data analysis, Writing- Original draft preparation, Writing- Reviewing and Editing

A.M. Khairul Islam: Data analysis, Writing- Original draft preparation, Writing- Reviewing and Editing

Koushik Majumder: Data analysis, Investigation

Fahmida Hoque Rimti: Writing- Original draft preparation, Visualization

Jyoti Das: Writing- Reviewing and Editing, Data curation

Mustari Nailah Tabassum: Writing- Reviewing and Editing

Arefin Naher Oishee: Methodology, Data curation

Tarannum Mahmood: Data curation, Writing- Original draft preparation

Monisha Paul: Data curation, Data analysis

Muntasrina Akhter: Methodology, Visualization

Alok Bijoy Bhadra: Conceptualization, Validation

Fariha Hoque Rimu: Data curation, Writing- Reviewing and Editing

Snahasish Chakraborty: Data curation, Investigation

Preetom Shom: Data curation, Visualization

Sirajum Monira Nosaibah: Methodology, Data curation

Md Ashikur Rahman: Data curation, Investigation

Ahmed Safwan khan: Conceptualization, Data curation

Anika Anjum: Data curation, Visualization

Sushmita khan: Data curation, Investigation

M. Mahbub Hossain: Methodology, Writing- Reviewing and Editing

Mohammad Delwer Hossain Hawlader: Supervision

Declarations:

Competing Interests

The authors have declared that no competing interests exist.

Financial Disclosure

The authors received no specific funding for this work.

References:

1.   Damon IK. Status of human monkeypox: Clinical disease, epidemiology and research.
Vaccine. 2011;29(SUPPL. 4).

2.   Parker S, Nuara A, Buller RML, Schultz DA. Human monkeypox: An emerging
zoonotic disease. Future Microbiol. 2007;2(1).

3.   Damon IK, Roth CE, Chowdhary V. Discovery of Monkeypox in Sudan. N Engl J
Med. 2006;355(9).

4.   Yinka-Ogunleye A, Aruna O, Dalhat M, Ogoina D, McCollum A, Disu Y, et al.
Outbreak of human monkeypox in Nigeria in 2017–18: a clinical and epidemiological
report. Lancet Infect Dis. 2019;19(8).

5.   Yong SEF, Ng OT, Ho ZJM, Mak TM, Marimuthu K, Vasoo S, et al. Imported
monkeypox, Singapore. Emerg Infect Dis. 2020;26(8).

6.   Rao AK, Schulte J, Chen T-H, Hughes CM, Davidson W, Neff JM, et al. Monkeypox
in a Traveler Returning from Nigeria — Dallas, Texas, July 2021 [Internet]. 2022.
Available from: https://wwwnc.cdc.gov/travel/destinations/traveler/none/nigeria?s_cid

7.   World Health Organization. WHO Director-General declares monkeypox outbreak a
public health emergency of international concern - PAHO/WHO | Pan American
Health Organization [Internet]. 2022 [cited 2023 Jul 26]. Available from:
https://www.paho.org/en/news/23-7-2022-who-director-general-declares-monkeypox-
outbreak-public-health-emergency

8.   Miura F, van Ewijk CE, Backer JA, Xiridou M, Franz E, de Coul EO, et al. Estimated
incubation period for monkeypox cases confirmed in the Netherlands, May 2022.
Eurosurveillance. 2022;27(24).

9.   Mileto D, Riva A, Cutrera M, Moschese D, Mancon A, Meroni L, et al. New
challenges in human monkeypox outside Africa: A review and case report from Italy.
Vol. 49, Travel Medicine and Infectious Disease. 2022.

10.   Bragazzi NL, Kong JD, Mahroum N, Tsigalou C, Khamisy-Farah R, Converti M, et al.
Epidemiological trends and clinical features of the ongoing monkeypox epidemic: A
preliminary pooled data analysis and literature review. J Med Virol. 2023;95(1).

11.   Khodakevich L, Jezek Z, Messinger D. Monkeypox virus: Ecology and public health
significance. Bull World Health Organ. 1988;66(6).

12.   Brown K, Leggat PA. Human monkeypox: Current state of knowledge and
implications for the future. Vol. 1, Tropical Medicine and Infectious Disease. 2016.

13.   Reynolds MG, Damon IK. Outbreaks of human monkeypox after cessation of
smallpox vaccination. Vol. 20, Trends in Microbiology. 2012.

14.   Ihekweazu C, Yinka-Ogunleye A, Lule S, Ibrahim A. Importance of epidemiological
research of monkeypox: is incidence increasing? Vol. 18, Expert Review of Anti-
Infective Therapy. 2020.

15.   World Health Organization. Mpox (monkeypox) [Internet]. 2023 [cited 2023 Jul 26].
Available from: https://www.who.int/news-room/fact-
sheets/detail/monkeypox?gclid=CjwKCAjw9NeXBhAMEiwAbaY4li6IGRnvkqzXu0h
ygnapvBTJt4leRNdyZoyq7PSnapHggCi2V-
3SUhoCU5EQAvD_BwE&fbclid=IwAR1xWm1fK0eEMRU9vJIua0fFhvl5pS5Egm_
u1EoKK9v4MKS171OkXK0uPhE

16.   Harapan H, Setiawan AM, Yufika A, Anwar S, Wahyuni S, Asrizal FW, et al.
Knowledge of human monkeypox viral infection among general practitioners: a cross-
sectional study in Indonesia. Pathog Glob Health. 2020;114(2).

17.   Alshahrani NZ, Algethami MR, Alarifi AM, Alzahrani F, Alshehri EA, Alshehri AM,
et al. Knowledge and Attitude Regarding Monkeypox Virus among Physicians in
Saudi Arabia: A Cross-Sectional Study. Vaccines. 2022;10(12).

18.   Riccò M, Ferraro P, Camisa V, Satta E, Zaniboni A, Ranzieri S, et al. When a
Neglected Tropical Disease Goes Global: Knowledge, Attitudes and Practices of
Italian Physicians towards Monkeypox, Preliminary Results. Trop Med Infect Dis.
2022;7(7).

19.   Alshahrani NZ, Alzahrani F, Alarifi AM, Algethami MR, Alhumam MN, Ayied HAM,
et al. Assessment of Knowledge of Monkeypox Viral Infection among the General
Population in Saudi Arabia. Pathogens. 2022;11(8).

20.   Bhuiyan AT, Nath SD, Das S, Das M, Ahsan M, Alam Ziad MIE, et al. Clinical
Observation of Hfnc Use Along With Non-rebreather Mask on Critical Covid-19

Diabetic Patients. Bioresearch Commun. 2023;9(2):1285–96.

21.   Wikipedia. 2022 Bangladeshi census [Internet]. 2022 [cited 2023 Jul 26]. Available
from: https://en.wikipedia.org/wiki/2022_Bangladeshi_census

22.   Centers for Disease Control and Prevention. Mpox | Poxvirus [Internet]. [cited 2023
Jul 26]. Available from: https://www.cdc.gov/poxvirus/mpox/index.html

23.   Nath SD, Chowdhury AS, Pinky SD, Akter KM, Nourin NA, Chowdhury T, et al.
Covariates of Knowledge, Attitude, Practice, and Burdens among the Caregivers of
Hypertensive Patients. Int J Hypertens. 2023;2023(4):1–11.

24.   Naing C, Ren WY, Man CY, Fern KP, Qiqi C, Ning CN, et al. Awareness of dengue
and practice of dengue control among the semi-urban community: A cross sectional
survey. J Community Health. 2011;36(6).

25.   Itrat A, Khan A, Javaid S, Kamal M, Khan H, Javed S, et al. Knowledge, awareness
and practices regarding dengue fever among the adult population of dengue hit
cosmopolitan. PLoS One. 2008;3(7).

26.   Syed M, Saleem T, Syeda UR, Habib M, Zahid R, Bashir A, et al. Knowledge,
attitudes and practices regarding dengue fever among adults of high and low
socioeconomic groups. J Pak Med Assoc. 2010;60(3).

27.   Daily Star. No monkeypox case detected in Bangladesh yet, BSMMU says [Internet].
[cited 2023 Jul 26]. Available from:
https://www.thedailystar.net/health/disease/news/no-Monkeypox-case-detected-
bangladesh-yet-bsmmu-says-3030726

28.   Cheema S, Maisonneuve P, Weber I, Fernandez-Luque L, Abraham A, Alrouh H, et al.
Knowledge and perceptions about Zika virus in a Middle East country. BMC Infect
Dis. 2017 Jul 26;17(1).

29.   World Health Organization. Multi-country monkeypox outbreak: situation update
[Internet]. 2022 [cited 2023 Jul 26]. Available from:
https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON392

30.   World Health Organization. WHO Advisory committee on variola virus research:
report of the nineteenth meeting, 1-2 November 2017, Geneva, Switzerland. World
Health Organization; 2018.

31.   Sherwat A, Brooks JT, Birnkrant D, Kim P. Tecovirimat and the Treatment of
Monkeypox — Past, Present, and Future Considerations. N Engl J Med. 2022;387(7).

32.   World Health Organization. Vaccines and immunization for monkeypox: Interim
guidance, 16 November 2022 [Internet]. 2022 [cited 2023 Aug 18]. Available from:
https://www.who.int/publications/i/item/WHO-MPX-Immunization

33.   Budd J, Miller BS, Manning EM, Lampos V, Zhuang M, Edelstein M, et al. Digital
technologies in the public-health response to COVID-19. Vol. 26, Nature Medicine.
2020.

34.   Souli D, Dilucca M. Knowledge, attitude and practice of secondary school students
toward COVID-19 epidemic in Italy: A cross selectional study. bioRxiv. 2020;2019.

35.   Aynalem YA, Akalu TY, Gebregiorgis BG, Sharew NT, Assefa HK, Shiferaw WS.
Assessment of undergraduate student knowledge, attitude, and practices towards
COVID-19 in Debre Berhan University, Ethiopia. PLoS One. 2021;16(5 May).

36.   Mohamed Ahmed Ayed M, abd Elaziem Mohamed A, Mohamed Mahmoud T,
Mohammed AbdElaziz S. Effect of Educational Intervention on Secondary School
Students' Knowledge, Practices and Attitudes Regarding COVID-19. Egypt J Heal
Care. 2021;12(2).

37.   Majerczak P, Strzelecki A. Trust, Media Credibility, Social Ties, and the Intention to
Share Information Verification in an Age of Fake News. Behav Sci (Basel).
2022;12(2).

38.   Islam FMA, Chakrabarti R, Dirani M, Islam MT, Ormsby G, Wahab M, et al.
Knowledge, attitudes and practice of diabetes in rural Bangladesh: The Bangladesh
Population based Diabetes and Eye Study (BPDES). PLoS One. 2014;9(10).

39.   Deepa M, Bhansali A, Anjana R, Pradeepa R, Joshi S, Joshi P, et al. Knowledge and
awareness of diabetes in urban and rural India: The Indian Council of Medical
Research India Diabetes Study (Phase I): Indian Council of Medical Research India
Diabetes 4. Indian J Endocrinol Metab. 2014;18(3).

40.   Rani PK, Raman R, Subramani S, Perumal G, Kumaramanickavel G, Sharma T.
Knowledge of diabetes and diabetic retinopathy among rural populations in India, and
the influence of knowledge of diabetic retinopathy on attitude and practice. Rural

Remote Health. 2008;8(3).

41. Wen F, Meng Y, Cao H, Xia J, Li H, Qi H, et al. Knowledge, attitudes, practices of primary and middle school students at the outbreak of COVID-19 in Beijing: A cross-sectional online study. medRxiv. 2020;(10).

42. Al ahdab S. A cross-sectional survey of knowledge, attitude and practice (KAP) towards COVID-19 pandemic among the Syrian residents. BMC Public Health. 2021;21(1).

43. Onchonga D. Monkeypox viral disease outbreak in non-endemic countries in 2022: What clinicians and healthcare professionals need to know. Saudi Pharm J. 2022;30(11).

44. Rahman M, Obaida-Nasrin S. Factors affecting acceptance of complete immunization coverage of children under five years in rural Bangladesh. Salud Publica Mex. 2010;52(2).

45. Fazle Rabbi AM. Mass Media Exposure and its Impact on Fertility: Current Scenario of Bangladesh. J Sci Res. 2012;4(2).

Figure 1: Awareness of the respondents about Mpox and their source of knowledge. In

Figure 1A, the pie chart shows if respondents were previously aware of Mpox or not. In

Figure 1B, the bar chart shows the sources of knowledge about Mpox, where the X-axis

denotes the percentage of participants (of those with awareness about Mpox) and the Y-axis

shows the sources.

Supplementary File 1: Questionnaire of the study