Title

Myopia is not a global epidemic: - what can we learn from a longitudinal study conducted in Sweden?

Authors

Pelsin Demir, P.D., MSc (*)
Karthikeyan Baskaran, K.B., PhD (*)
Pedro Lima Ramos, P.L.R., PhD (*, §)
Thomas Naduvilath, T.J.N., PhD (†)
Padmaja Sankaridurg, P.S., PhD (†)
Antonio Filipe Macedo, A.F.M., PhD (*, §)

* Medicine and Optometry, Linnaeus University, Kalmar, Sweden.
† Brien Holden Vision Institute, Sydney, NSW, Australia.
§ Department and Center of Physics – Optometry and Vision Science, University of Minho, Braga, Portugal.

Running title

Incidence of myopia in Swedish schoolchildren

Keywords:

incidence, myopic shift, myopia, environmental factors, parental myopia, refractive error

Abstract

Purpose

The aim of this longitudinal study was to investigate the incidence of myopia, incidence of myopic shift and associated risk factors in Swedish schoolchildren.

Methods

A longitudinal study was conducted between Jan-2019 and June-2021. Participants were Swedish schoolchildren aged 8-16 years. Myopia was defined as spherical equivalent refraction (SER)≤-0.50D. Myopic shift was defined as a minimum change in SER of -0.50D between measurements. The cumulative incidence and incidence rate was computed, Cox proportional hazards regression model and linear mixed models were used to modulate myopic shift and changes in SER.

Results

The study enrolled 128 participants, 86% Caucasian, 5.8% dropped-out during follow-up. The baseline prevalence of myopia was 10.0%, hyperopia was 48.0%, and emmetropia was 42.0%. The cumulative incidence of myopia during the two-years was 5.5%, incidence rate of myopia was 3.2 cases per 100 person-years. The cumulative incidence of myopic shift during the two-years was 21.0%, incidence rate of myopic shift was 12.4 cases per 100 person-years. Cox regression revealed that the probability of myopic shift reduced with "age" and increased with "axial length/corneal-curvature ratio". Myopic children at the baseline and children with two myopic parents showed a significant faster-paced SER change over time.

Conclusions

In the current study the incidence of myopia and myopic shift was low when compared with countries in East Asia. Parental myopia remains a critical factor to consider when predicting myopia progression. Further studies involving children from different ethnicities and incorporating longer follow-up period would further our understanding of the incidence of myopia in Swedish schoolchildren.

Introduction

The prevalence of myopia remains remarkably low in Scandinavia when compared with other parts of the globe. A recent study from Denmark found that the prevalence of myopia has been stable in the past 140 years[1]. Curiously, these findings have been corroborated by recent studies reporting 13.0% (CI95=8.7-18.3) myopia prevalence in Norway[2] and 10.0% (CI95= 4.4-14.9) in Sweden[3]. These findings are in sharp contrast with other regions of the globe where prevalence of myopia amongst children is estimated to be between 50.0% and 71.0%[4-6]. It remains unclear which factors can explain these contrasting results of prevalence of myopia.

Myopia is hereditary and that has been shown in many studies[14, 8, 15, 16]. Some studies showed that the odds ratio of having myopia in childhood with two myopic parents is around 3 when compared to children with no parental myopia.[18.] In addition, myopic children with two myopic parents also exhibited higher myopia (-2.33 D) when compared with children whose parents were not myopic (-1.13 D). In our previous study we also found significantly more negative refraction among children with two myopic parents compared with children whose parents were not myopic[3]. Therefore, it seems that there is a clear genetic susceptibility because parents with myopia tend to have children also with myopia. However, genetics may be only part of the picture and genetic susceptibility might be "activated" or "accelerated" when individuals are exposed to environmental risk factors.

Environmental factors such as near work and outdoor time have been associated with the incidence of myopia[15, 19-21]. For outdoor time, the most common result is that extended outdoor time may delay the onset of myopia[22]. There are currently conflicting results for near work. Some studies point that sustained near work is a risk factor for myopia development and progression while others have failed to confirm this association[15,19, 20, 21,23].

There is currently a lack of studies looking at the incidence of myopia and the progression of refractive errors in Swedish children. Likewise, there is a lack of studies investigating risk factors affecting myopia incidence and progression in Swedish schoolchildren. Having this information is relevant to understand the prevalence of myopia in Sweden and in Scandinavia and to generate new ideas to tackle the high

incidence of myopia in other parts of the globe. Given this gap in the knowledge, this longitudinal study aimed to assess the incidence of myopia, incidence of myopic shift and associated risk factors in Swedish schoolchildren.

Methods

This was longitudinal study involving a cohort of 128 children conducted in southern Sweden from January 2019 to June 2021. Readers are referred to our previously paper on baseline characteristics and recruitment procedures[3]. The study protocol complied with the tenets of the Declaration of Helsinki, informed consent was obtained the participants and their parents. The study received ethics approval from The Regional Committee for Medical Research Ethics in Linköping (Dnr 2018/423-31).

The follow-up period was 2 years and with a total of four visits: baseline, 0.5, 1 and 2 years. An identical comprehensive eye examination was repeated at every visit to determine changes in refractive error and changes in ocular parameters. Details of the methodology has been given in our previous publication[3]. In brief, each visit included measurements of: distance visual acuity, axial length and corneal curvature with noncontact optical coherence biometry (IOLMaster 500 (Carl Zeiss Meditec, AG, Jena, Germany, https://www.zeiss.com/corporate/int/home.html), hight, weight and cycloplegic refraction (NVision-K 5001autorefractor (Shin-Nippon, Rexxam, http://www.shin-nippon.jp/). To ensure paralysis of the ciliary muscle and complete dilation, accommodation and pupil response were verified with a RAF rule and a penlight respectively. Cycloplegic refraction was taken 30 minutes after installation of two drops of cyclopentolate 1% (Cyclogyl, Alcon, https://www.alcon.se/sv).

Questionnaires completed by parents were collected at baseline and follow-up assessments (given in Demir et al.[3]). The questions covered demographics, parental myopia, medical history, academic preferences of the child, education, living conditions, time spent on near work and time spent on outdoor activities after school, and reading habits. Questions were asked in Swedish and the answers were recorded in English. Parental myopia was confirmed by analysis of spectacle prescription.

Definitions and Statistical analysis

The definition of myopia was based on cycloplegic spherical equivalent refraction

(SER) of the right eye: SER= sphere + cylinder/2, cylinder with negative sign. A

participant with SER of -0.50 dioptre (D) or more negative was considered myopic.

Hyperopia was defined as SER +0.75 D or more positive[3]. Myopic shift was defined as

a change of -0.50 D or more in SER between visits (SER at ½ year was subtracted from

SER at baseline, then SER at 1 year was subtracted from SER at ½ year, and SER at 2

years was subtracted from SER at 1 year). Cumulative incidence was computed by

dividing the number of incident cases by the number at risk at baseline using the

equation:

Cumulative incidence = new cases / population at risk.

Incidence rate was computed using the equation:

Incidence rate= new cases during observation period / total person-time of observation

while at risk during study.

Cox proportional hazards regression model was performed to capture all possible

myopic shifts (repeated events) during the 2 years of follow-up. The model included

time-invariant covariates (e.g., "sex") and time-dependent covariates (e.g., "age") for

each participant. Independent variables tested are given in results, the descriptive

statistics in Table 1 and their contributions to the model in Table 2.

The effect of time, ametropia at baseline and parental myopia on SER was tested

using linear mixed models in SAS software PROC MIXED (Release: 3.8 Enterprise

Edition, SAS Institute Inc., Cary, NC, USA). Normalized SER was obtained by

subtracting the baseline SER from all other measures, that corresponds to a baseline

SER of 0 for all participants, that way it was possible to model SER progression. For

this analysis the normalized SER was defined as "dependent variable". Participants

were defined as "random factors". Explanatory factors or "fixed factors" were:

"ametropia" (Myopia, Emmetropia, Hyperopia), "parental myopia" (0,1,2 myopic

parents). Other factors and their interaction with time were tested. P-values were

adjusted for multiple comparisons using the Tukey-Kramer procedure. Means in the text

and shown in graphs are the estimated means (mean response for each factor, adjusted

for any other variables in the model) and their standard errors for the specified factors. Statistical significance was set at $p<0.05$.

Results

Sample characteristics

The cohort was formed of 128 participants, at baseline 70 (54.7%) were females and 58 males (45.3%), during the study seven participants dropped out meaning that 121 participants finished the study, 67 (55.4%) females and 54 (44.6%) males – the retention rate was 95%. The sample was predominantly Caucasian (86.0%), Table 1 summarizes longitudinal values of clinical and sociodemographic variables.

Incidence and prevalence

At baseline, the prevalence of myopia was 10.0% (CI95=4.4-14.9), hyperopia was 48.0% (CI95= 38.8-56.7), and emmetropia was 42.0% (CI95=33.5-51.2). The **cumulative incidence** of myopia during the two years of follow-up was 5.5% (CI95= 2.2-10.9), and the **incidence rate** of myopia was 3.2 cases per 100 person-years (CI95= 0.8-5.6). The **cumulative incidence** of myopic shift during the two years of follow-up was 21.0% (CI95= 14.4-29.2), and the **incidence rate** of myopic shift was 12.4 cases per 100 person-years (CI95= 7.7-17.1).

19 **Table 1. Summary of the key variables for baseline (0), 0.5, 1, and 2 years.**

	Baseline (n=128) Mean (SD) Median (IQR)	0.5 Years (n=126) Mean (SD) Median (IQR)	1 Years (n=114) Mean (SD) Median (IQR)	2 Years (n=121) Mean (SD) Median (IQR)
Age (Years)	12.0 (SD=2.5) 12.0 (IQR=5.0)	12.0 (SD=2.5) 12.5 (IQR=4.0)	12.5 (SD=2.5) 13.0 (IQR=4.0)	13.5 (SD=2.5) 14.0 (IQR=4.0)
SER (Diopters)	+0.65 (SD=1.2) +0.70 (IQR=1.0)	+0.70 (SD=1.3) +0.78 (IQR=1.1)	+0.66 (SD=1.2) +0.67 (IQR=1.1S)	+0.51 (SD=1.5) +0.64 (IQR=1.0)
AL (millimeters)	23.2 (SD=0.86) 23.2 (IQR=1.1)	23.2 (SD=0.87) 23.2 (IQR=1.1)	23.1 (SD=0.82) 23.1 (IQR=1.1)	23.3 (SD=0.92) 23.3 (IQR=1.2)
AL/CR	3.0 (SD=0.1) 3.0 (IQR= 0.1)	3.0 (SD=0.1) 3.0 (IQR= 0.1)	3.0 (SD=0.1) 3.0 (IQR= 0.1)	3.0 (SD=0.1) 3.0 (IQR= 0.1)
Near work (hours-per-day)	5.3 (SD=3.1) 4.6 (IQR=3.3)	4.8 (SD=2.6) 4.4 (IQR=3.2)	2.5 (SD=1.6) 2.2 (IQR=2.1)	2.9 (SD=1.8) 2.7 (IQR=2.4)
Outdoor time (hours-per-day)	2.6 (SD=2.2) 1.9 (IQR=2.1)	2.4 (SD=1.7) 1.8 (IQR=2.0)	1.6 (SD=1.0) 1.4 (IQR=1.3)	1.6 (SD=1.2) 1.3 (IQR=1.6)

SD= standard deviation, IQR= interquartile range, AL/CR = axial length/corneal-curvature ratio, AL = axial length

Cox proportional hazards model of myopic shift during follow-up

Table 2 (Initial Model) summarizes the initial Cox regression model with a full list

of predictors. AL/CR was the only independent predictor of the risk of myopic shift, age

was almost statistically significant. After performing the method of backward

elimination, the final model, given in Table 2 (Final Model), included AL/CR and age

as predictors for myopic shift.

Table 2. Summary of the Cox proportional hazards regression, at the top the initial
model and the bottom the final model. The event modulated was "myopic shift"
and multiple events were possible per subject. The null hypothesis was that all
regression coefficients in the model are equal to zero. When the p-value
(Pr>ChiSq) was less than 0.05 the parameters in the model were considered
different from zero, that is, independent predictors of the myopic shift.

Initial Model						
Parameter	**DF**	**Parameter Estimate**	**Standard Error**	**Chi-Square**	**Pr > ChiSq**	**Hazard Ratio**
AL*		1	-0.10770	0.29841	0.1303	0.7182	0.898
AL/CR*		1	8.56039	2.91571	8.6198	0.0033	5220.735
Age*		1	-0.16173	0.08690	3.4638	0.0627	0.851
Near work*		1	0.00129	0.09554	0.0002	0.9892	1.001
Outdoor time*		1	-0.11138	0.17256	0.4166	0.5186	0.895
Parental myopia**	0	1	-0.85186	0.57567	2.1897	0.1389	0.427
Parental myopia**	1	1	0.05503	0.43181	0.0162	0.8986	1.057
Sex**	Female	1	-0.19994	0.42042	0.2262	0.6344	0.819
Final Model						
Parameter	**DF**	**Parameter Estimate**	**Standard Error**	**Chi-Square**	**Pr > ChiSq**	**Hazard Ratio**
AL/CR		1	8.32555	2.04577	16.5620	<.0001	4128.009
Age		1	-0.19816	0.07839	6.3903	0.0115	0.820

* time-dependent covariates; ** time-invariant covariates

According to the final model in Table 2, "AL/CR" and Age had a statistically

significant effect on the hazard of the event. For each additional unit in age, the hazard

of myopic shift decreased by 100*(0.82-1) =18.0%. Hazard increased by 4128 for every

1 unit in AL/CR. In other words, for each additional 0.01 unit in AL/CR ratio, the

hazard of having an event of myopic shift increased by: $(e^{8.3255/100})$ -1= 8.7%. Figure 1

shows a survival function for three age category groups.

Figure 1. Kaplan-Meier graph assuming AL/CR ratio of a mean value of 2.98. for
different age categories. On the y-axis is the probability of having a myopic shift and on
the x-axis is the time in months since the start of the study. This graph shows different
age categories only to illustrate the effect of age; although, in the Cox regression age
was used as a continuous time-dependent covariate.

Longitudinal changes in SER and associated factors

There was a statistically significant effect of time on SER (F (3, 250) =16.3,
p<0.001). Table 3 summarizes pairwise comparisons with adjusted p-values of SER
between different time points.

Table 3. Summary of pairwise comparisons of SER between time points: baseline (0), 0.5, 1, and 2 years. The mean difference (M) was computed by subtracting SER at baseline in the column from SER in the row. SE= standard errors (SE) and p-values were adjusted with Bonferroni correction for multiple comparisons.

Time	0	0.5	1
0.5	M= 0.004 D (SE=0.04), p=0.063	-	-
1	**M= -0.16 D (SE= 0.05), p=0.005**	**M= -0.15 D (SE= 0.05), p=0.006**	-
2	**M= -0.27 D (SE= 0.04), p<0.001**	**M= -0.26 D (SE= 0.04), p<0.001**	M= -0.11 D (SE= 0.04), p=0.063

There was a significant interaction ametropia×time (F (2, 122) = 6.36, p=0.002), these results are summarized in Table 4 and Figure 2-A. There was a significant interaction time×parental myopia (F (2, 122) = 6.16, p=0.003), these results are summarized in Table 4 and Figure 2-B.

Figure 2. Multiple line graph showing results of the linear mixed model analysis. On the y-axis is the predicted change in SER and on the x-axis is the time in years with measurements performed at: baseline (0 years), 0.5 years, 1 year, 2 years. A) Progression of SER according to ametropia at baseline. B) Progression of SER according to parental myopia (none, one, or two myopic parents). Means in graphs are the estimated means (mean response for each factor, adjusted for any other variables in the model) and error bars are standard error of the mean.

Table 4. Summary of pairwise comparisons of SER changes between categories of ametropia on the left-hand side and parental myopia on the right-hand side at different time points: baseline (0), 0.5, 1, and 2 years. The mean difference (M) was computed by subtracting SER for the time in the first column from SER for the time in the first row. SE= standard errors (SE) and p-values were adjusted for multiple comparisons.

Ametropia	Time	0	0.5	1	Myopic parents	Time	0	0.5	1
Myopia	0.5	M= -0.10 D (SE= 0.10), p=0.286	-	-	none	0.5	M= -0.03 D (SE= 0.07), p=1.00	-	-
	1	**M= -0.30 D (SE= 0.12), p=0.011**	M= -0.20 D (SE= 0.12), p=0.085	-		1	M= 0.02 D (SE= 0.10), p=1.00	M= 0.05 D (SE= 0.01), p=1.00	-
	2	**M= -0.56 D (SE= 0.11), p<0.001**	**M= -0.46 D (SE= 0.10), p<0.001**	M= -0.26 D (SE= 0.11), p=0.365		2	M= -0.11 D (SE= 0.08), p=0.967	M= -0.10 D (SE= 0.07), p=0.993	M= -0.13 D (SE= 0.07), p=0.863
Hyperopia	0.5	M= 0.004 D (SE= 0.05), p=0.947	-	-	1	0.5	M= 0.11 D (SE= 0.06), p=0.866	-	-
	1	M= -0.10 D (SE= 0.06), p=0.931	M= -0.10 D (SE= 0.06), p=0.946	-		1	M= -0.14 D (SE= 0.07), p=0.746	**M= -0.24 D (SE= 0.07), p=0.029**	-
	2	M= -0.16 D (SE= 0.06), p=0.240	M= -0.16 D (SE= 0.05), p=0.153	M= -0.07 D (SE=0.06), p=0.989		2	M= -0.23 D (SE= 0.07), p=0.075	**M= -0.33 D (SE= 0.07), p<0.001**	M= -0.09 D (SE= 0.07), p=0.960
Emmetropia	0.5	**M= 0.09 D (SE= 0.05), p=0.046**	-	-	2	0.5	M= -0.10 D (SE= 0.08), p=0.991	-	-
	1	M= -0.08 D (SE= 0.05), p=0.941	**M= -0.18 D (SE= 0.05), p= 0.001**	-		1	**M= -0.37 D (SE= 0.10), p=0.006**	M= -0.28 D (SE= 0.10), p=0.110	-
	2	M= -0.08 D (SE= 0.05), p=0.956	**M= -0.17 D (SE= 0.05), p=0.020**	M= -0.01 D (SE= 0.05), p=1.00		2	**M= -0.47 D (SE= 0.09), p<0.001**	**M= -0.38 D (SE= 0.08), p<0.001**	M= -0.10 D (SE= 0.08), p=0.986

Discussion

The current study investigated the incidence of myopia and the incidence of myopic shift in a sample of Swedish schoolchildren. The study also investigated factors associated with myopia development and progression. Age and AL/CR showed to be independent predictors of the risk of myopic shift. Ametropia at baseline and parental myopia showed to be factors affecting progression of SER over time.

The two-year cumulative incidence of myopia and the incidence rate of myopia in the current study was consistent with the low baseline prevalence of myopia[3]. The findings of this study are consistent with studies from other Scandinavian countries[1,2,7-9]. A study from Denmark compared the myopia prevalence in 29 Danish studies between the years 1882 and 2018 and found no evidence of an increase in myopia numbers over the years[1]. The prevalence of myopia in Danish children aged 4.5-7 years ranged from 0.0% in year 2015-2016 to 8.9% in schoolchildren aged 7-10 years in year 1882. The same study showed that the prevalence of myopia was 7.2% in Danish children aged 10-14 years in year 1961-1962 and was 24.0% among children aged 11-14 years in year 1882[1]. In contrast with our findings, in East Asian countries the cumulative incidence of myopia ranges between 33.6% and 54%[10], and the annual incidence rate of myopia varies from 30 cases per 100-person years[11] to 31.7 cases per 100-person years[12]. The incidence of myopia tends to be higher in East Asians than in European Caucasians, a possible explanation for this is ethnicity. It is known that, for example, the prevalence of myopia is higher amongst British South Asian children than Caucasian children living in the UK despite their exposure to the same environment[13]. Overall, the incidence of myopia in the current study is lower than in East Asia and is in line with the expectations for Scandinavian populations.

In this study we report for the first time the incidence of the myopic shift in Swedish or Scandinavian children. Given the originality of our findings we can only compare our findings with other parts of the globe. Studies from East Asia have investigated the one-year cumulative incidence of myopic shift which varied between 51% and 74.6%[12,14,15]. These figures are considerably higher than in the current study. The incidence for myopia and myopic shift in our study were remarkable low compared with parts of the global where myopia is also more prevalent. It remains to be

understood which factors explain these differences, but ethnicity, environmental factors
and their interaction are expected to be relevant[16].

Survival analysis showed that age and AL/CR were independent predictors of
myopic shift. The results indicated that age is expected to reduce the probability of
myopic shift, that is. younger children were at a greater risk of having a myopic shift.
These findings are in line with other studies showing that younger age is associated with
a faster progression of myopia when compared to older ages[17-19]. Studies show a decline
in progression speed with increasing age in young myopes of both European and Asian
ethnicity[20,21]. An increase in the AL/CR ratio increased the probability of myopic shift.
The AL/CR ratio has a typical value of 3.0 in emmetropic eyes and more than 3.0 in
myopic eyes[22-25]. Our results are in line with other studies showing that increasing
AL/CR ratio is associated with a more myopic refractive error[25-28]. Recent studies have
shown that AL/CR ratio explains the total variance in refractive error better than AL
alone and is an influential variable for myopia detection in children[27,29]. Our results for
age give further evidence that is important to delay the onset of myopia[30] and our
findings for AL/CR are consistent with the often-successful results of orthokeratology
for myopia management[31].

The longitudinal analysis of changes in SER showed the expected trend in children,
that is, a change towards negative values. The progression was influenced by ametropia
and parental myopia. In the current study the mean change in SER was -0.30 D, that is
approximately -0.15 D per year. Reports from East Asia show a typical progression -
0.60 D per year[32,33] and that is 4-times our progression. These results explain the low
incidence of myopia in our study when compared to East Asia. Our findings showed
that participants with myopia at the baseline had a significant faster SER change during
the study. This results are in line with a recent study reporting myopia progression in
European children with myopia[8]. Another independent factor associated with
progression of SER was parental myopia. Children with two myopic parents had a faster
change towards myopia than the other two categories. Several studies have reported
similar results, showing that parental myopia is associated with a faster rate of myopia
progression[9,17,34].

The current study has two major strengths, the first strength was its longitudinal
design and the second was the low dropout rate. A possible limitation was the poor
recall of the information reported in the questionnaire. Even though, some studies have

shown that key answers tend to be reasonably accurate[35,36]. Another possible limitation

was the sampling method. Since the participants had to travel to the facilities of the

faculty where the study was based, it is unlikely that the sample was a random selection

of the population. This might have attracted overly concerned children and parents only,

but given the low myopia prevalence, this limitation seems to have had limited impact

on the findings. Further, the sample size limits the statistical power to find associations

between environmental factors and myopia. However, our findings are in line with the

most recent suggestions that environmental conditions such as educational stress at a

very young age may impact myopia onset and progression[37]. In the Swedish school

system children start formal school at 7 years of age which means less near work

pressure at very young age when compared to some parts of the world – this might be

the answer to the low incidence and prevalence of myopia reported in this study. Further

studies are warranted for investigating the effect of educational pressure on refractive

error development amongst Swedish children.

Conclusion

In this longitudinal study with a cohort formed mostly by Caucasian children living

in Sweden, the incidence of myopia and myopic shift was low when compared with

other parts of the globe. Younger children were at a greater risk of having a myopic

shift. Myopic children at the baseline and children with two myopic parents showed a

significant faster-paced refractive error change over time. Our results show that parental

myopia remains a critical confounder to consider when planning clinical trials for

myopia control interventions. Further studies are necessary involving children from

different ethnicities living in Sweden under the same environmental conditions to

investigate myopia prevalence and myopia progression patterns.

Acknowledgments

This study was supported by Specsavers Sweden AB, the faculty of Health

and Life Sciences, Linnaeus University, and Brien Holden Vision Institute.

ReferencesReferences

1. Hansen MH, Hvid-Hansen A, Jacobsen N, Kessel L. Myopia prevalence in Denmark - a review of 140 years of myopia research. *Acta Ophthalmol.* 2021;99(2):118-127.
2. Hagen LA, Gjelle JVB, Arnegard S, Pedersen HR, Gilson SJ, Baraas RC. Prevalence and Possible Factors of Myopia in Norwegian Adolescents. *Sci Rep.* 2018;8(1):13479.
3. Demir P, Baskaran K, Theagarayan B, Gierow P, Sankaridurg P, Macedo AF. Refractive error, axial length, environmental and hereditary factors associated with myopia in Swedish children. *Clin Exp Optom.* 2021;104(5):595-601.
4. Nakao SY, Miyake M, Hosoda Y, et al. Myopia Prevalence and Ocular Biometry Features in a General Japanese Population: The Nagahama Study. *Ophthalmology.* 2021;128(4):522-531.
5. Li Y, Liu J, Qi P. The increasing prevalence of myopia in junior high school students in the Haidian District of Beijing, China: a 10-year population-based survey. *BMC Ophthalmol.* 2017;17(1):88.
6. Guo Y, Duan JL, Liu LJ, et al. High myopia in Greater Beijing School Children in 2016. *PLoS One.* 2017;12(11):e0187396.
7. Pärssinen O, Kauppinen M, Viljanen A. The progression of myopia from its onset at age 8-12 to adulthood and the influence of heredity and external factors on myopic progression. A 23-year follow-up study. *Acta Ophthalmol.* 2014;92(8):730-739.
8. Polling JR, Klaver C, Tideman JW. Myopia progression from wearing first glasses to adult age: the DREAM Study. *Br J Ophthalmol.* 2021.
9. Pärssinen O, Kauppinen M. Associations of near work time, watching TV, outdoors time, and parents' myopia with myopia among school children based on 38-year-old historical data. *Acta Ophthalmol.* 2022;100(2):e430-e438.
10. Li L, Zhong H, Li J, Li CR, Pan CW. Incidence of myopia and biometric characteristics of premyopic eyes among Chinese children and adolescents. *BMC Ophthalmol.* 2018;18(1):178.
11. Wang SK, Guo Y, Liao C, et al. Incidence of and Factors Associated With Myopia and High Myopia in Chinese Children, Based on Refraction Without Cycloplegia. *JAMA Ophthalmol.* 2018;136(9):1017-1024.
12. Tsai DC, Fang SY, Huang N, et al. Myopia Development Among Young Schoolchildren: The Myopia Investigation Study in Taipei. *Invest Ophthalmol Vis Sci.* 2016;57(15):6852-6860.
13. Rudnicka AR, Owen CG, Nightingale CM, Cook DG, Whincup PH. Ethnic differences in the prevalence of myopia and ocular biometry in 10- and 11-year-old children: the Child Heart and Health Study in England (CHASE). *Invest Ophthalmol Vis Sci.* 2010;51(12):6270-6276.
14. Wu LJ, Wang YX, You QS, et al. Risk Factors of Myopic Shift among Primary School Children in Beijing, China: A Prospective Study. *Int J Med Sci.* 2015;12(8):633-638.
15. Jin P, Zou H, Xu X, et al. Longitudinal changes in choroidal and retinal thicknesses in children with myopic shift. *Retina.* 2019;39(6):1091-1099.
16. Pärssinen O, Soh ZD, Tan CS, Lanca C, Kauppinen M, Saw SM. Comparison of myopic progression in Finnish and Singaporean children. *Acta Ophthalmol.* 2021;99(2):171-180.
17. Jones-Jordan LA, Sinnott LT, Chu RH, et al. Myopia Progression as a Function of Sex, Age, and Ethnicity. *Invest Ophthalmol Vis Sci.* 2021;62(10):36.
18. Hyman L, Gwiazda J, Hussein M, et al. Relationship of age, sex, and ethnicity with myopia progression and axial elongation in the correction of myopia evaluation trial. *Arch Ophthalmol.* 2005;123(7):977-987.
19. Saw SM, Tong L, Chua WH, et al. Incidence and progression of myopia in Singaporean school children. *Invest Ophthalmol Vis Sci.* 2005;46(1):51-57.

20.   Donovan L, Sankaridurg P, Ho A, et al. Myopia progression in Chinese children is slower in summer than in winter. *Optom Vis Sci.* 2012;89(8):1196-1202.
21.   Morgan IG, Rose KA. Myopia: is the nature-nurture debate finally over? *Clin Exp Optom.* 2019;102(1):3-17.
22.   Grosvenor T, Scott R. Role of the axial length/corneal radius ratio in determining the refractive state of the eye. *Optom Vis Sci.* 1994;71(9):573-579.
23.   Ip JM, Huynh SC, Kifley A, et al. Variation of the contribution from axial length and other oculometric parameters to refraction by age and ethnicity. *Invest Ophthalmol Vis Sci.* 2007;48(10):4846-4853.
24.   Cheng D, Schmid KL, Woo GC. Myopia prevalence in Chinese-Canadian children in an optometric practice. *Optom Vis Sci.* 2007;84(1):21-32.
25.   Scheiman M, Gwiazda J, Zhang Q, et al. Longitudinal changes in corneal curvature and its relationship to axial length in the Correction of Myopia Evaluation Trial (COMET) cohort. *J Optom.* 2016;9(1):13-21.
26.   He X, Sankaridurg P, Naduvilath T, et al. Normative data and percentile curves for axial length and axial length/corneal curvature in Chinese children and adolescents aged 4-18 years. *Br J Ophthalmol.* 2021.
27.   He X, Zou H, Lu L, et al. Axial length/corneal radius ratio: association with refractive state and role on myopia detection combined with visual acuity in Chinese schoolchildren. *PLoS One.* 2015;10(2):e0111766.
28.   Jong M, Sankaridurg P, Naduvilath TJ, Li W, He M. The Relationship between Progression in Axial Length/Corneal Radius of Curvature Ratio and Spherical Equivalent Refractive Error in Myopia. *Optom Vis Sci.* 2018;95(10):921-929.
29.   Tideman JWL, Polling JR, Jaddoe VWV, Vingerling JR, Klaver CCW. Environmental Risk Factors Can Reduce Axial Length Elongation and Myopia Incidence in 6- to 9-Year-Old Children. *Ophthalmology.* 2019;126(1):127-136.
30.   Dhakal R, Shah R, Huntjens B, Verkicharla PK, Lawrenson JG. Time spent outdoors as an intervention for myopia prevention and control in children: an overview of systematic reviews. *Ophthalmic Physiol Opt.* 2022.
31.   Walline JJ, Lindsley KB, Vedula SS, et al. Interventions to slow progression of myopia in children. *Cochrane Database Syst Rev.* 2020;1(1):Cd004916.
32.   Fan DS, Lam DS, Lam RF, et al. Prevalence, incidence, and progression of myopia of school children in Hong Kong. *Invest Ophthalmol Vis Sci.* 2004;45(4):1071-1075.
33.   Tong L, Huang XL, Koh AL, Zhang X, Tan DT, Chua WH. Atropine for the treatment of childhood myopia: effect on myopia progression after cessation of atropine. *Ophthalmology.* 2009;116(3):572-579.
34.   Kurtz D, Hyman L, Gwiazda JE, et al. Role of parental myopia in the progression of myopia and its interaction with treatment in COMET children. *Invest Ophthalmol Vis Sci.* 2007;48(2):562-570.
35.   Breslin KM, O'Donoghue L, Saunders KJ. An investigation into the validity of self-reported classification of refractive error. *Ophthalmic Physiol Opt.* 2014;34(3):346-352.
36.   Ostrin LA, Sajjadi A, Benoit JS. Objectively Measured Light Exposure During School and Summer in Children. *Optom Vis Sci.* 2018;95(4):332-342.
37.   Ding X, Morgan IG, Hu Y, Yuan Z, He M. Exposure to the Life of a School Child Rather Than Age Determines Myopic Shifts in Refraction in School Children. *Investigative Ophthalmology & Visual Science.* 2022;63(3):15-15.