

## Title

**COVID-19 individual participant data meta-analyses. Can there be too many? Results from a rapid systematic review.**

## Authors

Lauren Maxwell\*, PhD

Heidelberger Institut für Global Health

Universitätsklinikum Heidelberg

Im Neuenheimer Feld 130/3

69120 Heidelberg, Germany

[lauren.maxwell@uni-heidelberg.de](mailto:lauren.maxwell@uni-heidelberg.de)

+49 06221 565344

ORCID: 0000-0002-0777-2092

\*Corresponding author

Priya Shreedhar, MSc

Heidelberger Institut für Global Health

Universitätsklinikum Heidelberg

Heidelberg, Germany

[priya.shreedhar@outlook.com](mailto:priya.shreedhar@outlook.com)

ORCID: 0000-0002-1920-2636

Brooke Levis, PhD

Centre for Clinical Epidemiology

Lady Davis Institute for Medical Research

Jewish General Hospital

Montreal, Canada

[b.levis@keele.ac.uk](mailto:b.levis@keele.ac.uk)

ORCID: 0000-0002-4310-3689

Sayali Arvind Chavan, MSc

Institute of Tropical Medicine and Public Health

Charité – Universitätsmedizin Berlin

Berlin, Germany

[sayalic94@gmail.com](mailto:sayalic94@gmail.com)

ORCID: 0000-0002-7908-9276

Shaila Akter, BDS

Heidelberger Institut für Global Health

Universitätsklinikum Heidelberg

Heidelberg, Germany

dr.shailaakter@gmail.com

ORCID: 0000-0002-4362-760X

Mabel Carabali, PhD

Département de Médecine Sociale et Préventive

École de Santé Publique, Université de Montréal

Montreal, Canada

mabel.carabali@umontreal.ca

ORCID: 0000-0002-9171-0483

## KEYWORDS

COVID-19, individual participant data meta-analysis, meta-analysis, data sharing

## OSF REGISTRATION NUMBER

10.17605/OSF.IO/93GF2

## DATA AVAILABILITY STATEMENT

A spreadsheet with comprehensive information on all planned or concluded IPD-MAs described in this review is available on Zenodo (10.5281/zenodo.6623480) under the Creative Commons Attribution 4.0 International (CC BY 4.0) license.

## ABSTRACT

### Background

Individual participant data meta-analyses (IPD-MAs), which include harmonising and analysing participant-level data from related studies, provide several advantages over aggregate data meta-analyses, which pool study-level findings. IPD-MAs are especially important for building and evaluating diagnostic and prognostic models, making them an important tool for informing the research and public health responses to COVID-19.

### Methods

We conducted a rapid systematic review of protocols and publications from planned, ongoing, or completed COVID-19-related IPD-MAs to identify areas of overlap and maximise data request and harmonisation efforts. We searched four databases using a combination of text and MeSH terms. Two independent reviewers determined eligibility at the title-abstract and full-text stage. Data were extracted by one reviewer into a pretested data extraction form and subsequently reviewed by a second reviewer. Data were analysed using a narrative synthesis approach. A formal risk of bias assessment was not conducted.

### Results

We identified 31 COVID-19-related IPD-MAs, including five living IPD-MAs and ten IPD-MAs that limited their inference to published data (e.g., case reports). We found overlap in study designs, populations, exposures, and outcomes of interest. For example, 26 IPD-MAs included RCTs; 17 IPD-MAs were limited to hospitalised patients. Sixteen IPD-MAs focused on evaluating medical treatments, including six IPD-MAs for antivirals, four on antibodies, and two that evaluated convalescent plasma.

### Conclusions

Collaboration across related IPD-MAs can leverage limited resources and expertise by expediting the creation of cross-study participant-level data datasets, which can, in turn, fast-track evidence synthesis for the improved diagnosis and treatment of COVID-19.

## BACKGROUND

The harmonisation and analysis of participant-level data and metadata for cross-study analyses, including individual participant data meta-analyses (IPD-MAs), can inform COVID-19 response through improved evaluation of diagnostic, preventative, and treatment measures. IPD-MAs have several analytic benefits over standard aggregate data meta-analyses when considering analyses of longitudinal data and the development and validation of clinical risk prediction tools.[1-3] IPD-MAs allow for joint consideration of study and subject-level heterogeneity to separate clinically relevant heterogeneity from heterogeneity related to study design or exposure and outcome ascertainment.[1-3] Separating clinically relevant from spurious heterogeneity is central to understanding whether observed differences in the risk of long COVID and COVID-19-related mortality are due to actual differences in exposure or immune response or due to study-level differences in selection, ascertainment, or residual confounding.

The implementation and management of IPD-MAs are resource-intensive.[1, 2, 4] Collecting the well-characterised metadata needed to appropriately describe included studies and cleaning and harmonising participant-level data from related studies require a significant investment of time and expertise from the primary studies and the IPD-MA management team.[2, 5] Additional barriers to sharing participant-level health-related data,[1] including fears of lost opportunities for publication and legal or ethical considerations, can prevent or slow down data sharing.[6-8] IPD-MAs are essential for informing research design, risk communication, and clinical practice for COVID-19. Given the significant resources needed to undertake an IPD-MA, identifying areas of overlap in exposures and outcomes of interest and inclusion criteria can foster cross-IPD-MA coordination to avoid duplication and maximise the utility of existing data.

## METHODS

We conducted a rapid systematic review to identify the protocols for or publications from planned, ongoing or completed COVID-19-related IPD-MAs. We described synergies across these efforts with a focus on study inclusion and exclusion criteria, including study population and study design, and exposure and outcomes of interest. We conducted a systematic search of four databases and protocol repositories, including Ovid Medline, the PROSPERO International Prospective Register of Systematic Reviews, the Open Science Foundation (OSF), and the Cochrane Database of Systematic Reviews, using a combination of MeSH (where applicable) and text terms (supplemental appendix table 1). We ran the searches on 2 June 2021, 29 October 2021, and 7 February 2022.

The protocol for this systematic review was developed per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)-Protocol statement guidelines.[9, 10] Before implementing the searches, we uploaded the systematic review protocol and search strategies to OSF (10.17605/OSF.IO/93GF2) after unsuccessfully trying to upload the protocol to the PROSPERO Registry of Systematic Reviews, which told our team that the systematic review of IPD-MAs was not a systematic review. This systematic review is reported in keeping with the 2020 PRISMA statement (supplemental appendix table 2).[11]

### Study selection and data extraction

Eligible protocols or published studies were IPD-MAs that planned to include or included participant-level COVID-19-related health data. IPD-MAs that only included social or psychological measures and

systematic reviews that were limited to aggregate measures rather than participant-level data from included studies were excluded. Two independent reviewers determined eligibility at the title abstract and full-text screening stages. One reviewer extracted data into a pre-piloted data extraction Google sheet. Data were subsequently reviewed by a second reviewer. Differences of opinion and discrepancies in data extraction were resolved through consensus.

## Analysis

We conducted a narrative synthesis of the results and summarise findings in a series of Sankey diagrams created in RStudio 1.4.1103. We did not include a formal risk of bias assessment as part of this rapid systematic review, as most IPD-MAs only had a protocol available for review at the time of data extraction.

## Patient and public involvement

Patients and the public were not directly involved in this systematic review; we used publicly available data for the analysis.

## RESULTS

We reviewed 116 full texts and identified 31 COVID-19-focused health-related IPD-MAs (see supplemental appendix figure 1 for the PRISMA flow diagram). The majority of IPD-MAs were identified through PROSPERO (n=21), followed by Ovid Medline (n=8) and OSF (n=2).[12, 13] No IPD-MAs were identified from the Cochrane Database of Systematic Reviews. The 31 ongoing or completed COVID-19 IPD-MAs are described in Table 1. As shown in the Sankey diagrams in Figures 1A–D, there were several areas of overlap in included study populations, designs, interventions, and outcomes of interest between ongoing or completed and static or living COVID-19-related IPD-MAs. Figures 1C–D limit inference to the 21 IPD-MAs that requested data from authors, which requires more effort than IPD-MAs of data included in publications.

## Study designs

Ten IPD-MAs included randomised controlled trials (RCTs), non-randomised intervention studies, or longitudinal observational studies; an additional 10 IPD-MAs were limited to RCTs only. Three IPD-MAs included RCTs and longitudinal or cross-sectional observational studies.[14-16] Two IPD-MAs had case reports and case series.[17, 18] One IPD-MA each was limited to case reports,[19] medical records,[12] and case series and longitudinal studies.[20] One IPD-MA included any study design;[21] two others included any study design other than case reports.[22, 23]

## Populations

More than half of the 31 IPD-MAs were conducted with data from hospitalised or intensive care unit (ICU) patients (n=17). Ten IPD-MAs included data from the general population, and two IPD-MAs were limited to children or adolescents.[23, 24] One IPD-MA was conducted with pregnant women[25] and one with older adults and health care workers.[26] Most IPD-MAs were not limited by geography (n=28). One IPD-MA was limited to studies in the US and Canada,[27] another to the US, Europe, and China,[28] and one to China.[29]

## Treatment or exposure

Sixteen IPD-MAs focused on the evaluation of medical treatments, including antivirals (n=6),[13, 16, 29-32] antibodies (n=4),[19, 24, 33, 34] angiotensin-converting-enzyme inhibitors (ACEIs) or angiotensin II receptor blockers (ARBs; n=2),[27, 35] convalescent plasma (n=2),[36, 37] COVID-19 vaccines (n=1),[38] and the Bacillus Calmette–Guérin (BCG)-vaccine (n=1).[26] One IPD-MA focused on extracorporeal membrane oxygenation (ECMO).[14] Two IPD-MAs evaluated any medical or mechanical intervention, including ECMO.[28, 39] One IPD-MA had frailty as the exposure.[40]




Figure 1. Sankey diagrams showing overlap between ongoing or completed and static or living COVID-19 IPD-MAs. **A.** Shows overlap between the focus, included study designs, and type of IPD-MA for all the ongoing or completed IPD-MAs. **B.** Shows overlap between the included study population, interventions/exposures, and outcomes of all the ongoing or completed IPD-MAs. **C.** Shows overlap between the focus, included study designs, and type of IPD-MA for only those IPD-MAs that requested data from authors. **D.** Shows overlap between the included study population, interventions/exposures, and outcomes of only those that requested data from authors.

ACEIs=angiotensin-converting-enzyme inhibitors. ARBs=angiotensin II receptor blockers. BCG=Bacillus Calmette–Guérin. ECMO=extracorporeal membrane oxygenation. GBS=Guillain–Barré syndrome. MIS-C= multisystem inflammatory syndrome in children. Obs=observational. RCTs=randomised controlled trials. RT-PCR=reverse transcription polymerase chain reaction.

## Outcomes

IPD-MAs shared a number of common outcomes, including rate of mechanical (n=8) or non-invasive ventilation (n=2),[31, 40] ECMO rate (n=2),[40, 41] rate of serious adverse events (SAEs) or adverse events (AEs; n=7), viral clearance or viral load (n=4),[13, 16, 30, 39] COVID-19 infection rate (n=3),[15, 26, 38] rate of hospitalization or rehospitalization (n=3)[22, 26, 30] or admittance to the ICU (n=6), time-to-hospital or ICU discharge (n=12), hospital discharge location (n=3),[21, 22, 40] time-to-clinical recovery (n=10), COVID-19 severity score (n=3),[27, 36, 37] quality of life-related measures (n=3),[13, 28, 38] and mortality (n=24). Areas of overlap in mortality measures included IPD-MAs that assessed in-hospital

mortality (n=7) and all-cause mortality (n=5). One IPD-MA assessed ICU mortality,[14] and another pregnancy-related mortality.[25] IPD-MAs that specified time-to-death, included: 14-day mortality (n=2),[27, 36] 28-day mortality (n=2),[13, 36] 30-day mortality (n=2),[12, 27] and 60-day mortality (n=1).[13] Individual IPD-MAs focused on the clinical presentation of long COVID-19[22] and COVID-19 in children;[23] time-to-recovery of smell or taste;[42] incidence of Guillain-Barre syndrome;[18] rate and spectrum of dermatological outcomes in COVID-19 patients;[17] viral load at of sample collection and its effect on the accuracy of RT-PCR;[20] and adverse birth outcomes and vertical transmission in pregnant women with COVID-19.[25]

### **Types of IPD-MAs**

Ten IPD-MAs were limited to published IPD, which means that the group conducting the IPD-MA did not contact authors to request data. Five were living IPD-MAs where datasets and related findings are regularly updated as evidence becomes available.[25, 26, 28, 36, 38] Living IPD-MAs included a real-time IPD-MA,[36] a network IPD-MA,[38] and an IPD-MA of IPD-MAs.[28] Living IPD-MAs focused on COVID-19 vaccines,[38] BCG vaccine,[26] any treatment,[28] convalescent plasma,[36] and issues of interest to perinatal populations.[25] Four of the five living IPD-MAs were limited to RCTs.[26, 28, 36, 38]

### **Availability of data from IPD-MAs**

Fifteen IPD-MAs were published when we submitted the manuscript for publication. Three published IPD-MAs made their data available through Github (n=1)[16] or the journal supplement (n=2).[21, 23] Two published IPD-MAs stated that interested researchers could request the dataset from the study team,[20, 28] five said that data would not be made available.[18, 19, 34, 38, 41] Five others did not include a statement related to data availability.[17, 32, 33, 36, 40] Three of the living IPD-MAs were published [28, 36, 38], although only one indicated that data could be requested from the study team.[28]

**Table 1** Overview of COVID-19-focused IPD-MAs

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
1	Angoulvant[24]	Initial treatment of multisystem inflammatory syndrome in children (MIS-C) and outcomes: a systematic review and meta-analysis of individual patient data; The International MIS-C Treatment Collaborative	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	Children or adolescents	RCTs and non-randomised intervention studies	IVIG plus glucocorticoids or glucocorticoids alone	Cardiovascular dysfunction; mortality; medical intervention rate; mechanical ventilation rate; clinical status; time-to-ICU discharge
2	Antwi-Amoabeng[19]	Clinical outcomes in COVID-19 patients treated with tocilizumab: An individual patient data systematic review	IPD-MA of published IPD only; Completed and published; Data not available	Pharmaceutical treatment or prophylaxis	Hospitalised patients	Case reports	Tocilizumab	In-hospital mortality; incidence of in-hospital complications; time-to-clinical recovery; inflammatory markers
3	Baral[35]	Individual patient data meta-analysis of renin-angiotensin-aldosterone system inhibitors in COVID-19	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	Hospitalised patients	RCTs and non randomised intervention studies, and longitudinal observational studies	ACEIs or ARBs	In-hospital mortality; ICU admission rate; mechanical ventilation rate; time-to-hospital discharge
4	Beyrouti[21]	Characteristics of intracerebral haemorrhage associated with COVID-19: a systematic review and pooled analysis of individual patient and aggregate data	IPD-MA of published IPD only (publication includes AD and IPD); Completed and published; Data included in the publication supplement	COVID-19 outcomes - intracerebral haemorrhage	Hospitalised patients	Any study design	N/A	Mortality, time-to-clinical recovery; discharge location

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
5	Campbell[22]	Predictors of COVID-19 outcomes: an individual participant meta-analysis	IPD-MA; Ongoing	COVID-19 outcomes - long COVID	General population	Any study design other than case reports	N/A	Clinical presentation of long COVID; mortality; rehospitalisation rate; time-to-hospital or ICU discharge, discharge location
6	Cao[29]	Comparative efficacy of treatments for patients infected with 2019 novel coronavirus: a systematic review and meta-analysis of individual patient data	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	General population	RCTs and non randomised intervention studies, and longitudinal observational studies	Lopinavir/Ritonavir	Mortality; time to clinical recovery
7	Cao[31]	Comparative effectiveness and safety of antiviral agents for patients with COVID-19: Protocol for a systematic review and individual patient data network meta-analysis	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	General population	RCTs and non randomised intervention studies, and longitudinal observational studies	Antiviral drugs alone or in any combination, including IFN- $\alpha$ , LPV/r, remdesivir, chloroquine, ribavirin, arbidol, and Xuebijing injection	Time to clinical recovery, all-cause mortality; vitals; mechanical ventilation rate; non-invasive ventilation rate; SAEs
8	Christophers[23]	Trends in Clinical Presentation of Children with COVID-19: A Systematic Review of Individual Participant Data	IPD-MA of published IPD only; Completed and published; Data included in the publication supplement	SARS-CoV-2 infection clinical presentation	Children or adolescents	Any study design other than case reports	N/A	Clinical presentation of COVID-19 in children
9	de Jong[12]	Clinical prediction models for	Living IPD-MA;	COVID-19	Hospitalised	EMRs	N/A	30-day and in-hospital

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
		mortality in COVID-19 patients: a living external validation and individual participant data meta-analysis (COVID-PRECISE)	Ongoing	outcomes - mortality	patients			mortality
10	Dominguez-Rodriguez[14]	Management of mechanical circulatory support during the COVID-19 pandemic: an individual patient data meta-analysis	IPD-MA; Ongoing	Non-pharmaceutical clinical treatment	ICU patients	RCTs and longitudinal and cross-sectional observational studies	ECMO	In-hospital and ICU mortality; time-to-hospital or ICU discharge; incidence of VAP, SAEs
11	Fontes[30]	Chloroquine/hydroxychloroquine for coronavirus disease 2019 (COVID-19) – a systematic review of individual participant data	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	Hospitalised patients	RCTs only	Chloroquine or hydroxychloroquine	COVID-19-related mortality; All-cause mortality; ARDS incidence; hospitalisation rate; ICU admission rate, time to clinical recovery; time to viral clearance, SAEs
12	Gastine[16]	A patient-level meta-analysis on SARS-CoV-2 viral dynamics to model response to antiviral therapies	IPD-MA; Completed and published; Data uploaded to GitHub	Pharmaceutical treatment or prophylaxis	General population	RCTs and longitudinal and cross-sectional observational studies	Antiviral medication	Viral load or clearance
13	Goldfeld[37]	Prospective individual patient data meta-analysis: Evaluating convalescent plasma for COVID-19	IPD-MA; Ongoing	Non-pharmaceutical clinical treatment	Hospitalised patients	RCTs only	Convalescent plasma	Mortality; time to hospital or ICU discharge; COVID-19 severity score
14	Harwood[41]	Which children and young	IPD-MA of	COVID-19	Hospitalised	Cohorts or other	N/A	In-hospital mortality;

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
		people are at higher risk of severe disease and death after hospitalisation with SARS-CoV-2 infection in children and young people: A systematic review and individual patient meta-analysis	published IPD only; Completed and published; Data not available	outcomes - long COVID or mortality	patients	longitudinal observational studies		mechanical ventilation rate; ECMO rate; ICU admission rate
15	Hasan[18]	Guillain-Barré syndrome associated with SARS-CoV-2 infection: A systematic review and individual participant data meta-analysis	IPD-MA of published IPD only; Completed and published; Data not available	COVID-19 outcomes - multiple	General population	Case reports and case series	N/A	Incidence of GBS
16	Hong[39]	Efficacy and safety of therapeutic treatments in patients with COVID-19: a network meta-analysis	IPD-MA; Ongoing	Any treatment	General population	RCTs and non randomised intervention studies, and longitudinal observational studies	Any medical or mechanical intervention	Mortality; time-to-hospital discharge; time-to-clinical recovery; viral load; AEs and SAEs
17	Juul[28]	Interventions for treatment of COVID-19. A living systematic review with individual patient data meta-analyses, aggregate data meta-analyses, trial sequential analyses, and network meta-analysis (The LIVING Project)	Living IPD-MA; Published and ongoing; Data available from emailing authors	Any treatment	Hospitalised patients	RCTs only	Any medical or mechanical intervention	All-cause mortality; ICU admission rate; renal replacement therapy rate; mechanical ventilation rate; QoL; AEs
18	Tan[42]	Prognosis of smell and taste recovery in COVID-19 patients: a systematic review and one-	IPD-MA; Ongoing	COVID-19 outcomes - neurologic	General population	Cohorts or other longitudinal observational	N/A	Time-to-recovery of smell or taste, time taken or extent of

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
		stage meta-analysis of individual patient time-to-event data			studies			improvement of smell or taste
19	Korang[38]	Vaccines to prevent COVID-19: a protocol for a living systematic review with network meta-analysis including individual patient data (The LIVING VACCINE Project)	Living IPD-MA; Published and ongoing; Data not available	Vaccine efficacy	General population (limited to those who were never infected with SARS-CoV-2)	RCTs only	Any COVID-19 vaccine	All-cause mortality; COVID-19 incidence; QoL
20	Lant[15]	Neurological associations of COVID-19 (COVID-Neuro): A protocol for a systematic review and meta-analysis of individual patient data	IPD-MA; Ongoing	COVID-19 outcomes - neurologic	Hospitalised patients	RCTs and longitudinal and cross-sectional observational studies	N/A	Infection rate; in-hospital mortality; time-to-ICU discharge; time-to-hospital discharge; mechanical ventilation rate; ICU admission rate
21	Lee[17]	Cutaneous manifestations of COVID-19: a systematic review and analysis of individual patient-level data	IPD-MA of published IPD only; Completed and published; No information on data availability	COVID-19 outcomes - cutaneous	General population	Case reports and case series	N/A	Rate and spectrum of dermatological outcomes
22	Ling[34]	Interleukin-6 receptor antagonists for severe coronavirus disease 2019: a meta-analysis of individual participant data from randomised controlled trials	IPD-MA of published IPD only; Completed and published; Data not available	Non-pharmaceutical clinical treatment	Hospitalised patients	RCTs only	IL-6 inhibitors	Time-to-clinical recovery; mortality

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
23	Mallett[20]	At what times during infection is SARS-CoV-2 detectable and no longer detectable using RT-PCR-based tests? A systematic review of individual participant data	IPD-MA of published IPD only; Completed and published; Data available from emailing authors	SARS-CoV-2 infection diagnosis	Hospitalised patients	Case series and longitudinal studies	N/A	Timing of sample collection for accurate SARS-CoV-2 diagnosis by RT-PCR
24	Simmons[32]	Sofosbuvir/daclatasvir regimens for the treatment of COVID-19: an individual patient data meta-analysis	IPD-MA; Completed and published; No information on data availability	Pharmaceutical treatment or prophylaxis	Hospitalised patients	RCTs and non-randomised intervention studies	Sofosbuvir/daclatasvir-based regimens	Clinical recovery within 14 days of randomisation; time-to-clinical recovery; all-cause mortality; time-to-hospital discharge; composite outcome of ICU admission or requirement for invasive mechanical ventilation
25	Smith[25]	Protocol for a sequential, prospective meta-analyses <sup>27</sup> to rapidly address priority perinatal COVID-19 questions	Living IPD-MA; Ongoing	COVID-19 outcomes - multiple	Pregnant women	RCTs and longitudinal observational studies	N/A	Adverse birth outcomes; pregnancy-related mortality and morbidity; vertical transmission rate of COVID-19
26	Speich[13]	Efficacy and safety of remdesivir in hospitalised patients with COVID-19: Systematic review and individual patient data meta-analysis of randomised trials	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	Hospitalised patients	RCTs only	Remdesivir	28- and 60-day mortality; mechanical ventilation rate; duration of mechanical ventilation; clinical

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
								status; time-to-clinical recovery; time-to-ICU or hospital discharge; QoL; Viral load or clearance; AEs or SAEs
27	Subramaniam[40]	Characteristics and Outcomes of Patients with Frailty Admitted to ICU with Coronavirus Disease 2019: An Individual Patient Data Meta-Analysis	IPD-MA; Completed and published; No information on data availability	COVID-19 outcomes - multiple	ICU patients	Cohorts or other longitudinal observational study only	Frailty	In-hospital mortality; time-to-hospital or ICU discharge; mechanical ventilation rate; non-invasive ventilation rate; ECMO rate; renal replacement therapy rate, vasoactive infusion rate; ICU bed occupancy; discharge location
28	Tasoudis[33]	Survival analysis of IL-6 inhibitors versus standard of care for COVID-19: a meta-analysis of individual patient data from randomised trials	IPD-MA of published IPD only; Completed and published; No information on data availability	Pharmaceutical treatment or prophylaxis	General population	RCTs only	IL-6 inhibitors	Mortality; ICU admission rate; hospital discharge rate; mechanical ventilation rate
29	Troxel[36]	Association of Convalescent Plasma Treatment With Clinical Status in Patients Hospitalised With COVID-19: A Meta-analysis	Living IPD-MA; Published and ongoing; No information on data availability	Non-pharmaceutical clinical treatment	Hospitalised patients	RCTs only	Convalescent plasma	14- and 28-day mortality; COVID-19 severity score
30	van	Anytime Live and Leading	Living IPD-MA;	Vaccine	Health care	RCTs only	BCG-vaccine	COVID-19 incidence;

	First author, last name	Title	Type, status, and availability of data for IPD-MA	Focus	Population	Study design	Treatment/exposure	Outcome(s)
	Werkhoven[26]	Interim* meta-analysis of the impact of BCG vaccine in health care workers and elderly during the SARS-CoV-2 pandemic (ALL-IN-META-BCG-CORONA)	Ongoing	efficacy	workers; elderly			hospitalisation rate; infection rate; time to clinical recovery; time to hospital discharge
31	Victory[27]	ACEi/ARB medications for hospitalised patients with COVID-19: an individual patient data (IPD)-based pooled analysis	IPD-MA; Ongoing	Pharmaceutical treatment or prophylaxis	Hospitalised patients	RCTs only	ACEIs or ARBs	COVID-19 severity score; time to hospital discharge; duration of mechanical ventilation; 14-day mortality; 30-day mortality, AEs or SAEs

ACEIs=angiotensin-converting-enzyme inhibitors. AD=aggregate data. AE, adverse event. ARBs=angiotensin II receptor blockers; ARDS=acute respiratory distress syndrome. BCG=Bacillus Calmette-Guérin. ECMO=extracorporeal membrane oxygenation. EMRs, electronic medical records. GBS=Guillain-Barré syndrome. ICU=intensive care unit. IL-6=Interleukin 6. IPD=individual participant data. IPD-MA=individual participant data meta-analysis. IVIG=intravenous immunoglobulins. MIS-C=multisystem inflammatory syndrome in children. N/A=not applicable. QoL=quality of life. RCT=randomized controlled trial. RT-PCR=reverse transcription polymerase chain reaction, SAE=serious adverse event. SARS-CoV-2=severe acute respiratory syndrome coronavirus. VAP=ventilator-associated pneumonia.

## DISCUSSION

IPD-MAs are an important tool for the rapid evidence generation needed to inform clinical practice, making them a vital part of the research response to emerging pathogens.[43] We conducted a rapid systematic review to identify ongoing or completed COVID-19-related IPD-MAs. There were many areas of overlap in the 31 COVID-19-related IPD-MAs, including in study design and population, exposure, and outcomes of interest. In particular, the 14 IPD-MAs that evaluated the same medical exposures (antivirals, antibodies, ACEIs and ARBs, and convalescent plasma) represent a missed opportunity to exploit synergies. Most IPD-MA protocols were registered on PROSPERO, which could flag these areas of overlap when researchers submit their protocol. IPD-MAs require a significant investment of time and expertise, both from the team conducting the IPD-MA and the groups contributing data to the IPD-MA. Rapidly identifying and exploiting shared inclusion criteria can help facilitate evidence generation and avoid unnecessary duplication of effort.

We identified at least 10 IPD-MAs that limited their analysis to data included in published reports. While IPD-MAs that are limited to published IPD have been conducted previously, the volume of the research response to COVID-19 coupled with the push for reproducibility and transparency have likely facilitated the rise in IPD-MAs of data that were included in the study publications. Almost half of the IPD-MAs of published data included case study or case series data (n=4/10; 40%).[17-20] Given that the utility of the IPD-MA is limited by the quality of the studies that contribute data,[2] findings from these rapidly produced IPD-MAs should be considered preliminary and updated when more detailed and less selective participant-level datasets become available. This finding is in keeping with a methodological review of published data that compared the methodological and reporting quality of COVID-19 and non-pandemic research and found a reduction in quality in the former.[44]

While we reviewed the protocols for all IPD-MAs, we could only identify the restriction to published IPD for those IPD-MAs that had published their analyses, which suggests a need to clarify inclusion criteria in IPD-MA protocols to specify the intent to limit inference to published IPD. Some of the unpublished studies identified in our review may be misclassified as having the classical approach to conducting an IPD-MA, which includes the challenges associated with requesting the data from the data producers.

Living IPD-MAs are regularly updated as more evidence becomes available, representing substantial investments. There was overlap in study design, exposure, and outcome measurements in several of the five living IPD-MAs and between the living IPD-MAs and static IPD-MAs, which represents an opportunity to share limited resources and expedite findings.

Only a few IPD-MAs of data received from authors had been published when this manuscript was submitted for publication (n=5/21; 24%), so we could not quantify the overlap in datasets across IPD-MAs that collected datasets from research teams which would be an important measure of cross-IPD-MA redundancy in efforts. Only three of the ten published IPD-MAs had made data available through a repository or the publication of supplementary materials,[16, 21, 23] which suggests a continued need to encourage data sharing.

Working collaboratively to harmonise and share data across related IPD-MAs would maximise limited resources and shorten the timeline to deliver results that best inform clinical and public health practice.

Testing the same hypotheses, especially with the same study designs or populations, represents a missed opportunity to evaluate novel hypotheses. Our findings support similar calls from a living review of COVID-19-related clinical trials and a scoping review of COVID-19-related data sharing platforms, which urged coordination across initiatives to reduce redundancies.[45] We propose the creation of a task force to identify concrete steps to enable cross-initiative collaboration and ensure that the harmonised participant-level data and study-related metadata correspond to the findable, accessible, interoperable, reusable (FAIR) principles for data resources.[46] These steps could include a cross-platform algorithm that uses natural language processing to alert researchers to similar initiatives at the time of protocol deposition.

We have listed additional resources facilitate the design and conduct of an IPD-MA in the context of an emerging pathogen in Table 2. The global scope and rapidly evolving nature of the pandemic underscore the need for more meta-collaborations that work to bring together data sharing efforts and cross-national analyses. The coordination of ongoing or planned IPD-MAs is a good starting place.

**Table 2** Resources for the conduct of an IPD-MA of an emerging pathogen

Focus	Description	Link and citation
Design & conduct of IPD-MAs	Website containing step-by-step guidance, videos, articles, and other material for the conduct of an IPD-MA	Richard Riley. Individual Participant Data (IPD) Meta-Analysis. Available at: <a href="http://www.ipdma.co.uk">www.ipdma.co.uk</a> .
Design & conduct of IPD-MAs	Virtual toolbox containing practical resources and information needed to carry out all stages of an IPD-MA	Levis B, Holden MA, Hattle M, Riley R, van Der Windt D, Campbell L, Fulton S, Ensor J. The Keele Toolbox for IPD Meta-analyses. Available at: <a href="https://www.keele.ac.uk/ipdmatoobox/">https://www.keele.ac.uk/ipdmatoobox/</a>
Planning & management of IPD-MAs	Virtual toolkit containing templates and resources for IPD-MA-related project management, data cleaning, and harmonisation-related tasks	Maxwell L, Shreedhar P, Carabali M, & Levis B. Templates and resources for individual participant data meta-analyses (IPD-MAs). Available at: <a href="https://doi.org/10.17605/OSF.IO/3FXVY">https://doi.org/10.17605/OSF.IO/3FXVY</a>
Creation & execution of harmonisation code	Open-source software packages, Mica & Opal, to facilitate harmonisation of IPD and dissemination of related metadata	Software available at: <a href="https://www.maelstrom-research.org">https://www.maelstrom-research.org</a> .
Guidance for IPD-MA in the context of prediction modelling	Framework and guidance documents regarding IPD-MAs of clinical prediction models	COVID-19 PRECISE. IPD Meta-Analysis. Available at: <a href="https://www.covprecise.org/ipdma/">https://www.covprecise.org/ipdma/</a> .

IPD-MAs=individual participant data meta-analyses; PRECISE=Precise Risk Estimation to optimise COVID-19 Care for Infected or Suspected patients in diverse sEttings.

## CONCLUSION

IPD-MAs are important for informed research and public health response to COVID-19. To identify areas of overlap, we conducted a rapid systematic review of completed or ongoing COVID-19 IPD-MAs. We identified 31 COVID-19-related IPD-MAs, including five living IPD-MAs, and found several areas of overlap in study designs, populations, exposures, and outcomes of interest. This review shows several potential areas of collaboration across related IPD-MAs which can leverage limited resources and expertise by expediting the creation of cross-study participant-level datasets. This, in turn, can fast-track evidence synthesis for the improved diagnosis and treatment of COVID-19.

## CONTRIBUTORS

LM, BL, and MC conceived of and designed the study. LM wrote the research protocol and developed the search strategy. LM, SA, and SC conducted the title abstract and full-text screening. LM, MC, PS, SA, and SC extracted and interpreted the data. LM wrote the first draft of the manuscript. All authors provided critical reviews of the manuscript. All authors had full access to all study data, take responsibility for data integrity and reliability of the analysis, and had final responsibility for the decision to submit for publication.

## COMPETING INTERESTS

None declared.

## FUNDING

ReCoDID project, funded by the EU Horizon 2020 research and innovation programme (grant agreement 825746) and the CIHR Institute of Genetics (grant agreement 01886-000) grant to LM.

## ROLE OF THE FUNDING SOURCE

The study's funders had no role in the study design, data collection, data analysis, data interpretation, or report writing. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit this manuscript for publication.

## REFERENCES

1. Tudur Smith C, Marcucci M, Nolan SJ, et al. Individual participant data meta-analyses compared with meta-analyses based on aggregate data. *Cochrane Database Syst Rev* 2016;9:MR000007.
2. Riley RD, Lambert PC, Abo-Zaid G. Meta-analysis of individual participant data: rationale, conduct, and reporting. *BMJ* 2010;340:c221.
3. Wilder-Smith A, Wei Y, Araújo TVBd, et al. Understanding the relation between Zika virus infection during pregnancy and adverse fetal, infant and child outcomes: a protocol for a systematic review and individual participant data meta-analysis of longitudinal studies of pregnant women and their infants and children. *BMJ Open* 2019;9:e026092.
4. Stewart LA, Tierney JF. To IPD or not to IPD?:Advantages and Disadvantages of Systematic Reviews Using Individual Patient Data. *Eval Health Prof* 2002;25:76–97.
5. Tierney JF, Vale C, Riley R, et al. Individual Participant Data (IPD) Meta-analyses of Randomised Controlled Trials: Guidance on Their Use. *PLoS Med* 2015;12:e1001855.
6. Schwalbe N, Wahl B, Song J, et al. Data Sharing and Global Public Health: Defining What We Mean by Data. *Front Digit Health* 2020;2:612339.
7. Jao I, Kombe F, Mwalukore S, et al. Research Stakeholders' Views on Benefits and Challenges for Public Health Research Data Sharing in Kenya: The Importance of Trust and Social Relations. *PLoS One* 2015;10:e0135545
8. van Panhuis WG, Paul P, Emerson C, et al. A systematic review of barriers to data sharing in public health. *BMC Public Health* 2014;14:1144.
9. Moher D, Shamseer L, Clarke M, et al. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
10. Shamseer L, Moher D, Clarke M, et al. Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;349:g7647.
11. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71.
12. de Jong VMT, Debray TPA. Clinical prediction models for mortality in COVID-19 patients: an external validation and individual participant data meta-analysis (COVID-PRECISE). OSF. 2021. Available at: <https://osf.io/96wgn/> (accessed June 9 2022).
13. Speich B, Schönenberger C, Griessbach A, et al. Efficacy and safety of remdesivir in hospitalized patients with COVID-19: Systematic review and individual patient data meta-analysis of randomized trials. OSF. 2021. Available at: <https://osf.io/7a4wf> (accessed June 9 2022).
14. Dominguez-Rodriguez A, Baez-Ferrer N, Abreu-Gonzalez P, et al. Management of mechanical circulatory support during the COVID-19 pandemic: an individual patient data meta-analysis. PROSPERO. 2020. Available at:  
[https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020203402](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020203402) (accessed June 9 2022).
15. Lant S, Cividini S, Singh B, et al. Neurological associations of COVID-19 (COVID-Neuro): A protocol for a systematic review and meta-analysis of individual patient data. PROSPERO. 2021. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020196542](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020196542) (accessed June 9 2022).
16. Gastine S, Pang J, Boshier FAT, et al. Systematic Review and Patient-Level Meta-Analysis of SARS-CoV-2 Viral Dynamics to Model Response to Antiviral Therapies. *Clin Pharmacol Ther* 2021;110:321–333.

17. Lee DS, Mirmirani P, McCleskey PE, et al. Cutaneous manifestations of COVID-19: a systematic review and analysis of individual patient-level data. *Dermatol Online J* 2020;26:13030/qt7s34p8rw.
18. Hasan I, Saif-Ur-Rahman KM, Hayat S, et al. Guillain-Barré syndrome associated with SARS-CoV-2 infection: A systematic review and individual participant data meta-analysis. *J Peripher Nerv Syst* 2020;25:335–343.
19. Antwi-Amoabeng D, Kanji Z, Ford B, et al. Clinical outcomes in COVID-19 patients treated with tocilizumab: An individual patient data systematic review. *J Med Virol* 2020;92:2516–2522.
20. Mallett S, Allen AJ, Graziadio S, et al. At what times during infection is SARS-CoV-2 detectable and no longer detectable using RT-PCR-based tests? A systematic review of individual participant data. *BMC Med* 2020;18:346.
21. Beyrouti R, Best JG, Chandratheva A, et al. Characteristics of intracerebral haemorrhage associated with COVID-19: a systematic review and pooled analysis of individual patient and aggregate data. *J Neurol* 2021;268:3105–15.
22. Campbell P, Ali M, Mason H, et al. Predictors of COVID-19 outcomes: an individual participant meta-analysis. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020224323](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020224323) (accessed June 9 2022).
23. Christophers B, Gallo Marin B, Oliva R, et al. Trends in Clinical Presentation of Children with COVID-19: A Systematic Review of Individual Participant Data. *Pediatr Res* 2022;91:494–501.
24. Angoulvant F, Levin M, McArdle A, et al. Initial treatment of multisystem inflammatory syndrome in children (MIS-C) and outcomes: a systematic review and meta-analysis of individual patient data; The International MIS-C Treatment Collaborative. PROSPERO. 2021. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42021292162](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021292162) (accessed June 9 2022).
25. Smith E. Protocol for a sequential, prospective meta-analyses (PMA) to rapidly addressing priority perinatal COVID-19 questions. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020188955](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020188955) (accessed June 9 2022).
26. van Werkhoven CH, Schure J, Bonten MJM, et al. Anytime Live and Leading Interim\* meta-analysis of the impact of Bacillus Calmette-Guérin vaccination in health care workers and elderly during the SARS-CoV-2 pandemic (ALL-IN-META-BCG-CORONA). PROSPERO. 2021. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42021213069](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021213069) (accessed June 9 2022).
27. Victory J, Freilich D, Hanley D, et al. ACEi/ARB medications for hospitalised patients with COVID-19: an individual patient data (IPD)-based pooled analysis. PROSPERO. 2022. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42021267770](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021267770) (accessed June 9 2022).
28. Juul S, Nielsen EE, Feinberg J, et al. Interventions for treatment of COVID-19: Second edition of a living systematic review with meta-analyses and trial sequential analyses (The LIVING Project). *PLoS One* 2021;16:e0248132.
29. Cao Y, Shan J, Gong Z, et al. Comparative efficacy of treatments for patients infected with 2019 novel coronavirus: a systematic review and meta-analysis of individual patient data. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020167038](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020167038) (accessed June 9 2022).

30. Fontes LE, Riera R, Martimbianco ALC, et al. Chloroquine/hydroxychloroquine for coronavirus disease 2019 (COVID-19) – a systematic review of individual participant data. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020178667](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020178667) (accessed June 9 2022).
31. Cao Y, Shan J, Gong Z, et al. Comparative effectiveness and safety of antiviral agents for patients with COVID-19: Protocol for a systematic review and individual-patient data network meta-analysis. *PLoS One* 2020;15:e0241947.
32. Simmons B, Wentzel H, Mobarak S, et al. Sofosbuvir/daclatasvir regimens for the treatment of COVID-19: an individual patient data meta-analysis. *J Antimicrob Chemother* 2021;76:286–291.
33. Tasoudis PT, Arvaniti CK, Adamou AT, et al. Interleukin-6 inhibitors reduce mortality in coronavirus disease-2019: An individual patient data meta-analysis from randomised controlled trials. *Eur J Intern Med* 2022;S0953-6205(22)00140-6.
34. Ling RR, Ramanathan K, Tan WQ, et al. Interleukin-6 Receptor Antagonists for Severe Coronavirus Disease 2019: A Meta-Analysis of Reconstructed Individual Participant Data from Randomised Controlled Trials. SSRN. 2021. Available at: <http://dx.doi.org/10.2139/ssrn.3844782> (accessed June 9 2022).
35. Baral R, Vassiliou V. Individual patient data meta-analysis of renin-angiotensin-aldosterone system inhibitors in COVID-19. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020208447](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020208447) (accessed June 9 2022).
36. Troxel AB, Petkova E, Goldfeld K, et al. Association of Convalescent Plasma Treatment With Clinical Status in Patients Hospitalised With COVID-19: A Meta-analysis. *JAMA Netw Open* 2022;5:e2147331.
37. Goldfeld KS, Wu D, Tarpey T, et al. Prospective individual patient data meta-analysis: Evaluating convalescent plasma for COVID-19. *Stat Med* 2021;40:5131–5151.
38. Korang SK, von Rohden E, Veroniki AA, et al. Vaccines to prevent COVID-19: a protocol for a living systematic review with network meta-analysis including individual patient data (The LIVING VACCINE Project). *PLoS One* 2022;17:e0260733.
39. Hong H, Halabi S. Efficacy and safety of therapeutic treatments in patients with COVID-19: a network meta-analysis. PROSPERO. 2020. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42020182500](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020182500) (accessed June 9 2022).
40. Subramaniam A, Anstey C, Curtis JR, et al. Characteristics and Outcomes of Patients With Frailty Admitted to ICU With Coronavirus Disease 2019: An Individual Patient Data Meta-Analysis. *Crit Care Explor* 2022;4:e0616.
41. Harwood R, Yan H, Talawila Da Camara N, et al. Which children and young people are at higher risk of severe disease and death after hospitalisation with SARS-CoV-2 infection in children and young people: A systematic review and individual patient meta-analysis. *EClinicalMedicine* 2022;44:101287.
42. Tan BKJ, Han R, Zhao JJ, et al. Prognosis & persistence of smell & taste dysfunction in COVID-19: a meta-analysis with parametric cure modelling of recovery curves. PROSPERO. 2021. Available at: [https://www.crd.york.ac.uk/prospero/display\\_record.php?ID=CRD42021283922](https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021283922) (accessed June 9 2022).

43. Zika Virus Individual Participant Data Consortium. The Zika Virus Individual Participant Data Consortium: A Global Initiative to Estimate the Effects of Exposure to Zika Virus during Pregnancy on Adverse Fetal, Infant, and Child Health Outcomes. *Trop Med Infect Dis* 2020;5:152.
44. Quinn TJ, Burton JK, Carter B, et al. Following the science? Comparison of methodological and reporting quality of covid-19 and other research from the first wave of the pandemic. *BMC Medicine* 2021;19:46.
45. Nguyen VT, Rivière P, Ripoll P, et al. Research response to coronavirus disease 2019 needed better coordination and collaboration: a living mapping of registered trials. *Journal of Clinical Epidemiology* 2021;130:107–16.
46. Wilkinson M, Dumontier M, Aalbersberg I, et al. The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data* 2016;3:160018.