1 **Title (108 characters): Childhood obesity is linked to putative neuroinflammation in brain white**
2 **matter, hypothalamus, and striatum**
3
4 **Running title (45 characters):** Brain microstructure in children with obesity
5

6 **Authors:** Zhaolong Li, BA[1, 2], Amjad Samara, MD[1, 3], Mary Katherine Ray, PhD[1], Jerrel Rutlin, BS[1],
7 Cyrus A. Raji, MD, PhD[3, 4], Joshua S. Shimony, MD, PhD[4], Peng Sun, PhD[4, a], Sheng-Kwei Song, PhD[4],
8 Tamara Hershey, PhD[1, 2, 3, 4, *], Sarah A. Eisenstein, PhD[1, 4]
9
Affiliations:
[1]Department of Psychiatry, Washington University in St. Louis School of Medicine, St. Louis, MO
63110, USA
[2]Department of Psychological and Brain Sciences, Washington University in St. Louis, St. Louis, MO
63130, USA
[3]Department of Neurology, Washington University in St. Louis School of Medicine, St. Louis, MO
63110, USA
[4]Mallinckrodt Institute of Radiology, Washington University in St. Louis School of Medicine, St. Louis,
MO 63110, USA
[a]Present address: Department of Imaging Physics, University of Texas MD Anderson Cancer Center,
Houston, TX 77030, USA
21
*Corresponding author:
Tamara Hershey, PhD
James S. McDonnell Professor of Cognitive Neuroscience
Department of Psychiatry and Mallinckrodt Institute of Radiology
Washington University in St. Louis School of Medicine
St. Louis, MO 63110
MSC 8134-0070-02
Phone: (314)-362-5593
Email: tammy@wustl.edu

31
32 **Keywords (5)**: diffusion MRI; hypothalamus; neuroinflammation; obesity; white matter
33
Word count in main text: 6080

Abbreviations
ABCD[®], Adolescent Brain Cognitive Development[SM]
ADC, apparent diffusion coefficient
ANOVA, analysis of variance
AUC, area-under-the-curve
BMI, body mass index
CI, confidence interval
DBSI, diffusion basis spectrum imaging
DTI, diffusion tensor imaging
DWI, diffusion-weighted image
FA, fractional anisotropy
FF, fiber fraction
FSL, FMRIB Software Library
FWE, family-wise error
ICV, intracranial volume
MRI, magnetic resonance imaging
NW, children with normal-weight
OB, children with obesity
OW, children with overweight
PDS, pubertal development stage
RF, restricted fraction
RNI, restricted normalized isotropic
RSI, restriction spectrum imaging
SD, standard deviation
SES, socioeconomic status
TBSS, tract-based spatial statistics
TFCE, threshold-free cluster enhancement
WC, waist circumference
WM, white matter

Abstract (200 words)

Neuroinflammation is both a consequence and driver of overfeeding and weight gain in rodent obesity

models. Advances in magnetic resonance imaging (MRI) enable investigations of brain microstructure

that suggests neuroinflammation in human obesity. To assess the convergent validity across MRI

techniques and extend previous findings, we used diffusion basis spectrum imaging (DBSI) to

characterize obesity-associated alterations in brain microstructure in 601 children (age 9-11 years) from

the Adolescent Brain Cognitive Development[SM] Study. Compared to children with normal-weight, greater

DBSI restricted fraction (RF), reflecting neuroinflammation-related cellularity, was seen in widespread

white matter in children with overweight and obesity. Greater DBSI-RF in hypothalamus, caudate

nucleus, putamen, and, in particular, nucleus accumbens, correlated with higher baseline body mass index

(BMI) and related anthropometrics. Comparable findings were seen in the striatum with a previously

reported restriction spectrum imaging (RSI) model. Gain in waist circumference over one and two years

related, at nominal significance, to greater baseline RSI-assessed restricted diffusion in nucleus

accumbens and caudate nucleus, and DBSI-RF in hypothalamus, respectively. Here we demonstrate that

childhood obesity is associated with microstructural alterations in white matter, hypothalamus, and

striatum. Our results also support the reproducibility, across MRI methods, of findings of obesity-related

putative neuroinflammation in children.

1. Introduction

Childhood obesity is a major growing health issue, affecting over 340 million children worldwide

in 2016 (World Health Organization, 2021). It is associated with expensive medical costs (Biener et al.,

2020), lower quality of life (Killedar et al., 2020), and elevated risk for health complications including

adult obesity, type 2 diabetes, and cardiovascular diseases (Liang et al., 2015; Simmonds et al., 2016).

Accumulating evidence also identifies childhood obesity as a risk factor for cognitive dysfunction and

Alzheimer's disease in late-life (Tait et al., 2022). Given the brain's prominent role in regulating feeding

and metabolism, it is essential to understand the relationship between obesity and brain health.

Determining which brain regions and networks might be involved in the development and maintenance of

childhood obesity could help identify targets for obesity prevention and intervention, thereby mitigating

short and long-term health consequences.

Obesity involves a chronic, low-grade, systemic inflammation affecting multiple organs (Gregor

& Hotamisligil, 2011). In rodent models of obesity, high-fat diets induce inflammation in the central

nervous system, or "neuroinflammation" (Baufeld et al., 2016; Buckman et al., 2013; De Souza et al.,

2005; Décarie-Spain et al., 2018; Valdearcos et al., 2017), which in turn cause memory deficits and

anxiodepressive behaviors (Beilharz et al., 2016; Décarie-Spain et al., 2018; Pistell et al., 2010). In

humans, post-mortem tissue analyses have revealed associations between obesity and increased gliosis in

multiple brain regions, including the hypothalamus, a key regulator of feeding and metabolism (Baufeld

et al., 2016; Schur et al., 2015). Aimed at assessing brain health *in vivo*, a number of magnetic resonance

imaging (MRI) studies have reported associations between obesity and altered brain structure. In adults,

higher body mass index (BMI) and visceral fat are consistently linked to lower cortical thickness and

smaller prefrontal and basal ganglia volumes (Fernández-Andújar et al., 2021; Gómez-Apo et al., 2021;

Raji et al., 2010; Willette & Kapogiannis, 2015), potentially due to neuronal loss consequent of obesity-

related neuroinflammation and/or microangiopathy (Gómez-Apo et al., 2021). These relationships are less

clear in children (Willette & Kapogiannis, 2015). Adult obesity has also been associated with

compromised white matter integrity, reflected by lower diffusion tensor imaging (DTI)-derived fractional

anisotropy (FA) and greater mean diffusivity of water, primarily in frontolimbic tracts and the corpus

callosum (Daoust et al., 2021; Kullmann et al., 2015; Verstynen et al., 2012). However, opposite findings

of greater white matter DTI-FA in obesity have also been noted (Birdsill et al., 2017; Carbine et al., 2020;

Dekkers et al., 2019), and the relationship between obesity and white matter integrity in children remains

unknown. Importantly, the standard single-tensor DTI model could be confounded by neuroinflammatory

processes such as cellularity and edema (Kullmann et al., 2016; Wang et al., 2011, 2015; Winklewski et

al., 2018), which may partially explain the mixed pattern of results.

In recent years, studies using multi-compartment diffusion MRI-based methods, though limited in

number, have yielded consistent observations of putative neuroinflammation in feeding-related brain

regions in obesity (Rapuano et al., 2020, 2022; Samara et al., 2020, 2021). The data-driven multi-tensor

diffusion basis spectrum imaging (DBSI) technique models diffusion-weighted signals as a linear

combination of discrete anisotropic tensors and isotropic diffusion spectra, enabling the *in vivo*

assessment of brain microstructure (Cross & Song, 2017; Wang et al., 2011, 2015). DBSI metrics, though

indirectly reflecting true anatomy, have been histopathologically validated as neuroinflammation-

sensitive using rodent and human neural tissue in multiple sclerosis (Chiang et al., 2014; Wang et al.,

2014; Wang et al., 2011, 2015), epilepsy (Zhan et al., 2018), and optic neuritis (Lin et al., 2017; Yang et

al., 2021). Notably, applying DBSI to adults with obesity, we previously observed microstructural

alterations in striatal and limbic regions that suggest cellularity, vasogenic edema, and lower apparent

axonal and dendritic densities (Samara et al., 2020, 2021), in line with the obesity-related

neuroinflammatory phenotype seen in animal and post-mortem human brain studies. In white matter

tracts, we found evidence of increased and widespread DBSI-assessed putative neuroinflammation in

young and middle-aged adults with obesity across two independent samples (Samara et al., 2020). DBSI

has not yet been used to characterize brain microstructure in childhood obesity. However, Rapuano et al.,

(2020) used restriction spectrum imaging (RSI), which in contrast to DBSI, models isotropic water

diffusion components based on the ratio of radial and axial diffusivities (Palmer et al., 2022; White et al.,

2013), and observed associations between greater purported striatal cellular density and higher baseline

and future waist circumference and BMI in children in the Adolescent Brain Cognitive Development[SM]

(ABCD) Study (Rapuano et al., 2020, 2022). Furthermore, using a non-diffusion method, namely

quantitative T2-weighted MRI, studies have reported that longer hypothalamic T2 relaxation time and

greater T2 signal intensity, both suggestive of reactive microglial and astrocytic gliosis, relate to higher

BMI in adults (Schur et al., 2015; Thaler et al., 2012) and children, including a subset from the ABCD

Study[®] (Sewaybricker et al., 2019; Sewaybricker, Kee, et al., 2021; Sewaybricker, Melhorn, et al., 2021).

Convergent findings amongst MRI methods in the same group of children would support the feasibility

and reliability of these techniques to assess putative neuroinflammation in childhood obesity.

141   In this study, we used the baseline ABCD Study[®] data from 601 children aged 9-11 years (see

section 2.1 for details on sample selection) to test the *a priori* hypotheses that 1) obesity-associated

microstructural alterations, including greater putative neuroinflammation-related cellularity (reflected by

greater DBSI restricted fraction (RF)) and lower axonal and dendritic densities (reflected by lower DBSI

fiber fraction (FF)) that we had observed in white matter and striatum in adults, and in one novel region

not yet assessed using diffusion MRI, i.e., the hypothalamus, would also be present in children, and that

2) greater hypothalamic and striatal cellularity (DBSI-RF) would relate to greater baseline waist

circumference and BMI metrics in children, similar to the RSI cellular density metric, namely restricted

normalized isotropic (RSI-RNI). We also explored associations between baseline DBSI and RSI metrics

in the hypothalamus and striatum and one and two-year longitudinal changes in anthropometrics. If our

results using DBSI are consistent to those in studies that used RSI and quantitative T2-weighted MRI,

they will support the use of non-invasive MRI-based methods to characterize obesity-related putative

neuroinflammation *in vivo* in humans, in the absence of histopathological validation.

2. Materials and methods

2.1. Participants

157   Participants were from the ABCD Study[®], a ten-year, 21-site study tracking brain development in

a diverse cohort of U.S. children and adolescents (Casey et al., 2018; Garavan et al., 2018; Jernigan et al.,

2018). Participants receive annual physical, sociocultural, and behavioral assessments, as well as

neuroimaging and bioassays every two years. Institutional review boards at study sites approved study

procedures; parents/caregivers provided written consent and children gave verbal assent. The ABCD

Study® 2.0.1 release included data from 11,875 participants at baseline and 4,951 participants at one-year

follow-up. In addition to the ABCD Study® inclusion/exclusion criteria (Garavan et al., 2018), we

excluded participants with 1) missing anthropometric or demographic data at baseline or one-year follow-

up; 2) current or past diagnosis of neurological (including cerebral palsy, brain tumor, stroke, aneurysm,

brain hemorrhage, intellectual disability, lead poisoning, muscular dystrophy, multiple sclerosis, and

others) and psychiatric (including schizophrenia, autism spectrum disorder, attention-deficit hyperactivity

disorder, and others) conditions and diabetes, similar to Rapuano et al., (2020); and 3) T1 or diffusion-

weighted images (DWIs) that did not pass quality control or had clinically significant incidental findings

(Hagler et al., 2019; Li et al., 2021). Also consistent with Rapuano et al., (2020), in order to maximize

harmonization of MRI data across sites, only scans performed on Siemens 3T Prisma platforms (Siemens

Healthineers AG, Erlangen, Germany) were included. As the ABCD Study® 4.0 release became available

during our study, we further included participants with complete data at two-year follow-up to extend

exploratory longitudinal analyses. Lastly, because head motion during MRI scans is known to interfere

with diffusion tensor model estimation and give spurious correlations (Ling et al., 2012; Yendiki et al.,

2014), we excluded participants with excessive head motion (defined as mean DWI framewise

displacement $\geq$ 2.5 mm) and covaried for mean head motion in statistical analyses.

Our inclusion/exclusion criteria selected a total of 1,613 qualifying participants (see

Supplementary Fig. 1 for flowchart). Age and sex-adjusted BMI percentiles at baseline were used to

classify participants by weight status (Kuczmarski et al., 2002), including 63 with underweight (BMI < 5th

percentile), 1,140 with normal-weight (NW; 5th to < 85th percentiles), 194 with overweight (OW; 85th to <

95th percentiles), and 216 with obesity (OB; $\geq$ 95th percentile). To achieve balanced group sizes as well as

reduce computational cost, we randomly selected 216 NW participants (matched to OB group size)

stratified by sex, and included all 194 OW and 216 OB participants. After neuroimaging processing, data

from 25 participants were excluded due to missing/incomplete T1 or DWI acquisition, missing field

maps, mismatch between field map and DWI dimensions, or missing/unclear DWI directions. The final

analytical sample therefore included 212 NW, 187 OW, and 202 OB participants, for a total $n = 601$.

Such sample size is similar to those in recent literature and should afford sufficient power to detect

obesity-related microstructural alterations (see **Supplementary Methods** for power analysis) (Jiang et al.,

2023; Sewaybricker, Kee, et al., 2021).

2.2. Obesity-related measures

Participant waist circumference (WC), weight, and height were measured at baseline and one and

two-year follow-ups (Barch et al., 2018). Raw BMI was calculated (weight$_{(lbs)}$/height$_{(in)}^2 \times 703$). BMI z-

scores corrected for age and sex were computed using the 2000 CDC growth charts (Kuczmarski et al.,

2002). These different measures were used to address the concern that a single index may be less

reflective of true adiposity and/or sensitive to fat gain in children (Cole et al., 2005; Taylor et al., 2000).

2.3. Neuroimaging

2.3.1. MRI acquisition

Details on T1 and DWI acquisition and harmonization across sites are published elsewhere

(Casey et al., 2018; Hagler et al., 2019). T1-weighted anatomical images were collected as a 3D T1-

weighted inversion prepared RF-spoiled gradient echo scan, with voxel resolution = 1 mm^3 isotropic. Spin

echo echo-planar imaging was used to acquire multi-shell DWIs with the following parameters: total

acquisition time = 7:31, repetition time = 4100 ms, time to echo = 88 ms, matrix size = $140 \times 140 \times 81$,

flip angle = 90°, acceleration factor = 3, and voxel resolution = 1.7 mm^3 isotropic. DWIs were imaged

with 7 b = 0 frames and 96 gradient directions (b's = 500, 1000, 2000, and 3000 s/mm^2 with 6, 15, 15,

and 60 directions, respectively).

2.3.2. DWI and DBSI processing

DWIs were corrected for susceptibility-induced distortion, eddy currents, and head motion using

FMRIB Software Library (FSL) *topup* and *eddy* (Smith et al., 2004). Multi-tensor DBSI maps were

estimated using an in-house script as previously described (Wang et al., 2011, 2015). Leveraging the

multi-shell DWI data, DBSI characterizes brain tissue microstructure by partitioning the total water

diffusion signal within each image voxel into isotropic and anisotropic compartments. DBSI modeling

produced maps of anisotropic fiber fraction (DBSI-FF; reflects axonal/dendritic density), isotropic

nonrestricted fraction (f(D) at apparent diffusion coefficient (ADC) > 0.3 μm^2/ms; reflects vasogenic

edema/tissue disintegration/extracellular water), and isotropic restricted fraction (DBSI-RF; f(D) at 0 <

ADC ≤ 0.3 μm^2/ms; reflects intracellular water/inflammation-related cellularity) (Chiang et al., 2014; Sun

et al., 2020; Wang et al., 2015). Details on DBSI model specification are provided in **Supplementary**

Methods. Notably, DBSI-FF and RF are consistently lower and greater, respectively, in adult obesity

(Samara et al., 2020, 2021) and serve as the neuroinflammation-related microstructural assessment in the

current study. DBSI maps were registered to T1 space first using *epi_reg* and a non-diffusion-weighted

image, then by applying the transformation matrix to individual maps using *applyxfm*.

2.3.3. Tract-based spatial statistics (TBSS)

Voxel-wise analyses of white matter DBSI-FF and RF were performed using TBSS (Smith et al.,

2006). The DTI model was fitted to preprocessed DWIs using FSL *dtifit*, and DTI-FA maps were eroded

by one voxel with end slices removed. Cleaned DTI-FA images were nonlinearly registered to the T1-

weighted image of a randomly selected NW participant, averaged, and assigned a threshold at FA > 0.2 to

create a white matter skeleton, onto which the DBSI-FF and RF maps were projected.

2.3.4. Segmentation of the striatum and hypothalamus

The nucleus accumbens, caudate nucleus, and putamen were segmented from T1-weighted

images using FSL *FIRST* (Patenaude et al., 2011). The hypothalamus was segmented using a novel,

automated algorithm developed with deep convolutional neural networks trained on adult data (Billot et

al., 2020). To assess the algorithm's accuracy in children, we compared automated and manual

hypothalamus segmentations in 20 participants (10 NW and 10 OB, randomly selected within each

group). Within this group, the automated and manual segmentations had good spatial overlap (mean Dice

similarity coefficient = 0.74, *SD* = 0.02, one-tailed $p < 0.001$ against the conventional threshold of 0.7)

and yielded highly correlated volumes ($r = 0.74$, $p < 0.001$). Neither spatial overlap nor volumetric

correlation between the automated and manual segmentations was different by weight group (NW vs. OB;

p's = 0.58 and 0.97). Although the automated segmentations had smaller volumes than manual

segmentations (*means* = 747 and 887 mm^3, $p < 0.001$), such volume reduction primarily excluded voxels

near the hypothalamic surface, reducing possible contamination of diffusion signal from neighboring

cerebrospinal fluid and vasculature (**Supplementary Fig. 2**). Also, the segmented volumes were

consistent with literature values (Neudorfer et al., 2020). Taken together, the automated algorithm reliably

produced hypothalamus segmentations comparable to manual segmentation. For each subcortical

structure, segmentations were visually inspected for accuracy before statistical analyses, and volume and

DBSI-FF and RF metrics were each extracted and combined/averaged between hemispheres.

2.4. Statistical analyses

All analyses, except for TBSS, were performed in R version 4.2.1 (R Core Team, 2013).

Differences in participant characteristics across NW, OW, and OB groups were assessed using analysis of

variance (ANOVA) or chi-square tests.

2.4.1. White matter

For TBSS, we excluded data from 28 randomly selected siblings, eliminating family dependency

confounds. Baseline DBSI-FF and RF in white mater tracts were compared amongst unrelated NW (n =

202), OW (n = 180), and OB (n = 191) participants using voxel-wise TBSS, first by ANOVAs for main

effects of group and second by t-tests for between-group comparisons. FSL *Randomize* (null distribution

built from 10,000 permutations; with recommended threshold-free cluster enhancement (TFCE)) was

used for these comparisons with spatial family-wise error (FWE) rate corrected at two-tailed $p \leq 0.05$)

(Winkler et al., 2014). Briefly, the raw statistical image was TFCE-transformed into an output image in

which voxel-wise TFCE scores were weighted sums of local clustered signals, such that larger TFCE

scores reflected magnitude of cluster-like spatial support greater than a given height (signal intensity) (Li

et al., 2017; Smith & Nichols, 2009). We specified the –T2 option in *Randomize* (2D optimization for

skeletonized data, cluster height weighted by $H = 2$, cluster extent weighted by $E = 1$, voxel connectivity

= 26). Voxel-wise analyses using TBSS and TFCE allowed for sensitive detection of regionally-specific

obesity-related DBSI-FF and RF effects in white matter, while stringently controlling for multiple

comparisons across space. Participant age, sex, race/ethnicity, parental education, household income,

parental marital status, pubertal development stage (PDS), mean head motion, and intracranial volume

(ICV) were covaried in TBSS. Group differences in white matter skeleton-average values of DBSI-FF

and RF were assessed with linear mixed-effects models using the *lme4* package (Bates et al., 2015),

where the same set of covariates plus weight group were fixed effects and site was the random effect.

2.4.2. Striatum and hypothalamus

DBSI-FF and RF outliers in the nucleus accumbens, caudate nucleus, putamen, and hypothalamus

that were $\pm$ 3 SD away from the mean were removed (**Supplementary Table 1**). One and two-year

changes in obesity-related measures (i.e., WC, BMI, and BMI *z*-scores) were calculated by subtracting

baseline from respective follow-up. Extreme BMI values (< 10 kg/m^2 or > 50 kg/m^2) and associated BMI

z-scores were removed, including 1 NW and 1 OB at one-year and 1 OW at two-year. Distributions for

obesity-related measures at and changes between all timepoints are shown in **Supplementary Fig. 3**.

Associations between DBSI metrics and baseline or future change in obesity-related measures

were assessed using linear mixed-effects models. Age (at baseline, one, or two-year), sex, race/ethnicity,

PDS (at baseline, one, or two-year), parental education, household income, parental marital status, mean

head motion, and ICV were covaried due to potential confounding (Lawrence et al., 2022; Li et al., 2023;

Palmer et al., 2022; Rapuano et al., 2020), and the random effect was family nested within sites. In

longitudinal models, baseline obesity-related measures were also covaried. As we had *a priori*

hypotheses, and the goal was to describe regionally-specific relations between tissue microstructure and

convergent obesity-related measures, multiple comparisons were corrected with each structure treated as a

family, at two-tailed $p = 0.05$ / (4 regions $\times$ 2 DBSI metrics) = 0.00625. Effect size estimates were

standardized β's with 95% confidence intervals (CIs) and partial R^2's. Models were checked for normality

of residuals, homoscedasticity, and low multicollinearity (variance inflation factors were ≤ 2.56). As there

were missing data following outlier removal, sample sizes varied and are reported in individual analyses.

2.4.3. Comparison between DBSI and RSI

Mean RSI restricted normalized isotropic (RSI-RNI) metrics in bilateral nucleus accumbens,

caudate nucleus, and putamen were obtained from the ABCD Study® tabulated dataset (Hagler et al.,

2019). The ABCD Study® segmented structures using FreeSurfer v5.3; the hypothalamus was not

specifically segmented and voxel-wise RSI-RNI maps were not available. RSI reflects cellularity as an

increase in the restricted isotropic (originating from intracellular water) diffusion signal, i.e., RSI-RNI

(Rapuano et al., 2020, 2022). Associations between RSI-RNI and baseline or future change in obesity-

related measures were evaluated using linear mixed-effects models, as in DBSI described in **section 2.4.2**.

To further compare model performance, DBSI-RF and RSI-RNI from the nucleus accumbens, caudate

nucleus, and putamen were each tested on classifying NW and OB participants using mixed-effects

logistic regression, with the same fixed and random effect covariates in linear models. Receiver operating

characteristic curves and areas-under-the-curve (AUCs) with 95% CIs were computed using the *pROC*

package, and AUCs from DBSI-RF and RSI-RNI were compared using DeLong's test (Robin et al.,

2011).

3. Results

3.1. Sample characteristics

Participant demographics, neuroimaging metrics, and obesity-related measures are described in

Table 1. Qualitatively, the OW and/or OB groups compared to the NW group had more non-White

participants, more advanced pubertal development, lower parental education, household income, and

proportion of married parents, higher baseline obesity-related measures, and greater one and two-year

gain in WC but decrease in BMI *z*-scores. Groups did not differ significantly in striatal or hypothalamic

volumes; these volumes were thus not covaried in addition to ICV in analyses.

Variable	Group (total $n = 601$)			p-value
	NW	OW	OB	
n	212 (35.3% of all)	187 (31.1% of all)	202 (33.6% of all)	N/A
Age (months)	121 ± 8	121 ± 7	120 ± 7	0.89
Sex				
Male	113 (53.3%)	98 (52.4%)	120 (59.4%)	0.31
Female	99 (46.7%)	89 (47.6%)	82 (40.6%)	
Race/ethnicity				
Asian	1 (0.5%)	1 (0.5%)	4 (2.0%)	
Black	16 (7.5%)	23 (12.3%)	27 (13.4%)	
Hispanic	25 (11.8%)	37 (19.8%)	54 (26.7%)	< 0.001 ***
White	156 (73.6%)	112 (59.9%)	99 (49.0%)	
Other	14 (6.6%)	14 (7.5%)	18 (8.9%)	
PDS category				
1	129 (60.8%)	76 (40.6%)	86 (42.6%)	
2	53 (25.0%)	49 (26.2%)	51 (25.2%)	< 0.001 ***
3+	30 (14.2%)	62 (33.2%)	65 (32.2%)	
Parental education				
No HS diploma	0 (0%)	1 (0.5%)	4 (2.0%)	
HS diploma/ GED	7 (3.3%)	5 (2.7%)	20 (9.9%)	
Some college	14 (6.6%)	28 (15.0%)	42 (20.8%)	< 0.001 **
Bachelor	97 (45.8%)	87 (46.5%)	84 (41.6%)	
Postgraduate	94 (44.3%)	66 (35.3%)	52 (25.7%)	
Household income				
< 50k	32 (15.1%)	45 (24.1%)	62 (30.7%)	
≥ 50k & < 100k	64 (30.2%)	74 (39.6%)	74 (36.6%)	< 0.001 ***
≥ 100k	116 (54.7%)	68 (36.4%)	66 (32.7%)	
Parental marriage				
Married	164 (77.4%)	128 (68.4%)	127 (62.9%)	0.005 **
Not married	48 (22.6%)	59 (31.6%)	75 (37.1%)	
Mean motion (mm)	1.23 ± 0.25	1.27 ± 0.31	1.27 ± 0.30	0.25
ICV (mm^3)	1539012 ± 128794	1569834 ± 151931	1559140 ± 144992	0.09
V$_{hypothalamus}$ (mm^3)	737 ± 90	740 ± 104	733 ± 108	0.77
V$_{nucleus\ accumbens}$ (mm^3)	1028 ± 204	1057 ± 225	1034 ± 204	0.36
V$_{caudate\ nucleus}$ (mm^3)	7902 ± 990	8038 ± 975	7853 ± 998	0.16
V$_{putamen}$ (mm^3)	10609 ± 1154	10691 ± 1162	10627 ± 1206	0.77
Obesity-related measures				
Baseline				
BMI (kg/m^2)	16.82 ± 1.39	20.92 ± 1.06	25.85 ± 3.41	< 0.001 ***
BMI percentile	48.51 ± 22.38	90.43 ± 2.90	97.63 ± 1.32	< 0.001 ***
BMI z-score	-0.06 ± 0.65	1.33 ± 0.17	2.05 ± 0.29	< 0.001 ***
WC (inch)	25.13 ± 2.27	29.16 ± 2.68	32.96 ± 4.10	< 0.001 ***
One-year				
BMI (kg/m^2)	17.61 ± 1.88	21.97 ± 2.24	27.16 ± 4.63	< 0.001 ***
BMI z-score	-0.01 ± 0.75	1.28 ± 0.47	1.98 ± 0.53	< 0.001 ***
WC (in)	25.93 ± 2.68	30.03 ± 2.82	34.89 ± 4.48	< 0.001 ***
Change (Δ) over one year				
ΔBMI (kg/m^2)	0.79 ± 1.29	1.05 ± 1.97	1.30 ± 3.57	0.12
ΔBMI z-score	0.04 ± 0.51	-0.05 ± 0.44	-0.08 ± 0.45	0.024 *
ΔWC (in)	0.80 ± 2.34	0.88 ± 2.66	1.94 ± 3.16	< 0.001 ***
Two-year				
BMI (kg/m^2)	18.43 ± 2.32	22.91 ± 2.48	28.11 ± 4.81	< 0.001 ***
BMI z-score	0.01 ± 0.86	1.28 ± 0.50	1.94 ± 0.54	< 0.001 ***
WC (in)	27.12 ± 3.01	31.34 ± 3.27	36.02 ± 4.98	< 0.001 ***
Change (Δ) over two years				
ΔBMI (kg/m^2)	1.62 ± 1.83	1.98 ± 2.27	2.25 ± 3.85	0.07
ΔBMI z-score	0.07 ± 0.65	-0.05 ± 0.48	-0.11 ± 0.48	0.002 **
ΔWC (in)	1.99 ± 2.72	2.18 ± 3.01	3.06 ± 3.72	0.001 **

Table 1. Participant demographics, brain volumes, and obesity-related measures. Statistics are
shown as mean ± standard deviation for continuous variables and count (frequency) for categorical data.
Variables were assessed at baseline unless otherwise noted. Comparisons were performed using one-way
analysis of variance or chi-squared tests as appropriate. The "Other" category under race/ethnicity
included participants who were parent/caregiver-identified as American Indian, Alaskan Native, Native
Hawaiian, other Pacific Islander, mixed, or otherwise not listed. Abbreviations: NW, children with
normal-weight; OW, with overweight; OB, with obesity; PDS, pubertal development stage; HS, high
school; GED, General Educational Development; ICV, intracranial volume; V, volume; BMI, body mass
index; WC, waist circumference. *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$.
320

3.2. Comparison of white matter DBSI metrics across groups

TBSS ANOVAs indicated significant main effects of weight group for both DBSI-FF and RF.

Follow-up TBSS *t*-tests showed that relative to NW, both OW and OB participants had significantly

lower DBSI-FF (reflecting lower axonal/dendritic density) and greater DBSI-RF (reflecting elevated

cellularity) in widespread white matter tracts (all FWE-corrected $p \leq 0.05$). Qualitatively, group

differences in both DBSI-FF and RF appeared more widespread throughout white matter tracts in the OB

vs. NW comparisons than in OW vs. NW comparisons (**Fig. 1**); nonetheless, white matter voxel-wise

DBSI-FF and RF were not significantly different between OB and OW groups (FWE-corrected $p >$

0.054). Consistent with voxel-wise comparisons, relative to NW, both OW and OB groups had lower

white matter average DBSI-FF (OW vs. NW, $\beta = -0.39$, 95% CI: -0.60 to -0.18, $p < 0.001$; OB vs. NW, β

= -0.33, 95% CI: -0.54 to -0.11, $p = 0.003$) and greater DBSI-RF (OW vs. NW, $\beta = 0.50$, 95% CI: 0.30 to

0.70, $p < 0.001$; OB vs. NW, $\beta = 0.36$, 95% CI: 0.16 to 0.56, $p < 0.001$), but these differences were not

significant between OW and OB (p's = 0.65 and 0.14 for DBSI-FF and RF comparisons).

DBSI Fiber Fraction (FF)

NW > OW   **NW > OB**

DBSI Restricted Fraction (RF)

NW < OW   **NW < OB**

Fig. 1. Voxel-wise comparisons of white matter DBSI metrics amongst unrelated children with normal-weight (NW; $n = 202$), overweight (OW, $n = 180$) and obesity (OB; $n = 191$). In each panel, axial images are shown from inferior (top left) to superior (bottom right). In *green*: white matter skeleton; in *blue, light blue*: NW > OB/OW at family-wise error (FWE) rate-corrected $p \leq 0.05$ and 0.01; in *red, yellow*: NW < OB/OW at FWE-corrected $p \leq 0.05$ and 0.01. Comparisons were adjusted for age, sex, race/ethnicity, parental education, household income, parental marital status, pubertal development stage, mean head motion, and intracranial volume. DBSI, diffusion basis spectrum imaging.

3.3. Associations between striatal and hypothalamic DBSI metrics and obesity-related measures

3.3.1. Baseline

Greater BMI at baseline was significantly associated with greater DBSI-RF in the hypothalamus ($\beta = 0.11$, 95% CI: 0.04 to 0.19, partial $R^2 = 0.014$, $p = 0.005$), nucleus accumbens ($\beta = 0.30$, 95% CI: 0.22 to 0.39, partial $R^2 = 0.096$, $p < 0.001$), caudate nucleus ($\beta = 0.18$, 95% CI: 0.08 to 0.27, partial $R^2 = 0.034$, $p < 0.001$), and the putamen ($\beta = 0.14$, 95% CI: 0.06 to 0.22, partial $R^2 = 0.021$, $p = 0.001$) (**Fig. 2A**). These results were consistent with WC and BMI *z*-scores as obesity-related measures. Further,

greater baseline BMI z-scores were significantly related to lower DBSI-FF in the hypothalamus (β = -

0.12, 95% CI: -0.20 to -0.05, partial R^2 = 0.016, p = 0.002) (**Fig. 2B**). Similar associations, at nominal but

not multiple comparison-adjusted significance, were seen between lower DBSI-FF in the hypothalamus

and WC (β = -0.08, p = 0.038) and BMI (β = -0.09, p = 0.024); in the nucleus accumbens and WC (β = -

0.10, p = 0.009); and in the putamen and BMI (β = -0.08, p = 0.048) and BMI z-scores (β = -0.08, p =

0.048). Detailed statistics for all models are reported in **Supplementary Table 2**. Follow-up analyses

revealed no DBSI metric by hemisphere interaction in relating to baseline obesity-related measures (i.e.,

no laterality effect; p's ≥ 0.30).

Beyond DBSI metrics, variables that were associated with greater obesity-related measures at

baseline included older age, lower parental education, and more advanced pubertal stage. As we did not

specifically power or hypothesize for demographics-related effects, these findings are exploratory and are

noted in **Supplementary Table 3**. In total, our linear-mixed effects models explained 18-25% of the

variance in baseline obesity-related measures.

Fig. 2. Significant associations **(A)** between baseline body mass index (BMI) and DBSI-RF in the
hypothalamus and striatum and **(B)** between baseline BMI z-scores and DBSI-FF in the hypothalamus in
children. BMI or BMI z-score residuals (adjusted for age, sex, race/ethnicity, parental education,
household income, parental marital status, pubertal development stage, mean head motion, intracranial
volume, and family nested by site) and DBSI metrics were standardized (std.). Standardized β regression
coefficients were reported with 95% confidence intervals (shaded). DBSI, diffusion basis spectrum
imaging.

3.3.2. One and two-year change

Greater DBSI-RF in the hypothalamus at baseline, at nominal significance not surviving multiple

comparison correction, predicted greater gain in WC over two years, accounting for baseline WC (β =

0.09, 95% CI: 0.01 to 0.18, partial R^2 = 0.008, p = 0.035; **Supplementary Fig. 4**). However, such effect

was not seen at one-year follow-up, or with changes in BMI or BMI z-scores as obesity-related measures

(p's = 0.79 and 0.52). Other one or two-year changes in obesity-related measures were not associated with

baseline DBSI metrics (**Supplementary Tables 4** and **5**).

3.4. Associations between striatal RSI-RNI and obesity-related measures

3.4.1. Baseline

Consistent with a previous study using ABCD Study® baseline data (n = 5,366; Rapuano et al.,

2020), greater baseline BMI was associated with higher RSI-RNI in the nucleus accumbens (β = 0.36,

95% CI: 0.27 to 0.44, partial R^2 = 0.125, p < 0.001), caudate nucleus (β = 0.15, 95% CI: 0.07 to 0.23,

partial R^2 = 0.025, p < 0.001), and putamen (β = 0.17, 95% CI: 0.09 to 0.26, partial R^2 = 0.030, p < 0.001)

in the current, smaller sample. Results were similar with WC and BMI z-scores. Detailed statistics for all

linear mixed-effects models are reported in **Supplementary Table 6**.

3.4.2. One and two-year change

Greater baseline RSI-RNI in the nucleus accumbens and caudate nucleus were respectively

associated, not surviving multiple comparison correction, with one-year gain in WC, accounting for

baseline levels (nucleus accumbens: β = 0.10, 95% CI: 0.00 to 0.20, partial R^2 = 0.008, p = 0.042; caudate

nucleus: β = 0.11, 95% CI: 0.02 to 0.19, partial R^2 = 0.011, p = 0.017; **Supplementary Fig. 5**). These

associations were not seen at two-year follow-up or with changes in BMI or BMI *z*-scores

(**Supplementary Tables 7** and **8**).

3.5. Comparison between DBSI and RSI on classifying NW and OB groups

395   Striatal DBSI and RSI metrics reflective of neuroinflammation-associated cellularity and cellular

396   density, i.e., DBSI-RF and RSI-RNI, showed similar sensitivity and specificity in classifying NW and OB

397   children (*p*'s ≥ 0.13; **Fig. 3**). Across the striatum, DBSI-RF was positively and strongly correlated with

398   RSI-RNI (*r*'s ≥ 0.60, *p*'s ≤ 0.001; **Supplementary Fig. 6**).

399
400   **Fig. 3.** Receiver operating characteristic curves comparing striatal DBSI restricted fraction (RF) and RSI
401   restricted normalized isotropic (RNI) performance in classifying children with normal-weight (NW) and
402   obesity (OB). AUC, area-under-the-curve; DBSI, diffusion basis spectrum imaging; RSI, restriction
403   spectrum imaging.

4. Discussion

4.1. Overview

Here we present both novel findings and support for the reproducibility of previous neuroimaging

studies that observed microstructural alterations suggestive of neuroinflammation in key feeding and

reward-related brain regions in childhood obesity. First, we demonstrate that elevated DBSI-assessed

cellularity, i.e., putative inflammatory marker, in the striatum relates to higher WC, BMI, and BMI *z*-

scores in 601 children aged 9 to 11 years from the ABCD Study®, reproducing observations made by

Rapuano et al., (2020) that used another diffusion-based RSI model in the same dataset. Quantitatively,

DBSI-RF and RSI-RNI were associated with obesity-related measures in similar magnitudes, and the two

methods exhibited comparable performance in classifying NW vs. OB children. Such convergence of

findings underpins the sensitivity and utility of diffusion MRI-based techniques in characterizing brain

microstructural alterations in obesity.

Second, we observed associations between obesity and increased purported cellularity consistent

with neuroinflammation in brain white matter tracts and hypothalamus, which were not assessed by

Rapuano et al., (2020). Our results in the hypothalamus align with reports of putative gliosis in this

region, assessed by quantitative T2 MRI, in both childhood and adult obesity (Schur et al., 2015;

Sewaybricker et al., 2019; Sewaybricker, Kee, et al., 2021; Sewaybricker, Melhorn, et al., 2021; Thaler et

al., 2012). Here, our findings add diffusion MRI-derived evidence of obesity-related putative

neuroinflammation in the hypothalamus in children. Furthermore, to our knowledge, our study is the first

to investigate and report overweight and obesity-associated white matter microstructural alterations in

children, in line with our earlier studies of DBSI-assessed putative white matter neuroinflammation in

adults (Samara et al., 2020). Collectively, our findings and those previously reported suggest that young

children manifest obesity-related differences in brain microstructure that are consistent with

neuroinflammation seen in animal and post-mortem human brain studies (Baufeld et al., 2016; Buckman

et al., 2013; De Souza et al., 2005; Décarie-Spain et al., 2018; Schur et al., 2015; Valdearcos et al., 2017).

Such brain differences may affect current and future susceptibility for weight gain and its comorbidities

including cognitive impairment, type 2 diabetes, and late-life dementia (Liang et al., 2015; Simmonds et

al., 2016; Tait et al., 2022).

4.2. Links between obesity, neuroinflammation, and brain function

The highly vascularized hypothalamus responds to feeding-related hormones, neuronal signals,

and nutrients derived from the bloodstream (Velloso & Schwartz, 2011). As a "metabolic sensor", the

hypothalamus is vulnerable to overfeeding and obesity-related elevations in peripheral pro-inflammatory

molecules including cytokines and saturated fatty acids (Jais & Brüning, 2017). Overfeeding also causes

the blood-brain barrier to break down, further enabling inflammatory factors to infiltrate brain tissue

(Guillemot-Legris et al., 2016; Guillemot-Legris & Muccioli, 2017; Stranahan et al., 2016). Our finding

that DBSI-assessed cellularity (DBSI-RF) in the hypothalamus is greater in childhood obesity is

consistent with the neuroinflammatory phenotype encompassing the recruitment, proliferation, and

activation of astrocytes and microglia (i.e., reactive gliosis) seen in this brain region in rodents fed with

high-fat diets (Buckman et al., 2013; De Souza et al., 2005). While such immune response may initially

be neuroprotective, chronic gliosis leads to dysregulated neuroinflammatory processes that disrupt

hypothalamic metabolic regulation and contribute to overfeeding, leptin and insulin-resistance, and

development of obesity (Gómez-Apo et al., 2021; Sochocka et al., 2017; Valdearcos et al., 2017).

Persistent neuroinflammation could also cause axonal damage and loss (Frischer et al., 2009; Kempuraj et

al., 2016), which may explain our observed association between obesity and lower DBSI-assessed

axonal/dendritic density (DBSI-FF).

The striatum plays a key role in reward processing and appetitive behavior (Stice et al., 2011).

Striatal activity, primarily dopamine neurotransmission, is influenced by homeostatic signals from the

hypothalamus and by circulating feeding-related hormones, both acting on receptors on midbrain

dopaminergic cells (Abizaid et al., 2006; Figlewicz, 2016; Hommel et al., 2006; King et al., 2011).

Altered dopamine neurotransmission has been noted in obesity (Geiger et al., 2009; Wang et al., 2001;

Wu et al., 2017). Beyond the hypothalamus, neuroinflammation in the striatum may further contribute to

obesogenic behavior. Indeed, our observation of heightened DBSI-assessed cellularity across the striatum

in childhood obesity matches the microstructural changes characteristic of diet-induced reactive gliosis in

the nucleus accumbens in rodents (Décarie-Spain et al., 2018; Molina et al., 2020). Taken together, MRI-

based assessments of hypothalamic and striatal microstructure by us and others consistently suggest

putative neuroinflammation in these regions in childhood obesity, in agreement with studies in rodent

models and human adults.

Longitudinally, greater DBSI-assessed cellularity in the hypothalamus weakly predicted two-year

gain in WC, aligning with a recent T2 MRI-based report of putative hypothalamic gliosis being associated

with weight gain in children (Sewaybricker, Kee, et al., 2021). Further, greater RSI-RNI in the nucleus

accumbens and caudate nucleus were linked to one-year WC gain, reproducing findings in Rapuano et al.,

(2020). However, these findings were at nominal but not multiple comparison-corrected significance, and

did not generalize across different obesity-related measures or MRI techniques. As our sample size was

not intended to power for the weaker longitudinal effects observed in Rapuano et al., (2020), these

findings require confirmation in larger studies involving more longitudinal observations as the ABCD

Study® continues to release data. Nonetheless, given evidence that striatal neuroinflammation causally

contribute to overfeeding in rodents (Décarie-Spain et al., 2018), plus emerging reports that putative

nucleus accumbens cellularity may mediate the relationships between eating behavior and obesity in both

adults and children (Rapuano et al., 2022; Samara et al., 2021), chronic neuroinflammation should be

evaluated as a potential contributing factor to obesity maintenance.

4.3. Brain microstructure in childhood vs. adult obesity

476   Overall, the pattern of our results in children agrees with DBSI-assessed microstructural

alterations seen in adult obesity (Ly et al., 2021; Samara et al., 2020, 2021). Obesity-associated decrease

in apparent axonal/dendritic density and increase in cellularity have been observed in white matter in both

adults and children. However, the pattern of results in the striatum differs by age. For example, greater

putative cellularity in the nucleus accumbens is associated with higher BMI and related metrics in

children, but such effect is absent in adults (Samara et al., 2021). Interestingly, it has been noted that in

adults, higher BMI is associated with smaller nucleus accumbens volumes (Dekkers et al., 2019; García-

García et al., 2020), whereas in children, such association is reversed (García-García et al., 2020;

Rapuano et al., 2017) or absent, as is in the current study and another analysis of the ABCD Study® data

(Adise et al., 2021). It is possible that as early reactive responses to obesity, striatal cellularity and gliosis

would manifest as microstructural but not volumetric alterations in children, while chronic

neuroinflammation would over time contribute to vasogenic edema and atrophy seen in adults (Dorrance

et al., 2014; Sochocka et al., 2017), as in multiple sclerosis (Kamholz & Garbern, 2005). Furthermore, as

executive control regions such as the prefrontal cortex mature later relative to the striatum (Spear, 2000),

striatal disruptions may lead to a more dysregulated reward system that influences obesogenic behavior

more strongly in children than in adults. As the ABCD Study® collects biennial neuroimaging scans in the

same participants from childhood through adulthood using harmonized MRI sequences, future research

should capitalize on this longitudinal dataset to delineate obesity-related brain microstructural changes

over development.

4.4. Comparison between DBSI and RSI findings

Although DBSI and RSI differ in their modeling of brain microstructure, their measures of

restricted water diffusion have been interpreted similarly such that the isotropic intracellular water

fraction (DBSI-RF and RSI-RNI) is thought to ultimately reflect the degree of neuroinflammation-related

immune cell infiltration or tissue cellularity (Cross & Song, 2017; Rapuano et al., 2020, 2022; Wang et

al., 2011, 2015). Indeed, in our study, DBSI and RSI-assessed striatal cellularity related similarly to

obesity-related measures and strongly with each other, and classified obesity status with comparable

performance. A true head-to-head comparison of the microstructural properties reflected by DBSI-RF and

RSI-RNI would however warrant a controlled phantom or immunohistological gold standard. In general,

the agreeing findings from DBSI and RSI highlight that diffusion MRI-based techniques are sensitive to

characterizing obesity-associated microstructural alterations in children, adding a novel neuroimaging

tool that assesses putative neuroinflammation *in vivo*.

4.5. Limitations

Limitations and future directions include, first, the lack of longitudinal timepoints besides one

and two-year follow-ups. It is possible that obesity-related neuroinflammation affects clinical and

behavioral outcomes on a timescale larger than two years. Second, as the ABCD Study® does not record

obesity duration, we could not assess when and to what extent brain microstructural changes occur

relative to obesity onset. Further research tracking children moving from normal-weight to obesity would

be useful. Third, as we focused on assessing associations between brain microstructure and obesity-

related measures, factors such as sex and socioeconomic status (SES) that likely impact child

development and complicate said associations, though controlled for in analyses, were not tested. In terms

of sex, girls have greater fat mass and more concentrated trunk adiposity than boys, even at similar BMIs

(Wisniewski & Chernausek, 2009). Further, though obesity is associated with elevated serum leptin levels

in both sexes, such effect is stronger in girls, who also demonstrate increases in leptin during puberty as

opposed to decreases in boys (Falorni et al., 1997). In terms of SES, socioeconomic adversity is a known

risk factor for childhood obesity (Hemmingsson, 2018; Vazquez & Cubbin, 2020), with physical

inactivity, unhealthy diet, and stress as proposed mediating mechanisms (Caprio et al., 2008;

Gebremariam et al., 2017; Hemmingsson, 2018; Mekonnen et al., 2020). Studies have also noted that girls

from disadvantaged neighborhoods are more susceptible to obesity compared to boys (Kranjac et al.,

2021), and that girls and boys experience differential dietary influences and weight expectations from

parents and peers (Caprio et al., 2008; Shah et al., 2020). Regarding brain microstructure, recent analyses

using the ABCD Study® data have shown that girls demonstrate greater RSI-assessed cell and neurite

density in white matter compared to boys (Lawrence et al., 2022), and lower SES interacts with greater

BMI in relating to putative white matter neuroinflammation and smaller brain volumes (Adise et al.,

2022; Dennis et al., 2022; Li et al., 2023). Collectively, these results suggest that there exist complex

associations between sex, sociocultural forces, and brain microstructure, and future research should adopt

an integrative framework to investigate how they may individually and interactively shape obesity

development. On a related note, we emphasize growing concerns that current practices of MRI acquisition

and quality control may inadvertently exclude participants in less accessible rural areas, from lower SES

families, and of racial/ethnic minorities (Ricard et al., 2023). The exclusion of neuroimaging data with

excessive head motion, in particular, poses a challenge in obesity research, as greater BMI is causally and

genetically linked to increased motion (Beyer et al., 2020). It is possible that our findings may not

generalize to children of all sociodemographic backgrounds, and confirmation in large samples of

marginalized populations is needed.

Finally, we note the limited interpretability of diffusion MRI-derived microstructural metrics.

While DBSI assessments have been histopathologically validated as neuroinflammation-sensitive in

inflammatory neurological diseases including human and rodent models of multiple sclerosis (Chiang et

al., 2014; Wang et al., 2011, 2015), and rodent optic neuritis (Lin et al., 2017; Yang et al., 2021),

validation remains ongoing for obesity. Although the cellularity and axonal density effects inferred from

DBSI-modeled water diffusivity agree with the neuroinflammatory phenotype seen in animal models and

human post-mortem brain of obesity (Baufeld et al., 2016; Buckman et al., 2013; De Souza et al., 2005;

Décarie-Spain et al., 2018; Schur et al., 2015; Valdearcos et al., 2017), we recognize that DBSI, as any

MRI technique, is an indirect marker of brain microstructure and could reflect neural development that

otherwise do not involve neuroinflammation (Palmer et al., 2022). On a related note, it is challenging to

determine whether feeding-related regions such as the hypothalamus and striatum are the only ones

involved in obesity-related neuroinflammation, since a true control region in which this phenomenon is

definitively absent has not been identified. Such limitation invites future research to evaluate

microstructure throughout gray matter as well as study potential interactions between gray and white

matter alterations in obesity. The confidence in the validity of MRI-based assessments of obesity-related

neuroinflammation could be explored with rodent models and/or human studies using positron emission

tomography methods for measuring neuroinflammatory indicators (e.g., astrocyte and microglia

activation).

5. Conclusions

With DBSI, we observed microstructural alterations in white matter, hypothalamus, and striatum

in children with overweight and obesity. Agreement between DBSI and RSI suggested that diffusion MRI

is a sensitive and useful tool for assessing obesity-related putative cellularity in children. Given that

childhood and adolescence involve substantial brain development, further longitudinal work is warranted

to elucidate how early changes in brain microstructure may contribute to obesity and its comorbidities in

the long run.

Funding: This work was supported by the National Institutes of Health (NIH) grants R01DK085575
(Hershey), T32DA007261-29 (Samara, Ray, Eisenstein), 1RF1AG072637-01 (Raji), KL2TR000450 in
partial form of Washington University Institute of Clinical and Translational Sciences Multidisciplinary
Clinical Research Career Development Program to (Ray, Raji), P30DK020579 in partial form of
Washington University Diabetes Research Center Pilot & Feasibility Award to (Eisenstein); the
Washington University Neuroimaging Laboratory Research Center Innovation Funds (Eisenstein), the
Radiological Society of North America Research Scholar Grant (Raji), the Mallinckrodt Institute of
Radiology Pilot Award (Eisenstein) and Summer Research Program (Li), the Society for Neuroscience
Trainee Professional Development Award (Li), the Washington University Summer Undergraduate
Research Award (Li), and Washington University McDonnell Center for Systems Neuroscience. The
ABCD Study® is supported by the NIH and federal partners (grants U01DA041048, U01DA050989,
U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987, U01DA041174,
U01DA041106, U01DA041117, U01DA041028, U01DA041134, U01DA050988, U01DA051039,
U01DA041156, U01DA041025, U01DA041120, U01DA051038, U01DA041148, U01DA041093,
U01DA041089, U24DA041123, and U24DA041147). A full list of supporters is available
at https://abcdstudy.org/federal-partners.html. The funders had no role in study design, data collection and
analysis, preparation of the manuscript, or decision to publish. The content is solely the responsibility of
the authors and does not necessarily represent the official views of the National Institutes of Health or
other funders.

Conflict of interest disclosures: Unrelated to this study, Dr. Cyrus A. Raji consults to Brainreader,
Neurevolution, Voxelwise, and the Pacific Neuroscience Foundation. Other authors have no conflict of
interest to disclose.

Author contributions (CRediT): Zhaolong Li, BA: Conceptualization, Data curation, Formal analysis,
Methodology, Software, Visualization, Writing – original draft, Writing – review & editing; **Amjad
Samara, MD:** Conceptualization, Data curation, Formal analysis, Software, Writing – Review & Editing;
Mary Katherine Ray, PhD: Conceptualization, Formal analysis, Writing – original draft, Writing –
Review & Editing; **Jerrel Rutlin, BS:** Data curation, Software; **Cyrus A. Raji, MD, PhD:** Writing –
review & editing; **Joshua S. Shimony, MD, PhD:** Writing – review & editing; **Peng Sun, PhD:**
Methodology, Writing – review & editing; **Sheng-Kwei Song, PhD:** Methodology, Writing – review &
editing; **Tamara Hershey, PhD:** Conceptualization, Methodology, Supervision, Writing – review &
editing; **Sarah A. Eisenstein, PhD:** Conceptualization, Formal analysis, Methodology, Software,
Supervision, Validation, Writing – original draft, Writing – review & editing

Acknowledgements: The authors wish to thank Richard Ni, Jonathan Koller, and Heather Lugar for their
assistance with neuroimaging analyses in this study.

Data availability statement: The ABCD Study® data are publicly available through the National
Institute of Mental Health Data Archive (https://nda.nih.gov/abcd). The ABCD Study® data used in this
report came from the ABCD Study® Data Release 2.0.1 (DOI: DOI 10.15154/1506087, July 2019) and
4.0 (DOI: 10.15154/1523041, October 2021). The p-code of script used to generate DBSI maps in this
study are available upon request, and the developers of DBSI are in the process of publishing an open-
source version of the scripts.

References

Abizaid, A., Liu, Z.-W., Andrews, Z. B., Shanabrough, M., Borok, E., Elsworth, J. D., Roth, R. H., Sleeman, M. W., Picciotto, M. R., Tschöp, M. H., Gao, X.-B., & Horvath, T. L. (2006). Ghrelin modulates the activity and synaptic input organization of midbrain dopamine neurons while promoting appetite. *The Journal of Clinical Investigation*, *116*(12), 3229–3239. https://doi.org/10.1172/JCI29867

Adise, S., Allgaier, N., Laurent, J., Hahn, S., Chaarani, B., Owens, M., Yuan, D., Nyugen, P., Mackey, S., Potter, A., & Garavan, H. P. (2021). Multimodal brain predictors of current weight and weight gain in children enrolled in the ABCD study ®. *Developmental Cognitive Neuroscience*, *49*, 100948. https://doi.org/10.1016/j.dcn.2021.100948

Adise, S., Marshall, A. T., Kan, E., Gonzalez, M. R., & Sowell, E. R. (2022). Relating neighborhood deprivation to childhood obesity in the ABCD study: Evidence for theories of neuroinflammation and neuronal stress. *Health Psychology : Official Journal of the Division of Health Psychology, American Psychological Association*. https://doi.org/10.1037/hea0001250

Barch, D. M., Albaugh, M. D., Avenevoli, S., Chang, L., Clark, D. B., Glantz, M. D., Hudziak, J. J., Jernigan, T. L., Tapert, S. F., Yurgelun-Todd, D., Alia-Klein, N., Potter, A. S., Paulus, M. P., Prouty, D., Zucker, R. A., & Sher, K. J. (2018). Demographic, physical and mental health assessments in the adolescent brain and cognitive development study: Rationale and description. *Developmental Cognitive Neuroscience*, *32*, 55–66. https://doi.org/https://doi.org/10.1016/j.dcn.2017.10.010

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software; Vol 1, Issue 1 (2015)* . https://www.jstatsoft.org/v067/i01

Baufeld, C., Osterloh, A., Prokop, S., Miller, K. R., & Heppner, F. L. (2016). High-fat diet-induced brain region-specific phenotypic spectrum of CNS resident microglia. *Acta Neuropathologica*, *132*(3), 361–375. https://doi.org/10.1007/s00401-016-1595-4

Beilharz, J. E., Maniam, J., & Morris, M. J. (2016). Short-term exposure to a diet high in fat and sugar, or liquid sugar, selectively impairs hippocampal-dependent memory, with differential impacts on inflammation. *Behavioural Brain Research*, *306*, 1–7. https://doi.org/https://doi.org/10.1016/j.bbr.2016.03.018

Beyer, F., Prehn, K., Wüsten, K. A., Villringer, A., Ordemann, J., Flöel, A., & Witte, A. V. (2020). Weight loss reduces head motion: Revisiting a major confound in neuroimaging. *Human Brain Mapping*, *41*(9), 2490–2494. https://doi.org/10.1002/hbm.24959

Biener, A. I., Cawley, J., & Meyerhoefer, C. (2020). The medical care costs of obesity and severe obesity in youth: An instrumental variables approach. *Health Economics*, *29*(5), 624–639. https://doi.org/https://doi.org/10.1002/hec.4007

Billot, B., Bocchetta, M., Todd, E., Dalca, A. V., Rohrer, J. D., & Iglesias, J. E. (2020). Automated segmentation of the hypothalamus and associated subunits in brain MRI. *NeuroImage*, *223*, 117287. https://doi.org/10.1016/j.neuroimage.2020.117287

Birdsill, A. C., Oleson, S., Kaur, S., Pasha, E., Ireton, A., Tanaka, H., & Haley, A. (2017). Abdominal obesity and white matter microstructure in midlife. *Human Brain Mapping*, *38*(7), 3337–3344. https://doi.org/10.1002/hbm.23576

Buckman, L. B., Thompson, M. M., Moreno, H. N., & Ellacott, K. L. J. (2013). Regional astrogliosis in the mouse hypothalamus in response to obesity. *Journal of Comparative Neurology*, *521*(6), 1322–

1333. https://doi.org/https://doi.org/10.1002/cne.23233

Caprio, S., Daniels, S. R., Drewnowski, A., Kaufman, F. R., Palinkas, L. A., Rosenbloom, A. L., &
Schwimmer, J. B. (2008). Influence of race, ethnicity, and culture on childhood obesity:
implications for prevention and treatment: a consensus statement of Shaping America's Health and
the Obesity Society. *Diabetes Care*, *31*(11), 2211–2221. https://doi.org/10.2337/dc08-9024

Carbine, K. A., Duraccio, K. M., Hedges-Muncy, A., Barnett, K. A., Kirwan, C. B., & Jensen, C. D.
(2020). White matter integrity disparities between normal-weight and overweight/obese adolescents:
an automated fiber quantification tractography study. *Brain Imaging and Behavior*, *14*(1), 308–319.
https://doi.org/10.1007/s11682-019-00036-4

Casey, B. J., Cannonier, T., Conley, M. I., Cohen, A. O., Barch, D. M., Heitzeg, M. M., Soules, M. E.,
Teslovich, T., Dellarco, D. V, Garavan, H., Orr, C. A., Wager, T. D., Banich, M. T., Speer, N. K.,
Sutherland, M. T., Riedel, M. C., Dick, A. S., Bjork, J. M., Thomas, K. M., … Dale, A. M. (2018).
The Adolescent Brain Cognitive Development (ABCD) study: Imaging acquisition across 21 sites.
Developmental Cognitive Neuroscience, *32*, 43–54.
https://doi.org/https://doi.org/10.1016/j.dcn.2018.03.001

Chiang, C. W., Wang, Y., Sun, P., Lin, T. H., Trinkaus, K., Cross, A. H., & Song, S. K. (2014).
Quantifying white matter tract diffusion parameters in the presence of increased extra-fiber
cellularity and vasogenic edema. *NeuroImage*, *101*, 310–319.
https://doi.org/10.1016/j.neuroimage.2014.06.064

Cole, T. J., Faith, M. S., Pietrobelli, A., & Heo, M. (2005). What is the best measure of adiposity change
in growing children: BMI, BMI %, BMI z-score or BMI centile? *European Journal of Clinical*
Nutrition, *59*(3), 419–425. https://doi.org/10.1038/sj.ejcn.1602090

Cross, A. H., & Song, S. K. (2017). "A new imaging modality to non-invasively assess multiple sclerosis
pathology." *Journal of Neuroimmunology*, *304*, 81–85.
https://doi.org/10.1016/j.jneuroim.2016.10.002

Daoust, J., Schaffer, J., Zeighami, Y., Dagher, A., García-García, I., & Michaud, A. (2021). White matter
integrity differences in obesity: A meta-analysis of diffusion tensor imaging studies. *Neuroscience*
and Biobehavioral Reviews, *129*, 133–141. https://doi.org/10.1016/j.neubiorev.2021.07.020

De Souza, C. T., Araujo, E. P., Bordin, S., Ashimine, R., Zollner, R. L., Boschero, A. C., Saad, M. J. A.,
& Velloso, L. A. (2005). Consumption of a fat-rich diet activates a proinflammatory response and
induces insulin resistance in the hypothalamus. *Endocrinology*, *146*(10), 4192–4199.
https://doi.org/10.1210/en.2004-1520

Décarie-Spain, L., Sharma, S., Hryhorczuk, C., Issa-Garcia, V., Barker, P. A., Arbour, N., Alquier, T., &
Fulton, S. (2018). Nucleus accumbens inflammation mediates anxiodepressive behavior and
compulsive sucrose seeking elicited by saturated dietary fat. *Molecular Metabolism*, *10*, 1–13.
https://doi.org/10.1016/j.molmet.2018.01.018

Dekkers, I. A., Jansen, P. R., & Lamb, H. J. (2019). Obesity, Brain Volume, and White Matter
Microstructure at MRI: A   Cross-sectional UK Biobank Study. *Radiology*, *291*(3), 763–
771. https://doi.org/10.1148/radiol.2019181012

Dennis, E., Manza, P., & Volkow, N. D. (2022). Socioeconomic status, BMI, and brain development in
children. *Translational Psychiatry*, *12*(1), 33. https://doi.org/10.1038/s41398-022-01779-3

Dorrance, A. M., Matin, N., & Pires, P. W. (2014). The effects of obesity on the cerebral vasculature.
Current Vascular Pharmacology, *12*(3), 462–472.

https://doi.org/10.2174/1570161112666140423222411

Falorni, A., Bini, V., Molinari, D., Papi, F., Celi, F., Stefano, G. Di, Berioli, M. G., Bacosi, M. L., &
Contessa, G. (1997). Leptin serum levels in normal weight and obese children and adolescents:
relationship with age, sex, pubertal development, body mass index and insulin. *International
Journal of Obesity*, *21*(10), 881–890. https://doi.org/10.1038/sj.ijo.0800485

Fernández-Andújar, M., Morales-García, E., & García-Casares, N. (2021). Obesity and Gray Matter
Volume Assessed by Neuroimaging: A Systematic Review. *Brain Sciences*, *11*(8).
https://doi.org/10.3390/brainsci11080999

Figlewicz, D. P. (2016). Expression of receptors for insulin and leptin in the ventral tegmental
area/substantia nigra (VTA/SN) of the rat: Historical perspective. *Brain Research*, *1645*, 68–70.
https://doi.org/https://doi.org/10.1016/j.brainres.2015.12.041

Frischer, J. M., Bramow, S., Dal-Bianco, A., Lucchinetti, C. F., Rauschka, H., Schmidbauer, M., Laursen,
H., Sorensen, P. S., & Lassmann, H. (2009). The relation between inflammation and
neurodegeneration in multiple sclerosis brains. *Brain*, *132*(5), 1175–1189.
https://doi.org/10.1093/brain/awp070

Garavan, H., Bartsch, H., Conway, K., Decastro, A., Goldstein, R. Z., Heeringa, S., Jernigan, T., Potter,
A., Thompson, W., & Zahs, D. (2018). Recruiting the ABCD sample: Design considerations and
procedures. *Developmental Cognitive Neuroscience*, *32*, 16–22.
https://doi.org/10.1016/j.dcn.2018.04.004

García-García, I., Morys, F., & Dagher, A. (2020). Nucleus accumbens volume is related to obesity
measures in an age-dependent fashion. *Journal of Neuroendocrinology*, *32*(12), e12812.
https://doi.org/10.1111/jne.12812

Gebremariam, M. K., Lien, N., Nianogo, R. A., & Arah, O. A. (2017). Mediators of socioeconomic
differences in adiposity among youth: a systematic review. *Obesity Reviews : An Official Journal of
the International Association for the Study of Obesity*, *18*(8), 880–898.
https://doi.org/10.1111/obr.12547

Geiger, B. M., Haburcak, M., Avena, N. M., Moyer, M. C., Hoebel, B. G., & Pothos, E. N. (2009).
Deficits of mesolimbic dopamine neurotransmission in rat dietary obesity. *Neuroscience*, *159*(4),
1193–1199. https://doi.org/https://doi.org/10.1016/j.neuroscience.2009.02.007

Gómez-Apo, E., Mondragón-Maya, A., Ferrari-Díaz, M., & Silva-Pereyra, J. (2021). Structural Brain
Changes Associated with Overweight and Obesity. *Journal of Obesity*, *2021*, 6613385.
https://doi.org/10.1155/2021/6613385

Gregor, M. F., & Hotamisligil, G. S. (2011). Inflammatory Mechanisms in Obesity. *Annual Review of
Immunology*, *29*(1), 415–445. https://doi.org/10.1146/annurev-immunol-031210-101322

Guillemot-Legris, O., Masquelier, J., Everard, A., Cani, P. D., Alhouayek, M., & Muccioli, G. G. (2016).
High-fat diet feeding differentially affects the development of inflammation in the central nervous
system. *Journal of Neuroinflammation*, *13*(1), 206. https://doi.org/10.1186/s12974-016-0666-8

Guillemot-Legris, O., & Muccioli, G. G. (2017). Obesity-Induced Neuroinflammation: Beyond the
Hypothalamus. In *Trends in Neurosciences* (Vol. 40, Issue 4, pp. 237–253). Elsevier Ltd.
https://doi.org/10.1016/j.tins.2017.02.005

Hagler, D. J., Hatton, S., Cornejo, M. D., Makowski, C., Fair, D. A., Dick, A. S., Sutherland, M. T.,
Casey, B. J., Barch, D. M., Harms, M. P., Watts, R., Bjork, J. M., Garavan, H. P., Hilmer, L., Pung,
C. J., Sicat, C. S., Kuperman, J., Bartsch, H., Xue, F., … Dale, A. M. (2019). Image processing and

analysis methods for the Adolescent Brain Cognitive Development Study. *NeuroImage*, *202*, 116091. https://doi.org/https://doi.org/10.1016/j.neuroimage.2019.116091

Hemmingsson, E. (2018). Early Childhood Obesity Risk Factors: Socioeconomic Adversity, Family Dysfunction, Offspring Distress, and Junk Food Self-Medication. *Current Obesity Reports*, *7*(2), 204–209. https://doi.org/10.1007/s13679-018-0310-2

Hommel, J. D., Trinko, R., Sears, R. M., Georgescu, D., Liu, Z.-W., Gao, X.-B., Thurmon, J. J., Marinelli, M., & DiLeone, R. J. (2006). Leptin Receptor Signaling in Midbrain Dopamine Neurons Regulates Feeding. *Neuron*, *51*(6), 801–810. https://doi.org/10.1016/j.neuron.2006.08.023

Jais, A., & Brüning, J. C. (2017). Hypothalamic inflammation in obesity and metabolic disease. *Journal of Clinical Investigation*, *127*(1), 24–32. https://doi.org/10.1172/JCI88878

Jernigan, T. L., Brown, S. A., & Dowling, G. J. (2018). The Adolescent Brain Cognitive Development Study. *Journal of Research on Adolescence : The Official Journal of the Society for Research on Adolescence*, *28*(1), 154–156. https://doi.org/10.1111/jora.12374

Jiang, F., Li, G., Ji, W., Zhang, Y., Wu, F., Hu, Y., Zhang, W., Manza, P., Tomasi, D., Volkow, N. D., Gao, X., Wang, G.-J., & Zhang, Y. (2023). Obesity is associated with decreased gray matter volume in children: a longitudinal study. *Cerebral Cortex*, *33*(7), 3674–3682. https://doi.org/10.1093/cercor/bhac300

Kamholz, J. A., & Garbern, J. Y. (2005). Neuronal cell injury precedes brain atrophy in multiple sclerosis. *Neurology*, *64*(1), 176 LP – 176. https://doi.org/10.1212/WNL.64.1.176

Kempuraj, D., Thangavel, R., Natteru, P. A., Selvakumar, G. P., Saeed, D., Zahoor, H., Zaheer, S., Iyer, S. S., & Zaheer, A. (2016). Neuroinflammation Induces Neurodegeneration. *Journal of Neurology, Neurosurgery and Spine*, *1*(1).

Killedar, A., Lung, T., Petrou, S., Teixeira-Pinto, A., Tan, E. J., & Hayes, A. (2020). Weight status and health-related quality of life during childhood and adolescence: effects of age and socioeconomic position. *International Journal of Obesity*, *44*(3), 637–645. https://doi.org/10.1038/s41366-020-0529-3

King, S. J., Isaacs, A. M., O'Farrell, E., & Abizaid, A. (2011). Motivation to obtain preferred foods is enhanced by ghrelin in the ventral tegmental area. *Hormones and Behavior*, *60*(5), 572–580. https://doi.org/https://doi.org/10.1016/j.yhbeh.2011.08.006

Kranjac, A. W., Boyd, C., Kimbro, R. T., Moffett, B. S., & Lopez, K. N. (2021). Neighborhoods matter; but for whom? Heterogeneity of neighborhood disadvantage on child obesity by sex. *Health & Place*, *68*, 102534. https://doi.org/https://doi.org/10.1016/j.healthplace.2021.102534

Kuczmarski, R. J., Ogden, C. L., Guo, S. S., Grummer-Strawn, L. M., Flegal, K. M., Mei, Z., Wei, R., Curtin, L. R., Roche, A. F., & Johnson, C. L. (2002). 2000 CDC Growth Charts for the United States: methods and development. *Vital and Health Statistics. Series 11, Data from the National Health Survey*, *246*, 1–190.

Kullmann, S, Schweizer, F., Veit, R., Fritsche, A., & Preissl, H. (2015). Compromised white matter integrity in obesity. *Obesity Reviews*, *16*(4), 273–281. https://doi.org/https://doi.org/10.1111/obr.12248

Kullmann, Stephanie, Callaghan, M. F., Heni, M., Weiskopf, N., Scheffler, K., Häring, H.-U., Fritsche, A., Veit, R., & Preissl, H. (2016). Specific white matter tissue microstructure changes associated with obesity. *NeuroImage*, *125*, 36–44. https://doi.org/https://doi.org/10.1016/j.neuroimage.2015.10.006

Lawrence, K. E., Abaryan, Z., Laltoo, E., Hernandez, L. M., Gandal, M. J., McCracken, J. T., & Thompson, P. M. (2022). White matter microstructure shows sex differences in late childhood: Evidence from 6797 children. *Human Brain Mapping*, *n/a*(n/a). https://doi.org/https://doi.org/10.1002/hbm.26079

Li, H., Nickerson, L. D., Nichols, T. E., & Gao, J.-H. (2017). Comparison of a non-stationary voxelation-corrected cluster-size test with TFCE for group-Level MRI inference. *Human Brain Mapping*, *38*(3), 1269–1280. https://doi.org/https://doi.org/10.1002/hbm.23453

Li, Y., Thompson, W. K., Reuter, C., Nillo, R., Jernigan, T., Dale, A., Sugrue, L. P., Brown, J., Dougherty, R. F., Rauschecker, A., Rudie, J., Barch, D. M., Calhoun, V., Hagler, D., Hatton, S., Tanabe, J., Marshall, A., Sher, K. J., Heeringa, S., … Brown, S. (2021). Rates of Incidental Findings in Brain Magnetic Resonance Imaging in Children. *JAMA Neurology*, *78*(5), 578–587. https://doi.org/10.1001/jamaneurol.2021.0306

Li, Z. A., Cai, Y., Taylor, R. L., Eisenstein, S. A., Barch, D. M., Marek, S., & Hershey, T. (2023). Associations between socioeconomic status and white matter microstructure in children: indirect effects via obesity and cognition. *MedRxiv*, 2023.02.09.23285150. https://doi.org/10.1101/2023.02.09.23285150

Liang, Y., Hou, D., Zhao, X., Wang, L., Hu, Y., Liu, J., Cheng, H., Yang, P., Shan, X., Yan, Y., Cruickshank, J. K., & Mi, J. (2015). Childhood obesity affects adult metabolic syndrome and diabetes. *Endocrine*, *50*(1), 87–92. https://doi.org/10.1007/s12020-015-0560-7

Lin, T.-H., Chiang, C.-W., Perez-Torres, C. J., Sun, P., Wallendorf, M., Schmidt, R. E., Cross, A. H., & Song, S.-K. (2017). Diffusion MRI quantifies early axonal loss in the presence of nerve swelling. *Journal of Neuroinflammation*, *14*(1), 78. https://doi.org/10.1186/s12974-017-0852-3

Ling, J., Merideth, F., Caprihan, A., Pena, A., Teshiba, T., & Mayer, A. R. (2012). Head injury or head motion? Assessment and quantification of motion artifacts in diffusion tensor imaging studies. *Human Brain Mapping*, *33*(1), 50–62. https://doi.org/10.1002/hbm.21192

Ly, M., Raji, C. A., Yu, G. Z., Wang, Q., Wang, Y., Schindler, S. E., An, H., Samara, A., Eisenstein, S. A., Hershey, T., Smith, G., Klein, S., Liu, J., Xiong, C., Ances, B. M., Morris, J. C., & Benzinger, T. L. S. (2021). Obesity and White Matter Neuroinflammation Related Edema in Alzheimer's Disease Dementia Biomarker Negative Cognitively Normal Individuals. *Journal of Alzheimer's Disease*, *79*, 1801–1811. https://doi.org/10.3233/JAD-201242

Mekonnen, T., Havdal, H. H., Lien, N., O'Halloran, S. A., Arah, O. A., Papadopoulou, E., & Gebremariam, M. K. (2020). Mediators of socioeconomic inequalities in dietary behaviours among youth: A systematic review. *Obesity Reviews*, *21*(7), e13016. https://doi.org/https://doi.org/10.1111/obr.13016

Molina, J., Joaquim, A., Bonamin, L. V., Martins, M. de F. M., Kirsten, T. B., Cardoso, C. V., Bernardi, M. M., & Bondan, E. F. (2020). Reduced astrocytic expression of GFAP in the offspring of female rats that received hypercaloric diet. *Nutritional Neuroscience*, *23*(6), 411–421. https://doi.org/10.1080/1028415X.2018.1512783

Neudorfer, C., Germann, J., Elias, G. J. B., Gramer, R., Boutet, A., & Lozano, A. M. (2020). A high-resolution in vivo magnetic resonance imaging atlas of the human hypothalamic region. *Scientific Data*, *7*(1), 305. https://doi.org/10.1038/s41597-020-00644-6

Palmer, C. E., Pecheva, D., Iversen, J. R., Hagler, D. J., Sugrue, L., Nedelec, P., Fan, C. C., Thompson, W. K., Jernigan, T. L., & Dale, A. M. (2022). Microstructural development from 9 to 14 years: Evidence from the ABCD Study. *Developmental Cognitive Neuroscience*, *53*, 101044.

https://doi.org/https://doi.org/10.1016/j.dcn.2021.101044

Patenaude, B., Smith, S. M., Kennedy, D. N., & Jenkinson, M. (2011). A Bayesian model of shape and
appearance for subcortical brain segmentation. *NeuroImage*, *56*(3), 907–922.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2011.02.046

Pistell, P. J., Morrison, C. D., Gupta, S., Knight, A. G., Keller, J. N., Ingram, D. K., & Bruce-Keller, A. J.
(2010). Cognitive impairment following high fat diet consumption is associated with brain
inflammation. *Journal of Neuroimmunology*, *219*(1–2), 25–32.
https://doi.org/10.1016/j.jneuroim.2009.11.010

R Core Team. (2013). *R: A language and environment for statistical computing*. R Foundation for
Statistical Computing.

Raji, C. A., Ho, A. J., Parikshak, N. N., Becker, J. T., Lopez, O. L., Kuller, L. H., Hua, X., Leow, A. D.,
Toga, A. W., & Thompson, P. M. (2010). Brain structure and obesity. *Human Brain Mapping*,
31(3), 353–364. https://doi.org/https://doi.org/10.1002/hbm.20870

Rapuano, K. M., Berrian, N., Baskin-Sommers, A., Décarie-Spain, L., Sharma, S., Fulton, S., Casey, B.
J., & Watts, R. (2022). Longitudinal Evidence of a Vicious Cycle Between Nucleus Accumbens
Microstructure and Childhood Weight Gain. *Journal of Adolescent Health*.
https://doi.org/https://doi.org/10.1016/j.jadohealth.2022.01.002

Rapuano, K. M., Laurent, J. S., Hagler, D. J., Hatton, S. N., Thompson, W. K., Jernigan, T. L., Dale, A.
M., Casey, B. J., & Watts, R. (2020). Nucleus accumbens cytoarchitecture predicts weight gain in
children. *Proceedings of the National Academy of Sciences*, *117*(43), 26977–26984.
https://doi.org/10.1073/pnas.2007918117

Rapuano, K. M., Zieselman, A. L., Kelley, W. M., Sargent, J. D., Heatherton, T. F., & Gilbert-Diamond,
D. (2017). Genetic risk for obesity predicts nucleus accumbens size and responsivity to real-world
food cues. *Proceedings of the National Academy of Sciences*, *114*(1), 160 LP – 165.
https://doi.org/10.1073/pnas.1605548113

Ricard, J. A., Parker, T. C., Dhamala, E., Kwasa, J., Allsop, A., & Holmes, A. J. (2023). Confronting
racially exclusionary practices in the acquisition and analyses of neuroimaging data. *Nature
Neuroscience*, *26*(1), 4–11. https://doi.org/10.1038/s41593-022-01218-y

Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., & Müller, M. (2011). pROC:
an open-source package for R and S+ to analyze and compare ROC curves. *BMC Bioinformatics*,
12(1), 77. https://doi.org/10.1186/1471-2105-12-77

Samara, A., Li, Z., Rutlin, J., Raji, C. A., Sun, P., Song, S.-K., Hershey, T., & Eisenstein, S. A. (2021).
Nucleus accumbens microstructure mediates the relationship between obesity and eating behavior in
adults. *Obesity*.

Samara, A., Murphy, T., Strain, J., Rutlin, J., Sun, P., Neyman, O., Sreevalsan, N., Shimony, J. S., Ances,
B. M., Song, S. K., Hershey, T., & Eisenstein, S. A. (2020). Neuroinflammation and White Matter
Alterations in Obesity Assessed by Diffusion Basis Spectrum Imaging. *Frontiers in Human
Neuroscience*, *13*. https://doi.org/10.3389/fnhum.2019.00464

Schur, E. A., Melhorn, S. J., Oh, S.-K., Lacy, J. M., Berkseth, K. E., Guyenet, S. J., Sonnen, J. A., Tyagi,
V., Rosalynn, M., De Leon, B., Webb, M. F., Gonsalves, Z. T., Fligner, C. L., Schwartz, M. W., &
Maravilla, K. R. (2015). Radiologic evidence that hypothalamic gliosis is associated with obesity
and insulin resistance in humans. *Obesity*, *23*(11), 2142–2148.
https://doi.org/https://doi.org/10.1002/oby.21248

Sewaybricker, L. E., Kee, S., Melhorn, S. J., & Schur, E. A. (2021). Greater radiologic evidence of hypothalamic gliosis predicts adiposity gain in children at risk for obesity. *Obesity*, *29*(11), 1770–1779. https://doi.org/https://doi.org/10.1002/oby.23286

Sewaybricker, L. E., Melhorn, S. J., Papantoni, A., Webb, M. F., Hua, J., Roth, C. L., Carnell, S., & Schur, E. A. (2021). Pilot multi-site and reproducibility study of hypothalamic gliosis in children. *Pediatric Obesity*, *16*(4), e12732. https://doi.org/https://doi.org/10.1111/ijpo.12732

Sewaybricker, L. E., Schur, E. A., Melhorn, S. J., Campos, B. M., Askren, M. K., Nogueira, G. A. S., Zambon, M. P., Antonio, M. A. R. G. M., Cendes, F., Velloso, L. A., & Guerra-Junior, G. (2019). Initial evidence for hypothalamic gliosis in children with obesity by quantitative T2 MRI and implications for blood oxygen-level dependent response to glucose ingestion. *Pediatric Obesity*, *14*(2), e12486. https://doi.org/https://doi.org/10.1111/ijpo.12486

Shah, B., Tombeau Cost, K., Fuller, A., Birken, C. S., & Anderson, L. N. (2020). Sex and gender differences in childhood obesity: contributing to the research agenda. *BMJ Nutrition, Prevention & Health*, *3*(2), 387–390. https://doi.org/10.1136/bmjnph-2020-000074

Simmonds, M., Llewellyn, A., Owen, C. G., & Woolacott, N. (2016). Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obesity Reviews*, *17*(2), 95–107. https://doi.org/https://doi.org/10.1111/obr.12334

Smith, S. M., Jenkinson, M., Johansen-Berg, H., Rueckert, D., Nichols, T. E., Mackay, C. E., Watkins, K. E., Ciccarelli, O., Cader, M. Z., Matthews, P. M., & Behrens, T. E. J. (2006). Tract-based spatial statistics: Voxelwise analysis of multi-subject diffusion data. *NeuroImage*, *31*(4), 1487–1505. https://doi.org/https://doi.org/10.1016/j.neuroimage.2006.02.024

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, *23*, S208–S219. https://doi.org/https://doi.org/10.1016/j.neuroimage.2004.07.051

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: Addressing problems of smoothing, threshold dependence and localisation in cluster inference. *NeuroImage*, *44*(1), 83–98. https://doi.org/https://doi.org/10.1016/j.neuroimage.2008.03.061

Sochocka, M., Diniz, B. S., & Leszek, J. (2017). Inflammatory Response in the CNS: Friend or Foe? *Molecular Neurobiology*, *54*(10), 8071–8089. https://doi.org/10.1007/s12035-016-0297-1

Spear, L. P. (2000). The adolescent brain and age-related behavioral manifestations. *Neuroscience & Biobehavioral Reviews*, *24*(4), 417–463. https://doi.org/https://doi.org/10.1016/S0149-7634(00)00014-2

Stice, E., Yokum, S., Burger, K. S., Epstein, L. H., & Small, D. M. (2011). Youth at Risk for Obesity Show Greater Activation of Striatal and Somatosensory Regions to Food. *The Journal of Neuroscience*, *31*(12), 4360 LP – 4366. https://doi.org/10.1523/JNEUROSCI.6604-10.2011

Stranahan, A. M., Hao, S., Dey, A., Yu, X., & Baban, B. (2016). Blood-brain barrier breakdown promotes macrophage infiltration and cognitive impairment in leptin receptor-deficient mice. *Journal of Cerebral Blood Flow and Metabolism : Official Journal of the International Society of Cerebral Blood Flow and Metabolism*, *36*(12), 2108–2121. https://doi.org/10.1177/0271678X16642233

Sun, P., George, A., Perantie, D. C., Trinkaus, K., Ye, Z., Naismith, R. T., Song, S.-K., & Cross, A. H. (2020). Diffusion basis spectrum imaging provides insights into MS pathology. *Neurology -*

Neuroimmunology Neuroinflammation, *7*(2), e655. https://doi.org/10.1212/NXI.0000000000000655

Tait, J. L., Collyer, T. A., Gall, S. L., Magnussen, C. G., Venn, A. J., Dwyer, T., Fraser, B. J., Moran, C.,
Srikanth, V. K., & Callisaya, M. L. (2022). Longitudinal associations of childhood fitness and
obesity profiles with midlife cognitive function: an Australian cohort study. *Journal of Science and*
Medicine in Sport, *25*(8), 667–672. https://doi.org/https://doi.org/10.1016/j.jsams.2022.05.009

Taylor, R. W., Jones, I. E., Williams, S. M., & Goulding, A. (2000). Evaluation of waist circumference,
waist-to-hip ratio, and the conicity index as screening tools for high trunk fat mass, as measured by
dual-energy X-ray absorptiometry, in children aged 3-19 y. *The American Journal of Clinical*
Nutrition, *72*(2), 490–495. https://doi.org/10.1093/ajcn/72.2.490

Thaler, J. P., Yi, C.-X., Schur, E. A., Guyenet, S. J., Hwang, B. H., Dietrich, M. O., Zhao, X., Sarruf, D.
A., Izgur, V., Maravilla, K. R., Nguyen, H. T., Fischer, J. D., Matsen, M. E., Wisse, B. E., Morton,
G. J., Horvath, T. L., Baskin, D. G., Tschöp, M. H., & Schwartz, M. W. (2012). Obesity is
associated with hypothalamic injury in rodents and humans. *The Journal of Clinical Investigation*,
122(1), 153–162. https://doi.org/10.1172/JCI59660

Valdearcos, M., Douglass, J. D., Robblee, M. M., Dorfman, M. D., Stifler, D. R., Bennett, M. L., Gerritse,
I., Fasnacht, R., Barres, B. A., Thaler, J. P., & Koliwad, S. K. (2017). Microglial Inflammatory
Signaling Orchestrates the Hypothalamic Immune Response to Dietary Excess and Mediates
Obesity Susceptibility. *Cell Metabolism*, *26*(1), 185-197.e3.
https://doi.org/10.1016/j.cmet.2017.05.015

Vazquez, C. E., & Cubbin, C. (2020). Socioeconomic Status and Childhood Obesity: a Review of
Literature from the Past Decade to Inform Intervention Research. *Current Obesity Reports*, *9*(4),
562–570. https://doi.org/10.1007/s13679-020-00400-2

Velloso, L. A., & Schwartz, M. W. (2011). Altered hypothalamic function in diet-induced obesity. In
International Journal of Obesity (Vol. 35, Issue 12, pp. 1455–1465).
https://doi.org/10.1038/ijo.2011.56

Verstynen, T. D., Weinstein, A. M., Schneider, W. W., Jakicic, J. M., Rofey, D. L., & Erickson, K. I.
(2012). Increased body mass index is associated with a global and distributed decrease in white
matter microstructural integrity. *Psychosomatic Medicine*, *74*(7), 682–690.
https://doi.org/10.1097/PSY.0b013e318261909c

Wang, G. J., Volkow, N. D., Logan, J., Pappas, N. R., Wong, C. T., Zhu, W., Netusll, N., & Fowler, J. S.
(2001). Brain dopamine and obesity. *Lancet*, *357*(9253), 354–357. https://doi.org/10.1016/S0140-
6736(00)03643-6

Wang, X., Cusick, M. F., Wang, Y., Sun, P., Libbey, J. E., Trinkaus, K., Fujinami, R. S., & Song, S.-K.
(2014). Diffusion basis spectrum imaging detects and distinguishes coexisting subclinical
inflammation, demyelination and axonal injury in experimental autoimmune encephalomyelitis
mice. *NMR in Biomedicine*, *27*(7), 843–852. https://doi.org/https://doi.org/10.1002/nbm.3129

Wang, Y., Sun, P., Wang, Q., Trinkaus, K., Schmidt, R. E., Naismith, R. T., Cross, A. H., & Song, S. K.
(2015). Differentiation and quantification of inflammation, demyelination and axon injury or loss in
multiple sclerosis. *Brain*, *138*(5), 1223–1238. https://doi.org/10.1093/brain/awv046

Wang, Y., Wang, Q., Haldar, J. P., Yeh, F.-C., Xie, M., Sun, P., Tu, T.-W., Trinkaus, K., Klein, R. S.,
Cross, A. H., & Song, S.-K. (2011). Quantification of increased cellularity during inflammatory
demyelination. *Brain : A Journal of Neurology*, *134*(Pt 12), 3590–3601.
https://doi.org/10.1093/brain/awr307

White, N. S., Leergaard, T. B., D'Arceuil, H., Bjaalie, J. G., & Dale, A. M. (2013). Probing tissue
microstructure with restriction spectrum imaging: Histological and theoretical validation. *Human*
Brain Mapping, *34*(2), 327–346. https://doi.org/https://doi.org/10.1002/hbm.21454

Willette, A. A., & Kapogiannis, D. (2015). Does the brain shrink as the waist expands? *Ageing Research*
Reviews, *20*, 86–97. https://doi.org/10.1016/j.arr.2014.03.007

Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M., & Nichols, T. E. (2014). Permutation
inference for the general linear model. *NeuroImage*, *92*, 381–397.
https://doi.org/https://doi.org/10.1016/j.neuroimage.2014.01.060

Winklewski, P. J., Sabisz, A., Naumczyk, P., Jodzio, K., Szurowska, E., & Szarmach, A. (2018).
Understanding the physiopathology behind axial and radial diffusivity changes-what do we Know?
In *Frontiers in Neurology* (Vol. 9, Issue FEB). Frontiers Media S.A.
https://doi.org/10.3389/fneur.2018.00092

Wisniewski, A. B., & Chernausek, S. D. (2009). Gender in childhood obesity: Family environment,
hormones, and genes. *Gender Medicine*, *6*, 76–85.
https://doi.org/https://doi.org/10.1016/j.genm.2008.12.001

World Health Organization. (2021). *Obesity and overweight*. https://www.who.int/en/news-room/fact-
sheets/detail/obesity-and-overweight

Wu, C., Garamszegi, S. P., Xie, X., & Mash, D. C. (2017). Altered Dopamine Synaptic Markers in
Postmortem Brain of Obese Subjects   . In *Frontiers in Human Neuroscience*   (Vol. 11, p. 386).
https://www.frontiersin.org/article/10.3389/fnhum.2017.00386

Yang, R., Lin, T.-H., Zhan, J., Lai, S., Song, C., Sun, P., Ye, Z., Wallendorf, M., George, A., Cross, A.
H., & Song, S.-K. (2021). Diffusion basis spectrum imaging measures anti-inflammatory and
neuroprotective effects of fingolimod on murine optic neuritis. *NeuroImage. Clinical*, *31*, 102732.
https://doi.org/10.1016/j.nicl.2021.102732

Yendiki, A., Koldewyn, K., Kakunoori, S., Kanwisher, N., & Fischl, B. (2014). Spurious group
differences due to head motion in a diffusion MRI study. *NeuroImage*, *88*, 79–90.
https://doi.org/10.1016/j.neuroimage.2013.11.027

Zhan, J., Lin, T.-H., Libbey, J. E., Sun, P., Ye, Z., Song, C., Wallendorf, M., Gong, H., Fujinami, R. S., &
Song, S.-K. (2018). Diffusion Basis Spectrum and Diffusion Tensor Imaging Detect Hippocampal
Inflammation and Dendritic Injury in a Virus-Induced Mouse Model of Epilepsy   . In *Frontiers in*
Neuroscience   (Vol. 12, p. 77). https://www.frontiersin.org/article/10.3389/fnins.2018.00077

984