1   Research Article
2

Evaluation of a microRNA-based Risk Classifier Predicting Cancer-Specific Survival in Renal Cell Carcinoma with Tumor Thrombus of the Inferior Vena Cava

6
Michael Kotlyar[a,1], Markus Krebs[a,b,1], Maximilian Burger[c], Hubert Kübler[a], Ralf Bargou[b], Susanne
Kneitz[d], Wolfgang Otto[c], Johannes Breyer[c], Daniel C. Vergho[c], Burkhard Kneitz[a], Charis Kalogirou[a]

[a] Department of Urology and Pediatric Urology, University Hospital Würzburg, 97080 Würzburg,

Germany

[b] Comprehensive Cancer Center Mainfranken, University Hospital Würzburg, 97080 Würzburg,

Germany

[c] Department of Urology, Caritas St. Josef and University of Regensburg Medical Center,

Regensburg, Germany

[d] Physiological Chemistry I, Biocenter, University of Würzburg, Würzburg, Germany

[1] both authors contributed equally and should be considered co-first authors.
17

Short Title: Prognostic Impact of microRNAs in Rena Cell Carcinoma with Vena Cava Infiltration

Corresponding Author:

Charis Kalogirou, MD

Department of Urology and Pediatric Urology, University Hospital Würzburg

Oberdürrbacher Str. 6

Würzburg, Bavaria, 97080, Germany

Tel: +49 931 201 32001

E-Mail: kalogirou_c@ukw.de

Keywords: kidney cancer, venous infiltration, cancer-specific survival, miR, risk stratification.

Abstract

<u>Introduction</u>: Clear cell renal cell carcinoma extending into the inferior vena cava (ccRCC^{IVC}) represents a clinical high-risk setting. However, there is substantial heterogeneity within this patient subgroup regarding survival outcomes. Previously, members of our group developed a microRNA(miR)-based risk classifier – containing miR-21, miR-126 and miR-221 expression – which significantly predicted cancer-specific survival (CSS) of ccRCC^{IVC} patients.

<u>Methods</u>: Examining a single-center cohort of tumor tissue from $n = 56$ patients with ccRCC^{IVC}, we measured expression levels of miR-21, miR-126 and miR-221 by qRT-PCR. Prognostic impact of clinicopathological parameters and miR expression were investigated via univariate and multivariate cox regression. Referring to the previously established risk classifier, we performed Kaplan Meier analyses for single miR expression levels and the combined risk classifier. Cut-off values and weights within the risk classifier were taken from the previous study.

<u>Results</u>: miR-21 and miR-126 expression were significantly associated with lymphonodal status at time of surgery, development of metastasis during follow-up, and cancer-related death. In Kaplan Meier analyses, miR-21 and miR-126 significantly impacted CSS in our cohort. Moreover, applying the miR-based risk classifier significantly stratified ccRCC^{IVC} according to CSS.

<u>Conclusion</u>: In our retrospective analysis, we successfully validated the miR-based risk classifier within an independent ccRCC^{IVC} cohort.

1. Introduction

In about 10 % of all cases, clear cell renal cell carcinomas (ccRCC) extend into the inferior vena cava (ccRCC^{IVC}) [1]–[3]. While constituting a high-risk setting in general, there still is substantial clinical heterogeneity within the ccRCC^{IVC} subgroup – with reported 5-year survival rates ranging from 37 % to 65 % for non-metastasized patients treated with nephrectomy and tumor thrombectomy [4]–[9]. Regarding this discrepancy, biomarkers are urgently needed to identify patients with a specifically high risk of cancer relapse [10], [11]. Potentially, this subgroup of ccRCC^{IVC} patients may also benefit from adjuvant systemic therapy and additional follow-up examinations.

MicroRNAs (miRs) as biomarker candidates are post-transcriptional regulators of gene expression in various cancer entities [12]–[14]. Regarding ccRCC, several studies demonstrated the prognostic impact of miR expression levels in tumor tissue [15]–[17]. Previously, Vergho et al. established a combined risk classifier for patients with ccRCC^{IVC} receiving nephrectomy and thrombectomy in curative intention [10]. Based on miR-21, miR-126 and miR-221 expression in tumor tissue, the risk classifier significantly stratified patients regarding cancer-specific survival in a

single-center cohort (*n* = 37) – with a 5-year cancer-specific survival (CSS) of 78 % vs. 18 % in the favorable and the unfavorable subgroup, respectively [10].

To further assess the miR-based risk classifier as a prognostic tool in ccRCC^IVC patients, we retrospectively evaluated it within an independent cohort (*n* = 56) from the Department of Urology, University of Regensburg (Regensburg, Germany). Cut-off values for miR expression levels as well as internal classifier weights were transferred from the previous pilot study [10], in order to test its transferability to independent study cohorts. Figure 1 illustrates the course of our study.

Fig. 1. Course of the study – using a microRNA (miR)-based risk classifier established previously [10], we examined the prognostic impact of miR-21, miR-126, and miR-221 expression in an independent cohort of clear cell renal cell carcinoma samples with infiltration of the inferior vena cava (ccRCC^IVC; *n* = 54). To assess the transferability of the miR-based risk classifier, cut-off value and weights were identical to the previous study.

2. Materials and Methods

2.1. Tumor Tissue Samples and Patients

Paraffin-embedded primary ccRCC^IVC tumor-samples of 56 subjects who underwent radical surgery were aggregated by the Department of Urology, University of Regensburg, Germany (1997–2006). A uropathologist selected sample-regions with > 90 % cancerous tissue. Follow-up data were collected by the Department of Urology, University of Regensburg (Regensburg, Germany). The study was approved by the local Ethics Committee (Regensburg: Nr. 08/108). Detailed characteristics of the study cohort are summarized in Table 1.

<u>2.2. RNA Extraction and qRT-PCR</u>

Using the RecoverAll^{TM} Total Nucleic Acid Isolation Kit (Thermo Fisher Scientific, Waltham, MA),

total RNA from paraffin-embedded samples was isolated according to the manufacturer's

instructions. RNA concentration and 260/280 ratio were analyzed by Spark® 10M (TECAN,

Männedorf, Switzerland). cDNA was synthesized from total RNA with stem-loop reverse

transcription primers (TaqMan microRNA Assay protocol, Applied Biosystems, Birchwood, UK).

TaqMan microRNA Assay kit was used to quantify miR expression according to manufacturer's

protocols. Samples showing a standard deviation > 0.5 were excluded (all reactions performed in

triplicates). Small nuclear RNA (RNU6b) expression was used for normalization of miRs relative

expression values. Samples with expression levels of RNU6B > 30 Ct were not included in further

analyses. Relative miR expressions were calculated using the ΔCt-method (ΔCt sample = Ct miR of

interest − Ct RNU6b). To calculate fold changes in miR expression between samples, the $2^{\Delta\Delta Ct}$

method (in this study, referred to as the ΔΔCt method) was used.

<u>2.3. Statistics and computational analysis</u>

To evaluate the validity of the miR-based risk classifier − (4.592 × ΔCt miR-21) + (-3.892 × ΔCt

miR-126) + (-1.938 × ΔCt miR-221) − weights and cut-off values (> 18.7 ΔCt = "unfavorable

subgroup", < 18.7 ΔCt = "favorable subgroup") were transferred from the pilot study [10] and

applied in order to stratify the ccRCC^{IVC} study cohort from Regensburg and perform Kaplan-Meier

analyses. Within the risk classifier formula, a negative factor indicates that higher expression levels

correlate with longer survival, while a positive factor correlates with shorter survival. For further

analysis, we transferred cut-off values for miR-21 (8.17 ΔCt), miR-126 (3.57 ΔCt) and miR-221

expression (1.84 ΔCt) to evaluate their predictive potential in the new cohort using the Kaplan-Meier

survival analysis. Kaplan-Meier, log-rank tests, univariate and multivariate Cox regression analysis

as well as partial effects on outcome analysis were performed using Python LifeLines Package

Version 0.27.0 [18]. Depending on the distribution of the data, Student's t-test or Mann-Whitney-

U-Test was used to analyze differences between miR expressions.

3. Results

Table 1 summarizes the basic clinical and pathological characteristics of our study cohort.

Detailed follow-up information were available for 54 of 56 ccRCC^{IVC} patients (96.4 %).

Table 1. Clinical and pathological patient characteristics (n = 56). Detailed follow-up information were available for 54 patients.

Characteristics	
Median Follow-up	94 (1 – 190) months
Median Age	67 (41 – 89) years
Sex	
female	22 (39.3 %)
male	34 (60.7 %)
Tumor Stage - pT3b	56 (100 %)
Fuhrman Grade	
G2	41 (73.2 %)
G3	15 (26.8 %)
Nodal Status	
N0	45 (80.4 %)
N+	11 (19.6 %)
Distant Metastasis during Follow-up	
M0	35 (62.5 %)
M1	21 (37.5 %)
Median Tumor Size	70 (18 – 225) mm
Overall survival	
yes	27 (48.2 %)
no	29 (51.8 %)
Cancer-related death	
yes	13 (23.2 %)
no	43 (76.8 %)

3.1. Association of miR-21, -126 and -221 Expression with Clinicopathological Characteristics

To investigate the impact of miR-21, -126 and -221 within our ccRCC^{IVC} cohort, we associated expression levels of miR-21, miR-126 and miR-221 to relevant clinical parameters. Figure 2 illustrates the results.

At time of surgery, 11 of 56 ccRCC^{IVC} patients (19.6 %) were diagnosed with nodal metastasis. In cases with positive nodal status, a trend towards up-regulation of miR-21 (p = 0.065) and a significant down-regulation of miR-126 (p < 0.001) were observed. For miR-221, no statistical association to nodal status could be shown.

During the follow-up period, distant metastasis emerged in 21 of 56 ccRCC^{IVC} patients (37.5 %). As shown in Figure 2b, we observed a significant up-regulation of miR-21 (p < 0.001) and down-regulation of miR-126 (p < 0.0001) as well as a trend towards downregulation for miR-221 (p = 0.06) in ccRCC^{IVC} samples of patients with metastasized disease.

13 of 56 patients with ccRCC^{IVC} (23.2 %) died during the follow-up period due to cancer (cancer-related death, CRD). Regarding miR expression levels, we found a significant up-regulation of miR-

21 (p < 0.0001) and a down-regulation of miR-126 (p < 0.001) in CRD cases. Instead, miR-221

expression did not show a statistically significant association to CRD in this analysis (p = 0.27).

Fig. 2. miR-21, -126 and -221 expression levels depending on lymphonodal status (a), distant metastases (b)

and cancer-related death (c). Significant changes between subgroups were calculated using unpaired Student's

t test or Mann-Whitney-U test. p < 0.05 *; p < 0.001**; p < 0.0001***.

<u>3.2. Cox regression Analysis</u>

Next, we performed a univariate Cox regression analysis to further assess the prognostic

potential of miR-21, -126, and -221 expression levels as predictors of CRD. Detailed follow-up

information were available for 54 of 56 cases with a median of 94 months.

As summarized in Table 2, miR-21 and miR-126 significantly predicted the occurrence of CRD in

our study cohort (p = 0.007 for miR-21, p = 0.00035 for miR-126). In contrast, miR-221 expression in

tumor tissue did not display significant prognostic potential (p = 0.28). Regarding further clinical

parameters, significant results were also observed for nodal involvement, metastatic status and

Fuhrman grade.

Table 2. Univariate and multivariate Cox regression of ccRCC[IVC] patients for miR expression levels and
clinicopathological parameters. 95 % Confidence intervals (CI) shown for Hazard ratios. *p* values were computed
using the chi-squared test. $p < 0.05$ *; $p < 0.001$ **; $p < 0.0001$ ***

	Cancer-related death			
	Univariate analysis		Multivariate analysis	
Parameters	Hazard Ratio (95 % CI)	*P* value	Hazard Ratio (95 % CI)	*P* value
miR-21	3.79 (1.55 – 9.26)	**0.003***	3.51 (1.2 – 10.2)	**0.02***
miR-126	0.19 (0.09 – 0.42)	**0.00035****	0.18 (0.075 – 0.43)	**0.0001****
miR-221	0.74 (0.46 – 1.19)	0.21	0.65 (0.38 – 1.13)	0.13
Age	0.98 (0.92 – 1.03)	0.42		
Sex	2.10 (0.58 – 7.65)	0.26		
Tumor size	1.01 (1.00 – 1.03)	0.07		
Fuhrman Grade	3.79 (1.27 – 11.33)	**0.02***		
Nodal status	6.70 (2.09 – 21.47)	**0.001***		
Distant metastasis	∞	NA		

Abbreviation: CI, confidence interval; ∞ / NA: In case of distant metastasis as clinical predictor of CRD, the coefficient was not estimable (positively infinite). Hazard ratio and p-value were therefore not shown.

Additionally, multivariate Cox regression for the three miR candidates yielded statistical
significance for miR-21 (*p* = 0.02, Hazard Ratio 3.51) and miR-126 expression (*p* = 0.0001, Hazard
Ratio 0.18) as predictors of CRD within our ccRCC[IVC] cohort. Despite a Hazard Ratio of 0.65, miR-221
expression again did not meet statistical significance as predictor of CRD (*p* = 0.13).

To further estimate the prognostic impact of single miR candidates from multivariate Cox
regression, we analyzed their partial effect on CSS for varying miR expression levels (Fig. 3). For miR-
21, higher relative expression levels were associated with lower CSS (Fig. 3a). In contrast, higher
expression levels of miR-126 as well as miR-221 were associated with higher CSS (Fig. 3b, c).

Fig. 3. (a–c) Partial effects of miR-21 (a), miR-126 (b) and miR-221 (c) expression levels on cancer-specific survival. The baseline represents median relative expression for each miRNA (miR-21 = 8.34 ΔCt, miR-126 = 4.0 ΔCt, miR-221 = 1.3 ΔCt). Blue line represents minimum and brown line the maximum relative expression of each miRNA in present study cohort. (d) Forest plot representing log Hazard Ratios (HR) from multivariate Cox regression of miR-21, -126, -221 for cancer-related death (CRD). p values were computed using the chi-squared test.

3.3. Kaplan Meier Analyses for single miR expression and the Risk Classifier

To investigate the prognostic transferability of single miRNAs, survival analyses using the cut-offs from Vergho et al. [10] were performed. Both, miR-21 and miR-126 showed a strong predictive significance in the Kaplan-Meier survival analysis (Fig. 4a, b). However, differences regarding CSS of miR-221 high vs. low expressing tumor specimens (Fig. 4c) did not reach statistical significance (p = 0.25).

To validate the predictive potential of the risk classifier (Fig. 4d), the cut-off level of 18.7 ΔCt (> 18.7 ΔCt = unfavorable subgroup, < 18.7 ΔCt = favorable subgroup) was used to stratify the validation cohort consisting of n = 54 ccRCC[IVC] tissue samples. Out of 13 CRD cases in total, the classifier correctly identified 12 patients who suffered from CRD as members of the unfavorable subgroup (92.3 %; > 18.7 ΔCt). Overall, a sensitivity of 92.3 % (CI 95 %: 62.1 % – 99.6 %) and a specificity of 61.0 % (CI 95 %: 44.5 % – 75.4 %) were reached. Difference in 5-years and 10-years CSS

was 100 % vs 70 % and 94 % vs 31 % between the favorable and the unfavorable subgroup,

respectively.

Fig. 4. Kaplan Meier survival analysis for CSS for external independent ccRCC[IVC] (*n* = 54) cohort from

Regensburg stratified by miR-21 (a), miR-126 (b) and miR-221 (c) expression levels. (d) Combined miR-based

risk classifier (miR-21, -126, -221) using identical cut-offs and weights from a previous publication [10]. *p* values

from log-rank tests are shown within each plot.

4. Discussion

ccRCCs infiltrating the inferior vena cava represent a clinically relevant high-risk subgroup.

Still, there is substantial clinical heterogeneity within this distinct subgroup – and biomarkers are

needed to assess the individual risk of progression. In general, adjuvant therapy with tyrosine kinase

inhibitors (TKI) or immune checkpoint blockers could be a promising therapeutic option after

nephrectomy – especially for patients suffering from high-risk RCC. However, European kidney

cancer guidelines currently do not contain strong recommendations towards adjuvant therapies due

to the mixed outcome in clinical trials [19]. For the TKI sunitinib, one trial displayed improved

disease-free survival (DFS) for patients – while showing no significant differences in overall survival

(OS) [20]. Additionally, another phase 3 trial did not detect significant survival effects for adjuvant

sunitinib or sorafenib in nonmetastatic high-risk renal cell carcinoma [21]. Due to the sobering TKI

results, research efforts were mainly shifted towards immune checkpoint blockers. For the PD-1

(Programmed Cell Death Protein 1) inhibitor pembrolizumab, KEYNOTE-564 trial detected improved

progression-free survival (PFS) in an adjuvant setting after nephrectomy [22].

4.1. Evaluating a miR-based Risk Classifier for RCC with Infiltration of the Vena Cava

To estimate the individual risk of patients suffering from ccRCC[IVC], members of our research

group have established a risk classifier based on the tissue expression of miR-21, miR-126, and miR-

221 [10]. Former cohort contained tumor tissue of n = 37 patients undergoing surgery at the

University Hospital of Würzburg, Germany. In this study, we externally validated the prognostic

potential of the miR-based risk classifier. Therefore, we examined an independent cohort of ccRCC[IVC]

from the University of Regensburg, Germany (n = 56). To test the transferability and usability of the

classifier within an external tissue cohort, we applied identical cut-off values and weights as in the

previous pilot study.

Regarding clinicopathological characteristics of our study cohort, low miR-126 expression was

significantly associated with a positive lymphonodal status at time of surgery. Moreover, occurrence

of metastases during follow-up was significantly associated with higher miR-21 and lower miR-126

levels. Tumor tissue from patients suffering from CRD was also characterized by a significant

upregulation of miR-21 and a downregulation of miR-126. Within univariate cox regression, miR-21

and miR-126 showed prognostic significance regarding cancer-specific survival (CSS). Lower levels

of miR-221 in tumor tissue and its association with CRD did not reach statistical significance. Beyond

miR expression, Fuhrman grade, lymphonodal status and occurrence of metastasis emerged as

prognostically relevant. Next, we added a multivariate Cox regression for the three miR candidates.

Again, this study identified miR-21 and miR-126 expression to significantly impact CRD. Kaplan Meier

analyses based on cut-off values determined previously by Vergho et al. [10] confirmed the

significant influence of miR-21 as well as miR-126 expression levels on CSS. Finally, applying the miR-

based classifier using identical cut-off values and weights split patients in two groups. Of note, the

classifier nearly stratified the study cohort in two halves – with n = 26 patients belonging to the

favorable and n = 28 patients belonging to the unfavorable subgroup. Regarding the substantial

difference in CSS between both groups, adjuvant therapies appear promising especially for the

unfavorable subgroup of ccRCC[IVC] patients.

4.2. Functional Roles of miR-21, miR-126, and miR-221 in Cancer

After confirming the prognostic potential of the miR classifier using the validation cohort from

Regensburg, we were interested in previously reported functions of these miRs in RCC and other

malignancies. For miR-21, several researchers demonstrated oncogenic effects in various cancers,

including RCC [23], [24]. Among the prominent miR-21 target genes are key players of apoptosis

induction like PDCD4 (Programmed Cell Death 4) [25] and genes like PTEN (Phosphatase and Tensin

Homolog) [23]. Latter is an established tumor suppressor gene best known for regulating PI3K/Akt

signaling. In contrast to miR-21, miR-126 acts as a tumor suppressor in tumor tissue, e. g. by

targeting ROCK1 (Rho Associated Coiled-Coil Containing Protein Kinase 1) [26] and VEGFA (Vascular

Endothelial Growth Factor A) [27]. For miR-221, oncogenic versus protective functions appear to

depend on the underlying cancer entity, as researchers demonstrated both roles [28]–[30]. For RCC,

a downregulation of miR-221 appears well in line with previous publications. Specifically, miR-221

is reported to regulate KDR (Kinase Insert Domain Receptor) – also known as VEGFR2 (Vascular

Endothelial Growth Factor Receptor 2) – in ccRCC [31] and prostate cancer [32], thereby regulating

the sensitivity towards sunitinib. In summary, among diverse tumorigenic functions of these miRs,

all three candidates prominently influence angiogenesis-related pathways (so-called

AngiomiRs) [33], [34]. Given that not all ccRCCs depend on angiogenic signaling to the same degree

[35], it is tempting to assume that the unfavorable subgroup identified by our risk classifier could

benefit from adjuvant anti-angiogenic therapy.

4.3. Limitations and Future Directions

Our study has several limitations. Leaving aside the special RCC subgroup investigated here,

sample size of our study is relatively small. Moreover, we purposely did not adjust cut-off values and

individual miR weights determined previously in order to check the transferability of the risk

classifier to external tissue cohorts. More research is needed – ideally in a prospective setting – to

further validate the risk classifier in a clinical setup and to elucidate whether sub-classification of

ccRCC^{IVC} is able to identify patients most susceptible towards adjuvant therapy.

5. Conclusions

While RCC extending into the inferior vena cava represents a high-risk setting, there is still

substantial clinical heterogeneity within this patient subgroup. Previously, Vergho et al. established

a miR-based risk classifier - containing miR-21, miR-126 and miR-221 expression – which significantly

predicted CSS for patients from this subgroup. To validate this classifier, we examined its impact on

an external and independent patient cohort. Using identical cut-off values for single miRs and

identical weights within the classifier, we confirmed a highly significant risk stratification within the

new cohort. Patients with an unfavorable constellation according to the miR-based classifier could

especially benefit from adjuvant therapy and continuous follow-up examinations.

Statement of Ethics

All procedures performed in studies involving human participants were in accordance with the

ethical standards of the institutional and/or national research committee and with the 1964 Helsinki

declaration and its later amendments or comparable ethical standards. The study was approved by

the local Ethics Committees (Regensburg: Nr. 08/108, Würzburg: Nr. 136/08).

271
272 **Conflict of Interest Statement**

273 The authors have no conflicts of interest to declare.

274
275 **Funding Sources**

276 Markus Krebs was funded by a personal grant from Else-Kröner-Foundation (Else Kröner Integrative

277 Clinician Scientist College for Translational Immunology, University Hospital Würzburg, Germany).

278
Author Contributions

Conceptualization, M.Ko., M.Kr., B.K., D.V. and C.K.; methodology, M.Kr., B.K., D.V. and C.K.;

software, M.Ko., S.K. and B.K.; validation, M.Kr., M.B., J.B., H.K., D.V., B.K. and C.K.; formal analysis,

M.Ko., S.K. and C.K.; investigation, M.Ko., M.Kr., W.O., J.B., S.K., B.K. and C.K; resources, J.B. and

B.K.; data curation, M.Ko., B.K. and C.K.; writing—M.Kr., M.Ko., J.B., B.K. and C.K.; writing—review

and editing, M.Ko., M.Kr., M.B., H.K., W.O., J.B., S.K., D.V., B.K. and C.K.; visualization, M.Ko. and

M.Kr.; supervision, M.B., H.K., W.O., D.V., B.K. and C.K.; project administration, M.Kr., B.K. and C.K.;

funding acquisition, M.B. and H.K.

References

[1] B. Ljungberg, R. Stenling, B. Osterdahl, E. Farrelly, T. Aberg, and G. Roos, 'Vein Invasion in Renal
Cell Carcinoma: Impact on Metastatic Behavior and Survival', *Journal of Urology*, vol. 154, no.
5, pp. 1681–1684, Nov. 1995, doi: 10.1016/S0022-5347(01)66749-1.

[2] W. Hoehn and P. Hermanek, 'Invasion of Veins in Renal Cell Carcinoma -Frequency, Correlation
and Prognosis', *Eur Urol*, vol. 9, no. 5, pp. 276–280, 1983, doi: 10.1159/000474103.

[3] A. C. Novick and D. M. Cosgrove, 'Surgical Approach for Removal of Renal Cell Carcinoma
Extending into the Vena Cava and the Right Atrium', *Journal of Urology*, vol. 123, no. 6, pp. 947–
950, Jun. 1980, doi: 10.1016/S0022-5347(17)56207-2.

[4] A. Haferkamp *et al.*, 'Renal Cell Carcinoma With Tumor Thrombus Extension Into the Vena Cava:
Prospective Long-Term Followup', *Journal of Urology*, vol. 177, no. 5, pp. 1703–1708, May 2007,
doi: 10.1016/j.juro.2007.01.039.

[5] M. L. Blute, B. C. Leibovich, C. M. Lohse, J. C. Cheville, and H. Zincke, 'The Mayo Clinic experience
with surgical management, complications and outcome for patients with renal cell carcinoma
and venous tumour thrombus', *BJU Int*, vol. 94, no. 1, pp. 33–41, Jul. 2004, doi: 10.1111/j.1464-
410X.2004.04897.x.

[6] D. C. Vergho, A. Loeser, A. Kocot, M. Spahn, and H. Riedmiller, 'Tumor thrombus of inferior vena
cava in patients with renal cell carcinoma – clinical and oncological outcome of 50 patients after
surgery', *BMC Res Notes*, vol. 5, no. 1, p. 264, Dec. 2012, doi: 10.1186/1756-0500-5-264.

[7] A. Q. Haddad *et al.*, 'Oncologic Outcomes Following Surgical Resection of Renal Cell Carcinoma
with Inferior Vena Caval Thrombus Extending Above the Hepatic Veins: A Contemporary
Multicenter Cohort', *Journal of Urology*, vol. 192, no. 4, pp. 1050–1056, Oct. 2014, doi:
10.1016/j.juro.2014.03.111.

[8] T. Klatte *et al.*, 'Prognostic Factors for Renal Cell Carcinoma With Tumor Thrombus Extension',
Journal of Urology, vol. 178, no. 4, pp. 1189–1195, Oct. 2007, doi: 10.1016/j.juro.2007.05.134.

[9] A. Berczi, T. Flasko, T. Szerafin, B. Thomas, Z. Bacso, and C. Berczi, 'Surgical Management and
Outcome of Renal Cell Carcinoma with Inferior Vena Cava Tumor Thrombus', *Urol Int*, vol. 99,
no. 3, pp. 267–271, 2017, doi: 10.1159/000464108.

[10] D. C. Vergho *et al.*, 'Impact of miR-21, miR-126 and miR-221 as Prognostic Factors of Clear Cell
Renal Cell Carcinoma with Tumor Thrombus of the Inferior Vena Cava', *PLoS ONE*, vol. 9, no. 10,
p. e109877, Oct. 2014, doi: 10.1371/journal.pone.0109877.

[11] P. J. Spachmann *et al.*, 'Impact of E-Cadherin and β-Catenin as Prognostic Factor in Renal Cell
Carcinoma with Tumor Thrombus of the Vena Cava', *Urol Int*, vol. 102, no. 4, pp. 413–420, 2019,
doi: 10.1159/000497279.

[12] V. Ambros, 'microRNAs', *Cell*, vol. 107, no. 7, pp. 823–826, Dec. 2001, doi: 10.1016/S0092-
8674(01)00616-X.

[13] A. Esquela-Kerscher and F. J. Slack, 'Oncomirs — microRNAs with a role in cancer', *Nat Rev
Cancer*, vol. 6, no. 4, pp. 259–269, Apr. 2006, doi: 10.1038/nrc1840.

[14] Y. W. Kong, D. Ferland-McCollough, T. J. Jackson, and M. Bushell, 'microRNAs in cancer
management', *The Lancet Oncology*, vol. 13, no. 6, pp. e249–e258, Jun. 2012, doi:
10.1016/S1470-2045(12)70073-6.

[15] M. S. Zaman *et al.*, 'Up-Regulation of MicroRNA-21 Correlates with Lower Kidney Cancer
Survival', *PLoS ONE*, vol. 7, no. 2, p. e31060, Feb. 2012, doi: 10.1371/journal.pone.0031060.

[16] D. Vergho *et al.*, 'Combination of expression levels of miR-21 and miR-126 is associated with
cancer-specific survival in clear-cell renal cell carcinoma', *BMC Cancer*, vol. 14, no. 1, p. 25, Dec.
2014, doi: 10.1186/1471-2407-14-25.

[17] F. Spadaccino *et al.*, 'The Ambivalent Role of miRNAs in Carcinogenesis: Involvement in Renal
Cell Carcinoma and Their Clinical Applications', *Pharmaceuticals*, vol. 14, no. 4, p. 322, Apr.
2021, doi: 10.3390/ph14040322.

[18] C. Davidson-Pilon, 'lifelines: survival analysis in Python', *JOSS*, vol. 4, no. 40, p. 1317, Aug. 2019,
doi: 10.21105/joss.01317.

[19] J. Bedke *et al.*, '2021 Updated European Association of Urology Guidelines on the Use of
Adjuvant Pembrolizumab for Renal Cell Carcinoma', *European Urology*, vol. 81, no. 2, pp. 134–
137, Feb. 2022, doi: 10.1016/j.eururo.2021.11.022.

[20] A. Ravaud *et al.*, 'Adjuvant Sunitinib in High-Risk Renal-Cell Carcinoma after Nephrectomy', *N
Engl J Med*, vol. 375, no. 23, pp. 2246–2254, Dec. 2016, doi: 10.1056/NEJMoa1611406.

[21] N. B. Haas *et al.*, 'Adjuvant sunitinib or sorafenib for high-risk, non-metastatic renal-cell
carcinoma (ECOG-ACRIN E2805): a double-blind, placebo-controlled, randomised, phase 3 trial',
Lancet, vol. 387, no. 10032, pp. 2008–2016, May 2016, doi: 10.1016/S0140-6736(16)00559-6.

[22] T. K. Choueiri *et al.*, 'Adjuvant Pembrolizumab after Nephrectomy in Renal-Cell Carcinoma', *N*
Engl J Med, vol. 385, no. 8, pp. 683–694, Aug. 2021, doi: 10.1056/NEJMoa2106391.

[23] C. Kalogirou *et al.*, 'Metformin-Derived Growth Inhibition in Renal Cell Carcinoma Depends on
miR-21-Mediated *PTEN* Expression', *Urol Int*, vol. 96, no. 1, pp. 106–115, 2016, doi:
10.1159/000441011.

[24] D. Bautista-Sánchez *et al.*, 'The Promising Role of miR-21 as a Cancer Biomarker and Its
Importance in RNA-Based Therapeutics', *Molecular Therapy - Nucleic Acids*, vol. 20, pp. 409–
420, Jun. 2020, doi: 10.1016/j.omtn.2020.03.003.

[25] X. Li *et al.*, 'MicroRNA-21 (miR-21) Post-Transcriptionally Downregulates Tumor Suppressor
PDCD4 and Promotes Cell Transformation, Proliferation, and Metastasis in Renal Cell
Carcinoma', *Cell Physiol Biochem*, vol. 33, no. 6, pp. 1631–1642, 2014, doi: 10.1159/000362946.

[26] G.-M. Zhang *et al.*, 'MicroRNA-126 inhibits tumor cell invasion and metastasis by
downregulating ROCK1 in renal cell carcinoma', *Molecular Medicine Reports*, vol. 13, no. 6, pp.
5029–5036, Jun. 2016, doi: 10.3892/mmr.2016.5160.

[27] T. Sasahira *et al.*, 'Downregulation of miR-126 induces angiogenesis and lymphangiogenesis by
activation of VEGF-A in oral cancer', *Br J Cancer*, vol. 107, no. 4, pp. 700–706, Aug. 2012, doi:
10.1038/bjc.2012.330.

[28] M. Garofalo, C. Quintavalle, G. Romano, C. M. Croce, and G. Condorelli, 'miR221/222 in Cancer:
Their Role in Tumor Progression and Response to Therapy', *CMM*, vol. 12, no. 1, pp. 27–33, Jan.
2012, doi: 10.2174/156652412798376170.

[29] F. Li, J.-W. Xu, L. Wang, H. Liu, Y. Yan, and S.-Y. Hu, 'MicroRNA-221-3p is up-regulated and serves
as a potential biomarker in pancreatic cancer', *Artificial Cells, Nanomedicine, and*
Biotechnology, vol. 46, no. 3, pp. 482–487, Apr. 2018, doi: 10.1080/21691401.2017.1315429.

[30] B. Kneitz *et al.*, 'Survival in patients with high-risk prostate cancer is predicted by miR-221, which
regulates proliferation, apoptosis, and invasion of prostate cancer cells by inhibiting IRF2 and
SOCS3', *Cancer Res.*, vol. 74, no. 9, pp. 2591–2603, May 2014, doi: 10.1158/0008-5472.CAN-13-
1606.

[31] H. W. Z. Khella *et al.*, 'miR-221/222 Are Involved in Response to Sunitinib Treatment in
Metastatic Renal Cell Carcinoma', *Mol. Ther.*, vol. 23, no. 11, pp. 1748–1758, Nov. 2015, doi:
10.1038/mt.2015.129.

[32] M. Krebs *et al.*, 'miR-221-3p Regulates VEGFR2 Expression in High-Risk Prostate Cancer and
Represents an Escape Mechanism from Sunitinib In Vitro', *JCM*, vol. 9, no. 3, p. 670, Mar. 2020,
doi: 10.3390/jcm9030670.

[33] Y. M. Salinas-Vera *et al.*, 'AngiomiRs: MicroRNAs driving angiogenesis in cancer (Review)', *Int J*
Mol Med, vol. 43, no. 2, pp. 657–670, Feb. 2019, doi: 10.3892/ijmm.2018.4003.

[34] S. Wang and E. N. Olson, 'AngiomiRs—Key regulators of angiogenesis', *Current Opinion in*
Genetics & Development, vol. 19, no. 3, pp. 205–211, Jun. 2009, doi: 10.1016/j.gde.2009.04.002.

[35] A. Marquardt *et al.*, 'Subgroup-Independent Mapping of Renal Cell Carcinoma—Machine
Learning Reveals Prognostic Mitochondrial Gene Signature Beyond Histopathologic
Boundaries', *Front. Oncol.*, vol. 11, p. 621278, Mar. 2021, doi: 10.3389/fonc.2021.621278.

386