

Clinical predictors of treatment outcome in adult Tuberculous Meningitis patients at the Kundiawa General Hospital.

Stanley Aiwara Jnr¹, Isi H. Kevau², David Linge³, Izzard Aglua⁴.

1. Internal Medicine Registrar (MBBS, MMedII), Port Moresby General Hospital.
2. Professor in Cardiology & Specialist Physician, Internal Medicine Department (PhD, MMed, MBBS). Port Moresby General Hospital.
3. Professor in Internal Medicine (PhD, MMed, MBBS). Port Moresby General Hospital.
4. Internal Medicine Registrar & Clinical Research Coordinator (MBBS, MPH) - Sir Joseph Nombri Memorial-Kundiawa General Hospital.

Abstract

Background

Tuberculous meningitis is the most severe form of extrapulmonary TB and accounted for 5 % of 10 million TB cases globally in the WHO 2018 report, with mortality as high as 19.35% in children and 30% in adults. Clinicians in resource-poor settings are often challenged with limited diagnostic and therapeutic options for optimal patient care, and often rely on clinical parameters for diagnosis, treatment, monitoring and outcome prediction.

Aim

This study was done to identify potential clinical predictors of i) length of in-hospital stay, and ii) death or discharge at 28 days, amongst Melanesian adults with TB meningitis at the Kundiawa General Hospital in Simbu Province of Papua New Guinea.

Method

A retrospective observational study was conducted on 65 Melanesian adults with TB meningitis at the Kundiawa General Hospital in Simbu province between 2015 and 2019.

Result

High case fatality (48%) and mortality rates (2.22 per 100,000 per year) for TB Meningitis in this study. Even higher case fatality of 90.9% with HIV co-infection. Parameters associated with higher 28-day mortality included admission GCS (P=0.039, 95% CI), HIV-TBM co-infection (P=0.049, 95% CI), positive fluid balance 24 hours after admission (P=0.061, 95% CI) and male gender (P=0.057, 95% CI).

Conclusion

Study showed high case fatality (48%) and mortality rates (2.22 per 100,000 per year) for adult TB Meningitis, with even higher case fatality (90.9%) for HIV-TB co-infection. Admission GCS, positive fluid balance 24 hours after admission, HIV-TB co-infection and male gender were associated with higher 28-day mortality.

Background

Tuberculous meningitis (TBM) is the most severe form of extrapulmonary Tuberculosis (TB) and accounted for up to 5 % of 10 million TB cases reported globally by WHO in 2018, with a high mortality and morbidity ^{1,2}. TB meningitis studies have reported mortalities as high as 19.35% in children and 30% in adults, with an associated high burden of neurological disability after treatment ³⁻⁵. Mortality was as high as 80% in studies from Sub-Saharan Africa and Asia ⁵.

The ongoing challenge for effective treatment with optimal outcome is further compounded in resource-poor settings in which clinicians are often faced with limited diagnostic and therapeutic options that impede timely and appropriate treatment ⁶⁻⁷. Clinicians often rely on clinical parameters for diagnosis, treatment, monitoring and outcome, to inform management decisions as well as discussions on disease condition and likely outcome with patients and relatives.

This observational study was therefore done to explore the role of certain clinical and cerebrospinal fluid (CSF) parameters in predicting treatment outcome amongst Melanesian adults with TB Meningitis at the Simbu Provincial Hospital in Papua New Guinea. The primary outcome measure assessed was death or discharge at 28 days, with length-of-hospitalization as the secondary outcome measure. Potential predictors of outcome assessed included duration of illness, duration of TB treatment, admission Glasgow Coma Score (GCS), mean arterial pressure (MAP) and plasma sodium on admission, fluid balance status 24 hours after admission, HIV-TBM co-infection and CSF findings (polymorphs, lymphocytes and acid-fast bacilli [AFB]), besides age and gender^{8,9}.

Aim

To identify potential clinical predictors of i) length of in-hospital stay, and ii) death or discharge at 28 days, amongst Melanesian adults with TB meningitis at the Kundiawa General Hospital in Simbu Province.

Method

A retrospective observational study was conducted on a total of 65 Melanesian adults with TB meningitis at the medical ward of the Sir Joseph Nombri Memorial, Kundiawa General Hospital in Simbu province between 2015 and 2019.

A modified Thwaites Criteria using clinical, radiological and CSF findings was employed for the diagnosis of Tuberculous Meningitis ^{10, 11}, whereby a patient with at least four of the following thematic features was diagnosed as a case of TBM:

i) meningitis symptoms (headache, fever, photophobia, vomiting, fitting) \geq 2 weeks, ii) personal or family history or symptoms of TB, iii) neck stiffness \pm Kernig's or Brudzinski's sign, iv) altered conscious status, v) focal neurological symptoms, vi) regional lymphadenopathy or other extrapulmonary TB features, vii) consistent chest radiograph and viii) CSF findings (AFB positive, lymphocyte predominance, elevated glucose with low protein). Of the four diagnostic features, at least one of TB and one of meningitis was required to make a TB meningitis diagnosis. Focal neurological symptoms included cranial nerve palsies and CSF AFB was assessed by polymerase chain reaction (PCR).

Study data was collected retrospectively from admission notes in patients' information chart using a semi-structured document review form by one investigator at a monthly interval over a 5-year period. Raw data was entered onto an excel spread sheet and imported into Stata 15.1 for analysis. Summary statistics and multivariate regression analysis were performed on variables of interest for possible relationships between predictor and outcome variables. Logistic regression was applied for 28-day mortality given its dichotomous nature, whilst linear regression was applied for length of in-hospital stay as it is numerical. The primary outcome measure assessed was death or discharge at 28 days, with length-of-hospitalization as the secondary outcome measure. Parameters assessed included duration of illness (days), duration of TB treatment (days), admission

Glasgow Coma Score (GCS), mean arterial pressure (MAP) and plasma sodium on admission, fluid balance status 24 hours after admission, HIV-TBM co-infection and CSF findings (polymorphs, lymphocytes and acid-fast bacilli [AFB]), besides age and gender ^{8,9}. The retrospective nature of the study limited its ability to effectively assess neurological deficit (neuro-motor, cognitive) as an outcome measure, owing to data insufficiency.

65 patients, over 13 years of age and diagnosed with TB meningitis (per criteria) with or without HIV infection were included for this study. All study subjects received a course of dexamethasone (2-8 mg iv/oral/NGT) during in-hospital treatment.

Results

▪ Death at 28 days

Case fatality for TB meningitis was 31/65, 48%.

Mortality rate ~2.22 per 100,000 adults per year in the province as at 2019 population data. Estimated adult population in Simbu province at 2019 was 278, 971 (2011 census data: total provincial population 376 973, 37% children under 15, and population growth rate 2.2% for PNG) ¹².

For an estimated population of 400,000 adults, 9 would have died from TB meningitis every year, at the given mortality rate of 2.22 per 100,000 per year, for the 5-year study period.

Predictors of death or discharge at 28 days.

Logistic regression for age, gender, duration of illness, duration of TB treatment, length of hospitalization, HIV-TBM co-infection, admission GCS, admission MAP, admission sodium, fluid balance status 24 hours after admission, CSF polymorphs, lymphocytes and positive AFB (PCR) on death at 28 days was performed.

Table 1. showing result from multi-variate regression analysis on outcome indicators for death at 28 days.

Mortality at 28 days	Odds Ratio	Std. Error	z	P> z	95% Confidence Interval	
Age	1.042368	.0551968	0.78	0.433	.9396088	1.156365
Gender	37.58772	71.72348	1.90	0.057	.8929247	1582.257
Illness duration	1.012295	.010914	1.13	0.257	.9911287	1.033914
Treatment duration	.8142552	.3854285	-0.43	0.664	.3219908	2.059101
Admission GCS	.5197426	.1648529	-2.06	0.039	.2791276	.9677741
Admission MAP	1.007134	.0421572	0.17	0.865	.9278065	1.093245
Admission sodium	1.108749	.1014481	1.13	0.259	.9267235	1.326528
Positive Fluid Balance	2844.575	12056.71	1.88	0.061	.701787	1.15e+07
HIV-TB	73.63644	246.7033	1.28	0.199	.103591	52343.6
CSF polymorphs	1.037084	.0326724	1.16	0.248	.9749837	1.103139
CSF lymphocytes	1.005336	.0097699	0.55	0.584	.9863688	1.024668
CSF PCR	.0845687	.2571376	-0.81	0.417	.0002183	32.76074
Length of hospitalization	1.002112	.0112889	0.19	0.851	.9802287	1.024484
_cons	2.43e-10	3.82e-09	-1.41	0.159	9.79e-24	6010.404
Observations	65					
Prob > chi2	0.0007					
R ²	0.6166					

- Statistically significant association between admission GCS (P=0.039, 95% CI) and 28-day mortality, male gender (P=0.057, 95% CI) and 28-day mortality, and a marginally significant association between positive fluid balance 24 hours after admission (P=0.061, 95% CI) and 28-day mortality, after controlling for other variables. Overall regression model was significant (Prob>chi2 = 0.0007).

Table 2. showing result from regression analysis for admission GCS, HIV-TBM coinfection, male gender and fluid balance 24 hours after admission on 28-day mortality.

Mortality at 28 days	Odds Ratio	Std. Error	z	P> z	95% Confidence Interval	
Male	2.525294	1.710047	1.37	0.171	.6697397	9.521774
Admission GCS	.7146273	.099275	2.42	0.016	.5442919	.9382688
HIV-TBM	9.439305	10.77502	1.97	0.049	1.007601	88.42835
Positive Fluid Balance	8.594234	7.913265	2.34	0.019	1.414005	52.2352
_cons	7.026309	13.62728	1.01	0.315	.1569827	314.487
Observations	65					
Prob > chi2	0.0000					
R ²	0.3108					

- Statistically significant, though weak (R²=0.3108), association between admission GCS (P=0.016, 95% CI), HIV-TBM coinfection (P=0.049, 95% CI), positive fluid balance (P=0.019, 95% CI) and death at 28 days, after controlling for each other. Overall regression model was significant (Prob>chi2 = 0.0000).

Table 3. Summary of Predictor Variables for 28-day Mortality


Predictors		Died	Discharged	Total
Admission GCS	≤ 5	1	0	1
	6 - 8	5	1	6
	9 - 11	9	5	14
	12 - 13	10	5	15
	14 - 15	7	22	29
Gender	Male	20	18	38
	Female	12	15	27
24-Hour fluid balance	Positive	15	6	21
	Negative	17	27	44
HIV-TBM co-infection	Positive	10	1	11
	Negative	22	32	54
				N = 65

▪ **Length of in-hospital stay**

1 day to 473 days. 50% < 28 days, 90% < 60days. Mean admission duration of 36.62 days.

Table 4. summary statistics for duration of hospitalization.

Summary Statistics – Length of in-hospital Stay		
	Percentiles	Days
	1%	1
	5%	1
	10%	2
	25%	6
	50%	26
	75%	43
	90%	59
	95%	119
	99%	473
Observations	65	
Mean	36.62	
Std. Deviation	63.04	



Predictors of length of in-hospital stay

Table 5. shows results from multivariate linear regression for potential determinants of duration of in-hospital stay.

Length of hospitalization	Coefficient	Std. Error	t	P> t	95% Confidence Interval	
Age	-.3961081	.5600288	-0.71	0.482	-1.517127	.7249113
Male	1.779555	15.49925	0.11	0.909	-29.24556	32.80467
HIV- TBM coinfection	-44.58576	20.599	-2.16	0.035	-85.81913	-3.352386
Positive fluid balance	-34.33084	15.98143	-2.15	0.036	-66.32114	-2.340533
CSF AFB	92.1679	24.695	3.73	0.000	42.73548	141.6003
Admission GCS	-.8063363	3.020443	-0.27	0.790	-6.852411	5.239738
_cons	67.46652	47.93152	1.41	0.165	-28.47884	163.4119
observations	65					
Prob > F	0.0062					
R ²	0.26					

- Statistically significant, though weak (R²=0.26), association between positive CSF AFB (P=0.000, 95% CI), HIV-TBM Co-infection (P=0.036, 95% CI), positive fluid balance (P=0.036, 95% CI) and length of in-hospital stay after controlling for each other. Overall regression model was significant (Prob>F = 0.0062).

Discussion

The high case fatality (48%) and mortality rates (2.22 per 100,000 per year) for TB Meningitis in this study concur with similar studies from resource-constrained settings around the world¹³⁻¹⁵. Apart from the diagnostic and therapeutic limitations, parameters inherent to the biology of *Mycobacterium tuberculosis* including virulence and persistence, together with the site and nature of disease condition, may account for similar higher mortalities for TB meningitis observed even in resource-rich settings¹⁶⁻¹⁸.

At the estimated mortality rate of 2.22 per 100,000 adults per year in the province, 9 adults would have died from TB meningitis every year for a population of around 400,000 adults for the 5-year study period in the province.

The clinical parameters associated with 28-day mortality from this study were admission GCS, HIV-TBM co-infection, positive fluid balance status in the first 24 hours of treatment and male gender.

A multivariate regression analysis (Table 1) showed a statistically significant association between admission GCS ($P=0.039$, 95% CI), male gender ($P=0.057$, 95% CI) and 28-day mortality after controlling for possible confounders. The strength of the association between the predictor and outcome variables was however weak ($R^2=0.3108$) in the model, owing largely to the small sample size (65), and therefore limiting the ability of the study to conclusively quantify the observed associations in terms of the relevant measure of association in odds ratio (OR). Hence, whilst it may be statistically sound to state that there were observed associations between predictor and outcomes variables of interest, any meaningful discussion of the OR (strength of these associations) is limited. A regression coefficient of $R^2 = 0.31$ in the model means that a unit change in any of the predictor variables will entail only a 31% corresponding change in the outcome variable (28-day mortality), which is far less than at least 75% ($R^2 \geq 0.75$) which is deemed necessary to represent a reliably strong association between predictor and outcome variables. Therefore, larger, prospective cohort studies (RCT) would be ideal to further explore and clearly ascertain the potential associations observed in this study.

55.4% of study subjects had a GCS of 13 or lower on admission, which inadvertently affected their progress on treatment and subsequent outcome of higher mortality at 28 days. Depressed conscious status has long been identified as an independent prognostic factor for poor outcomes in TB, and other forms of chronic meningitis around the world¹⁹. Delayed presentation to health care facilities with late treatment initiation continue to contribute to disease severity on admission, as reflected in this study by a high proportion of patients with lower admission GCS, and with subsequent poor outcomes in developing countries^{20, 21}.

The marginally significant association between positive fluid balance ($P=0.061$, 95% CI) 24 hours after admission and 28-day mortality observed, after controlling for confounding, highlights the role of fluid and electrolyte balance in treatment outcomes. A positive fluid balance after 24 hours of treatment was observed in only 25% of the study subjects but was significantly associated with higher mortality at 28 days, which warrants a closer attention by clinicians to fluid and electrolytes status during in-patient treatment of patients with TB meningitis and associated raised intracranial pressures^{22, 23}. TB Meningitis and raised intracranial pressure can cause syndromes of salt wasting or excess anti-diuretic hormone action (SIADH) with resulting hyponatremia which in turn entails functional neurological derangement with poor outcomes²⁴⁻²⁷.

TBM-HIV Co-infection is another independent predictor of mortality reported in several similar studies, with higher mortalities compared to non-HIV subjects with TB meningitis²⁸⁻³⁰. The co-infection rate in this study was relatively low at 16.92% but with a significantly high case fatality of 90.9%. There was no statistically significant association between HIV-TBM co-infection ($P=0.199$, 95% CI) and 28-day mortality in the multivariate regression after controlling for other variables (Table 1). However, in the model regressing identified predictors (admission GCS, gender, fluid balance 24 hours after admission and HIV-TB co-infection) in table 2, there was a statistically

significant association between HIV-TBM co-infection and 28-day mortality ($P=0.049$, 95% CI). This disparity may be attributable to potential confounding and/or underpowering of the study owing to the small sample size of 65 (type 2 error).

The duration of hospitalization ranged from 1 to 473 days, with a mean duration of 36.62 days. 90% of subjects stayed more than two weeks but less than 60 days (2 months) in the hospital, which is characteristic of chronic diseases and highlights the challenge of considerations in longer in-hospital care.

In the regression model for identified predictors (Table 5), there was a weak ($R^2=0.25$) but statistically significant association between positive CSF AFB ($P=0.000$, 95% CI), TBM-HIV Co-infection ($P=0.036$, 95% CI), positive fluid balance ($P=0.036$, 95% CI) and length of in-hospital stay after controlling for each other. Again, this may be attributable to confounding and the small sample size.

The weak associations observed in regression models ($R^2 < 0.75$) limit the value of assessed parameters as reliable predictors of length of hospitalization.

Conclusion

This study showed a relatively high case fatality (48%) and mortality rates (2.22 per 100,000 per year) for TB Meningitis amongst Melanesian adults admitted to the Kundiawa General Hospital. The case fatality for HIV-infected individuals with TB meningitis was much higher at 90.9%. Ninety percent of the subjects stayed for more than 2 weeks but less than 2 months in the hospital.

Admission GCS, positive fluid balance 24 hours after admission, HIV-TB co-infection and male gender were associated with higher 28-day mortality, and therefore can serve as useful clinical predictors of mortality in the care of TB meningitis patients, especially in resource-constrained settings. A positive CSF AFB can also serve as a useful indicator for duration of hospitalization amongst TB meningitis patients.

Prospective cohort studies with larger sample sizes are needed to further explore and ascertain the associations observed in this study.

Ethics

Ethical consideration and approval was granted by the interim Institutional Review Committee of the SiPHA-Clinical Research Center.

Acknowledgement

I would like to extend a sincere thank you to individuals and institutions without whose help and support this study would not have been possible, including:

- My spouse and colleague Dr. Jacqueline Pahun
- Sir Joseph Nombri Memorial-Kundiawa General Hospital administration and ward staff: Dr. Francis Wandj, Dr. Raymond Saulep, Dr. Cassius Maingu, Dr. Kassi Ken, Dr. Izzard Agua, Sr. Moro Drua and the nursing staff.
- Internal Medicine Department, UPNG SMHS & PGMH: Dr. Alexander Maha, Prof. David Linge, Prof. Sir Isi. H. Kevau.

Abbreviations

TBM-Tuberculous Meningitis, TB-Tuberculosis, WHO-World Health Organization, GCS-Glasgow Coma Scale, CSF-Cerebrospinal Fluid, PCR-Polymerase Chain Reaction, CI-Confidence Interval, AFB-Acid Fast Bacilli, NGT-Nasogastric Tube, SiPHA-Simbu Provincial Health Authority.

Keywords: Tuberculosis Meningitis, TB meningitis, prognostic indicators, clinical predictors, Melanesian adults, Papua New Guinea.

References

1. World Health Organization, 2014. World Health Organization Global Tuberculosis Report 2013.
2. Thao, L.T.P., Heemskerk, A.D., Geskus, R.B., Mai, N.T.H., Ha, D.T.M., Chau, T.T.H., Phu, N.H., Chau, N.V.V., Caws, M., Lan, N.H. and Thu, D.D.A., 2018. Prognostic models for 9-month mortality in tuberculous meningitis. *Clinical Infectious Diseases*, 66(4), pp.523-532.
3. Chiang, S.S., Khan, F.A., Milstein, M.B., Tolman, A.W., Benedetti, A., Starke, J.R. and Becerra, M.C., 2014. Treatment outcomes of childhood tuberculous meningitis: a systematic review and meta-analysis. *The Lancet Infectious Diseases*, 14(10), pp.947-957.
4. Thwaites, G.E. and Hien, T.T., 2005. Tuberculous meningitis: many questions, too few answers. *The Lancet Neurology*, 4(3), pp.160-170.
5. Stadelman, A.M., Ellis, J., Samuels, T.H., Mutengesa, E., Dobbin, J., Ssebambulidde, K., Rutakingirwa, M.K., Tugume, L., Boulware, D.R., Grint, D. and Cresswell, F.V., 2020, August. Treatment outcomes in adult tuberculous meningitis: a systematic review and meta-analysis. In *Open forum infectious diseases* (Vol. 7, No. 8, p. ofaa257). US: Oxford University Press.
6. World Health Organization, 2007. *Improving the diagnosis and treatment of smear-negative pulmonary and extrapulmonary tuberculosis among adults and adolescents: recommendations for HIV-prevalent and resource-constrained settings* (No. WHO/HTM/TB/2007.379). World Health Organization.
7. Getahun, H., Harrington, M., O'Brien, R. and Nunn, P., 2007. Diagnosis of smear-negative pulmonary tuberculosis in people with HIV infection or AIDS in resource-constrained settings: informing urgent policy changes. *The Lancet*, 369(9578), pp.2042-2049.
8. Wen, L., Li, M., Xu, T., Yu, X., Wang, L. and Li, K., 2019. Clinical features, outcomes and prognostic factors of tuberculous meningitis in adults worldwide: systematic review and meta-analysis. *Journal of neurology*, 266(12), pp.3009-3021.
9. Li, K., Tang, H., Yang, Y., Li, Q., Zhou, Y., Ren, M., Long, X., Shen, W., Hu, R., Wang, X. and Zeng, K., 2017. Clinical features, long-term clinical outcomes, and prognostic factors of tuberculous meningitis in West China: a multivariate analysis of 154 adults. *Expert review of anti-infective therapy*, 15(6), pp.629-635.
10. Sunbul, M., Atilla, A., Esen, S., Eroglu, C. and Leblebicioglu, H., 2005. Thwaites' diagnostic scoring and the prediction of tuberculous meningitis. *Medical Principles and Practice*, 14(3), pp.151-154.
11. Thwaites, G.E., Bang, N.D., Dung, N.H., Quy, H.T., Oanh, D.T.T., Thoa, N.T.C., Hien, N.Q., Thuc, N.T., Hai, N.N., Lan, N.T.N. and Lan, N.N., 2004. Dexamethasone for the treatment of tuberculous meningitis in adolescents and adults. *New England Journal of Medicine*, 351(17), pp.1741-1751.
12. Allen, B., 2014. Papua New Guinea national census 2011: Rates of population change in local level government areas.
13. Soria, J., Metcalf, T., Mori, N., Newby, R.E., Montano, S.M., Huaroto, L., Ticona, E. and Zunt, J.R., 2019. Mortality in hospitalized patients with tuberculous meningitis. *BMC infectious diseases*, 19(1), pp.1-7.
14. Hosoglu, S., Geyik M, F., Balik, I., Aygen, B., Erol, S., Aygencel T, G., Mert, A., Saltoglu, N., Dokmetas, I., Felek, S. and Sunbul, M., 2002. Predictors of outcome in patients with tuberculous meningitis. *The International Journal of Tuberculosis and Lung Disease*, 6(1), pp.64-70.
15. Woldeamanuel, Y.W. and Girma, B., 2014. A 43-year systematic review and meta-analysis: case-fatality and risk of death among adults with tuberculous meningitis in Africa. *Journal of neurology*, 261(5), pp.851-865.
16. Smith, I., 2003. Mycobacterium tuberculosis pathogenesis and molecular determinants of virulence. *Clinical microbiology reviews*, 16(3), pp.463-496.

17. Davis, A.G., Rohlwick, U.K., Prout, A., Figaji, A.A. and Wilkinson, R.J., 2019. The pathogenesis of tuberculous meningitis. *Journal of leukocyte biology*, 105(2), pp.267-280.
18. Thee, S., Basu Roy, R., Blázquez-Gamero, D., Falcón-Neyra, L., Neth, O., Noguera-Julian, A., Lillo, C., Galli, L., Venturini, E., Buonsenso, D. and Götzinger, F., 2021. Treatment and Outcome in Children with Tuberculous Meningitis: A Multicenter Pediatric Tuberculosis Network European Trials Group Study. *Clinical Infectious Diseases*.
19. Yasar, K.K., Pehlivanoglu, F. and Sengoz, G., 2010. Predictors of mortality in tuberculous meningitis: a multivariate analysis of 160 cases. *The International journal of tuberculosis and lung disease*, 14(10), pp.1330-1335.
20. Nataprawira, H.M., Ruslianti, V., Solek, P., Hawani, D., Milanti, M., Anggraeni, R., Memed, F.S. and Kartika, A., 2016. Outcome of tuberculous meningitis in children: the first comprehensive retrospective cohort study in Indonesia. *The International Journal of Tuberculosis and Lung Disease*, 20(7), pp.909-914.
21. Sheu, J.J., Yuan, R.Y. and Yang, C.C., 2009. Predictors for outcome and treatment delay in patients with tuberculous meningitis. *The American journal of the medical sciences*, 338(2), pp.134-139.
22. Brancusi, F., Farrar, J. and Heemskerk, D., 2012. Tuberculous meningitis in adults: a review of a decade of developments focusing on prognostic factors for outcome. *Future microbiology*, 7(9), pp.1101-1116.
23. Donovan, J., Rohlwick, U.K., Tucker, E.W., Hiep, N.T.T., Thwaites, G.E., Figaji, A.A. and Tuberculous Meningitis International Research Consortium, 2019. Checklists to guide the supportive and critical care of tuberculous meningitis. *Wellcome Open Research*, 4.
24. Harrigan, M.R., 2001. Cerebral salt wasting syndrome. *Critical care clinics*, 17(1), pp.125-138.
25. Ganong, C.A. and Kappy, M.S., 1993. Cerebral salt wasting in children: the need for recognition and treatment. *American Journal of Diseases of Children*, 147(2), pp.167-169.
26. Palmer, B.F., 2003. Hyponatremia in patients with central nervous system disease: SIADH versus CSW. *Trends in Endocrinology & Metabolism*, 14(4), pp.182-187.
27. Cotton, M.F., Donald, P.R., Schoeman, J.F., Van Zyl, L.E., Aalbers, C. and Lombard, C.J., 1993. Raised intracranial pressure, the syndrome of inappropriate antidiuretic hormone secretion, and arginine vasopressin in tuberculous meningitis. *Child's Nervous System*, 9(1), pp.10-15.
28. Garg, R.K. and Sinha, M.K., 2011. Tuberculous meningitis in patients infected with human immunodeficiency virus. *Journal of neurology*, 258(1), pp.3-13.
29. Thinyane, K.H., Motsemme, K.M. and Cooper, V.J.L., 2015. Clinical presentation, aetiology, and outcomes of meningitis in a setting of high HIV and TB prevalence. *Journal of tropical medicine*, 2015.
30. Cecchini, D., Ambrosioni, J., Brezzo, C., Corti, M., Rybko, A., Perez, M., Poggi, S. and Ambroggi, M., 2009. Tuberculous meningitis in HIV-infected and non-infected patients: comparison of cerebrospinal fluid findings. *The International Journal of Tuberculosis and Lung Disease*, 13(2), pp.269-271.