Increased Levels of N-Lactoylphenylalanine After Exercise are Related to Adipose Tissue Loss During Endurance Training in Humans With Overweight and Obesity

Lac-Phe and Exercise-Induced Fat Loss in Humans

Miriam Hoene[a], Xinjie Zhao[b], Jürgen Machann[c,d,e], Andreas L. Birkenfeld[c,d,f], Martin Heni[a,c,d,g], Andreas Peter[a,c,d], Andreas Niess[h,i], Anja Moller[c,d,f], Rainer Lehmann[a,c,d], Guowang Xu[b,*], Cora Weigert[a,c,d, *]

[a]Institute for Clinical Chemistry and Pathobiochemistry, Department for Diagnostic Laboratory Medicine, University Hospital Tübingen

[b]CAS Key Laboratory of Separation Science for Analytical Chemistry, Dalian Institute of Chemical Physics, Chinese Academy of Sciences, Dalian, China

[c]Institute for Diabetes Research and Metabolic Diseases of Helmholtz Zentrum München at the University of Tübingen, Tübingen, Germany

[d]German Center for Diabetes Research (DZD), Neuherberg, Germany

[e]Section on Experimental Radiology, Department of Diagnostic and Interventional Radiology, University Hospital Tübingen, Tübingen, Germany

[f]Department of Internal Medicine IV, University Hospital Tübingen, Tübingen, Germany

[g]Division of Endocrinology and Diabetology, Department of Internal Medicine 1, University Hospital Ulm, Ulm, Germany

[h]Department of Sports Medicine, University Hospital Tübingen, Tübingen, Germany

[i]Interfaculty Research Institute for Sports and Physical Activity, University of Tübingen, Tübingen, Germany

* These authors contributed equally to this work and share senior authorship

Correspondig author: Miriam Hoene, miriam.hoene@med.uni-tuebingen.de

Abstract

Objective:

The exercise-inducible metabolite N-Lactoylphenylalanine (Lac-Phe) has recently been shown to reduce food intake and adipose tissue mass in mice. We addressed whether Lac-Phe could have a similar function in humans.

Methods:

Sedentary subjects with overweight and obesity completed an 8-week supervised endurance exercise intervention (n=22). Before and after the intervention, blood plasma levels of Lac-Phe were determined by UHPLC-MS in the resting state and immediately after an acute endurance exercise test. Adipose tissue and muscle volume were quantified by MRI.

Results:

Acute exercise caused a pronounced increase in Lac-Phe, both before and after the intervention. Higher levels of Lac-Phe after acute exercise were associated with a greater reduction in abdominal subcutaneous and, to a lower degree, visceral adipose tissue during the intervention.

Conclusions:

Lac-Phe produced during physical activity could boost weight loss in humans with obesity and overweight, possibly by transmitting or enhancing the appetite-suppressing effects of lactate. Exercise-induced Lac-Phe could be employed to predict and, potentially, improve the effectiveness of lifestyle interventions in subjects with overweight and obesity.

Trial Registration

Clinicaltrials.gov

NCT03151590

22 May 2017

https://clinicaltrials.gov/ct2/show/NCT03151590

Keywords

N-Lactoylphenylalanine (Lac-Phe), lactic acid and lactate; sedentary subjects with overweight and obesity; prevention of type 2 diabetes and cardiometabolic diseases; subcutaneous and visceral abdominal adipose tissue; biomarker for the prediction and personalization of exercise-based lifestyle interventions; acute endurance exercise and training

Abbreviations

ACN, acetonitrile; BMI, body mass index; MeOH, methanol; MRI, magnetic resonance imaging, UHPLC-MS, ultra high-performance liquid chromatography-mass spectrometry.

Introduction

The incidence of type 2 diabetes and related cardio-metabolic diseases is increasing worldwide. Physical activity and weight loss are two important pillars of diabetes prevention [1]. There is, however, a large variability in the effectiveness of exercise-based lifestyle interventions regarding therapeutic targets such as improvement of blood glucose control or reduction of adipose tissue mass that cannot satisfactorily be explained up to now [2–4].

The pronounced changes in skeletal muscle and whole-body energy metabolism induced by physical activity cause an increase in the concentration of a multitude of small metabolites in the circulation. A growing number of these metabolites have been found to not only reflect metabolic spillover but also play an essential role in mediating crosstalk between different cells and organs and thereby, the beneficial effects of physical activity [5–7].

N-Lactoylphenylalanine (Lac-Phe) is a pseudo-dipeptide generated from lactate and phenylalanine that has been reported to increase in a particularly pronounced fashion during and shortly after physical exercise [8,9]. A potential role in humans has not been reported yet, however, Lac-Phe has recently been shown to lower body weight and adipose tissue mass in mice rendered obese by a high-fat diet [9]. This prompted us to assess whether differences in the reduction in adipose tissue volume during an exercise intervention in humans with overweight and obesity could be attributed to differences in the exercise-induced production of Lac-Phe.

Methods

Study design and participants

All participants gave written informed consent. The study was approved by the ethics committee of the University of Tübingen and registered at Clinicaltrials.gov (NCT03151590). Details of the study protocol including recruitment and exclusion criteria have been published recently [10].

In brief, healthy subjects with <120 min of physical activity per week and a BMI >27 kg/m^2 completed an 8-week supervised exercise intervention flanked by two acute exercise visits performed as follows: Blood was collected in the morning in the fasted state, 45 min before the commencement of exercise. In the meantime, the participants received a standardized breakfast (1 bun, 20 g butter, 1 slice of cheese, 150 g apple puree, water). Subsequently, they performed 30 min of bicycle ergometer exercise at the heartrate corresponding to 80% of their individual VO$_2$peak. A second blood sample was collected 5 min after this bout of exercise. EDTA blood samples were immediately placed on ice, processed within 30 min and plasma stored at -80 °C.

The training intervention consisted of three times per week 1 hour supervised endurance exercise, 30 minutes each cycling and walking at 80% VO$_2$peak. The target heartrate was maintained throughout the intervention. Determination of VO$_2$peak and magnetic resonance imaging (MRI)-based quantification of adipose and lean tissue in arms and legs have been described [10,11]. Abdominal adipose tissue was segmented into visceral and non-visceral, which mainly consists of subcutaneous adipose tissue, using an automated procedure [11]. After excluding one subject with newly diagnosed autoimmune thyreoiditis, complete blood sample sets from 22 out of 26 subjects were available for metabolomics analysis.

Quantification of plasma metabolites

The plasma levels of Lac-Phe, Lactate and Phe were determined using ultra high-performance liquid chromatography-mass spectrometry (UHPLC-MS). 50 µL of plasma were mixed with 250 µL of MeOH, vortexed 30 s and centrifuged for 20 min at 16,000 g, 4 °C. The supernatant was vacuum-dried in aliquots of 200 µl. Dry samples were resuspended in 50 µL 25% ACN/water. The analysis was performed on a Vanquish UHPLC coupled to a Q Exactive (both Thermo Fisher Scientific, Waltham, USA) operated in negative ion mode as previously described with slight modifications [12]. The separation was performed on a 2.1×100 mm ACQUITYTM UPLC HSS 1.8 µm T3 column (Waters, Milford, MA, USA). The mobile phases were (A) 6.5 mM ammonium bicarbonate in water and (B) 6.5 mM ammonium bicarbonate in 95% MeOH/water (B). The elution started with 2% B for 1 min, linearly changed to 100% B within 20 min, reverted back to 2% B, and equilibrated for 2.9 min (flow rate 0.35 mL/min, column temperature 50 °C). The Q Exactive was set to 140,000 resolution and full scan mode, mass scan range was 70-1050 m/z. Nitrogen sheath gas and nitrogen auxiliary gas were set at flow rates of 45 and 10 AU. Capillary and aux gas heater temperatures were 300 °C and 350 °C, respectively. The spray voltage was 3.00 kV. Parallel reaction monitoring was used to obtain high-resolution MS/MS spectra of Lac-Phe (m/z = 236.0928) with a resolution of 17500 and a collision energy of 30 eV. The internal standard d5-Phe (0.8 µg/mL in extraction solvent) was used to normalize signal intensities.

Statistical analysis

Statistical analyses were performed using JMP 16 (SAS Institute Inc, Cary, North Carolina, USA). Longitudinal comparisons were performed using paired t-tests. Multiple linear regression analyses were performed on log-transformed data and adjusted for sex, age, baseline values of the respective tissue compartment or BMI, or change in muscle volume, as indicated. Normal distribution of the residuals was confirmed with the Shapiro-Wilk test in all analyses. A p-value < 0.05 was considered statistically significant.

Results

Lac-Phe was identified by LC-MS/MS (Fig. 1A) and could be quantified in all samples (Fig. 1B). Acute exercise caused a significant increase in plasma Lac-Phe levels, both before and after the 8-week training intervention (Fig. 1B). Training had no effect on Lac-Phe concentrations in the resting state or after the acute bout of exercise (Fig. 1B), which was performed at the same relative intensity before and after the intervention. Plasma levels of Lac-Phe exhibited a significant correlation with phenylalanine (Fig. 1C, R^2=0.35) and a very strong correlation with lactate (Fig. 1D, R^2=0.82).

The training intervention resulted in an improvement in VO$_2$peak and an increase in lean tissue, i.e. muscle mass, in the legs (Tab. 1). At the same time, BMI and abdominal subcutaneous and visceral adipose tissue were decreased (Tab. 1). The decrease in subcutaneous adipose tissue was inversely correlated to the plasma concentration of Lac-Phe after acute exercise, both before (Fig. 1E) and after the intervention (Fig. 1F) and also after adjustment for sex, age, and adipose tissue baseline values (results of multiple regression analyses shown in Tab. 2). The decrease in visceral

adipose tissue was inversely correlated to the Lac-Phe concentration after the final acute exercise
test and tended to be correlated after the first acute exercise test (Tab. 2).

Plasma lactate levels after acute exercise exhibited a similar, but slightly weaker correlation to
the change in abdominal adipose tissue (Tab. 2). This association only reached statistical
significance for the subcutaneous depot after the pre-training acute exercise bout but not for
visceral adipose tissue (Tab. 2). Furthermore, lactate levels after acute exercise were positively
correlated to the increase in muscle mass of the lower extremities during the intervention.
Importantly, Lac-Phe levels after acute exercise still exhibited an inverse correlation to the change
in subcutaneous adipose tissue when additionally adjusting for the change in leg muscle volume
in the multiple linear regression models (p=0.020, standardized Beta coefficient=-0.56 for the pre-
training and a trend of p=0.088, Beta=-0.42 for the post-training exercise bout).

No significant correlation of Lac-Phe or lactate could be observed with the changes in BMI or in
the lean tissue of the arms (Tab. 2). As expected, the latter was not increased by the training
scheme, i.e. by cycling and treadmill exercise (Tab. 1).

Discussion

The metabolite Lac-Phe is produced during physical exercise and has recently gained attention as
a mediator of weight loss in mice [9] but its relevance and function in humans remain to be
demonstrated. We provide a first clue by showing that higher levels of Lac-Phe after exercise are
related to a greater reduction in abdominal subcutaneous and, to a lower extent, visceral adipose
tissue in subjects with obesity during a supervised exercise intervention.

Lac-Phe produced during physical exercise has been shown to reduce obesity by lowering food
intake in mice [4]. Food intake has also been identified as an important factor for weight loss in
humans with obesity [13,14]. While food intake has not been systematically assessed in our study,
it seems plausible that higher levels of Lac-Phe did cause a greater transient suppression of
hunger after each exercise session that contributed to a negative energy balance.

A reduction of food intake following physical exercise may seem counter-intuitive at first sight.
However, in humans, a compensatory increase in calorie intake only occurs with a delay in the
range of days and more pronouncedly in lean subjects [15] while in subjects with obesity, physical
activity may even reduce food intake by restoring the physiologic regulatory circuitry [13,14].
Acutely, physical activity causes a repression of hunger that serves the purpose of preserving
blood flow to skeletal muscle and that correlates with the circulating concentrations of lactate
[16]. Studies of lactate administration have supported an appetite-suppressing effect of this
metabolite and suggested different potential mechanisms of action [16,17]. Since lactate drives
the formation of Lac-Phe [8], which then peaks after lactate [9], it could be speculated that Lac-
Phe mediates a more sustained appetite-suppressing signal than lactate itself. One mode of
action for Lac-Phe could be via G protein-coupled receptors in the brain, the central regulator of
food intake [18,19].

Independent of its signalling function, Lac-Phe could serve as a biomarker to predict and,
potentially, optimize the individual response to exercise-based lifestyle interventions. This is
particularly relevant given the large variability in the extent to which different subjects benefit

from a given exercise scheme [2–4]. Exercise intensity is usually personalized, e.g. based on the
individual VO_2peak as in our study, but parameters to determine the ideal intensity or modality
of exercise that are most suitable for an individual have not been available up to now. Lac-Phe
could serve as such a parameter since subjects exhibiting higher levels after an exercise test
achieved a greater reduction in adipose tissue mass despite having exercised at the same relative
VO_2peak. In addition to modifying the exercise scheme, future studies could also aim at boosting
the Lac-Phe response in order to enhance weight loss and thereby prevent type 2 diabetes and
related cardiometabolic diseases.

Acknowledgements

187 The manuscript has been published as a preprint (DOI: 10.1101/2022.09.07.22279536).

Ethics Approval and Consent to Participate

189 All participants gave written informed consent. The study was approved by the ethics committee
190 of the University of Tübingen and registered at Clinicaltrials.gov (NCT03151590).

Conflict of Interests

192 Miriam Hoene, Xinjie Zhao, Jürgen Machann, Andreas L. Birkenfeld, Martin Heni, Andreas Peter,
193 Andreas Niess, Anja Moller, Rainer Lehmann, Guowang Xu, and Cora Weigert declare no
194 competing interests.

Author Contributions

MHo analyzed and interpreted the data and wrote and edited the manuscript. XZ performed LC-
MS analyses. JM performed whole-body MRI. ALB, MHe, AP and, AN provided scientific guidance
and contributed to the discussion. AM designed the study and analyzed anthropometric data. RL
provided scientific guidance and experimental design and contributed to the discussion. GX
provided scientific guidance and experimental design, contributed to the discussion and reviewed
the manuscript. CW designed the study, supervised the whole project, contributed to the
discussion, reviewed the manuscript and is the guarantor of the study. All authors approved the
final version of the manuscript.

Funding

205 This work was supported by the Sino-German Center for Research Promotion (M-0257), the key
206 foundation from the National Natural Science Foundation of China (21934006), the German
207 Federal Ministry of Education and Research (BMBF) (DZD e.V., 01GI0925), the German Diabetes
208 Association and the University of Tübingen.

Data Availability Statement

210 The data will only be made available to interested researchers upon reasonable request as far as
211 privacy and consent of research participants are not compromised.

References

1. Delahanty LM. Weight loss in the prevention and treatment of diabetes. Prev Med. 2017 Nov;104:120–3.

2. O'Donoghue G, Kennedy A, Andersen GS, Carr B, Cleary S, Durkan E, et al. Phenotypic Responses to a Lifestyle Intervention Do Not Account for Inter-Individual Variability in Glucose Tolerance for Individuals at High Risk of Type 2 Diabetes. Front Physiol. 2019;10:317.

3. Magalhães JP, Hetherington-Rauth M, Júdice PB, Correia IR, Rosa GB, Henriques-Neto D, et al. Interindividual Variability in Fat Mass Response to a 1-Year Randomized Controlled Trial With Different Exercise Intensities in Type 2 Diabetes: Implications on Glycemic Control and Vascular Function. Front Physiol. 2021;12:698971.

4. Solomon TPJ. Sources of Inter-individual Variability in the Therapeutic Response of Blood Glucose Control to Exercise in Type 2 Diabetes: Going Beyond Exercise Dose. Front Physiol. 2018;9:896.

5. Reddy A, Bozi LHM, Yaghi OK, Mills EL, Xiao H, Nicholson HE, et al. pH-Gated Succinate Secretion Regulates Muscle Remodeling in Response to Exercise. Cell. 2020 Oct 1;183(1):62-75.e17.

6. Maurer J, Hoene M, Weigert C. Signals from the Circle: Tricarboxylic Acid Cycle Intermediates as Myometabokines. Metabolites. 2021 Jul 23;11(8):474.

7. Yang YR, Kwon KS. Potential Roles of Exercise-Induced Plasma Metabolites Linking Exercise to Health Benefits. Front Physiol. 2020 Dec 3;11:602748.

8. Jansen RS, Addie R, Merkx R, Fish A, Mahakena S, Bleijerveld OB, et al. N-lactoyl-amino acids are ubiquitous metabolites that originate from CNDP2-mediated reverse proteolysis of lactate and amino acids. Proc Natl Acad Sci U S A. 2015 May 26;112(21):6601–6.

9. Li VL, He Y, Contrepois K, Liu H, Kim JT, Wiggenhorn AL, et al. An exercise-inducible metabolite that suppresses feeding and obesity. Nature. 2022 Jun;606(7915):785–90.

10. Hoffmann C, Schneeweiss P, Randrianarisoa E, Schnauder G, Kappler L, Machann J, et al. Response of Mitochondrial Respiration in Adipose Tissue and Muscle to 8 Weeks of Endurance Exercise in Obese Subjects. J Clin Endocrinol Metab. 2020 Nov 1;105(11):dgaa571.

11. Würslin C, Machann J, Rempp H, Claussen C, Yang B, Schick F. Topography mapping of whole body adipose tissue using a fully automated and standardized procedure. J Magn Reson Imaging. 2010 Feb;31(2):430–9.

245   12.   Zhao X, Zeng Z, Chen A, Lu X, Zhao C, Hu C, et al. Comprehensive Strategy to Construct In-
246   House Database for Accurate and Batch Identification of Small Molecular Metabolites. Anal
247   Chem. 2018 Jun 19;90(12):7635–43.

248   13.   Martins C, Morgan L, Truby H. A review of the effects of exercise on appetite regulation: an
249   obesity perspective. Int J Obes. 2008 Sep;32(9):1337–47.

250   14.   King NA, Hopkins M, Caudwell P, Stubbs RJ, Blundell JE. Individual variability following 12
251   weeks of supervised exercise: identification and characterization of compensation for
252   exercise-induced weight loss. Int J Obes. 2008 Jan;32(1):177–84.

253   15.   Melzer K, Kayser B, Saris WHM, Pichard C. Effects of physical activity on food intake. Clinical
254   Nutrition. 2005 Dec 1;24(6):885–95.

255   16.   McCarthy SF, Islam H, Hazell TJ. The emerging role of lactate as a mediator of exercise-
256   induced appetite suppression. Am J Physiol Endocrinol Metab. 2020 Oct 1;319(4):E814–9.

257   17.   Schultes B, Schmid SM, Wilms B, Jauch-Chara K, Oltmanns KM, Hallschmid M. Lactate
258   infusion during euglycemia but not hypoglycemia reduces subsequent food intake in healthy
259   men. Appetite. 2012 Jun;58(3):818–21.

260   18.   Kullmann S, Kleinridders A, Small DM, Fritsche A, Häring HU, Preissl H, et al. Central nervous
261   pathways of insulin action in the control of metabolism and food intake. Lancet Diabetes
262   Endocrinol. 2020 Jun;8(6):524–34.

263   19.   Lund J, Clemmensen C, Schwartz TW. Outrunning obesity with Lac-Phe? Cell Metab. 2022
264   Aug 2;34(8):1085–7.

Table 1. Anthropometric, fitness and metabolic data

	Untrained	Trained	p-value
Sex	14 female / 8 male		-
Age [years]	30 ± 8.9 (19 – 59)		-
VO$_2$peak$_{ergo}$/body mass [mL/(kg*min)]	25.0 ± 4.2 (18.3 – 32.3)	26.5 ± 4.7 (16.0 – 34.9)	0.042*
BMI [kg/m²]	31.7 ± 4.5 (27.5 – 45.5)	31.3 ± 4.7 (26.3 – 45.2)	0.006*
Subcutaneous abdominal adipose tissue [L]	15.3 ± 5.9 (8.4 – 32.2)	14.7 ± 6.1 (7.2 – 33.1)	0.006*
Visceral adipose tissue [L]	3.53 ± 1.65 (0.81 – 7.26)	3.38 ± 1.57 (0.94 – 6.68)	0.012*
Lean tissue legs [L]	17.9 ± 4.1 (12.3 – 27.5)	18.2 ± 3.9 (13.2 – 27.7)	0.034*
Lean tissue arms [L]	9.73 ± 1.96 (7.31 – 14.27)	9.85 ± 2.26 (7.19 – 14.42)	0.551
Glucose fasting [mmol/L]	5.09 ± 0.40 (4.61 – 6.00)	5.02 ± 0.40 (4.33 – 5.61)	0.336

269   N = 22, VO$_2$peak N=21; mean ± SD (range of values); * $p < 0.05$.

271   **Table 2.** Association of the increased Lac-Phe and lactate levels after acute exercise with the
272   training response

Training fold change	Lac-Phe untrained acute	Lac-Phe trained acute	Lactate untrained acute	Lactate trained acute
Subcutaneous abdominal adipose tissue [L]	Beta = -0.62 p = 0.004*	Beta = -0.52 p = 0.028*	Beta = -0.60 p = 0.008*	Beta = -0.39 p = 0.102
Visceral adipose tissue [L]	Beta = -0.42 p = 0.075	Beta = -0.48 p = 0.037*	Beta = -0.23 p = 0.372	Beta = -0.37 p = 0.123
BMI [kg/m²]	Beta = -0.25 p = 0.279	Beta = -0.15 p = 0.538	Beta = -0.13 p = 0.600	Beta = 0.07 p = 0.784
Lean tissue legs [L]	Beta = 0.37 p = 0.079	Beta = 0.22 p = 0.357	Beta = 0.42 p = 0.047*	Beta = 0.47 p = 0.036*
Lean tissue arms [L]	Beta = 0.14 p = 0.584	Beta = 0.08 p = 0.758	Beta = 0.27 p = 0.279	Beta = 0.08 p = 0.759

273   Multiple linear regression analyses with adjustments for sex, age and baseline values of the respective tissue
274   compartment or BMI. N=22, Beta, standardized beta coefficient, * $p < 0.05$.

Figure 1: A: Structure of N-Lactoylphenylalanine (Lac-Phe) and identification by LC-MS/MS.
B: Plasma levels of Lac-Phe in subjects with overweight and obesity (n=22) before (resting) or
immediately after (acute) a 30-min endurance exercise session in the untrained state or after an
8-week exercise intervention (trained). **C-D**: Correlation of plasma Lac-Phe with plasma
phenylalanine (**C**) and lactate (**D**). **E-F**: Correlation of the fold change in subcutaneous abdominal
adipose tissue volume during the exercise intervention with the concentration of Lac-Phe after
acute exercise before (**E**) and after the training intervention (**F**). R^2, standardized beta coefficient
(Beta) and p-value are shown for the simple linear regression (values from multiple regression
presented in Table 2). AU, arbitrary units.