Improved Robustness of SARS-CoV-2 Whole-Genome Sequencing from Wastewater with a Nonselective Virus Concentration Method

Emily Segelhurst,[1] Jonathan E. Bard,[2,3,6] Annemarie N. Pillsbury,[1] Natalie A. Lamb,[2] Chonglin Zhu,[1] Alyssa Pohlman,[2] Amanda Boccolucci,[2] Jamaal Emerson,[4] Brandon J. Marzullo,[2,3] Donald A. Yergeau,[2] Norma J. Nowak,[2,3] Ian M. Bradley,[1,5] Jennifer A. Surtees,[3,4,6] Yinyin Ye[1,*]

[1]Department of Civil, Structural and Environmental Engineering, University at Buffalo, Buffalo, New York 14260, United States

[2]UB Genomics and Bioinformatics Core, University at Buffalo, Buffalo, New York 14203, United States

[3]Department of Biochemistry, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, New York 14203, United States

[4]Department of Microbiology and Immunology, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, New York 14203, United States

[5]Research and Education in Energy, Environmental and Water (RENEW) Institute, University at Buffalo, Buffalo, New York 14260, United States

[6]Genetics, Genomics and Bioinformatics Graduate Program, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, New York 14203, United States

---

*Corresponding author: yinyinye@buffalo.edu

Tel. (716) 645-4002

ABSTRACT

The sequencing of human virus genomes from wastewater samples is an efficient method for tracking viral transmission and evolution at the community level. However, this requires the recovery of viral nucleic acids of high quality. We developed a reusable tangential-flow filtration system to concentrate and purify viruses from wastewater for whole-genome sequencing. A pilot study was conducted with 94 wastewater samples from four local sewersheds, from which viral nucleic acids were extracted, and the whole genome of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was sequenced using the ARTIC V4.0 primers. Our method yielded a high probability (0.9) of recovering complete or near-complete SARS-CoV-2 genomes (>90% coverage at 10× depth) from wastewater when the COVID-19 incidence rate exceeded 33 cases per 100 000 people. The relative abundances of sequenced SARS-CoV-2 variants followed the trends observed from patient-derived samples. We also identified SARS-CoV-2 lineages in wastewater that were underrepresented or not present in the clinical whole-genome sequencing data. The developed tangential-flow filtration system can be easily adopted for the sequencing of other viruses in wastewater, particularly those at low concentrations.

KEYWORDS

SARS-CoV-2, COVID-19, whole-genome sequencing, tangential-flow filtration, wastewater-based epidemiology

SYNOPSIS

The tangential-flow filtration method extracts viral nucleic acids of high enough quality from wastewater for robust and successful whole-genome sequencing.

GRAPHIC FOR TABLE OF CONTENTS (TOC)

INTRODUCTION

The global spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has facilitated the emergence of genome mutations, resulting in new lineages that further threaten public health.[1] Whole-genome sequencing (WGS) of clinical samples is a powerful method for tracking the spread of various SARS-CoV-2 lineages,[2] but this can be expensive, slow, and subject to biased sampling. Furthermore, the frequency of clinical testing has declined, aided by the availability of rapid at-home testing kits. WGS of SARS-CoV-2 in wastewater samples circumvents the issues as a high-throughput and more comprehensive strategy that monitors a larger portion of the population.[3-6] Successful WGS of SARS-CoV-2 in wastewater samples relies on the depth (the number of times a nucleotide is sequenced)[7] and breadth of genome coverage (hereafter, coverage; the percentage of nucleotide positions sequenced to a given depth).[7] Because SARS-CoV-2 lineages differ by only a few mutations, greater depth and coverage are needed to quantify lineage abundance and detect low-frequency mutations.[8, 9]

Different virus concentration methods have been developed for polymerase chain reaction (PCR)-based detection of viral genes from wastewater.[10-13] However, the nucleic acids recovered with these methods may not be suitable for WGS—PCR assays typically target <1% of the genome. Sequencing inhibitors, such as humic acids, and background nucleic acids from prokaryotes and eukaryotes are likely to reduce the genome coverage of SARS-CoV-2. Indeed, near-complete (>90%) SARS-CoV-2 genomes are not recovered from most samples (>85%) when aluminum hydroxide is used to directly precipitate viruses without separating viruses from other microorganisms in wastewater.[14] The removal of wastewater solids before the Amicon/Centricon ultra-centrifugal concentration[15] or electronegative membrane filtration[16] can improve sequencing

success, but the quality of the extracted genomes can vary and random sequencing failures can

occur.[17]

We report a reusable and nonselective tangential-flow filtration system to concentrate and

purify viruses from wastewater for WGS. We evaluated the robustness of the method to recover

complete or near-complete SARS-CoV-2 genomes using the samples collected from four local

sewersheds over six months. We subsequently compared the findings with those from patient-

derived clinical samples at the County level. Finally, the timing of lineage detection was compared

among clinical WGS, wastewater WGS, and quantitative reverse transcription polymerase chain

reactions (RT-qPCR).

MATERIALS AND METHODS

Wastewater sample collection

Time- or flow-weighted 24-hr composite influent samples were collected every 1 or 2

weeks between October 1 and December 16, 2021, and between January 18 and April 12, 2022,

from three wastewater treatment plants covering four sewersheds that serve ~80% of the total

population of Erie County (New York): Tonawanda, Kenmore-Tonawanda, Amherst, and Bird

Island (**Table S1**; **Figure S1**). Influent was collected by an autosampler every 30 min, and the

samples were kept at 4 °C in high-density polyethylene bottles that were cleaned with 10% bleach.

The samples were transported on ice packs and stored at 4 °C until they were processed.

Virus concentration from wastewater

Wastewater samples (125 mL) were centrifuged in sterile bottles at $10\,000 \times g$ for 15 min

at 4 °C to remove large wastewater solids, prokaryotic and eukaryotic cells, and other debris. The

supernatant was concentrated in a Vivaflow laboratory cross flow cassette system (Sartorius)

equipped with a 30-kDa Hydrosart ultrafilter membrane (Sartorius) at a feedline flow rate of 8.5

mL/min (**Figure S2**). The residual liquid in the ultrafilter membrane and connected tubing was

blown with air into the sample reservoir. The concentrate (~25 mL) in the reservoir was then

collected, overlayed with 5 mL of a 20% (v/v) sucrose solution and ultracentrifuged at 100 000 ×

g for 45 min at 4 °C in an SW 32 Ti rotor using an Optima XE series centrifuge (Beckman Coulter).

Immediately after the ultracentrifugation, the pellet was resuspended in 200 µL phosphate-

buffered saline (pH 7.4; Gibco) and stored at −80 °C ~two weeks until nucleic acid extraction.

After use, the ultrafilter membranes were immediately flushed with ~100 mL of 0.5 M

NaOH solution preheated to 55 °C at a feedline flow rate of 8.5 mL/min. The NaOH solution was

then recirculated for 20–30 min at the same flow rate. The cleaned membrane was stored in 0.5 M

NaOH at 4 °C. Cleaning efficiency was verified on six random days by processing 125 mL of

autoclaved Milli-Q water with the same concentration procedure. The absence of SARS-CoV-2

genes through washed filters were confirmed by RT-qPCR (**Figure S3**).

Nucleic acid extraction and RT-qPCR

Viral nucleic acids were extracted from the concentrated samples with QIAamp viral RNA

mini kits (Qiagen) and eluted in 60 µL AVE buffer (Qiagen) per the manufacturer's instructions.

One-step RT-qPCR was performed to quantify the SARS-CoV-2 N gene according to the CDC

N2 assay[18] and to determine the presence/absence of specific S gene mutations, including WT493-

498, Q493R and Q498R, delH69/V70, and delL24/P25/P26 and A27S as variant determinants for

Delta, generic Omicron, Omicron BA.1, and Omicron BA.2, respectively, following methods

published previously (**Table S2**).[19, 20] The limit of detection for the SARS-CoV-2 N gene was

determined to be 1.8 gene copies/µL, and the limit of quantification was determined to be 5 gene

copies/µL following a method as described previously (**Figure S4**).[21] A checklist of the MIQE

(<u>m</u>inimum <u>i</u>nformation for publication of <u>q</u>uantitative real-time PCR <u>e</u>xperiments) guidelines[22] is

provided in **Table S3**. All RT-qPCR assays were conducted in duplicates on a CFX96 Touch real-

time PCR detection system (Bio-Rad), and the threshold cycle (C_T) value was determined using a

CFX Maestro Software (Version 4.0, Bio-Rad). To remove PCR inhibition, the extracted nucleic

acids were diluted 5- or 10-fold in nuclease-free water before RT-qPCR analyses (**Figure S5**).

Detailed procedures of RT-qPCR inhibition test, reaction compositions, thermocycling conditions,

and determination of the limits of detection and quantification are described in the Supporting

Information.

SARS-CoV-2 whole-genome sequencing (WGS)

The quality of the extracted nucleic acids was assessed with the Agilent Fragment

Analyzer. WGS of SARS-CoV-2 was performed following a modified ARTIC protocol[23] using

the V4.0 nCOV-2019 amplicon panel (IDT). Briefly, 8 µL of nucleic acid extracts was reverse

transcribed to cDNA with random hexamers using the Invitrogen SuperScript IV first-strand

synthesis system (Thermo Fisher Scientific) according to the manufacturer's protocol. The cDNA

was amplified with two primer pools using Q5 hot-start high-fidelity 2× master mix (New England

BioLabs) with the following parameters: 98 °C for 30 s, 25 cycles of 98 °C for 15 s and 65 °C for

5 min. The resulting amplicons from both primer pools were combined. Excess primers and

reagents were removed with 1× AMPure XP beads (Beckman Coulter), and the amplicons were

eluted in EB buffer (Qiagen). The total eluted volume of amplicons was input to generate libraries

using the NEBNext Ultra II DNA library prep kit (New England BioLabs) without fragmentation

per the manufacturer's protocol. Individual samples were then barcoded and pooled for

quantification using the sparQ Universal Library Quant kit (QuantaBio) for Illumina sequencing.

The libraries were diluted to 10 pM and sequenced on an Illumina MiSeq instrument (V3

chemistry, PE300) with 1% PhiX as internal control.

Bioinformatic analysis

The Illumina sequencing output files were demultiplexed using bcl2fastq (v2.20.0.422) to

convert into FASTQ files. Initial quality control was performed through FastQ Screen.[24] Samples

were then processed using the UB Genomics and Bioinformatics Core SARS-CoV-2 analysis

pipeline (https://github.com/UBGBC/fastq-to-consensus). Briefly, adapters were trimmed before

sequencing reads were aligned to the SARS-CoV-2 reference genome (MN908947.2) using the

BWA-MEM algorithm.[25] Variants, insertions, and deletions were then called using BCFtools

(v.1.10.2),[26] requiring a minimum depth per nucleotide position of $10\times$ or $50\times$ and generating VCF

file outputs along with a final consensus FASTA file for each input sample. The resulting VCF

and depth mpileup files were used as input into Freyja to perform lineage composition analysis

(https://github.com/andersen-lab/Freyja).[5] The coverage (--covcut) was specified at $10\times$ or $50\times$,

and only confirmed lineages were reported. The Freyja pipeline was selected because of its high

accuracy and efficiency.[5]

Patient-derived SARS-CoV-2 data

The daily cases of COVID-19 during the sampling periods in the studied sewersheds were

extracted from the Erie County Wastewater Monitoring Dashboard.[27] The COVID-19 incidence

rates (per 100 000 population in the sewershed) were then calculated as 7-day rolling averages.

The patient-derived WGS data at the County level were downloaded from the Global Initiative on

Sharing All Influenza Data (GISAID) database[28] with the location as "North America/USA/New

York/Erie County," collection date from "2021-10-01" to "2022-04-30", and low coverage

excluded (GISAID Identifier: EPI_SET_220906wr). Most of these samples were sequenced at

University at Buffalo.[29] On each collection date, the relative abundances of SARS-CoV-2 lineages

in the patient samples were calculated, and the number of patient samples that were collected for

sequencing were smoothed as 7-day rolling averages.

Statistical analysis

The probability of recovering >90% coverage of the SARS-CoV-2 genome at 10× depth at

a given RT-qPCR C_T value or COVID-19 incidence rate was predicted by binary logistic

regression analysis. Data were visualized with Prism 9.2.0 (GraphPad Software) and the ggplot2[30]

package in R version 4.2.1.[31]

Data availability

Raw sequencing data are available in NCBI Sequence Read Archive (SRA) under the

BioProject ID: PRJNA877272. Codes for analyzing SARS-CoV-2 lineage in wastewater are

available at https://github.com/UBGBC/fastq-to-consensus.

RESULTS AND DISCUSSION

Coverage and depth of SARS-CoV-2 genomes from wastewater

Complete or near-complete SARS-CoV-2 genomes (>90% coverage) at 10× depth were

recovered from 68% (64/94) of the wastewater samples (**Table S4**). This success rate is 53% higher

than that in a SARS-CoV-2 wastewater sequencing study that applied aluminum hydroxide

flocculation and the ARTIC protocol.[14] A binary logistic regression analysis predicted a 0.9

probability of sequencing success (>90% coverage at 10× depth) when the C_T value of the SARS-

CoV-2 N gene was 31.8 (corresponding to ~3500 gene copies loaded for sequencing) (**Figure 1A**).

For comparison, at the 0.9 probability of sequencing success, C_T values of SARS-CoV-2 gene

ranged from <26 to ~29 with other workflows in previous pilot studies (**Figure S6**).[14-17] Our

workflow demonstrates higher coverage of SARS-CoV-2 genome at higher C_T values (lower gene

levels).

Most genomic regions were sequenced at a minimum depth of 50× (**Figure S7**). We noted

that three regions were often sequenced at lower depth (<10×): nucleotides 21750 to 22250 in the

S gene (amplicons 72 and 73) and 26750 to 27000 in the M gene (amplicon 89) (**Figure S7**).

Similar amplicon dropout issues were previously reported for a few amplicons using the ARTIC

V3 panel.[32, 33] We used ARTIC V4.0 primers, which address the V3 coverage issues but do not

include all the mutations in SARS-CoV-2 Delta and Omicron lineages in the primer design.[34] Other

wastewater factors may also explain the low-depth coverage within certain regions of SARS-CoV-

2 genome. First, the low concentrations of the SARS-CoV-2 genome in wastewater (<10^5 gene

copies/mL)[35] may have contributed to the lower sequencing depth. The presence of primer dimers

compete for primer-template interactions, resulting in reduced amplification efficiency.[36] Also,

mutations of SARS-CoV-2 genomes identified in wastewater but not in patient samples could

influence primer binding at some sites.[3, 6, 37] Our results suggest that up-to-date primer design and

optimization is critical to sequence emerging variants from wastewater.

Robustness of SARS-CoV-2 WGS from wastewater

Influence of sewersheds. The rates of sequencing success were similar for samples

collected from the four studied sewersheds: 70.8% (17/24) for Tonawanda, 72.0% (18/25) for

Kenmore-Tonawanda, 72.7% (16/22) for Amherst, and 65.2% (15/23) for Bird Island (**Table S4**).

Although the Bird Island sewershed serves a population that is larger than the other three (**Table**

S1), similar average C_T values were obtained for the SARS-CoV-2 N gene (33.5 for Tonawanda,

33.4 for Kenmore-Tonawanda, 34.2 for Amherst, and 34.1 for Bird Island) (**Table S4**), indicating

that similar genome levels were available for sequencing. Notably, the Bird Island wastewater

treatment plant collects wastewater exclusively through combined sewer systems (**Table S1**). A

large fraction of sequencing inhibitors, such as humic acids and heavy metals, in the stormwater

runoff may slightly reduce the WGS success rate.[38, 39]

Influence of COVID-19 incidence. The probability of sequencing success was >0.9 when

the COVID-19 incidence rate was >33/100 000 persons but decreased to 0.75 when the incidence

rate was 25.2/100 000 persons (**Figure 1B**). A previous study reported a probability of 0.75 to

quantify SARS-CoV-2 gene in wastewater by RT-qPCR when the 14-day COVID-19 case rate

was 152/100 000 persons.[40] In that study, viruses were concentrated from solids-removed influent

samples by 10-kDa Centricon ultra-filters.[40] We are not aware of other wastewater sequencing

studies that have calculated SARS-CoV-2 WGS success rates with regard to the incidence rate,

but our results suggest that SARS-CoV-2 WGS with the viruses recovered by tangential-flow

filtration method can be successfully applied at low COVID-19 incidence rates.

Figure 1. Coverage of the SARS-CoV-2 genome at 10× depth and the probability of sequencing

success (>90% coverage at 10× depth) with regard to C_T value of the SARS-CoV-2 N gene

measured with RT-qPCR (A) and the COVID-19 incidence rate (rolling 7-day average of cases

per 100 000 people) (B). The probability of sequencing success was calculated according to a

binary logistic regression analysis.

Lineage distributions estimated by wastewater and clinical data

Most of our wastewater samples were sequenced to an average depth of >50× per 250-nucleotide-region across the genome (**Figure S7**). Although the estimated lineage abundances at sequencing depths of 10× and 50× were very similar (**Figure S8**), we only reported lineage distributions estimated by the WGS data filtered at a depth of 50×.

Nine dominant groups of lineages were identified in the wastewater samples: AY.* (Delta), B.1.617.2 (Delta), B.1.2, B.1.1.529 (Omicron), BA.1/BA.1.1 (Omicron), BA.1.1.16 (Omicron), other BA.1.* (Omicron), BA.2.12/BA.2.12.1 (Omicron), and other BA.2.* (Omicron). The prevalence of Delta AY.* lineages in wastewater in October–December 2021 was coincident with the lineages observed via clinical surveillance (**Figure 2**). Two Omicron infection waves in 2022 (BA.1 in January/February and BA.2 in March/April) predicted from the wastewater data aligned with clinical data (**Figure 2**). Remarkably, wastewater sequencing revealed lineages that were underrepresented or not present in the clinical data. Specifically, the B.1.2 lineage was abundant in the Kenmore-Tonawanda sewershed in October but was not detected in patient samples from that time (**Figure 2A**). Moreover, in January/February of 2022, the BA.1/BA.1.1 lineages were prevalent in patient samples (**Figure 2A**), whereas the BA.1.1.16 lineage dominated in the wastewater samples (**Figure 2B–E**). Notably, the genome sequences of the BA.1/BA.1.1 and BA.1.1.16 lineages are similar, which might affect the predictions of their relative abundance.

The reason that B.1.2 lineage identified in wastewater samples but largely absent from patient samples is unclear. The Freyja pipeline identified B.1.2 on the basis of a mutation spectrum, consisting of 13 nucleotide substitutions (A18424G, A23403G, C10319T, C1059T, C14408T, C21304T, C241T, C27964T, C28472T, C28869T, C3037T, G25563T, and G25907T),

three of which (C1059T, C21304T, and G25563T) overlap the mutations in highly abundant

Delta lineages circulating at the time. Furthermore, the B.1.2 mutation spectrum does not share a

strong correlation with any of the mutation spectra of Delta lineages (**Figure S9**). This is less

likely a misassignment by the pipeline. Taken together, our findings suggest that the WGS of

patient samples can miss SARS-CoV-2 lineages circulating in the community. Those lineages

may not be clinically relevant, but future research is needed to understand their roles in viral

evolution and lineage emergence.

Figure 2. Distributions of dominant SARS-CoV-2 lineages estimated from Erie County clinical

data deposited on GISAID (A) and wastewater sequencing data (B–E) during the sampling periods.

The solid black lines represent the rolling 7-day average numbers of clinical samples collected for

sequencing (A) and COVID-19 incidence rates (cases per 100 000 people) in each sewershed (B–

E).

Early detection of variants in wastewater

The values of early detection of SARS-CoV-2 variants in wastewater depend on the delays in clinical analysis.[41] The workflow optimization for a quick turnaround of WGS analysis is beyond the scope of this study. Here, we only compare the detection of emerging SARS-CoV-2 variants based on the date of collection for both wastewater and patient samples.

The Omicron BA.1 was first identified in Amherst and Bird Island wastewater with a relative abundance of 1-2% in the second week of December, which is ~1 week later than the patient-derived samples (**Figure 3**). The detection of Omicron BA.2 from wastewater samples in March was similarly delayed (**Figure 3**). Note that some BA.2 patient-derived samples were collected in January and February, but there was no Omicron BA.2 outbreak until March. The poor detection of Omicron BA.2 in wastewater may be attributable to the relatively low coverage of the SARS-CoV-2 genome from wastewater during mid-February and early March (<65%; Table S3). Nevertheless, our findings suggest that wastewater is a relevant proxy for patient samples. Wastewater WGS should be considered in the face of practical delays in clinical sequencing for early detection of variants.

We found that RT-qPCR assays were more sensitive than wastewater WGS for detecting Omicron BA.1-specific mutations, but wastewater WGS was more sensitive for Omicron BA.2-specific mutations (**Table S4**). However, RT-qPCR assays are less sensitive than digital PCR assays,[42] which were reported to detect variant-specific mutations earlier than genetic sequencing.[9] Interestingly, our RT-qPCR assays detected S:delH69/V70 mutations (for Omicron BA.1) in the Kenmore-Tonawanda wastewater samples collected in October 2021. Although WGS also detected these mutations, other Omicron-specific mutations in the S gene (Q493R and Q498R)

296   were not detected at the time. Wastewater WGS may thus be more precise than PCR assays for

297   early detection of SARS-CoV-2 variants.

Figure 3. Detection of Omicron BA.1 and BA.2 in wastewater and clinical samples. The relative

abundance of Omicron was predicted by Freyja pipeline with the wastewater sequencing data

filtered at 50× depth. The relative abundance of BA.1 (top) is the sum of relative abundances of

B.1.1.529, BA.1, BA.1.1, and other BA.1.*. The relative abundance of BA.2 (bottom) comprises

BA.2.12, BA.2.12.1, and other BA.2.*. The number of clinical samples collected for sequencing

was counted on every collection date.

Environmental implications

Our research demonstrates that tangential-flow filtration to concentrate viruses from wastewater samples enables extraction of nucleic acids of high-enough quality for stable performance of SARS-CoV-2 WGS. Complete or near-complete genomes at a depth of $10\times$ were sequenced from 68% (64/94) of the wastewater samples. Moreover, the tangential-flow filtration method improves the WGS success at lower COVID-19 incidence rates. Our results report a 0.9 probability of WGS success when the COVID-19 incidence rate exceeded 33/100 000 persons. Furthermore, WGS of SARS-CoV-2 in wastewater revealed lineages underrepresented or not detected from patient samples. Future studies are needed to advance wastewater WGS data interpretation and streamline the workflow of wastewater WGS to shorten the turnaround time for early detection.

The developed reusable and likely cost-effective tangential-flow filtration method is readily applicable to other sequencing assays that requires high-quality viral nucleic acids from wastewater. Given the flexibility of concentrating large volumes of wastewater, the tangential-flow filtration system can be further optimized for genomic surveillance of low-abundance viruses in wastewater.

SUPPORTING INFORMATION

The supporting information is available free of charge at http://pubs.acs.org.

Inhibition test; RT-qPCR assay details and Limit of detection (LOD) and limit of quantification (LOQ) determination; Details of four sewersheds in this study; Table S2. Details of RT-qPCR assay for quantifying SARS-CoV-2 N gene and S gene mutations; Check list of RT-qPCR experiments according to the MIQE (minimum information for

publication of quantitative real-time PCR experiments) guidelines; Summary of wastewater samples sequenced in this study; Geographic locations of four sewersheds in Erie County, New York; Schematic of tangential-flow filtration system; SARS-CoV-2 N gene levels in wastewater and blank test samples; Determination of limit of detection (LOD; left) and limit of quantification (LOQ; right) of SARS-CoV-2 N gene by RT-qPCR; RT-qPCR inhibition test of SARS-CoV-2 N gene in the wastewater nucleic acid extracts; Comparison of SARS-CoV-2 genome coverage from wastewater with different workflows; Heat map of read depths of SARS-CoV-2 whole-genome sequencing from wastewater samples; Comparison of relative abundances of SARS-CoV-2 lineages estimated by the sequencing data at depths of $10\times$ and $50\times$; Pair-wise pearson correlation of B.1.2 lineage mutaton spectrum against all other lineage mutation spectra.

ACKNOWLEDGEMENTS

This work was funded by Erie County Department of Health (ECDOH) award 91287 to Y.Y. and I.M.B. and award 91279 to J.A.S.. We thank undergraduate students Vicky Huang and Danya Hanin for processing the wastewater samples in the lab. We also thank Joseph L. Fiegl for coordinating the wastewater sample collection. The GISAID data used in the manuscript (Identifier: EPI_SET_220906wr) is composed of 4952 individual genome sequences published in GISAID's EpiCoV database. To view the contributors of each individual sequence with details, visit doi.org/10.55876/gis8.220906wr. Most of the samples (4653/4952) were sequenced at the University at Buffalo. We are grateful for the sequencing work of scientists and healthcare professionals who generated the remaining sequences from the region and made them publicly accessible through GISAID.

REFERENCES

(1) Walensky, R. P.; Walke, H. T.; Fauci, A. S. SARS-CoV-2 variants of concern in the United States-challenges and opportunities. *JAMA* **2021**, *325* (11), 1037-1038. DOI: 10.1001/jama.2021.2294.

(2) Oude Munnink, B. B.; Nieuwenhuijse, D. F.; Stein, M.; O'Toole, A.; Haverkate, M.; Mollers, M.; Kamga, S. K.; Schapendonk, C.; Pronk, M.; Lexmond, P.; et al. Rapid SARS-CoV-2 whole-genome sequencing and analysis for informed public health decision-making in the Netherlands. *Nat Med* **2020**, *26* (9), 1405-1410. DOI: 10.1038/s41591-020-0997-y.

(3) Crits-Christoph, A.; Kantor, R. S.; Olm, M. R.; Whitney, O. N.; Al-Shayeb, B.; Lou, Y. C.; Flamholz, A.; Kennedy, L. C.; Greenwald, H.; Hinkle, A.; et al. Genome Sequencing of Sewage Detects Regionally Prevalent SARS-CoV-2 Variants. *mBio* **2021**, *12* (1), e02703-e02720. DOI: 10.1128/mBio.02703-20.

(4) Nemudryi, A.; Nemudraia, A.; Wiegand, T.; Surya, K.; Buyukyoruk, M.; Cicha, C.; Vanderwood, K. K.; Wilkinson, R.; Wiedenheft, B. Temporal Detection and Phylogenetic Assessment of SARS-CoV-2 in Municipal Wastewater. *Cell Rep Med* **2020**, *1* (6), 100098. DOI: 10.1016/j.xcrm.2020.100098.

(5) Karthikeyan, S.; Levy, J. I.; De Hoff, P.; Humphrey, G.; Birmingham, A.; Jepsen, K.; Farmer, S.; Tubb, H. M.; Valles, T.; Tribelhorn, C. E.; et al. Wastewater sequencing reveals early cryptic SARS-CoV-2 variant transmission. *Nature* **2022**, *609*, 101-108. DOI: 10.1038/s41586-022-05049-6.

(6) Smyth, D. S.; Trujillo, M.; Gregory, D. A.; Cheung, K.; Gao, A.; Graham, M.; Guan, Y.; Guldenpfennig, C.; Hoxie, I.; Kannoly, S.; et al. Tracking cryptic SARS-CoV-2 lineages detected in NYC wastewater. *Nature Commun* **2022**, *13* (1), 1-9. DOI: 10.1038/s41467-022-28246-3.

(7) Sims, D.; Sudbery, I.; Ilott, N. E.; Heger, A.; Ponting, C. P. Sequencing depth and coverage: key considerations in genomic analyses. *Nat Rev Genet* **2014**, *15* (2), 121-132. DOI: 10.1038/nrg3642.

(8) Valieris, R.; Drummond, R. D.; Defelicibus, A.; Dias-Neto, E.; Rosales, R. A.; Tojal da Silva, I. A mixture model for determining SARS-Cov-2 variant composition in pooled samples. *Bioinformatics* **2022**, *38* (7), 1809-1815. DOI: 10.1093/bioinformatics/btac047.

(9) Lou, E. G.; Sapoval, N.; McCall, C.; Bauhs, L.; Carlson-Stadler, R.; Kalvapalle, P.; Lai, Y.; Palmer, K.; Penn, R.; Rich, W.; et al. Direct comparison of RT-ddPCR and targeted amplicon sequencing for SARS-CoV-2 mutation monitoring in wastewater. *Sci Total Environ* **2022**, 155059. DOI: 10.1016/j.scitotenv.2022.155059.

(10) Graham, K. E.; Loeb, S. K.; Wolfe, M. K.; Catoe, D.; Sinnott-Armstrong, N.; Kim, S.; Yamahara, K. M.; Sassoubre, L. M.; Grijalva, L. M. M.; Roldan-Hernandez, L.; et al. SARS-CoV-2 RNA in wastewater settled solids is associated with COVID-19 cases in a large urban sewershed. *Environ Sci Technol* **2021**, *55* (1), 488-498. DOI: 10.1021/acs.est.0c06191.

(11) LaTurner, Z. W.; Zong, D. M.; Kalvapalle, P.; Gamas, K. R.; Terwilliger, A.; Crosby, T.; Ali, P.; Avadhanula, V.; Santos, H. H.; Weesner, K.; et al. Evaluating recovery, cost, and throughput of different concentration methods for SARS-CoV-2 wastewater-based epidemiology. *Water Res* **2021**, *197*, 117043. DOI: 10.1016/j.watres.2021.117043.

(12) Zheng, X.; Deng, Y.; Xu, X.; Li, S.; Zhang, Y.; Ding, J.; On, H. Y.; Lai, J. C. C.; In Yau, C.; Chin, A. W. H.; et al. Comparison of virus concentration methods and RNA extraction methods for SARS-CoV-2 wastewater surveillance. *Sci Total Environ* **2022**, *824*, 153687. DOI: 10.1016/j.scitotenv.2022.153687.

(13) Peccia, J.; Zulli, A.; Brackney, D. E.; Grubaugh, N. D.; Kaplan, E. H.; Casanovas-Massana, A.; Ko, A. I.; Malik, A. A.; Wang, D.; Wang, M.; et al. Measurement of SARS-CoV-2 RNA in wastewater tracks community infection dynamics. *Nat Biotechnol* **2020**, *38* (10), 1164-1167. DOI: 10.1038/s41587-020-0684-z.

(14) Perez-Cataluna, A.; Chiner-Oms, A.; Cuevas-Ferrando, E.; Diaz-Reolid, A.; Falco, I.; Randazzo, W.; Giron-Guzman, I.; Allende, A.; Bracho, M. A.; Comas, I.; et al. Spatial and temporal distribution of SARS-

CoV-2 diversity circulating in wastewater. *Water Res* **2022**, *211*, 118007. DOI:
10.1016/j.watres.2021.118007.
(15) Fontenele, R. S.; Kraberger, S.; Hadfield, J.; Driver, E. M.; Bowes, D.; Holland, L. A.; Faleye, T. O. C.;
Adhikari, S.; Kumar, R.; Inchausti, R.; et al. High-throughput sequencing of SARS-CoV-2 in wastewater
provides insights into circulating variants. *Water Res* **2021**, *205*, 117710. DOI:
10.1016/j.watres.2021.117710.
(16) Bar-Or, I.; Weil, M.; Indenbaum, V.; Bucris, E.; Bar-Ilan, D.; Elul, M.; Levi, N.; Aguvaev, I.; Cohen, Z.;
Shirazi, R.; et al. Detection of SARS-CoV-2 variants by genomic analysis of wastewater samples in Israel.
Sci Total Environ **2021**, *789*, 148002. DOI: 10.1016/j.scitotenv.2021.148002.
(17) Izquierdo-Lara, R.; Elsinga, G.; Heijnen, L.; Munnink, B. B. O.; Schapendonk, C. M. E.; Nieuwenhuijse,
D.; Kon, M.; Lu, L.; Aarestrup, F. M.; Lycett, S.; et al. Monitoring SARS-CoV-2 circulation and diversity
through community wastewater sequencing, the Netherlands and Belgium. *Emerg Infect Dis* **2021**, *27*
(5), 1405-1415. DOI: 10.3201/eid2705.204410.
(18) Lu, X.; Wang, L.; Sakthivel, S. K.; Whitaker, B.; Murray, J.; Kamili, S.; Lynch, B.; Malapati, L.; Burke, S.
A.; Harcourt, J.; et al. US CDC real-time reverse transcription PCR panel for detection of Severe Acute
Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* **2020**, *26* (8), 1654. DOI:
10.3201/eid2608.201246.
(19) Lee, W. L.; Gu, X.; Armas, F.; Wu, F.; Chandra, F.; Chen, H.; Xiao, A.; Leifels, M.; Chua, F. J. D.; Kwok,
G. W. C.; et al. Quantitative detection of SARS-CoV-2 Omicron BA.1 and BA.2 variants in wastewater
through allele-specific RT-qPCR. *medRxiv* **2022**. DOI: 10.1101/2021.12.21.21268077.
(20) Lee, W. L.; Imakaev, M.; Armas, F.; McElroy, K. A.; Gu, X. Q.; Duvallet, C.; Chandra, F.; Chen, H. J.;
Leifels, M.; Mendola, S.; et al. Quantitative SARS-CoV-2 Alpha variant B.1.1.7 tracking in wastewater by
allele-specific RT-qPCR. *Environ Sci Technol Lett* **2021**, *8* (8), 675-682. DOI: 10.1021/acs.estlett.1c00375.
(21) He, H.; Zhou, P.; Shimabuku, K. K.; Fang, X.; Li, S.; Lee, Y.; Dodd, M. C. Degradation and deactivation
of bacterial antibiotic resistance genes during exposure to free chlorine, monochloramine, chlorine
dioxide, ozone, ultraviolet light, and hydroxyl radical. *Environ Sci Technol* **2019**, *53* (4), 2013-2026. DOI:
10.1021/acs.est.8b04393.
(22) Bustin, S. A.; Benes, V.; Garson, J. A.; Hellemans, J.; Huggett, J.; Kubista, M.; Mueller, R.; Nolan, T.;
Pfaffl, M. W.; Shipley, G. L.; et al. The MIQE guidelines: minimum information for publication of
quantitative real-time PCR experiments. *Clin Chem* **2009**, *55* (4), 611-622. DOI:
10.1373/clinchem.2008.112797.
(23) *nCoV-2019 sequencing protocol V.1*. 2020. https://www.protocols.io/view/ncov-2019-sequencing-
protocol-bp2l6n26rgqe/v1?version_warning=no (accessed 2020-01-22).
(24) Wingett, S. W.; Andrews, S. FastQ Screen: A tool for multi-genome mapping and quality control.
F1000Research **2018**, *7*, 1338. DOI: 10.12688/f1000research.15931.2.
(25) Li, H.; Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **2009**, *25* (14), 1754-1760. DOI: 10.1093/bioinformatics/btp324.
(26) Danecek, P.; Bonfield, J. K.; Liddle, J.; Marshall, J.; Ohan, V.; Pollard, M. O.; Whitwham, A.; Keane, T.;
McCarthy, S. A.; Davies, R. M.; et al. Twelve years of SAMtools and BCFtools. *Gigascience* **2021**, *10* (2).
DOI: 10.1093/gigascience/giab008 From NLM Medline.
(27) *Erie County SARS-CoV-2 wastewater monitoring dashboard*.
https://erieny.maps.arcgis.com/apps/dashboards/a95853269eec489ea59e5b71571f2e76 (accessed
2022-04-15).
(28) Elbe, S.; Buckland-Merrett, G. Data, disease and diplomacy: GISAID's innovative contribution to
global health. *Glob Chall* **2017**, *1* (1), 33-46. DOI: 10.1002/gch2.1018.
(29) Lamb, N. A., Bard, J.E., Pohlman, A., Boccolucci, A., Yergeau, D.A., Marzullo, B.J., Pope, C., Burstein,
G., Tomaszewski, J., Nowak, N.J. and Surtees, J.A. Genomic Surveillance of SARS-CoV-2 in Erie County,
New York. *medRxiv* **2021**. DOI: 10.1101/2021.07.01.21259869.

(30) Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag New York, 2016.

(31) Team, R. C. *RStudio: Integrated Development for R. RStudio*; PBC, Boston, MA, 2020.

(32) Jahn, K.; Dreifuss, D.; Topolsky, I.; Kull, A.; Ganesanandamoorthy, P.; Fernandez-Cassi, X.; Bänziger,
C.; Devaux, A. J.; Stachler, E.; Caduff, L.; et al. Early detection and surveillance of SARS-CoV-2 genomic
variants in wastewater using COJAC. *Nat Microbiol* **2022**, 1-10. DOI: 10.1038/s41564-022-01185-x.

(33) Lambisia, A. W.; Mohammed, K. S.; Makori, T. O.; Ndwiga, L.; Mburu, M. W.; Morobe, J. M.; Moraa,
E. O.; Musyoki, J.; Murunga, N.; Mwangi, J. N.; et al. Optimization of the SARS-CoV-2 ARTIC Network V4
Primers and Whole Genome Sequencing Protocol. *Front Med (Lausanne)* **2022**, *9*, 836728. DOI:
10.3389/fmed.2022.836728.

(34) Clark, C. R.; Hardison, M. T.; Houdeshell, H. N.; Vest, A. C.; Whitlock, D. A.; Skola, D. D.; Koble, J. S.;
Oberholzer, M.; Schroth, G. P. Evaluation of an optimized protocol and Illumina ARTIC V4 primer pool for
sequencing of SARS-CoV-2 using COVIDSeq™ and DRAGEN™ COVID Lineage App workflow. *bioRxiv* **2022**.
DOI: 10.1101/2022.01.07.475443.

(35) Mantilla-Calderon, D.; Huang, K. Y.; Li, A. J.; Chibwe, K.; Yu, X. Q.; Ye, Y.; Liu, L.; Ling, F. Q. Emerging
investigator series: meta-analyses on SARS-CoV-2 viral RNA levels in wastewater and their correlations
to epidemiological indicators. *Environ Sci: Wat Res Technol* **2022**, *8* (7), 1391-1407. DOI:
10.1039/d2ew00084a.

(36) Itokawa, K.; Sekizuka, T.; Hashino, M.; Tanaka, R.; Kuroda, M. Disentangling primer interactions
improves SARS-CoV-2 genome sequencing by multiplex tiling PCR. *PLoS One* **2020**, *15* (9), e0239403.
DOI: 10.1371/journal.pone.0239403.

(37) Agrawal, S.; Orschler, L.; Lackner, S. Metatranscriptomic analysis reveals SARS-CoV-2 mutations in
wastewater of the Frankfurt metropolitan area in southern Germany. *Microbiol Resour Announc* **2021**,
10 (15), e00280-00221. DOI: 10.1128/MRA.00280-21.

(38) Grant, S. B.; Rekhi, N. V.; Pise, N. R.; Reeves, R. L.; Matsumoto, M.; Wistrom, A.; Moussa, L.; Bay, S.;
Kayhanian, M. A review of the contaminants and toxicity associated with particles in stormwater runoff.
Terminology **2003**, *2* (2), 1-173.

(39) Schrader, C.; Schielke, A.; Ellerbroek, L.; Johne, R. PCR inhibitors - occurrence, properties and
removal. *J Appl Microbiol* **2012**, *113* (5), 1014-1026. DOI: 10.1111/j.1365-2672.2012.05384.x.

(40) Tiwari, A.; Lipponen, A.; Hokajarvi, A. M.; Luomala, O.; Sarekoski, A.; Rytkonen, A.; Osterlund, P.; Al-
Hello, H.; Juutinen, A.; Miettinen, I. T.; et al. Detection and quantification of SARS-CoV-2 RNA in
wastewater influent in relation to reported COVID-19 incidence in Finland. *Water Res* **2022**, *215*,
118220. DOI: 10.1016/j.watres.2022.118220.

(41) Larsen, D. A.; Wigginton, K. R. Tracking COVID-19 with wastewater. *Nat Biotechnol* **2020**, *38* (10),
1151-1153. DOI: 10.1038/s41587-020-0690-1.

(42) Ahmed, W., Smith, W.J., Metcalfe, S., Jackson, G., Choi, P.M., Morrison, M., Field, D., Gyawali, P.,
Bivins, A., Bibby, K. and Simpson, S.L. Comparison of RT-qPCR and RT-dPCR platforms for the trace
detection of SARS-CoV-2 RNA in wastewater. *ACS ES&T Water* **2022**. DOI: 10.1021/acsestwater.1c00387.

484