1   **Anemia among Pregnant Women in Cambodia: A Descriptive Analysis of Temporal and Geospatial**

2   **Trends and Logistic Regression-Based Examination of Factors Associated with Anemia in Pregnant**

3   **Women**

4   Samnang Um[1][¶], Heng Sopheab[1], An Yom[1], Jonathan A. Muir[2][&]

5   1 The National Institute of Public Health, Tuol Kork District, Phnom Penh, Cambodia, 2 The Global

6   Health Institute, Emory University, Atlanta, Georgia, United States.

7   * umsamnang56@gmail.com

8   [¶]These authors contributed equally to this work.

9   [&]These authors also contributed equally to this work.

ABSTRACT

Anemia is a major public health problem for thirty-two million pregnant women worldwide. Anemia during pregnancy is a leading cause of child low birth weight, preterm birth, and perinatal/neonatal mortality. Pregnant women are at higher risk of anemia due to micronutrient deficiencies, hemoglobinopathies, infections, socio-demographic and behavioral factors. This study aimed to: 1) assess temporal and geospatial trends of anemia in Cambodia and 2) identify factors associated with anemia among pregnant women aged 15-49 years old in Cambodia. We analyzed data from the Cambodia Demographic and Health Survey (CDHS) for 2005, 2010, and 2014. Data were pooled across the three survey years for all pregnant women aged 15-49 years. Survey weights were applied to account for the complex survey design of the CDHS. Descriptive statistics were estimated for key sociodemographic characteristics of the study population. We used logistic regressions to assess factors associated with anemia among pregnant women aged 15-49 years old. Anemia in pregnant women aged 15-49 in Cambodia decreased from 56% in 2005 to 53% in 2014. With the highest in Preah Vihear and Stung Treng provinces (74.3%), in Kratie province (73%), and in Prey Veng (65.4%) in 2005, 2010, and 2014 respectively. Compared to pregnant women from the wealthiest households, women from poorest households were more likely to have anemia (AOR = 2.8; 95% CI: 1.76-4.9). Pregnant women from coastal regions were almost twice as likely of having anemia (AOR = 1.9; 95% CI: 1.2-3.0). Pregnant women were more likely anemic if they were in their 2nd trimester (AOR = 2.6; 95% CI: 1.9-3.6) or 3rd trimester (AOR = 1.6 95% CI: 1.2-2.3). Anemia remains highly prevalent among pregnant women in Cambodia. Public health interventions and policies to alleviate anemia should be prioritized and shaped to address these factors.

Keywords: Anemia, Cambodia, Pregnant Women, Maternal Anemia, CDHS

Introduction

Anemia during pregnancy occurs in both developed and developing countries and is associated with abnormal outcomes in pregnancy; for example, greater risk for low birth weight, preterm birth, and perinatal/neonatal mortality [1]. According to the World Health Organization (WHO), anemia is defined as hemoglobin (Hb) levels less than 11.0 g/dL in pregnant women [2]. In 2019, it was estimated that anemia affected approximately 37% (32 million) of pregnant women worldwide and as many as half of all pregnant women in low-income and middle-income countries were diagnosed with anemia [3]. In Southeast Asia countries, 48% of pregnant women were anemic [4]. These patterns are reflected in Cambodia where the prevalence of anemia is higher among pregnant women compared with non-pregnant women (i.e., 53.2% compared to 45%) [5].

Anemia in pregnant women has severe consequences on health as well as social and economic development [6]. contributing to low physical activity and increases maternal morbidity and mortality, especially among those with severe anemia [1] and those living in developing countries [7]. The WHO estimates that 20% of maternal deaths are attributable to anemia [2]. In Southeast Asia, half of all maternal deaths were due to anemia in 2016 [8]. In Cambodia, anemia is affected 51.1% of pregnant women aged 15-49 years old in 2019 [9]. Despite significant economic development in Cambodia over the past two decades, the prevalence of anemia in women has not substantially declined.

Pregnant women with anemia are twice as likely to die during or shortly after pregnancy compared to those without anemia [8]. In many countries, anemia varies by socioeconomic factors such as education, household wealth status, occupation, and residence [10]. Pregnant women aged 30 and older are more likely to develop anemia than younger aged pregnant women [11-13]. Risk of

anemia was greater among women living in the lowest wealth quintile, with limited education [14],

and rural area [12,13]. Similarly, repeated childbearing, inadequate water hygiene and sanitation

status, and parasitic infection increases the risk of anemia in pregnant women [15]. Factors associated

with anemia during pregnancy in developing countries include nutritional deficiencies of iron,

vitamin B12, folate, and parasitic diseases (for example, malaria) [6,16,17]. Compared to women who

initiated antenatal care (ANC) in the first trimester, the odds of having anemia in pregnancy were

significantly higher among pregnant women who initiated ANC in the second trimester (AOR=2.71)

and third trimester (AOR=5.01) [18].

The prevalence of anemia in Cambodia has slowly decreased in both pregnant and non-

pregnant women. To further reduce anemia in Cambodia, improved understanding of its geographical

distribution and associated risk factors is required. Enhanced understanding will help identify

populations at greater risk for anemia and prioritize geographic areas for targeted interventions. To

our knowledge, there are no published peer-reviewed studies that assessed social and demographic

factors associated with anemia among pregnant women in Cambodia over time. One prior study on

anemia patterns and risk factors utilized data from the CHDS 2014 [14]. This study included all

women of reproductive age 15-49 years and pooled DHS data from seven South and Southeast Asian

countries. An additional study aimed to describe trends in nutritional status of women of

reproductive age in Cambodia over four nationally representative Demographic Health Surveys

(2000, 2005, 2010, and 2014) and to assess factors associated with nutritional disparities [19]. Finally,

a research abstract published in association with a research conference in Cambodia referenced using

CDHS 2014; this was a preliminary version of the present study [20]. Given the paucity of scholarship

addressing this health concern in Cambodia, in this study we aimed to:

1. Assess temporal and geospatial trends of anemia among pregnant women in Cambodia

2. Identify factors associated with anemia among pregnant women in Cambodia

Understanding these factors may further support policy development and programs with more

effective strategies and interventions in Cambodia to reduce the prevalence of anemia among

pregnant women and associated health risks such as maternal mortality.

Materials and methods

Data

To analyze trends and factors associated with anemia in pregnant women in Cambodia, we

used existing women's data from the 2005, 2010, and 2014 CDHS. The CDHS is a nationally

representative population-based household survey that is regularly conducted roughly every 5 years.

The survey typically uses two-stage stratified cluster sampling to collect the samples from all

provinces that divided into sampling domains. In the first stage, clusters, or enumeration areas (EAs)

that represent the entire country are randomly selected from the sampling frame using probability

proportional to cluster size (PPS). The second stage then involves the systematic sampling of

households listed in each cluster or EA. Interviews were then conducted with women aged 15-49 in

selected households. Using the women's survey questionnaire, variables collected by the CDHS

include births to women aged 15-49, sociodemographic characteristics, household assets that are used

to calculate a household wealth index, health-related indicators and nutritional status, number of

ANC visits and other pregnancy/delivery indicators for past births, and involvement in household

decision making using questionnaires. It further includes height and weight measurements of women

and children, hemoglobin level, malaria test, and vitamin A level [5]. We restricted our sample to

pregnant women with recorded hemoglobin level; which resulted in a total sample size of 1,629

(1,567 weighted) pregnant women aged 15-49 (Fig 1).

Fig 1. Selection process and the final sample size from the 2005, 2010, and 2014 CDHS.

Measurements

Anemia is the outcome variable used in the study. For women in CDHS, hemoglobin was

measured by collecting capillary blood from a finger prick with the HemoCue 201+analyzer [19]. The

original variable for anemia level in the CHDS was recorded as a categorical variable with "not

anemic", "mild anemia", "moderate anemia", and "severe anemia". Then, the original variable was

recoded into the dichotomous variable **Anemia**, with mild, moderate, and severe anemia coded as

Anemic = 1 and non-Anemic = 0.

Right-hand-sided variables included women's and household characteristics. Women's age was

categorized into an ordinal level variable with 15-20 years = 1, 21-31 years = 2 (reference category),

and 32-49 years = 3. Women's marital status was coded into a dichotomous variable with Married = 1,

Not Married = 0 (includes divorce, single, widowed, and separated). Women's education was coded as

an ordinal level variable with Higher = 1 (reference category), Secondary = 2, Primary = 3, No

Education = 4. Women's occupation was coded as a categorical variable with Professional = 1

(reference category; included clerical, technical, managerial, professional, sales, and services),

Agriculture/Manual = 2 (included skilled and unskilled manual work), Not Working = 3 (1

observation had a missing value). Household level wealth was measured using a wealth index that

was coded as an ordinal level variable with richest =1 (reference category), richer = 2, middle = 3,

poorer = 4, and poorest = 5 (we opted to use the wealth index that was provided by each respective

survey year of the CDHS as opposed to calculating a wealth index across the pooled data. This

decision was based on prior research that found wealth status designation was comparable across the

CDHS provided indices and an index based on data pooled across years) [20,21].

Pregnancy duration (trimester) was coded as an ordinal level variable with $1^{st} = 1$ (*reference*

category), $2^{nd} = 2$, and $3^{rd} = 3$. BMI was coded as an ordinal level variable with Overweight or Obese =

1 (reference category), Normal = 2, and Underweight = 3. Prior births was coded as an ordinal level

variable with Less than 3 Births = 1 (reference category), 3-5 Births = 2, and 6 or more Births = 3 (3

observations had a missing value). Tobacco use was coded as a dichotomous variable with No Tobacco

Use = 0 and Tobacco Use = 1 (possible types of tobacco use included cigarettes, cigars, and chewing

tobacco; 2 observations had a missing value).). Healthcare barriers was coded as a dichotomous

variable with No Barriers = 0 and 1 or more barriers = 1 (possible barriers reported included distance,

money, and waiting time; 1 observation had a missing value).

Residence was a dichotomous variable representing urban vs. rural place of residence with

Rural =1, Urban = 0. Phnom Penh was a dichotomous variable representing residence in the

provincial area surrounding Phnom Penh vs. other regions of residence with Other Regions =1,

Phnom Penh = 0. Cambodia's domains/provinces were regrouped for analytic purposes into a

categorical variable with 4 geographical regions that were coded as Plains = 1 (reference category),

Tonle Sap = 2, Costal/sea = 3, and Mountains = 4. Health facility visit and aborted pregnancy were

coded as dichotomous variables representing whether the respondent had visited a health facility in

the past year or had ever terminated a pregnancy; results are not presented due to lack of statistical

significance in unadjusted analyses.

Analytic strategy

Data cleaning and analyses were performed using STATA version 16 (Stata Corp 2019, College

Station, TX) [22]. Additional visualization was performed using R version 4.2.0 (code available upon

request); the complex survey design of the CHDS was accounted for using the "survey" package

[23,24]. Sample weights and complex survey design were accounted for in both descriptive and

logistic regression analyses. To convey temporal and geographic trends in anemia prevalence,

weighted estimates of overall and region-specific temporal trends in anemia prevalence were

visualized using the ggplot2 package in R [25]. In addition, a temporal series of maps illustrating

provincial variation in the prevalence of anemia were also prepared by our team using ArcGIS

software version 10.8 [26]. The underlying shapefiles of Cambodia provinces were obtained from the

publicly available Spatial Data Repository associated with the DHS website at

[https://spatialdata.dhsprogram.com/covariates/] [27]. Data quality checks and minor data cleaning

followed standard procedures in preparation for statistical analysis [28].

Cross tabulation chi-square tests were used to describe and provide preliminary assessment of

associations between independent variables of interest including maternal demographic, household

characteristics, geographical regions, and health-related) and anemia status. Unadjusted models and

several pathway specific models were examined prior to fitting the final model (results from the pathway

specific models are available upon request). Variables were selected for inclusion in the final adjusted

logistic regression if their unadjusted association with anemia had theoretical justification and/or a

significance level of p-value ≤ 0.15 [12]. Correlation among independent variables was assessed using

a combination of cross tabulation chi-square tests and Cramer's V scores; all results suggested minimal

to moderate correlation between independent variables (all Crammer's v scores were lower than .5,

most were lower than .3). Further consideration of whether to include a combination of variables was

based on broader evaluations of model fit, which we assessed using Pseudo-R^2, Deviance, AIC, and BIC

scores. The Pseudo-R^2 value was the largest and the Deviance, AIC, BIC scores were the smallest for the

fully adjusted model, indicating that this model had superior fit compared to other models.

Unadjusted logistic regression was used to analyze the magnitude effect of associations between

anemia and maternal bio-demographic and household characteristics, geographical regions, and

health-related factors. Results are reported as Odds Ratios (OR) with 95% confidence intervals (CI).

Adjusted logistics regression was then used to assess independent factors associated with anemia after

adjusting for other potential confounding factors in the model. Results from the final adjusted model

are reported as adjusted odds ratios (AOR) with 95% confidence intervals and corresponding p-

values. Results from the final adjusted logistic regression model were considered as statistically

significant based on a p-value less than 0.05 and 95% confidence intervals. Evaluations of effect

modification were not statistically significant. Observations with missing values were omitted from

the regression analyses.

This study was approved by the National Ethical Committee for Health Research (Ref: 225

NECHR) Cambodia and the Institutional Review Board (IRB) of ICF in Rockville, Maryland, USA.

The CDHS data are publicly accessible and were made available to us upon request to the DHS

Program website at (https://dhsprogram.com/data/available-datasets.cfm) [29]. Written consents

were obtained from all participants before the interview with participants in the CDHS.

Results

A little over 60% of pregnant women included in the analysis were aged 21-30 years, and

almost 98% were married (Table 1). Approximately 17% had no schooling, 52% had a primary

education, 29% had a secondary education, and only 2.6% had a higher education. Roughly 45%

were from poorest or poorer household wealth quintiles. Pregnancy duration was rather evenly

distributed with 28% of pregnant women in their 1^{st} trimester, 37% in their 2^{nd} trimester, and 36% in

their third trimester. Slightly over 50% of pregnant women had a normal BMI, with another 33%

were either overweight or obese. About 80% reported having fewer than 3 prior births. Only about

6% of pregnant women reported using tobacco. A majority (85%) of them resided in rural areas

(Table 1).

Table 1. Descriptive characteristics of anemia prevalence among study population

Variables	2005					2010					2014					2005-2014				
	Anemic		Not Anemic		P value	Anemic		Not Anemic		P value	Anemic		Not Anemic		P value	Anemic		Not Anemic		P value
	Freq.	%	Freq.	%		Freq.	%	Freq.	%		Freq.	%	Freq.	%		Freq.	%	Freq.	%	
Age (Years)																				
15-20	54	57.5	40	42.5	0.159	52	63.6	30	36.4	0.003	76	58.8	53	41.2	0.386	182	59.7	123	40.3	0.002
21-31	134	51.4	127	48.6		150	47.2	169	52.8		189	51.0	182	49.0		474	49.8	477	50.2	
32-49	84	63.9	47	36.1		44	67.6	21	32.4		62	54.1	53	45.9		190	61.1	121	38.9	
Marital Status																				
Married	269	56.2	210	43.8	0.270	237	52.4	215	47.6	0.215	318	52.8	284	47.2		824	64.3	709	35.7	0.232
Not Married	2	36.4	4	63.6		10	69.5	4	30.5		10	72.0	4	28.0		22	53.8	12	46.2	
Education																				
Higher	5	74.9	2	25.1	0.633	1	13.6	6	86.4	0.107	9	33.7	17	66.3	0.025	15	36.7	25	63.3	0.025
Secondary School	48	53.4	42	46.6		57	47.4	63	52.6		116	48.5	123	51.5		221	49.2	228	50.8	
Primary School	148	54.2	125	45.8		143	54.6	119	45.4		170	59.5	116	40.5		461	56.2	359	43.8	
No Education	71	61.0	45	39.0		46	59.3	32	40.7		33	50.8	32	49.2		150	58.0	109	42.0	
Occupation																				
Professional	35	43.9	44	56.1	0.045	43	48.6	46	51.4	0.504	46	39.9	70	60.1	0.021	124	43.7	160	56.3	0.010
Agricultural/Manual labor	235	58.0	170	42.0		157	52.5	142	47.5		185	59.1	128	40.9		577	56.7	440	43.3	
Not Working	2	100.0	0	0.0		46	59.3	32	40.7		96	51.7	90	48.3		145	54.4	122	45.6	
Wealth Index																				
Richest	27	40.2	40	59.8	0.002	25	34.8	47	65.2	0.012	51	40.9	73	59.1	0.001	103	39.0	160	61.0	<0.001
Richer	43	43.4	57	56.6		37	51.7	34	48.3		64	51.8	60	48.2		145	48.9	151	51.1	

	n	%	n	%	p	n	%	n	%	p	n	%	n	%	p	n	%	n	%	p
Middle	51	57.0	39	43.0		58	51.4	55	48.6		40	39.3	62	60.7		150	49.0	156	51.0	
Poorer	66	60.5	43	39.5		50	52.7	45	47.3		99	66.3	50	33.7		215	60.9	138	39.1	
Poorest	84	70.4	35	29.6		77	66.7	38	33.3		73	63.4	42	36.6		234	66.9	116	33.1	
Pregnancy Duration																				
1st Trimester	53	45.5	64	54.5	0.028	53	37.3	89	62.7	0.001	75	43.8	96	56.2	0.003	182	42.1	250	57.9	<0.001
2nd Trimester	112	65.0	60	35.0		108	65.1	58	34.9		150	63.7	86	36.3		371	64.5	204	35.5	
3rd Trimester	106	54.3	89	45.7		85	54.2	72	45.8		102	49.1	106	50.9		294	52.4	267	47.6	
BMI																				
Overweight or obese	67	45.1	71	54.9	0.042	72	43.7	66	56.3	0.524	27	47.8	29	52.2	0.081	72	45.6	86	54.4	0.005
Normal	182	61.4	114	38.6		152	55.1	124	44.9		184	58.3	132	41.7		518	58.4	370	41.6	
Underweight	23	48.3	28	51.7		23	52.0	29	48.0		117	47.8	127	52.2		255	49.1	265	50.9	
Prior Births																				
Less than 3 Births	190	54.9	156	45.1	0.154	196	50.7	191	49.3	0.207	280	51.4	264	48.6	0.065	666	52.1	612	47.9	0.017
3-5 Births	61	54.4	51	45.6		40	63.5	23	36.5		44	68.3	21	31.7		145	60.5	95	39.5	
6 and more Births	21	77.0	6	23.0		11	65.8	6	34.2		4	56.5	3	43.5		35	70.6	14	29.4	
Tobacco Use																				
No	250	54.6	208	45.4	0.015	234	52.6	211	47.4	0.564	320	52.7	288	47.3	0.021	805	53.2	707	46.8	0.027
Yes	22	78.6	6	21.4		13	60.4	9	39.6		7	100.0	0	0.0		42	74.1	15	25.9	
Healthcare Barriers																				
No Barriers	43	48.4	45	51.6	0.215	50	44.3	63.0	55.7	0.073	81	49.7	82	50.3	0.342	174	47.7	191	52.3	0.019
1 or more Barriers	226	57.3	169	42.7		197	55.7	156.0	44.3		246	54.5	205	45.5		669	55.8	530	44.2	
Place of Residence																				
Urban	23	42.7	32	57.3	0.029	28	39.3	43	60.7	0.025	50	46.1	58	53.9	0.095	101	43.2	133	56.8	<0.001
Rural	248	57.7	182	42.3		219	55.4	176	44.6		278	54.7	230	45.3		745	55.9	588	44.1	

The overall prevalence of anemia among pregnant women decreased slightly from 56% in 2005

to 53% in 2014 (Fig 2). This decline varied across the four geographical regions with the highest

prevalence of anemia in the coastal region in 2010 (66%). Plain region had the lowest prevalence of

anemia across all survey years with prevalence levels in 2005 lower than the prevalence levels of

other regions in 2014. Anemia was highest among pregnant women in Preah Vihear and Stung Treng

provinces (74.3%), in Kratie province (73%), and in Prey Veng (65.4%) for 2005, 2010 and 2014

respectively (Fig 3). Pregnant women living in Phnom Penh had the lowest prevalence of anemia in

2005 and 2010 (i.e., 33.3% in 2005, 30.0% in 2010. In 2014, the lowest prevalence of anemia among

pregnant women was observed in Takeo province (23.8%).

Fig 2. Overall and regional trends of anemia among pregnant women by survey year.

Fig 3. **Geographical distribution of anemia among pregnant women aged 15-49 years old.** Map created
using ArcGIS software version 10.8 [26]. Shapefiles for administrative boundaries in Cambodia are
publicly accessible through DHS website at [https://spatialdata.dhsprogram.com/covariates/] [27].
CDHS   data   are   publicly   accessible   through   the   DHS   website   at
(https://dhsprogram.com/data/available-datasets.cfm) [28].
215
Regarding socio-demographic factors associated with anemia in pregnancy in unadjusted

logistic regression analysis, pregnant women were more likely to have anemia if they were aged 15-

20 (OR= 1.5; 95% CI: 1.1-2.0) or aged 31-49 (OR= 1.6; 95% CI: 1.1-2.2) compared to pregnant

women aged 21-30 (Fig 4). Pregnant women with no formal education were more likely to have

anemia compared to women with a higher education (2.4; 95% CI: 1.1-5.1) as were women with only

a primary level education (OR = 2.2; 95% CI: 1.1-4.5). Women working in agricultural or manual

labor fields were more likely to have anemia compared to women working in professional fields (OR

= 1.7; 95% CI: 1.2-2.3). Juxtapose pregnant women from the richest wealth quintile, pregnant women

with lower wealth were more likely to have anemia; for example, women from the poorest quintile

were over 3 times as likely to have anemia (OR = 3.2; 95% CI: 2.1-4.7) and women from the poorer

quintile were over 2 times as likely to have anemia (OR = 2.4; 95% CI: 1.6-3.6).

Fig 4. Factors associated with anemia among pregnant women aged 15-49 years old in unadjusted

logistic regression analysis (n=1,567). Forest plots were generated using the *forest model* package in

R.

Health indicators associated with anemia in unadjusted logistic regression analysis included

pregnancy duration, BMI, number of prior births, tobacco use, and barriers to healthcare. Pregnant

women were more likely to have anemia if they were in their 2nd trimester (OR = 2.5; 95% CI: 1.8-

3.4) or their 3rd trimester (OR = 1.5; 95% CI: 1.1-2.1) compared to pregnant women in their 1st

trimester. A normal BMI was also associated with anemia during pregnancy compared to a BMI

indicating that a pregnant woman was overweight or obese (OR = 1.5; 95% CI: 1.1-1.9). Compared to

having had fewer than 3 births, pregnant women with 6 or more prior births were more likely to

have anemia (2.2; 95% CI: 1.2-4.2). Pregnant women using 1 type of tobacco were more likely to

have anemia compared to pregnant women not using tobacco (1.9; 95% CI: 1.1-3.2); using 2 or more

types of tobacco was likewise positively associated with having anemia (17.5; 95% CI: 3.7-83.3).

Finally, reporting 1 or more barriers to accessing healthcare was associated with anemia during

pregnancy (1.4; 95% CI: 1.1-1.8).

Geographic regions of residence were likewise associated with a woman having anemia

during pregnancy. Pregnant women from rural areas were more likely to have anemia compared to

pregnant women from urban areas (1.7; 95% CI: 1.3-2.2). Likewise, pregnant women living outside of

the area surrounding Phnom Penh were more likely to have anemia compared to pregnant women

from the Phnom Penh area (1.6; 95% CI: 1.1-2.3). Coastal regions were positively associated with

anemia compared to plains regions (1.9; 95% CI: 1.2-2.9) as were mountain regions (1.43; 95% CI:

1.04-1.98).

In the final adjusted logistic model (**Fig 5**), factors independently associated with anemia

included age, wealth index, pregnancy duration, BMI, and geographical regions. Pregnant women

were more likely to have anemia if they were aged 31-49 (AOR = 1.6; 95% CI: 1.0-2.3) compared to

pregnant women aged 21-30. Compared to pregnant women from the richest wealth quintile,

pregnant women from the poorest quintile were almost 3 times as likely to have anemia (AOR = 2.8;

95% CI: 1.76-4.9) as were women from the poorer quintile (AOR = 2.2; 95% CI: 1.3-3.9). Pregnant

women were more likely anemic if they were in their 2nd trimester (AOR = 2.6; 95% CI: 1.9-3.6) or

3rd trimester (AOR = 1.6 95% CI: 1.2-2.3) compared to their 1st trimester. Pregnant women with a

normal BMI were more likely of developing anemia (AOR = 1.4; 95% CI: 1.0-1.8) compared to

pregnant women who were overweight or obese. Pregnant women living in coastal regions had

higher odds having anemia (AOR = 1.9; 95% CI: 1.2-3.0) compared to those living in plain region.

Statistically significant associations were further visualized as predicted probabilities (see Figs 6–9 in

the supplement).

Fig 5. Factors independently associated with anemia among pregnant women aged 15-49 years old in

adjusted logistic regression analysis (n=1,567). Forest plots were generated using the forest model

package in R.

Discussion

Prevalence of anemia among pregnant women in Cambodia decreased slightly from 56% in

2005 to 53% in 2014. This prevalence level was relatively higher than many Southeast Asian

countries during the same time period [30]; for example, in Thailand the prevalence of anemia among

pregnant women ranged from 31% in 2005 to 31.9% in 2016 and prevalence of anemia among

pregnant women in Vietnam ranged from 33.8% in 2005 to 28.2% in 2015. The elevated prevalence

of anemia among pregnant women in Cambodia may be explained by factors related to dietary habits,

nutrition deficiencies in micronutrients such as iron or vitamin A, genetic hemoglobin disorders such

as the high prevalence of thalassemia in Cambodia (e.g., prevalence rang 50 to 60%) [27], lack of

medical treatments, inflammation [31], and malaria infections, and helminths; these potential

pathways require further evaluation [16]. Interventions to help mitigate anemia in Cambodia include

the promotion of a diversified diet; recommendations that all pregnant women attend at least 4 ANC

visit, receive a standard dose of 30–60 mg iron and 400 µg folic acid beginning as soon as possible

during gestation, and iron-containing supplements no later than the first trimester of pregnancy;

prevention and treatment of malaria; use of insecticide-treated bed nets; helminth prevention and

control; delayed cord clamping; and increased birth spacing [32]. However, some have noted that

factors addressed by these interventions cannot fully account for the elevated prevalence of anemia

[31] and data from the 2000-2014 Cambodia DHS indicated that the percentage of pregnant women

attended four or more antenatal care visits has increased dramatically, from 9% in 2000 to 76% in

2014. However proportion was lowest reported by women resided in rural, plateau, and coastal

region, lowest families income, limited education, and first pregnant [9], over 76% took standard iron

and folic acid supplements during pregnancy and after delivery through the national health system,

while only 65% of pregnant women took iron supplements living in lowest wealth households, which

lowest in Preah Vihea/Stung Treng (63.3%) and Mondol Kiri/Ratanakiri (56%) [5]. In the last 10

years, Cambodia has also piloted weekly iron and folic acid supplementation for women of

reproductive age. However, despite evidence for an impact on iron status during pregnancy, factors

such as cost and need for behavioral change have hampered implementing this policy on a large scale

[5]. In contrast, food fortification efforts, such as iron-fortified fish sauce, are currently underway

[33] and 72% of women received medications to combat parasites [5]; hence the interest in better

understanding associations between anemia and demographic, economic, and social factors as well as

health-seeking behaviors across different parts of the country [34].

Our results suggest independent associations between anemia in pregnant women and age,

wealth, pregnancy duration, BMI, and region of residence. Older aged pregnant women were more

likely to have anemia compared to young pregnant women, which is consistent with previous studies

[11-13] and may reflect biological and physical disadvantages with increased maternal age; for

example, developing gestational diabetes, high blood pressure during pregnancy, a mother having

multiple pregnancies, and enduring the collective impact of exhausting labor-related complications

[35-37]. Consistent with studies from Rwanda [38], Ethiopia [39], and Uganda [40], pregnant women

were also more likely to have anemia if they were from the poorest wealth quintiles with the patterns

of association between anemia and wealth exhibiting a dose response to increased poverty. Lower

socioeconomic status is often associated with negative consequences regarding consumption of a

healthy diet, greater chance of infection diseases, and less access to healthcare services [42]. In

contrast, individuals from higher socioeconomic groups have a comparative advantage that enables

them to purchase sufficient food with greater variety and quality, thus protecting against anemia [43].

In addition, pregnant women living in the coastal regions of residence were more likely have anemia

compared to women living in plains regions. These differences in the prevalence of anemia might be

due to variations in socioeconomic status, attention given to focused antenatal care and

supplementation of iron sulfate throughout the pregnancy, dietary patterns, sample size, and

geographical and lifestyle variations [44]. Finally, pregnant women with a normal BMI were more

likely of have anemia compared to pregnant women who were overweight or obese, which is

surprising given prior studies suggesting that higher BMI is negatively associated with iron status

among pregnant women [45]. This association may be related to nutrition; e.g., it may be that

overweight or obese women have a higher iron intake compared to normal weigh women [33].

However, the level of iron content of foods consumed by pregnant women in Cambodia remains

unclear, hence future research should employ a nutrition survey for pregnant women and evaluate

these potential associations directly.

This study was conducted using pooled Demographic and Health surveys for 2005, 2010, and

2014, which enabled us to perform a descriptive analysis of temporal and geographical trends in

anemia among pregnant women and also analyze potential risk factors using nationally representative

data and generalize our findings to pregnant women throughout Cambodia. However, using cross-

sectional data limits our ability to assess causal relationship among the associated factors that we

identified. Using secondary data that was gathered to collect health and demographic data across a

broad variety of variables related to child and maternal health enabled us to evaluate some of the

factors likely associated with anemia among pregnant women, but other factors such as diet,

nutritional supplements (Vitamin B12, folate); malaria infection, non-communicable diseases such as

hypertension and diabetes, and awareness about anemia were excluded from the analysis due to data

limitations. Finally, the most recent round of CDHS data was gathered in 2014, which may no longer

reflect present prevalence of anemia among pregnant women in Cambodia. Future research should

further evaluate these relationships and the general prevalence of anemia among pregnant

Cambodian women once additional data becomes available (data gathering for the next round of

Cambodia DHS data is underway, but delayed due to COVID-19). With additional data, future

research should consider a formal spatial analysis for enhancing the scientific rigor of geospatial

trends [46].

Our findings suggest that the high prevalence of anemia among pregnant women in Cambodia

declined only slightly over the 10-year period between 2005 and 2014. This is concerning given that

anemia during pregnancy is a public health problem that should be actively addressed to mitigate the

proximal and distal health risks associated with the condition. We identified age, socioeconomic

status, higher BMI, and regions of residence as potential risk factors associated with anemia during

pregnancy. Public health practitioners and policy makers should consider demographic groups

represented by these characteristics for better targeting interventions to address anemia among

pregnant women in Cambodia.

Acknowledgments

The authors would like to thank the National Institute of Public Health for allowing us to take part in Sabbatical for Writing Program 2022. We also sincerely thank the DHS-ICF, who approved the data used for this paper.

Authors' contributions

Samnang Um contributed to the conceptualization, methodology, data analysis, writing the original draft, review, and editing; **Heng Sopheab** contributed to writing, review, and editing the manuscript; **An Yom** contributed to writing, review, and editing the manuscript; and **Jonathan A. Muir** contributed to methodology, data analysis, writing, review, and editing the manuscript. All authors read and approved the final manuscript.

Data availability

Our study used 2005, 2010, and 2014 Cambodia Demographic and Health Survey (CDHS) datasets. The DHS data are publicly available from the website: https://dhsprogram.com.

Funding

The authors received no funding for this study.

Competing interests

The authors have declared that no competing interests exist.

Abbreviations

ANC, Antenatal Care; AOR, Adjusted Odds Ratio; CDHS, Cambodian Demographic Health Survey; EA, Enumeration areas; Hb, Hemoglobin; WHO, World Health Organization

References

1.   Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, et al. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis, 2. The American journal of clinical nutrition. 2016;103(2):495-504.

2.   WHO. The global prevalence of anaemia in 2011. Geneva: World Health Organization. 2015.

3.   Stevens GA, Paciorek CJ, Flores-Urrutia MC, Borghi E, Namaste S, Wirth JP, et al. National, regional, and global estimates of anaemia by severity in women and children for 2000–19: a pooled analysis of population-representative data. The Lancet Global Health. 2022;10(5):e627-e39.

4.   WHO U. Global Nutrition Targets 2025: Breastfeeding policy brief (WHO/NMH/NHD14. 7). Geneva: World Health Organization. 2014.

5.   NIS. National Institute of Statistics. Directorate General for HC. International ICF . Cambodia Demographic and Health Survey 2014. Phnom Penh, Cambodia: National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF International; 2015. 2014.

6.   McLean E, Cogswell M, Egli I, Wojdyla D, De Benoist B. Worldwide prevalence of anaemia, WHO vitamin and mineral nutrition information system, 1993–2005. Public health nutrition. 2009;12(4):444-54.

7.   Elzahrani S. Prevalence of iron deficiency anemia among pregnant women attending antenatal clinics at Al-Hada Hospital. Canadian Journal on Medicine. 2012;3(1):10-4.

8.   Daru J, Zamora J, Fernández-Félix BM, Vogel J, Oladapo OT, Morisaki N, et al. Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: a multilevel analysis. The Lancet Global Health. 2018;6(5):e548-e54.

9.   Mallick L, Courtney Allen, and Rathavuth Hong. . Trends in Maternal and Child Health inCambodia, 2000-2014. Further Analysis of the 2000, 2005, 2010, and 2014 Cambodia Demographic andHealth Surveys. DHS Further Analysis Reports No. 106. Rockville, Maryland, USA: ICF. Available from: https://www.researchgate.net/publication/323150392_Trends_in_Maternal_and_Child_Health_in_Cambodia_2000-2014 [accessed Mar 30 2023]. 2016.

10.   Balarajan Y, Ramakrishnan U, Özaltin E, Shankar AH, Subramanian S. Anaemia in low-income and middle-income countries. The lancet. 2011;378(9809):2123-35.

11.   Woldegebriel AG, Gebregziabiher Gebrehiwot G, Aregay Desta A, Fenta Ajemu K, Berhe AA, Woldearegay TW, et al. Determinants of anemia in pregnancy: findings from the Ethiopian health and demographic survey. Anemia. 2020;2020.

12.   Lin L, Wei Y, Zhu W, Wang C, Su R, Feng H, et al. Prevalence, risk factors and associated adverse pregnancy outcomes of anaemia in Chinese pregnant women: a multicentre retrospective study. BMC pregnancy and childbirth. 2018;18(1):1-8.

13.   Wu Y, Ye H, Liu J, Ma Q, Yuan Y, Pang Q, et al. Prevalence of anemia and sociodemographic characteristics among pregnant and non-pregnant women in southwest China: a longitudinal observational study. BMC Pregnancy and ChildBirth. 2020;20(1):1-10.

14.   Sunuwar DR, Singh DR, Chaudhary NK, Pradhan PMS, Rai P, Tiwari K. Prevalence and factors associated with anemia among women of reproductive age in seven South and Southeast Asian countries: Evidence from nationally representative surveys. PloS one. 2020;15(8):e0236449.

15.   Harding KL, Aguayo VM, Namirembe G, Webb P. Determinants of anemia among women and children in Nepal and Pakistan: An analysis of recent national survey data. Maternal & child nutrition. 2018;14:e12478.

16.   Charles CV, Summerlee AJ, Dewey CE. Anemia in Cambodia: Prevalence, etiology and research needs. Asia Pacific journal of clinical nutrition. 2012;21(2):171-81.

17.   Charles CV, Dewey CE, Daniell WE, Summerlee AJ. Iron-deficiency anaemia in rural Cambodia: community trial of a novel iron supplementation technique. European journal of public health. 2011;21(1):43-8.

18.   Tibambuya BA, Ganle JK, Ibrahim M. Anaemia at antenatal care initiation and associated factors among pregnant women in West Gonja District, Ghana: a cross-sectional study. The Pan African medical

journal. 2019;33:325. Epub 2019/11/07. doi: 10.11604/pamj.2019.33.325.17924. PubMed PMID: 31692871; PubMed Central PMCID: PMCPmc6815505.

19.   Croft T, Marshall A, Allen C. Guide to DHS Statistics. Rockville, Maryland, USA: ICF; 2018. 2018.

20.   Muir JA. Societal Shocks as Social Determinants of Health: The Ohio State University; 2021.

21.   Muir JA. Another mHealth? Examining Motorcycles as a Distance Demolishing Determinant of Health Care Access in South and Southeast Asia. Journal of Transport & Health. 2018;11:153-66.

22.   Cunningham SA, Muir JA. Data Cleaning. In: Nichols AN, Edlund J, editors. The Cambridge Handbook of Research Methods and Statistics for the Social and Behavioral Sciences: Volume 1: Building a Program of Research. 1: Cambridge University Press; 2023.

23.   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2022.

24.   Lumley T. Package 'survey'. Available at the following link: https://cran r-project org. 2020.

25.   Wickham H. ggplot2: elegant graphics for data analysis: springer; 2016.

26.   Desktop A. Release 10.8. 1.(2020). ESRI (Environmental Systems Research Institute), Redlands, CA.

27.   Karakochuk CD, Whitfield KC, Barr SI, Lamers Y, Devlin AM, Vercauteren SM, et al. Genetic hemoglobin disorders rather than iron deficiency are a major predictor of hemoglobin concentration in women of reproductive age in rural Prey Veng, Cambodia. The Journal of nutrition. 2015;145(1):134-42.

28.   Cunningham, S., & Muir, J. (2023). Data Cleaning. In A. Nichols & J. Edlund (Eds.), The Cambridge Handbook of Research Methods and Statistics for the Social and Behavioral Sciences: Volume 1: Building a Program of Research (Cambridge Handbooks in Psychology, pp. 443-467). Cambridge: Cambridge University Press. doi:10.1017/9781009010054.022.

29.   Asamrew N, Endris AA, Tadesse M. Level of patient satisfaction with inpatient services and its determinants: a study of a specialized hospital in Ethiopia. Journal of environmental and public health. 2020;2020.

30.   WHO. World Health Organization-Global Health Observatory Data Repository, Prevalence of anaemia in non-pregnant women (aged 15-49) (%). Available at [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/anaemia-in-non-pregnant-women-prevalence-(-)]. Accessed March 2023.

31.   Wieringa FT, Dahl M, Chamnan C, Poirot E, Kuong K, Sophonneary P, et al. The High Prevalence of Anemia in Cambodian Children and Women Cannot Be Satisfactorily Explained by Nutritional Deficiencies or Hemoglobin Disorders. Nutrients. 2016;8(6):348. PubMed PMID: doi:10.3390/nu8060348.

32.   Barrett B. Nutrition Profile of Cambodia: A Review of Current Conditions and Strategies to Prevent Macronutrient and Micronutrient Deficiencies. 2020.

33.   Theary C, Panagides D, Laillou A, Vonthanak S, Kanarath C, Chhorvann C, et al. Fish sauce, soy sauce, and vegetable oil fortification in Cambodia: where do we stand to date? Food and nutrition bulletin. 2013;34(2_suppl1):S62-S71.

34.   Charles CV, Dewey CE, Hall A, Hak C, Channary S, Summerlee AJ. Anemia in Cambodia: A cross-sectional study of anemia, socioeconomic status and other associated risk factors in rural women. Asia Pacific journal of clinical nutrition. 2015;24(2):253-9.

35.   Mayo Clinic Press. Pregnancy after 35: Healthy moms, healthy babies. Available at: https://www.mayoclinic.org/healthy-lifestyle/getting-pregnant/in-depth/pregnancy/art-20045756. Accessed on March 31, 2022.

36.   Londero AP, Rossetti E, Pittini C, Cagnacci A, Driul L. Maternal age and the risk of adverse pregnancy outcomes: a retrospective cohort study. BMC pregnancy and childbirth. 2019;19(1):1-10.

37.   Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, Bommarito K, Madden T, Olsen MA, et al. Maternal age and risk of labor and delivery complications. Maternal and child health journal. 2015;19(6):1202-11.

38.   Habyarimana F, Zewotir T, Ramroop S. Spatial distribution and analysis of risk factors associated with Anemia among women of reproductive age: case of 2014 Rwanda demographic and health survey data. The Open Public Health Journal. 2018;11(1).

39.   Kibret KT, Chojenta C, D'Arcy E, Loxton D. Spatial distribution and determinant factors of anaemia among women of reproductive age in Ethiopia: a multilevel and spatial analysis. BMJ open. 2019;9(4):e027276.

40.   Nankinga O, Aguta D. Determinants of Anemia among women in Uganda: further analysis of the Uganda demographic and health surveys. BMC Public Health. 2019;19(1):1-9.

41.   Soofi S, Khan GN, Sadiq K, Ariff S, Habib A, Kureishy S, et al. Prevalence and possible factors associated with anaemia, and vitamin B 12 and folate deficiencies in women of reproductive age in Pakistan: analysis of national-level secondary survey data. BMJ open. 2017;7(12):e018007.

42.   Arpey NC, Gaglioti AH, Rosenbaum ME. How socioeconomic status affects patient perceptions of health care: a qualitative study. Journal of primary care & community health. 2017;8(3):169-75.

43.   Teshale AB, Tesema GA, Worku MG, Yeshaw Y, Tessema ZT. Anemia and its associated factors among women of reproductive age in eastern Africa: A multilevel mixed-effects generalized linear model. Plos one. 2020;15(9):e0238957.

44.   Balarajan YS, Fawzi WW, Subramanian S. Changing patterns of social inequalities in anaemia among women in India: cross-sectional study using nationally representative data. BMJ open. 2013;3(3):e002233.

45.   Zimmermann MB, Hurrell RF. Nutritional iron deficiency. The lancet. 2007;370(9586):511-20.

46.   Sunuwar DR, Singh DR, Adhikari B, Shrestha S, Pradhan PMS. Factors affecting anaemia among women of reproductive age in Nepal: a multilevel and spatial analysis. BMJ Open. 2021;11(3):e041982. Epub 2021/03/31. doi: 10.1136/bmjopen-2020-041982. PubMed PMID: 33782019; PubMed Central PMCID: PMCPmc8009228.

Supplemental table

Table 2. Prevalence of Anemia in the 19 domains in CDHS 2005, CDHS 2010 and 2014

Province	CDHS 2005 (n=486)	CDHS 2010 (n=467)	CDHS 2014 (n=615)
	% of anemia	% of anemia	% of anemia
Banteay Meanchey	62.0	37.0	30.0
Kampong Cham/Tbong Khmum	56.0	50.0	44.0
Kampong Chhnang	53.0	67.0	63.0
Kampong Speu	61.0	48.0	56.0
Kampong Thom	62.0	57.0	62.0
Kandal	53.0	49.0	50.0
Kratie	53.0	78.0	66.0
Phnom Penh	34.0	31.0	46.0
Prey Veng	50.0	60.0	69.0
Pursat	56.0	41.0	52.0
Siem Reap	64.0	56.0	53.0
Svay Rieng	45.0	61.0	64.0
Takeo	64.0	40.0	26.0
Otdar Meanchey	78.0	30.0	59.0
Battambang/Pailin	55.0	58.0	50.0
Kampot/Kep	62.0	76.0	61.0
Preah Sihanouk/Koh Kong	76.0	48.0	65.0

Preah Vihear/Stung Treng	73.0	61.0	59.0
Mondul Kiri/Ratanak Kiri	56.0	49.0	42.0

Supplemental figures

Fig 6. Predicted probability of anemia among pregnant women aged 15-49 years old (n=1,567). Visualized using "ggplot2" package in R.

Fig 7. Predicted probability of anemia among pregnant women aged 15-49 years old (n=1,567). Visualized using "ggplot2" package in R.

Fig 8. Predicted probability of anemia among pregnant women aged 15-49 years old (n=1,567). Visualized using "ggplot2" package in R.

Fig 9. Predicted probability of anemia among pregnant women aged 15-49 years old (n=1,567). Visualized using "ggplot2" package in R.

Total number of women aged 15-49 years

    CDHS 2014: 17,578

    CDHS 2010: 18,754

    CDHS 2005: 16,823

Excluded those who were not pregnant and without hemoglobin collection

    CDHS 2014: 16,323

    CDHS 2010: 18,284

    CDHS 2005: 16,919

Final sample size included in the data analyzed for the study

    CDHS 2014: 659 (615 weighted cases)

    CDHS 2010: 470 (467 weighted cases)

    CDHS 2005: 500 (486 weighted cases)

Variable	Odds Ratio		p
Age			
21–30		Reference	
15–20		1.49 (1.11, 2.01)	0.009
31–49		1.58 (1.14, 2.19)	0.006
Marital Status			
Married		Reference	
Not Married		1.55 (0.77, 3.13)	0.224
Education			
Higher		Reference	
Secondary		1.67 (0.80, 3.49)	0.172
Primary		2.21 (1.08, 4.53)	0.030
No Education		2.38 (1.11, 5.11)	0.026
Occupation			
Professional		Reference	
Agricultural/Manual		1.69 (1.26, 2.28)	<0.001
Not Working		1.53 (1.00, 2.36)	0.052
Wealth Index			
Richest		Reference	
Richer		1.50 (1.02, 2.19)	0.037
Middle		1.50 (1.03, 2.18)	0.032
Poorer		2.43 (1.63, 3.63)	<0.001
Poorest		3.15 (2.11, 4.71)	<0.001
Pregnancy Duration			
1st Trimester		Reference	
2nd Trimester		2.50 (1.85, 3.38)	<0.001
3rd Trimester		1.51 (1.09, 2.09)	0.013
BMI			
Overweight or Obese		Reference	
Normal		1.44 (1.11, 1.88)	0.007
Underweight		0.86 (0.54, 1.38)	0.536
Prior Births			
Less than 3 Births		Reference	
3–5 Births		1.41 (0.98, 2.02)	0.066
6 or more Births		2.21 (1.15, 4.24)	0.017
Tobacco Use			
No Tobacco Use		Reference	
Tobacco Use		2.05 (1.21, 3.45)	0.007
Healthcare Barriers			
No Barriers		Reference	
1 or More Barriers		1.41 (1.08, 1.86)	0.013
Residence			
Urban		Reference	
Rural		1.67 (1.26, 2.20)	<0.001
Phnom Penh			
Phnom Penh		Reference	
Other Regions		1.61 (1.10, 2.35)	0.014
Regions			
Plains		Reference	
Tonle Sap		1.25 (0.93, 1.68)	0.132
Coastal		1.86 (1.20, 2.89)	0.006
Mountains		1.43 (1.04, 1.98)	0.027
Year			
2014		Reference	
2010		0.99 (0.73, 1.34)	0.939
2005		1.12 (0.83, 1.50)	0.459

Variable	Odds Ratio		p
Age			
21–30		Reference	
15–20		1.35 (0.97, 1.88)	0.071
31–49		1.55 (1.03, 2.33)	0.036
Education			
Higher		Reference	
Secondary		1.08 (0.51, 2.28)	0.835
Primary		1.10 (0.52, 2.35)	0.799
No Education		0.79 (0.34, 1.84)	0.580
Occupation			
Professional		Reference	
Agricultural/Manual		1.20 (0.84, 1.71)	0.320
Not Working		1.42 (0.89, 2.26)	0.140
Wealth Index			
Richest		Reference	
Richer		1.32 (0.82, 2.13)	0.252
Middle		1.39 (0.86, 2.26)	0.183
Poorer		2.21 (1.27, 3.86)	0.005
Poorest		2.81 (1.62, 4.86)	<0.001
Pregnancy Duration			
1st Trimester		Reference	
2nd Trimester		2.60 (1.88, 3.58)	<0.001
3rd Trimester		1.62 (1.15, 2.27)	0.006
BMI			
Overweight or Obese		Reference	
Normal		1.36 (1.00, 1.84)	0.049
Underweight		1.05 (0.64, 1.72)	0.854
Prior Births			
Less than 3 Births		Reference	
3–5 Births		1.10 (0.71, 1.68)	0.676
6 or more Births		1.32 (0.57, 3.06)	0.521
Tobacco Use			
No Tobacco Use		Reference	
Tobacco Use		1.26 (0.69, 2.28)	0.455
Healthcare Barriers			
No Barriers		Reference	
1 or More Barriers		1.09 (0.80, 1.48)	0.603
Residence			
Urban		Reference	
Rural		0.99 (0.67, 1.47)	0.977
Regions			
Plains		Reference	
Tonle Sap		1.07 (0.79, 1.45)	0.645
Coastal		1.87 (1.17, 2.97)	0.009
Mountains		1.14 (0.79, 1.64)	0.484
Year			
2014		Reference	
2010		1.02 (0.74, 1.41)	0.909
2005		1.07 (0.76, 1.51)	0.683
	0.5 1 2		