

1 **Title: A cross-sectional study of low birth satisfaction among Iranian**
2 **postpartum women during COVID-19 epidemics' fifth wave**

3

4 **Short title: birth satisfaction during COVID-19 pandemic**

5 Forough Mortazavi ^{1*}, Maryam Mehrabadi²

6 ¹Associate Professor, Non-Communicable Diseases Research Center, Sabzevar University of
7 Medical Sciences, Sabzevar, Iran Email: frmortazavi@yahoo.com

8 <https://orcid.org/0000-0001-8190-0569>

9 ²MSc of Midwifery, Health Chancellery, Sabzevar University of Medical Sciences, Sabzevar, Iran

10 Corresponding Information: Forough Mortazavi, Non-Communicable Diseases Research Center,
11 Sabzevar University of Medical Sciences, Pardis Building, Towhidshahr Blvd, Sabzevar, Iran

12 Email: frmortazavi@yahoo.com Tel: +98(051)44018306

13

14 **Abstract**

15 **Background:** Birth dissatisfaction may increase the risk for postpartum depression and requests
16 for an elective cesarean for the next birth. The outbreak of COVID-19 pandemic has had a
17 considerable impact on the healthcare systems and their users in many aspects. We investigated
18 predictors of birth satisfaction in a sample of Iranian postpartum women during the COVID-19
19 epidemics' fifth wave.

20 **Methods:** This cross-sectional study was conducted on 601 postpartum women admitted to
21 postpartum wards of Mobini maternity hospital using a convenience sampling method between 2
22 Aug and 18 September 2021. We collected data on socio-demographic, obstetric, labor and birth,
23 and psychological variables. We used the general linear model and multiple linear regression
24 analyses to determine predictors of birth satisfaction.

25 **Results:** The mean birth satisfaction score was 28.6 ± 7.3 . The percentages of mothers who gave
26 birth by elective and emergency cesarean were 19.5% and 10.8%, respectively. Overall predictors
27 of birth satisfaction were emergency cesarean [-7.463(-9.310, -5.616), instrumental birth [-3.571(-
28 6.907, -0.235)], episiotomy [-2.227 (-3.591, -0.862)], Entonox analgesia [-1.548(-2.726, -0.371)],
29 Well-being score < 50 [-1.812(-3.146, -0.478)], fear of COVID -19 [-1.216(-2.288,, -0.144)], low
30 satisfaction with pregnancy -2.539(-3.952, -1.127) and low satisfaction with spouse's support [-
31 2.419(-4.598, -0.240)].

32 **Conclusions:** During the pandemic, fear of COVID -19, low level of well-being, low satisfaction
33 with pregnancy and low satisfaction with spouse's support as well as women's experience of
34 emergency cesarean, instrumental birth, episiotomy, and Entonox analgesia, are exerting negative

35 influences on birth satisfaction. To improve birth satisfaction and thus maternal mental health
36 interventions to lower fear of contracting COVID -19 and reduce rates of episiotomy, emergency
37 cesarean, and instrumental birth are recommended.

38 **Keywords:** patient satisfaction, postpartum period, natural childbirth, cesarean section, cross-
39 sectional study, Iran, COVID-19

40

41

42 **Introduction**

43 Childbirth is a turning point in a woman's life that can be either a wonderful experience or a
44 traumatic one. Although the primary goal of childbirth is the birth of a healthy infant from a healthy
45 mother, women would generally like it to be a positive and satisfying experience (1). Birth
46 satisfaction refers to the mother's overall assessment of childbirth and the extent to which she
47 perceived it as a positive and satisfying experience (2). Studies in developed countries report that
48 7–10% of women have had a negative birth experience (3).

49 Several studies have found positive association between birth dissatisfaction and a number of
50 outcomes including postpartum depression (2-4), difficulties in mother-baby bonding, poor
51 maternal care of the baby and abstaining from exclusive breastfeeding (3). Findings of a study in
52 New York City hospitals indicate that women with lower birth satisfaction were more likely to
53 experience higher levels of anxiety, stress, and depressive symptoms during the postpartum period
54 (4). Birth dissatisfaction may also increase requests for an elective cesarean for the next birth (5)
55 and the likelihood of mothers delaying their next pregnancy (6). Dissatisfaction with vaginal birth
56 among women in Hong Kong after their first childbirth resulted in 23.8% of them changing their
57 preferred mode of birth from vaginal to elective cesarean (5).

58 Several factors have been found to contribute to dissatisfaction with childbirth including
59 emergency cesarean (5, 7, 8), elective cesarean, higher family income, use of epidural analgesia
60 (5), induction, unwanted pregnancy (7), primiparity, low level of well-being, low satisfaction with
61 pregnancy, severe fear of childbirth, long time between hospital admission and giving birth (8),
62 and lack of support from partner (7, 8). Also, childbirth satisfaction has been found to be associated
63 with planned childbirth (9), mothers' acquaintance with labor, moderate labor pain (10), women's

64 participation in deciding matters that affect them (11), and empathy and respectful behavior of
65 caregivers (11, 12).

66 The COVID-19 pandemic has led to increasing rates of anxiety, depression, and stress among
67 women during pregnancy and childbirth (13-15). Stress was related to constrained antenatal care
68 during the pandemic, worries about being infected with the virus, the possibility of negative effects
69 of the infection on the baby, and worry about the health of loved ones (16). Fear of COVID-19 has
70 been found to be a common problem during the pandemic with a significant impact on women's
71 well-being (13). It had a statistically significant relationship with depression, suicidal tendencies,
72 and poor psychological quality of life in pregnant women (17). Studies have also found that, just
73 as in the pre-pandemic period, dissatisfaction with childbirth during COVID-19 pandemic is
74 associated with postpartum depression (15, 18). The pandemic has also impacted the organization
75 and the practices of health care provision in the antenatal, intrapartum, and postnatal periods. It
76 has also impacted the attitudes and the responses of healthcare users. Factors that might have been
77 impacted include pregnancy intention, care providers' behavior towards women, more restrictive
78 hospital rules, the likelihood of experiencing a hassle-free pregnancy, and request for elective
79 cesarean. It may also have increased households' economic concerns. Therefore, it would not be
80 unreasonable to expect lower levels of birth satisfaction in women giving birth during COVID-19
81 pandemic (15).

82 Like many other countries, Iran has experienced high rates of cesarean (19) and maternal requests for
83 cesarean (20) in recent decades. At the same time, the decreasing total fertility rate (TFR) has been
84 increasingly viewed as a threat to the future development of the country and thus a cause for concern (21).
85 Because birth dissatisfaction may increase requests for an elective cesarean for the next birth and the
86 likelihood of mothers delaying their next pregnancy, there is concerns that the COVID-19 epidemic with

87 its negative impact on birth satisfaction may undermine the Health Transformation Plan (HTP), an initiative
88 launched by the Iranian government for the reform of the healthcare system in 2014 (22). HTP includes a
89 number of guidelines and measures with regard to maternity care aiming to promote maternal experience
90 of childbirth. Thus, we hypothesize that maternal birth experience and childbirth satisfaction might have
91 been affected by COVID-19 pandemic. In the present study, we investigated the extent to which fear of
92 COVID -19, maternal well-being status, and labor and birth factors predicted lower birth satisfaction during
93 the COVID-19 epidemics' fifth wave in Sabzevar, Northeast Iran.

94 **Materials and Methods**

95 **Design, participants, and data collection**

96 This cross-sectional study was conducted from Aug 2 to September 18, 2021. Women were
97 recruited in the postpartum and surgery wards of Mobini Hospital affiliated with Sabzevar
98 University of Medical Sciences. As a matter of hospital policy, they were usually hospitalized for
99 the first 24 hours and 48 hours after vaginal delivery and cesarean, respectively. Women who had
100 vaginal deliveries completed the 45-item study questionnaire on the morning of the day after
101 childbirth when they felt they were ready for the task. In the cesarean group, they filled out the
102 scales on the morning of the second day after birth. Sampling was done using the convenience
103 method. Women who gave birth to a healthy, live infant were included in the study. We excluded
104 women with mental illness under treatment, women with severe postpartum complications, women
105 with infants admitted to the intensive care unit, those with preterm birth, and COVID-positive
106 women. Women admitted for birth in the hospital who showed mild signs of COVID were tested
107 and those with positive results were isolated from other women in labor and postpartum. Women
108 with severe disease were transferred to another facility dedicated to COVID-19 patients. The
109 average annual birth rate in the hospital which was 5898 before the COVID -19 outbreak,

110 decreased to 5245 in the first year after the outbreak. There were 4020 births in the first 9 months
111 of the second year of the pandemic. A research colleague, who was a midwifery graduate,
112 identified eligible mothers and, if they agreed to participate in the study, presented them with
113 written consent forms and anonymous questionnaires. We trained the midwife on how to present
114 the questionnaire and collect the data. Demographic, social, and obstetric information of the
115 mothers were extracted from their medical records and recorded by the colleague.

116 **Ethical consideration**

117 The Ethics Committee of Sabzevar University of Medical Sciences has reviewed and approved
118 this study (approval number: IR.MEDSAB.REC.1399.103). All procedures were performed in
119 accordance with the guidelines of Sabzevar University of Medical Sciences, which is in
120 accordance with the Declaration of Helsinki. Women who consented to participate in the study
121 signed an informed consent form and were assured about the confidentiality of their information.
122 After recording the socio-demographic information of the women in the questionnaire, the midwife
123 asked women to fill out the scales. The questionnaires and scales were anonymous.

124 **Instruments**

125 Interview form

126 A three-part questionnaire was completed by the midwife. The first part contained questions on
127 socio-demographic characteristics (including age, education, residency, and job). The second part
128 consisted of obstetrical information (such as parity, attending prenatal classes, history of chronic
129 disease, the desirability of pregnancy, poor obstetric history, infant gender, history of abortion,
130 and complicated pregnancy). The third part consisted of labor and birth information (such as mode
131 of birth, induced or spontaneous labor pain, pain relief method during labor and birth, having a

132 private midwife at birth, gestational age, birth weight, admission to birth duration, and
133 episiotomy/tear repair).

134 The level of satisfaction with pregnancy was assessed based on the extent of health problems
135 experienced during pregnancy using a five-point Likert scale from 1 to 5 (1 = dissatisfied to 5 =
136 very satisfied). Women's satisfaction with husband's emotional/financial support and
137 marital/sexual satisfaction were evaluated with the same scale. A question on receiving fundal
138 pressure was also included. Women rated their household income as 1= insufficient or 2=
139 sufficient. (Recorded in a supplementary file 1)

140 To measure birth satisfaction, there are two scales developed for measuring birth experience: the
141 Birth Satisfaction Scale–Revised (BSS-R) and the Childbirth Experience Questionnaire (CEQ2)
142 (23). We examined both scales and found similar results with regard to their ability to explain birth
143 satisfaction variance. We present our results on the CEQ2 in a supplementary file.

144 The Birth Satisfaction Scale–Revised (BSS-R)

145 The Birth Satisfaction Scale–Revised (BSS-R), created by Martin & Martin in 2014, is regarded
146 as the best instrument for measuring women's birth satisfaction (24). The BSS-R includes 10-items
147 in three factors: stress experienced during labor, women's personal attributes, and quality of care
148 provision. Participants were asked to rate each item on a four-point Likert scale which ranges from
149 0-4 (0 = strongly disagree to 4 = strongly agree). The BSS-R showed acceptable internal
150 consistency (Cronbach's alpha = 0.79). Its total score ranges from zero to 40. Mortazavi and
151 colleagues translated the scale into Persian. The validity study of the Persian BSS-R indicted that
152 it has three dimensions which are identical to those proposed by the developers of the original

153 scale. The reliability of the Persian BSS-R has been confirmed (Cronbach's alpha = 0.76) (25). In
154 the present study, we calculated a Cronbach's alpha value of 0.734.

155 The World Health Organization's Well-Being Index (WHO-5 Well-Being Index)

156 The 5-item World Health Organization Well-Being Index (WHO-5) assesses the emotional well-
157 being of individuals over the preceding two weeks. It has five items which are rated using a 6-
158 point Likert scale where zero represents 'having good feelings at no time' and five represent 'having
159 good feelings all the time' (26). The scale's total score ranges from 0 to 25 which is converted to a
160 scale of 0 to 100. The scale is used in screening programs for depression with a score of 50 as the
161 cut-off point. Individuals with scores less than fifty should be referred for further assessment. The
162 validity and reliability of the Persian version of WHO-5 in pregnant women were confirmed and
163 its unidimensionality and reliability was proven (Cronbach's alpha = 0.85) (27). In our study, we
164 calculated a Cronbach's alpha value of 0.889.

165 Fear of COVID-19 Scale (FCV-19S)

166 The Fear of COVID-19 Scale (FCV-19S) is one of the most widely used scales for assessing fear
167 of COVID-19. It was developed by Ahorsu et al. (2020) for the purpose of assessing fear of
168 COVID-19. The scale was translated into Persian and the validity study for the scale was conducted
169 using a sample of Iranian students (28). It is a unidimensional instrument consisting of 7 items
170 rated on a 5-point Likert-type scale (1 to 5 points). The total score ranges from 7 to 35 with higher
171 scores indicating a higher level of fear of COVID-19. A Cronbach's alpha value of 0.88 was
172 reported in the original study on the scale (28). In our study, we calculated a Cronbach's alpha
173 value of 0.928.

174 **Data analysis**

175 We used the SPSS version 18 to analyze the data. Descriptive statistics were used to characterize
176 the participants. We evaluated the normal distribution of birth satisfaction scores and other
177 quantitative variables using skewness and kurtosis. We used the general univariate linear model to
178 identify independent variables with a significant impact on birth satisfaction scores. Then,
179 variables with $p < 0.25$ in the univariate linear regression were entered into five separate multiple
180 linear regression analyses by the backward-LR method. These analyses enabled us to determine
181 the demographic, obstetric, labor and birth, psychological, and overall predictors of birth
182 dissatisfaction. We checked linear regression assumptions. The normality of residuals was verified
183 and collinearity statistics indicated no multicollinearity (tolerance < 1 and variance inflation factor
184 < 2). The effect sizes for the entire model and main predictors of birth satisfaction scores were
185 calculated.

186 **Results**

187 Of the 676 women who gave birth in Mobini hospital during the study period, 67 women were
188 excluded from the study of whom two women were COVID positive, 27 had a preterm birth, 35
189 women had an infant in NICU, four women experienced severe postpartum hemorrhage, and eight
190 women did not consent to participate in the study. Overall, 601 women participated in the study.
191 The mean value of age (year), education (years), gestational age (week), birth weight (gr), and
192 admission to delivery duration (hour) were 28.7 ± 6.6 , 11.1 ± 4.1 , 39.1 ± 1.2 , 3250 ± 465 , and 7.2 ± 8.0 ,
193 respectively. The mean scores of FVC-19S, WHO-5, and birth satisfaction were 14.7 ± 7.5 ,
194 69.5 ± 26.8 , 67.5 ± 13.0 , and 28.6 ± 7.3 , respectively. Sixty-five point nine percent (65.9%) of the
195 participants were multiparous. The correlation between birth satisfaction total scores and the

196 duration between admission to hospital and giving birth was 0.247 ($p < 0.001$). Sample
 197 characteristics and the results of general linear models for the relationship between the BSS-R
 198 scores and independent variables are presented in table 1.

199 Table 1. Sample characteristics, mean (SD) of birth satisfaction scores, and the results of general
 200 linear models on birth satisfaction scores (N = 601).

Demographic variables	N (%)	M (SD) of birth satisfaction scores [€]	Mean difference (95% CI)	P
Age (years)				
≤ 20	70 (11.6)	28.3 ± 7.6	Ref	
21-34	405 (67.4)	28.7 ± 7.3	0.42 (-1.44, 2.27)	0.659
≥ 35	126 (21.0)	28.4 ± 7.2	0.12 (-2.01, 2.25)	0.912
Educational level (years)				
Primary school	94 (15.6)	29.9 ± 6.7	-2.36 (-4.59, -0.13)	0.034*
High school	335 (55.7)	28.8 ± 7.6	-1.15 (-3.19, 0.88)	0.518
University	172 (28.6)	27.6 ± 6.9	Ref	
Job				
Housewife	547 (91.0)	28.8 ± 7.1	Ref	0.086†
Employed	54 (9.0)	27.0 ± 8.7	-1.79 (-3.82, 0.025)	
Household income				
Insufficient	107 (17.8)	27.9 ± 8.6	-0.90 (-2.42, 0.63)	0.249

Sufficient	494 (82.2)	28.8 ± 7.0	Ref	
Resident				
Urban	411 (68.4)	28.1 ± 7.2	-1.49 (-2.74, 0.24)	0.019*†
Rural	190 (31.6)	29.6 ± 7.3	Ref	
Obstetric variables				
Parity				
Primipara	205 (34.1)	26.9 ± 7.4	-2.62 (-3.83, -1.40)	<0.001***†
Multipara	396 (65.9)	29.6 ± 7.0	Ref	
History of abortion				
Yes	176 (29.3)	28.3 ± 7.4	-0.38 (-1.66, 0.90)	0.558
No	425 (70.7)	28.7 ± 7.2	Ref	
Infant gender				
Female	315 (52.4)	28.8 ± 6.9	0.48 (-0.69, 1.64)	0.425
Male	286 (47.6)	28.4 ± 7.7	Ref	
Complicated pregnancy				
Yes	123 (20.5)	28.1 ± 8.0	-0.66 (-2.11, 0.79)	0.371
No	478 (79.5)	28.7 ± 7.1	Ref	
Poor obstetric history				
Yes	51 (8.5)	26.9 ± 8.2	-1.86 (-3.95, 0.23)	0.081†
No	550 (91.5)	28.8 ± 7.2	Ref	
Chronic disease				
Yes	62 (10.3)	27.8 ± 8.4	-0.91 (-2.83, 1.01)	0.352

No	539 (89.7)	28.7 ± 7.1	Ref	
Attending prenatal class				
Yes	206 (34.3)	27.8 ± 7.3	-1.27 (-2.49, -0.04)	0.042*†
No	395 (65.7)	29.0 ± 7.2	Ref	
Labor and Birth				
Gestational age at birth (week)				
< 38	91 (15.1)	29.2 ± 7.9	1.07 (-0.81, 2.96)	0.263
38-40	354 (58.9)	28.7 ± 7.2	0.55 (-0.83, 1.92)	0.438
> 40	156 (26.0)	28.1 ± 7.0	Ref	
Birth weight (gr)				
2500-3999	536 (89.2)	28.8 ± 7.2	Ref	
< 2500	24 (4.0)	27.3 ± 8.6	-1.55 (-4.52, 1.43)	0.308
≥ 4000	41 (6.8)	26.3 ± 7.4	-2.50 (-4.80, -0.19)	0.034*†
Mode of birth				
Elective cesarean	117 (19.5)	30.6 ± 5.9	Ref	
Emergency cesarean	65 (10.8)	23.0 ± 7.9	-7.59 (-9.70, -5.48)	<0.001***†
Vaginal delivery	391 (65.1)	29.1 ± 7.0	-1.49 (-2.93, -0.05)	0.043*†
VBAC [‡]	11 (1.8)	29.0 ± 5.7	-1.07 (-5.92, 2.70)	0.464
Instrumental delivery	17 (2.8)	24.1 ± 8.1	-6.49 (-10.04, -2.94)	<0.001***†
Labor pain				
Spontaneous	306 (50.9)	29.8 ± 7.4	Ref	
Induced	179 (29.8)	26.9 ± 7.6	-1.92 (-3.25, -0.60)	0.004***†

Elective cesarean	116 (19.3)	30.7 ± 5.9	1.86 (0.33, 3.34)	0.018*†
Having a private midwife				
Yes	150 (25.0)	28.3 ± 7.2	Ref	
No	336 (55.9)	28.0 ± 7.6	-0.30 (-1.70, 1.10)	0.669
Elective cesarean	115 (19.1)	30.7 ± 6.0	-2.34 (-4.10, 0.59)	0.009**
Fundal pressure				
Yes	308 (51.2)	28.4 ± 7.3	-1.99 (-3.57, -0.42)	0.013*†
No	111 (18.5)	30.4 ± 6.4	Ref	
Cesarean	182 (30.3)	27.9 ± 7.6	-2.48 (-4.19, -0.77)	0.005**
Suturing				
Yes – spontaneous tear	193 (32.1)	29.7 ± 7.1	-1.12 (-3.02, 0.77)	0.245†
Yes – episiotomy	143 (24.8)	27.0 ± 7.3	-3.83 (-5.80, -1.85)	<0.001***†
Cesarean	182 (30.3)	27.9 ± 7.6	-2.89 (-4.81, -0.98)	0.003**
No	77 (12.8)	30.8 ± 5.9	Ref	
Pain relief method‡				
Cesarean-general analgesia	24 (4.0)	29.2 ± 6.7	-1.16 (-4.38, 2.05)	0.478
Cesarean -Spinal analgesia	158 (26.3)	27.7 ± 7.8	-2.63 (-4.41, -0.84)	0.004***†
Vaginal birth-Entonox	291 (48.4)	28.3 ± 7.1	-2.01 (-3.62, -0.40)	0.015*†
Vaginal birth -Hot water shower/massage/birth ball	22 (3.7)	29.9 ± 7.9	-0.38 (-3.71, 2.96)	0.825
Nothing	106 (17.6)	30.3 ± 6.7	Ref	
Admission to birth duration (hour)				
< 5	289 (48.1)	30.0 ± 6.4	Ref	

≥ 5	312 (51.9)	27.4 \pm 7.8	-2.60 (-3.75, -1.46)	<0.001***†
Psychological variables				
Desirability of pregnancy				
Wanted	378 (62.9)	28.3 \pm 7.0	Ref	
Unplanned	74 (12.3)	27.8 \pm 9.0	-0.474 (-2.29, 1.34)	0.607
Unwanted	149 (24.8)	29.7 \pm 7.0	1.42 (0.04, 2.80)	0.044*†
Fear of COVID-19				
Low fear (< the median score)	280 (46.6)	29.3 \pm 7.6	Ref	
High fear (\geq the median score)	321 (53.4)	28.0 \pm 6.9	-1.27 (-2.44, -0.11)	0.032*†
Well-being index (WHO-5 score)				
< 50	139 (23.1)	26.7 \pm 7.1	-2.48 (-3.85, -1.11)	<0.001***†
≥ 50	462 (76.9)	29.2 \pm 7.3	Ref	
Satisfaction with pregnancy				
Low satisfied	133 (22.1)	26.6 \pm 8.0	-3.23 (-4.99, -1.46)	<0.001***†
Moderately satisfied	139 (23.1)	27.8 \pm 6.9	-2.00 (-3.75, -0.27)	0.017*†
Satisfied/very satisfied	329 (54.7)	29.8 \pm 6.9	Ref	
Perceived marital/sexual satisfaction				
Low satisfied	46 (7.6)	28.0 \pm 9.3	-0.72 (-2.93, 1.49)	0.521
Moderately satisfied	91 (15.1)	28.1 \pm 6.7	-0.70 (-2.34, 0.94)	0.401
Satisfied/very satisfied	464 (77.3)	28.8 \pm 7.2	Ref	
Satisfaction with spouse's support				
Low satisfied	40 (6.7)	25.7 \pm 9.2	-3.15 (-5.49, -0.81)	0.008***†

Moderately satisfied	74 (12.3)	28.6 ± 7.0	-0.30 (-2.07, 1.48)	0.742
Satisfied/very satisfied	487 (81.0)	28.9 ± 7.1	Ref	
<p>†variables entered in the multiple linear regression analysis , ‡vaginal birth after cesarean, §NVD group was included in the analysis,</p> <p>¶higher mean scores are indicative of higher birth satisfaction, the highest and lowest possible scores (0-40), * <0.05, **<0.01, ***<0.001</p>				

201

202 In table 2, the results of multiple linear regression analysis for the birth satisfaction scores are

203 presented. Also, effect sizes for the main predictors of birth satisfaction are included in table 2.

204 Obstetric and labor and birth predictors of birth satisfaction were primiparity [B = -2.650, CI (-

205 3.861, -1.439)], birth weight (gr) \geq 4000 [-2.351(-4.521, -0.180)], emergency cesarean [-7.332 (-

206 9.276, -5.389)], and episiotomy [-2.466 (-3.853, -1.079)]. Psychological predictors of birth

207 satisfaction were unwanted pregnancy [2.292 (.958, 3.625)], Well-being score < 50 [-1.742 (-3.145, -

208 0.338)], low satisfaction with spouse's support [-2.523 (-4.828, -0.219)], and low [-2.910 (-4.400,

209 -1.419)] and moderate [-2.168 (-3.580, -0.755)] satisfaction with pregnancy. Overall predictors of

210 birth satisfaction were emergency cesarean [-7.463(-9.310, -5.616), instrumental delivery [-3.571(-

211 6.907, -0.235)], episiotomy [-2.227 (-3.591, -0.862)], Entonox analgesia [-1.548(-2.726, -0.371)], Well-

212 being score < 50 [-1.812(-3.146, -0.478)], fear of COVID -19 [-1.216(-2.288,, -0.144)], low satisfaction

213 with pregnancy -2.539(-3.952, -1.127) and low satisfaction with spouse's support [-2.419(-4.598, -

214 0.240)]. The overall proportion of the variance in birth satisfaction explained by all variables is

215 17.4% (effect size = 0.174). Labor and birth variables explained 12.2% of the variance in birth

216 satisfaction.

217 Table 2. Results of multiple linear regression analysis on the birth satisfaction scores.

	Coefficients		Collinearity Statistics
--	--------------	--	-------------------------

	Beta	B (95.0% CI)	P	Tolerance	VIF
Demographic predictors					
Primary school vs. University	0.073	1.464 (-0.280, 3.209)	0.100	0.836	1.196
Urban vs. rural	-0.082	-1.280 (-2.600, 0.041)	0.057	0.891	1.123
Insufficient income	-0.078	-1.490 (-3.068, 0.089)	0.064	0.920	1.086
Obstetrics predictors					
Primiparity	-0.173	-2.650 (-3.861, -1.439)	<0.001***	0.999	1.001
Poor obstetric history	-0.077	-1.998 (-4.057, 0.062)	0.057	0.999	1.001
Labor and birth predictors					
Birth weight (gr) \geq 4000	-0.082	-2.351(-4.521, -0.180)	0.034*	0.996	1.004
Instrumental birth	-0.072	-3.170 (-6.600, 0.258)	0.070	0.924	1.083
Emergency cesarean	-0.313	-7.332 (-9.276, -5.389)	<0.001**	0.819	1.221
Induced labor pain	-0.069	-1.102 (-2.360, 0.156)	0.086	0.902	1.109
Episiotomy repair	-0.146	-2.466 (-3.853, -1.079)	0.001**	0.832	1.202
Vaginal birth-Entonox	-0.078	-1.142 (-2.385, 0.101)	0.072	0.774	1.293
Psychological predictors					
Unwanted pregnancy	0.136	2.292 (0.958, 3.625)	0.001**	0.958	1.044
Well-being score < 50	-0.101	-1.742 (-3.145, -0.338)	0.015*	0.907	1.103

High fear of COVID -19 (scores \geq 13)	-0.070	-1.016 (-2.151, -0.119)	0.079	0.991	1.009
Low satisfaction with spouse's support	-0.086	-2.523 (-4.828, -0.219)	0.032*	0.963	1.039
Low satisfaction with pregnancy	-0.166	-2.910 (-4.400, -1.419)	<0.001***	0.830	1.205
Moderate satisfaction with pregnancy	-0.126	-2.168 (-3.580, -0.755)	0.003**	0.896	1.116
Overall predictors					
Emergency cesarean (ref: elective cesarean)	-0.319	-7.463(-9.310, -5.616)	<0.001***	0.853	1.172
Instrumental birth	-0.081	-3.571(-6.907, -0.235)	0.036*	0.918	1.090
Episiotomy repair	-0.132	-2.227 (-3.591, -0.862)	0.001**	0.809	1.237
Vaginal birth-Entonox	-0.106	-1.548(-2.726, -0.371)	0.010*	0.811	1.233
Well-being score < 50	-0.105	-1.812(-3.146, -0.478)	0.008**	0.887	1.127
High fear of COVID -19	-0.083	-1.216(-2.288, -0.144)	0.026*	0.981	1.019
Low satisfaction with pregnancy	-0.145	-2.539(-3.952, -1.127)	<0.001***	0.816	1.225
Low satisfaction with spouse's support	-0.083	-2.419(-4.598, -0.240)	0.030*	0.952	1.051
Primary school vs. university	0.069	1.372(-0.112, 2.856)	0.070	0.966	1.035

Unwanted pregnancy	0.075	1.259(-0.028, 2.547)	0.055	0.908	1.101
Moderate satisfaction with pregnancy	-0.076	-1.319 (-2.660, 0.023)	0.054	0.877	1.140
Birth weight (gr) \geq 4000	-0.070	-2.022(-4.154, 0.110)	0.063	0.971	1.029
* <0.05, **<0.01, ***<0.001, Adjusted R square for the first regression analysis= 1.2%, Adjusted R square for the second regression analysis = 3.2%, , Adjusted R square for the third regression analysis = 12.2%, Adjusted R square for the fourth regression analysis = 7.5%, Adjusted R square for the fifth regression analysis = 17.4%, method: backward					

218
219
220 We investigated the relationship between the presence of a private midwife during labor and fear
221 of COVID-19 in the case of women who had a vaginal delivery. Women who were accompanied
222 by a private midwife reported a higher level of fear of COVID-19 than those without a private
223 midwife ($p = 0.044$). The presence of a private midwife during labor had no relationship with birth
224 satisfaction scores.

225 In table 3, the distribution of psychological variables including birth satisfaction, well-being, and
226 fear of COVID-19 according to parity and household income in the pandemic period are presented.
227 The BSS-R mean scores was higher in multiparas than primiparas ($p < 0.001$). The FVC-19S and
228 WHO-5 mean scores were not different between multiparas and primiparas ($p > 0.05$). They were
229 also not different between women with sufficient and insufficient income ($p > 0.05$).

230 Table 3. Mean scores of scales used in the study for different income and parity groups

		BSS-R [†]	WHO-5 [‡]	FCV-19S [€]
Parity				
	Primiparas	26.8 \pm 7.4	71.5 \pm 25.3	14.2 \pm 7.1
	Multiparas	29.4 \pm 7.0	68.4 \pm 27.4	14.9 \pm 7.7

P		<0.001***	0.159	0.299
Income				
	Sufficient	28.7 ± 6.9	70.1 ± 26.2	14.6 ± 7.2
	Insufficient	27.8 ± 8.5	66.3 ± 29.3	15.0 ± 8.6
P		0.314	0.183	0.634
†birth satisfaction scale-revised, ‡WHO-5 well-being index, € Fear of COVID-19 scale * <0.05, **<0.01, ***<0.001				

231

232 Discussion

233

234 We investigated the predictors of birth satisfaction in early postpartum during the COVID-19

235 epidemics' fifth wave in Iran. During the early pandemic, women could not have a companion in

236 the postpartum ward. Also, childbirth preparatory classes were closed and midwives were reluctant

237 to offer their services as private midwives. But after one month the situation improved. Virtual

238 preparatory classes became available and private midwives resumed their services supporting

239 women at birth. The fifth wave of the epidemic, caused by the delta variant, resulted in the highest

240 number of infections and deaths in comparison with previous waves in the country. Although a

241 lockdown was imposed, access to health care facilities was not limited and pregnant women could

242 visit hospitals. They were also allowed to have a private midwife in labor and a companion during

243 postpartum. At this juncture in the COVID-19 pandemic in Iran, vaccination of pregnant women

244 had not yet been begun but almost all midwives had been infected and all had been vaccinated.

245 We explored the role of variables related to labor and birth and also psychological variables in

246 birth satisfaction. Our results indicate that emergency cesarean, instrumental birth, episiotomy,

247 Entonox analgesia, low well-being score, high fear of COVID -19, low satisfaction with pregnancy

248 and a low satisfaction with spouse's support are predictors of lower levels of birth satisfaction.

249 The overall proportion of the variance in birth satisfaction explained by all variables is 17.4%.
250 Labor and birth variables explained 12.2% of the variance in birth satisfaction.

251 Our results are in line with those of Preis and colleagues (29). They found that in the pandemic
252 period, established predictors of low birth satisfaction such as nulliparity, mode of birth, social
253 support, and labor and birth complications explained 35% of the variance in birth satisfaction.
254 According to the same study, pandemic-related variables including maternal concerns about
255 preparation for birth and restrictions on the number of family members allowed to accompany a
256 birthing mother explained 3% of the variance in birth satisfaction (29).

257 We found that fear of contracting COVID-19 during the fifth wave of the disease was a predictor
258 of lower birth satisfaction. But we expected it to be a stronger predictor of lower birth satisfaction
259 than our results suggest. Fear of COVID-19 may induce higher anxiety before admission to the
260 hospital because of uncertainty and unfamiliarity with the hospital setting and the higher risk of
261 contracting COVID. But during labor and birth, uncertainty and unfamiliarity are diminished and
262 women feel quite relieved when they are about to be discharged from the hospital in a few hours.

263 Previous studies have found that the level of stress experienced during pregnancy and childbirth
264 was significantly associated with birth dissatisfaction (15, 18). We found no difference in the levels
265 of fear of COVID-19 between parity or income groups. This implies that all mentioned groups of
266 women experienced fear of COVID-19. In contrast, the results of a previous study indicate that fear
267 of COVID-19 was associated with parity and stage of pregnancy (30).

268 We found that compared to elective cesarean, emergency cesarean and instrumental birth were
269 better predictors of lower levels of birth satisfaction. This result may be explained by delays and
270 shortcomings in the implementation of policies and guidelines promoting physiological vaginal

271 birth in our setting. Results from previous studies indicated that cesarean was associated with lower
272 birth satisfaction in comparison with physiological vaginal birth (4, 8, 18, 29, 31).

273 In our study, women's low level of well-being was a predictor of lower birth satisfaction. We
274 found similar result in our previous study (8). Comparisons of well-being according to parity and
275 income indicate that in the pandemic period, well-being mean scores are not different between
276 parity and income groups. This implies that the pandemic has influenced well-being in both these
277 groups. In the present study, having a pregnancy that is stressful and characterized by hassle was
278 a psychological predictor of birth satisfaction. Also, low satisfaction with spouse's support could
279 predict birth satisfaction. These results are in agreement with the findings of previous studies (8,
280 32).

281 We found that vaginal birth by episiotomy rather than spontaneous tear is a predictor of birth
282 satisfaction. In our sample, 24.8 and 32.1 percent of the women experienced episiotomy and
283 spontaneous tear, respectively. Our study was conducted in a mother friendly hospital and so
284 episiotomy is not performed as a routine procedure; however, because it is a training hospital,
285 newly assigned obstetrics residents may perform episiotomy. In a study in Tehran, Iran, vaginal
286 birth by episiotomy accompanied by tear was a predictor of birth satisfaction (33).

287 Our results indicate that receiving Entonox analgesia is a predictor of lower birth satisfaction.
288 Entonox analgesia is used for vaginal birth. Our result is not in accord with those reported by
289 Fumagalli and colleagues. Their findings indicate that none of the intrapartum interventions was
290 associated with birth satisfaction (34).

291 In our study, having a stressful, hassled pregnancy was a predictor of lower birth satisfaction. Also
292 low satisfaction with spouse's support could predict lower birth satisfaction. These results are in
293 agreement with findings of our previous study (8). In a study on 225 postpartum women in Khaf,
294 Iran, childbirth experience improved with lower hassle and an increased sense of uplift (32).

295 We found that giving birth to an infant with high birth weight is a predictor of lower birth
296 satisfaction. High birth weight may cause long labor and higher pain levels which are associated
297 with lower levels of birth satisfaction.

298 We found that none of the socio-demographic variables were related to maternal birth satisfaction.
299 This is in line with the study by Fumagalli et al. (34) and our previous study (8). Results from the
300 present study indicate that primiparity is an obstetrical predictor of lower birth satisfaction. This
301 is in agreement with the results of several previous studies (18, 33, 34). Satisfaction with childbirth
302 is related to three factors: women's attributes, quality of care received, and stress. Several studies
303 have shown that multiparas experience lower levels of stress and fear of childbirth because of their
304 previous birth experience. Also, giving birth to the first child is usually more difficult than the
305 second or third child. That the satisfaction with current birth is higher in multiparas than primiparas
306 may be the result of their previous experience which decreases their fear of birth and also makes
307 delivery easier and more comfortable.

308 We found no relationship between the presence of a private midwife during labor and the birth
309 satisfaction score. It seems that the role of private midwives in improving birth satisfaction is not
310 as significant as commonly believed and needs further evaluation. Childbirth preparatory classes
311 have become increasingly popular among Iranian pregnant women. In prenatal visits, pregnant
312 women receive the option to register for free childbirth preparatory classes. During the pandemic,

313 virtual classes were held by midwives who work in private practices or public health centers. After
314 completing the course, participants may opt to have a private midwife during labor, birth, and early
315 postpartum. Private midwives are permitted to support women during active labor, birth, and early
316 postpartum. Most of them do not have permission to perform vaginal exams, delivery of the baby,
317 and other interventions. Obstetrics residents or midwifery students under the supervision of
318 mentors usually perform the deliveries.

319 We found that women who had a private midwife during labor and birth reported a higher level of
320 fear of COVID-19. A possible reason for this may be that during the pandemic, women with higher
321 levels of fear had hired private midwives to receive better care. But because this is a cross-sectional
322 study, we cannot conclude a causal relationship between the two factors. In a study in Russia, the
323 rate of having a support person at labor decreased from 58% in the pre-pandemic period to 27% in
324 the pandemic due to the COVID-19 restrictions (35). Results of a study in the US indicate that the
325 number of support persons during birth predicted birth satisfaction (29).

326 **Limitations and strengths**

327 The main limitation of this study is due to its cross-sectional design which makes it difficult to
328 establish cause and effect relationships between some variables. The strong points of this study
329 are the large sample size. We investigated birth satisfaction in the early hours after birth; so, in
330 comparison with studies that were conducted several months after birth, our results are relatively
331 more precise. This study was performed using a sample of postpartum women who gave birth in a
332 conventional birth setting. Its findings cannot be generalized to populations who give birth in
333 labor-delivery-recovery settings where women are isolated from other parturient and therefore are
334 less worried about contacting COVID-19.

335 **Implications for future research**

336 Data collection for this study was completed before the vaccination of pregnant women against
337 COVID-19 had started. We recommend that further studies be undertaken in Iran to explore the
338 effects of vaccination on birth satisfaction. According to our findings, hiring a private midwife
339 was not associated with higher birth satisfaction. We believe that this finding merits further
340 exploration and so we recommend that qualitative studies be conducted to explore perspectives of
341 women and midwives on the role of private midwives.

342 **Conclusions**

343 We found that fear of COVID -19 is a predictor of lower birth satisfaction. Our findings also
344 indicate that variables related to labor and birth were predictors of birth satisfaction. These
345 variables which were responsible for a large part of the birth satisfaction during the pandemic
346 include the following: emergency cesarean, instrumental birth, Entonox analgesia, episiotomy, low
347 level of well-being, low satisfaction with pregnancy, and low satisfaction with husband's support.
348 Based on our findings we recommend a number interventions to increase birth satisfaction and
349 thus maternal mental health. Chief among these are interventions to lower fear of contracting
350 COVID -19 and to reduce rates of episiotomy, emergency cesarean, and instrumental birth.

351 **Acknowledgements**

352 We would like to thank pregnant women who participated in this study.

353 **References**

- 354 1. Bryanton J, Gagnon AJ, Johnston C, Hatem M. Predictors of Women ' s Perceptions of
355 the Childbirth Experience. JOGNN. (2008)37 24-34.doi:10.1111/J.1552-6909.2007.00203.x

- 356 2. Karlström A, Nystedt A, Hildingsson I. The meaning of a very positive birth experience:
357 focus groups discussions with women. *BMC pregnancy and childbirth*.
358 (2015)15(1):251.doi:10.1186/s12884-015-0683-0
- 359 3. Bell AF, Andersson E, Goding K, Vonderheid SC. The birth experience and maternal
360 caregiving attitudes and behavior: A systematic review. *Sexual & reproductive healthcare* :
361 official journal of the Swedish Association of Midwives. (2018)16:67-
362 77.doi:10.1016/j.srhc.2018.02.007
- 363 4. Janevic T, Maru S, Nowlin S, McCarthy K, Bergink V, Stone J, et al. Pandemic Birthing:
364 Childbirth Satisfaction, Perceived Health Care Bias, and Postpartum Health During the COVID-
365 19 Pandemic. *Matern Child Health J*. (2021)25(6):860-9.doi:10.1007/s10995-021-03158-8
- 366 5. Pang MW, Leung TN, Lau TK, Hang Chung TK. Impact of first childbirth on changes in
367 women's preference for mode of delivery: follow-up of a longitudinal observational study. *Birth*
368 (Berkeley, Calif). (2008)35(2):121-8.doi:10.1111/j.1523-536X.2008.00225.x
- 369 6. Preis H, Tovim S, Mor P, Grisaru-Granovsky S, Samueloff A, Benyamini Y. Fertility
370 intentions and the way they change following birth- a prospective longitudinal study. *BMC*
371 *pregnancy and childbirth*. (2020)20(1):228.doi:10.1186/s12884-020-02922-y
- 372 7. Waldenström U, Hildingsson I, Rubertsson C, Rådestad I. A negative birth experience:
373 prevalence and risk factors in a national sample. *Birth (Berkeley, Calif)*. (2004)31(1):17-
374 27.doi:10.1111/j.0730-7659.2004.0270.x
- 375 8. Mortazavi F, Mehrabadi M. Predictors of low birth satisfaction among Iranian
376 postpartum women: A cross-sectional study. *Nursing Open*. (2022)9(1):604-
377 13.<https://doi.org/10.1002/nop2.1104>
- 378 9. Tadele M, Bikila D, Fite RO, Obsa MS. Maternal satisfaction towards childbirth Service
379 in Public Health Facilities at Adama town, Ethiopia. *Reproductive Health*.
380 (2020)17(1):60.doi:10.1186/s12978-020-00911-0
- 381 10. Jafari E, Mohebbi P, Mazloomzadeh S. Factors Related to Women's Childbirth
382 Satisfaction in Physiologic and Routine Childbirth Groups. *Iran J Nurs Midwifery Res*.
383 (2017)22(3):219-24.doi:10.4103/1735-9066.208161
- 384 11. Takacs L, Seidlerova JM, Sulova L, Hoskovcova SH. Social psychological predictors of
385 satisfaction with intrapartum and postpartum care - what matters to women in Czech maternity
386 hospitals? *Open medicine (Warsaw, Poland)*. (2015)10(1):119-27.doi:10.1515/med-2015-0022

- 387 12. Pantoja L, Weeks FH, Ortiz J, Cavada G, Foster J, Binfa L. Dimensions of childbirth care
388 associated with maternal satisfaction among low-risk Chilean women. *Health Care for Women*
389 *International*. (2020)41(1):89-100.doi:10.1080/07399332.2019.1590360
- 390 13. Mortazavi F, Ghardashi F. The lived experiences of pregnant women during COVID-19
391 pandemic: a descriptive phenomenological study. *BMC pregnancy and childbirth*.
392 (2021)21(1):193.doi:10.1186/s12884-021-03691-y
- 393 14. Mollard E, Kupzyk K, Moore T. Postpartum stress and protective factors in women who
394 gave birth in the United States during the COVID-19 pandemic. *Women's health (London,*
395 *England)*. (2021)17:17455065211042190.doi:10.1177/17455065211042190
- 396 15. Mariño-Narvaez C, Puertas-Gonzalez JA, Romero-Gonzalez B, Peralta-Ramirez MI.
397 Giving birth during the COVID-19 pandemic: The impact on birth satisfaction and postpartum
398 depression. *International Journal of Gynecology & Obstetrics*. (2021)153(1):83-
399 8.doi.org/10.1002/ijgo.13565
- 400 16. Mortazavi F, Mehrabadi M, KiaeeTabar R. Pregnant women's well-being and worry
401 during the COVID-19 pandemic: a cross-sectional study. *BMC pregnancy and childbirth*.
402 (2021)21(1):59.doi:10.1186/s12884-021-03548-4
- 403 17. Ahorsu DK, Imani V, Lin CY, Timpka T, Broström A, Updegraff JA, et al. Associations
404 Between Fear of COVID-19, Mental Health, and Preventive Behaviours Across Pregnant
405 Women and Husbands: An Actor-Partner Interdependence Modelling. *Int J Ment Health Addict*.
406 (2020):1-15.doi:10.1007/s11469-020-00340-x
- 407 18. Urbanová E, Škodová Z, Bašková M. The Association between Birth Satisfaction and the
408 Risk of Postpartum Depression. *Int J Environ Res Public Health*.
409 (2021)18(19).doi:10.3390/ijerph181910458
- 410 19. Gibbons L, Belizán JM, Lauer JA, Betrán P, Merialdi M, Althabe F. The Global Numbers
411 and Costs of Additionally Needed and Unnecessary Caesarean Sections Performed per Year:
412 Overuse as a Barrier to Universal Coverage Geneva: WHO 2010.
- 413 20. Darsareh F, Aghamolaei T, Rajaei M, Madani A. Determinants of caesarean birth on
414 maternal demand in the Islamic Republic of Iran: a review. *Eastern Mediterranean Health*
415 *Journal*. (2017)23(6):441-8.doi: 10.26719/2017.23.6.441

- 416 21. Cincotta R, Sadjadpour K. IRAN IN TRANSITION: The Implications of the Islamic
417 Republic's Changing Demographics. In: Peace. CEfI, editor. Washington, DC: Carnegie
418 Endowment for International Peace; (2017).
- 419 22. Moradi-Lakeh M, Vosoogh-Moghaddam A. Health sector evolution plan in Iran; equity
420 and sustainability concerns. *Int J Health Policy Manag.* (2015)4(10):637-40.doi:
421 10.15171/ijhpm.2015.160
- 422 23. Walker KF, Dencker A, Thornton JG. Childbirth experience questionnaire 2: Validating
423 its use in the United Kingdom. *European Journal of Obstetrics & Gynecology and Reproductive*
424 *Biology: X.* (2020)5:100097.<https://doi.org/10.1016/j.eurox.2019.100097>
- 425 24. Martin CJH, Martin CR. Development and psychometric properties of the Birth
426 Satisfaction Scale-Revised (BSS-R) Midwifery. (2014)30:610-
427 9.doi.org/10.1016/j.midw.2013.10.006
- 428 25. Mortazavi F, Mehrabadi M, Hollins Martin CJ, Martin CR. Psychometric properties of
429 the birth satisfaction scale-revised (BSS-R) in a sample of postpartum Iranian women. *Health*
430 *Care for Women International.* (2020):1-16.doi:10.1080/07399332.2020.1802464
- 431 26. About the WHO-5.: WHO; 2020 [Available from: [https://www.psykiatri-regionh.dk/who-](https://www.psykiatri-regionh.dk/who-5/about-the-who-5/Pages/default.aspx)
432 [5/about-the-who-5/Pages/default.aspx](https://www.psykiatri-regionh.dk/who-5/about-the-who-5/Pages/default.aspx). Accessed 15 Dec 2020.
- 433 27. Mortazavi F, Mousavi SA, Chaman R, Khosravi A. Validation of the World Health
434 Organization-5 Well-Being Index; assessment of maternal well-being and its associated factors.
435 *Turk psikiyatri dergisi = Turkish journal of psychiatry.* (2015)26(1):1-7
- 436 28. Ahorsu DK, Lin C-Y, Imani V, Saffari M, Griffiths MD, Pakpour AH. The Fear of
437 COVID-19 Scale: Development and Initial Validation. *International Journal of Mental Health*
438 *and Addiction.* (2020).doi:10.1007/s11469-020-00270-8
- 439 29. Preis H, Mahaffey B, Heiselman C, Lobel M. The impacts of the COVID-19 pandemic
440 on birth satisfaction in a prospective cohort of 2,341 U.S. women. *Women and birth : journal of*
441 *the Australian College of Midwives.* (2021).doi:10.1016/j.wombi.2021.10.004
- 442 30. Giesbrecht GF, Rojas L, Patel S, Kuret V, MacKinnon AL, Tomfohr-Madsen L, et al.
443 Fear of COVID-19, mental health, and pregnancy outcomes in the pregnancy during the COVID-
444 19 pandemic study: Fear of COVID-19 and pregnancy outcomes. *Journal of affective disorders.*
445 (2022)299:483-91.doi.org/10.1016/j.jad.2021.12.057

- 446 31. Kahalon R, Preis H, Benyamini Y. Who benefits most from skin-to-skin mother-infant
447 contact after birth? Survey findings on skin-to-skin and birth satisfaction by mode of birth.
448 *Midwifery*. (2021)92:102862.doi:10.1016/j.midw.2020.102862
- 449 32. Khalife-Ghaderi F, Amiri-Farahani L, Haghani S, Hasanpoor-Azghady SB. Examining
450 the experience of childbirth and its predictors among women who have recently given birth.
451 *Nursing Open*. (2021)8(1):63-71.doi.org/10.1002/nop2.603
- 452 33. Nahae J, Mohammad-Alizadeh-Charandabi S, Abbas-Alizadeh F, Martin CR, Hollins
453 Martin CJ, Mirghafourvand M, et al. Pre- and during-labour predictors of low birth satisfaction
454 among Iranian women: a prospective analytical study. *BMC pregnancy and childbirth*.
455 (2020)20(1):408.doi:10.1186/s12884-020-03105-5
- 456 34. Fumagalli S, Colciago E, Antolini L, Riva A, Nespoli A, Locatelli A. Variables related to
457 maternal satisfaction with intrapartum care in Northern Italy. *Women and Birth*.
458 (2021)34(2):154-61.doi.org/10.1016/j.wombi.2020.01.012
- 459 35. Yakupova V, Suarez A, Kharchenko A. Birth Experience, Postpartum PTSD and
460 Depression before and during the Pandemic of COVID-19 in Russia. *Int J Environ Res Public*
461 *Health*. (2021)19(1).doi:10.3390/ijerph19010335

462

463