1 Detailed evaluation of chromatic pupillometry and full-field stimulus testing to assess ultra-
2 low vision in retinitis pigmentosa
3

4 Midori Yamamoto[1,6], Take Matsuyama[1,6], Tadao Maeda[1], Seiji Takagi[2],
5 Naohiro Motozawa[1], Daiki Sakai[1,3,4], Yasuhiko Hirami[1,3],Akiko Maeda[1],Yasuo Kurimoto[1,3],
6 Masayo Takahashi[1,5],Michiko Mandai[1]
7

[1] Department of Ophthalmology , Kobe City Eye Hospital, Kobe, Japan
[2] Department of Ophthalmology, Toho University Graduate School of Medicine, Tokyo,
Japan
[3] Department of Ophthalmology, Kobe City Medical Center General Hospital, Kobe, Japan
[4] Department of Surgery, Division of Ophthalmology, Kobe University Graduate School of
Medicine, Kobe, Japan
[5]Vision Care Inc., Kobe, Japan
[6]First authors (equal contribution)
16

17 Meeting Presentation: The 74[th] Annual congress of Japan clinical Opthalmology,2020
18

19 Financial Support: Japan Agency for Medical Research and Development (AMED) under
20 grant number JP22bm0204002
21

22 Conflict of Interest: No conflicting relationship exists for any author
23

24 Running head: Multimodal approach to evaluate visual function in patients with low vision
25

26 Keywords: retinitis pigmentosa, FST, chromatic pupillometry
27

Address correspondence to:
Takeshi Matsuyama and Michiko Mandai
Kobe City Eye Hospital, 2-1-8 minamimachi, minatojima,
chuo-ku, kobe, 650- 0047, Japan
Tel: +81-78-381-9876
Fax: +81-78-381-9705
E-mail: michiko_mandai@kcho.jp
35
36

Abbreviations

CF	Counting fingers	
FST	Full-field stimulus testing	
HM	Hand motion	
IOP	Intraocular pressure	
IRD	Inherited retinal disease	
LP	Light perception	
iPSC	induced pluripotent cell	
NLP	Non light perception	
OCT	Optical coherence tomography	
RP	Retinitis pigmentosa	
RPE	Retinal pigment epithelium	

Abstract

Purpose

Novel therapeutic options, such as regenerative medicine and gene therapy are now emerging as viable treatment options for patients with severe visual impairments, such as retinitis pigmentosa (RP). Gradable assessment of patients' visual function is essential to consider treatment options and to evaluate treatment outcomes, however, evaluation of visual function in advanced low vision patients is often challenging due to patients' poor and sometimes unpredictable responses. In this study, we attempted to accurately assess the visual capabilities and disease stage in RP patients with visual acuity of 0.01 or lower, using chromatic pupillometry and full-field stimulus testing (FST), combined with retinal structural features, as determined by spectral-domain OCT.

Design

Retrospective analysis of visual function tests in 84 eyes of 43 patients with advanced RP with visual acuity of 0.01 and lower visited Kobe City Eye Hospital from 2019 to 2021.

Methods: Eighty-four eyes of 43 patients with advanced RP with visual acuity of 0.01 and lower were evaluated by chromatic pupillometry, FST, BCVA (best-corrected visual acuity), and OCT (optical coherence tomography). Hierarchical (multilevel) Bayesian modeling was used to estimate individual eye's pupil response and FST, taking into account the ambiguity and randomness often observed in ultra-low vision patients. Using the estimated ability

obtained from each test, the correlation between each test and retinal thickness was further

analyzed to make a comprehensive assessment of the data.

Results

FST and pupillometry measurements were moderately correlated with visual acuity, but

exhibited a wide range of values within the same visual acuity groups. FST was not

correlated with central retinal thickness at CF/HM VA range and seemed to reflect overall

remaining photoreceptor function including peripheral retina. Pupillometry seemed to

distinguish different levels inner retinal function.

Conclusion

The combination of pupillometry and FST allowed for graded evaluation of visual function

within patients grouped in the same visual acuity groups in advanced RP patients with ultra-

low vision.

Introduction

Retinitis pigmentosa (RP) is an inherited retinal degeneration with a reported prevalence of

1 in 4000 and is one of the leading causes of severe visual dysfunction worldwide.[1] In typical

retinitis pigmentosa, the preceding loss of rod photoreceptor cells leads to night blindness

and concentric visual field constriction, followed by a progressive loss of cone photoreceptor

cells resulting in deterioration of central vision, sometimes leading to blindness.

As with many neurodegenerative diseases, there have been no established treatments to

restore visual function once it is lost. However, in recent years, the development of retinal

cell-based therapies and gene therapies have been put in clinical trials for severe retinal

diseases. These new therapies conceptually allow for high expectations that they will not only

inhibit the disease progression but also improve visual function.

Optogenetic intervention, which aims to restore light responsiveness by ectopic expression of

light activated proteins in inner retinal cells, is an alternative gene therapy approach which

have also been applied to human patients[2]. Regenerative medicine is a replacement therapy

for damaged or lost cells to regenerate visual function. We reported a proof-of-concept for

pluripotent stem cell (PSC)-derived retinal organoid transplantation in animal models of RP

by confirming host-graft synapse formation, restoration of RGC light responses after

transplantation, and also by behavioral improvement[3,4]. Based on these POC, we recently

conducted a clinical trial for RP patients using human-induced pluripotent stem cell (iPSC)-

derived retinal organoids (jRCTa050200027)[5].

Development and clinical application of these new therapies mandate an accurate and

gradable assessment of patients' visual function, since they target specific points of the visual

pathway and disease stage. Unfortunately, patients with "severe visual dysfunction", for

example patients with a visual acuity lower than 0.01, which are the primary candidates for

these new treatments, are often grouped together into broad categories. A more detailed

assessment of which part of the visual pathway have a residual function, and to what extent,

will inform the best treatment options. If the therapeutic concept is to enhance photoreceptor

function through gene therapy or other means, it is important to determine the degree of

photoreceptor survival in the retina. For photoreceptor transplantation, it is crucial that

secondary and tertiary neurons that transmit signals from photoreceptors are functioning, or

can become functional following photoreceptor transplantation. Optogenetics and artificial

eye approaches would require a functioning neural pathway from retinal ganglion cells to the

visual cortex. A more accurate assessment of visual function and disease stage will help

determine the indications and methods of treatment. Additionally, the same methodology

could be utilized to reliably determined the effectiveness of the treatment and assess the

extent of improvement in visual functions, if any.

To assess visual function in late-stage RP patients with low vision, a number of functional
tests have been reported including full-field stimulus testing（FST）[6–8], and chromatic
pupillometry[9–12]. FST is a subjective test using the full field light stimuli, which can
theoretically separate the cone and rod photoreceptor responses by using different
wavelengths of the stimulus. Chromatic pupillometry is a noninvasive and objective method
to study pupillary responses induced by cone, rod photoreceptors and melanopsin-containing
intrinsically-photosensitive retinal ganglion cells (ipRGCs), which can be distinguished by
the intensity and wavelength (color) of light stimuli. [11,13,14] Light information for pupillary
light reflex is relayed to the brain through a pathway separate from the conventional visual
pathway, with optic tract fibers terminating at the pretectal nucleus in the midbrain, and not
at the LGN of the thalamus.[15–20] Selected stimulus conditions can produce pupil responses
that reflect phototransduction primarily mediated by rods, cones, or melanopsin/ipRGCs. We
conducted a detailed analysis of FST and chromatic pupillometry on advanced RP patients
with ultra-low vision. We then examined the correlation between the results of these tests and
conventional visual acuity (VA) test, as well as OCT observation. Our results indicate that
chromatic pupillometry and FST together with OCT can be used complementarily to more
accurately evaluate the visual function in advanced RP patients and make better assessments
on estimated pathological status of patients.
142

Methods

Patients

This retrospective observational study protocol was approved by the institutional review
board of Kobe City Medical Center General Hospital (approval number: E18006). The study
was conducted in accordance with the terms of the Declaration of Helsinki. Written informed
consent was obtained from all participants after an explanation of the nature and possible
consequences of the study. We included the patients with both RP and RP-related diseases
who visited the Kobe City Eye Hospital, Kobe, Japan between 2019 and 2021. Inclusion
criteria for patients were that the visual acuity was equal to or lower than 20/2000.
Examinations were performed in 84 eyes of 43 subjects (Table 1) and all patients underwent
a series of ophthalmic examinations that included the following: best-corrected visual acuity
(BCVA), intraocular pressure (IOP), fundus examination, spectral-domain optical coherence
tomography (SD-OCT), chromatic pupillometry and FST.
156

Visual acuity measurements

BCVA was tested using a standard Landolt C acuity chart. The visual acuities were converted
to logMAR to allow for statistical analysis. Visual acuity of 0.01 were assigned logMAR visual

acuity of 2.0, counting fingers (CF) 2.6, hand motion (HM) 2.9, light perception (LP) 3.1,

and no light perception (NLP) 3.4, following the convention of Jhonson et al.[21]

SD-OCT evaluation

The SD-OCT images were obtained by using the Heidelberg Spectralis (Heidelberg

Engineering, Spectralis, Heidelberg, Germany). The thickness of total retinal thickness (TRT)

and the thickness of Ganglion Cell Complex (GCC), Inner Nuclear Layer (INL), and

photoreceptor layer (photo) were measured at four cardinal points 2,000 μm from the fovea

in horizontal and vertical scans. All measurements were performed by using with "caliper"

function of the Heidelberg instrument. TRT was defined as the distance between the signal

peak at the vitreoretinal interface (the internal limiting membrane: ILM) and the posterior

boundary of the major signal peak that corresponds to the basal retinal pigment

epithelium/Bruch's membrane complex (RPE/BrM). GCC was defined as the three

innermost retinal layers: the nerve fiber layer, the ganglion cell layer, and the inner plexiform

layer. INL was defined as the distance between the basal Inner Plexiform Layer (IPL) and the

OPL. Lastly, the photoreceptor layer was defined as the region between Outer Plexiform

Layer (OPL)and RPE/BrM.

Chromatic pupillometry

Chromatic pupillometry and FST measurements were performed using an Espion system with

a ColorDome light stimulator in a completely dark room. The protocol for chromatic

pupillometry consisted of the following four steps. For each step measurements were repeated

5 times. The first step (rod) consisted of a -3 log cd/m^2 blue light stimulus following 10

minutes of dark adaptation. The second step (cone1) consisted of a 1 log cd/m^2 red light

stimulus on 0.1cd/m^2 blue background light following 2 minutes of light adaptation. The third

step (cone2) consisted of a 3 cd/m^2 white stimulus light on 1 cd/m^2 white background light

following 2 minutes of light adaptation. Lastly, the fourth step (mela) consisted of a blue light

stimulus of 150 cd/m^2 without background light following 2 minutes of dark adaptation. Rod

and cone responses are observed as transient responses immediately after light stimulation

whereas melanopsin response is observed as a sustained pupil response. Each test was

performed on both eyes simultaneously, and changes in pupil diameter were recorded with an

infrared camera in the color dome™.

Full-field stimulus testing (FST)

FST was examined using the same equipment as chromatic pupillometry. After 45 minutes

of dark adaptation following mydriatic eye drops in both eyes, blue (448nm), white

(590nm), green (530nm), and red (627nm) light stimuli were used to examine the rod and
cone functions. The non-tested eye was shielded so that one eye was tested at a time.
Subjects were given a button box (Diagnosys LLC, MA, USA) with a Yes/No button and
answered whether they perceived light in the color dome™ after a beep. Thresholds to the
light stimulus were measured for each color while also taking into account false negatives
and false positives.
202
Statical analyses
We used full Bayesian statistical inference with MCMC sampling for statistical modeling
using Rstan (Stan Development Team. 2017. RStan: the R interface to Stan. R package
version 2.21.5. http://mc-stan.org). Since most data consists of measurements gathered
from both eyes of subjects, we incorporated terms for the patient and eye bias to account for
the natural hierarchical structure of the data. Using these and other relevant parameters,
such as color of stimulus light, we obtained posterior estimates from linear regression
models. We show the posterior distributions of parameters of interest, which indicate the
probability for the value of the parameter given the data, with 95% compatibility intervals
(confidence intervals) and mean indicated. Note that posterior distributions do not
represent a simple pooling (average) of data for a particular set of predictor combinations,
rather they represent the effect of predictors while considering the data as a whole. Detailed
explanation of the statistical models, as well as the Stan code for the models are provided in
a GitHub repository (https://github.com/matsutakehoyo/Clinical-Data-Analysis).
217
218 **Results**
219
Robust hierarchical modeling of FST makes better threshold estimates
FST measurements were analyzed using a multilevel (hierarchical) logistic model as
illustrated in Figure 1A. We estimated patients' ability to detect light taking into
consideration the effect of light intensity, (x_i, in dB), patient (β_{pat}), eye (β_{eye}), and
stimulus color (β_{clr}) as predictors in our model. We also allowed eye biases to vary for
different colors (eye and color interaction, denoted eyeXclr). Figure 1B shows the posterior
distribution of model parameters. Overall, despite the presence of a large patient to patient
variation (β_{pat}), most of the patients showed little variation between the responses by right
(RE) and left eyes (LE). There is a clear effect for the color of the stimulus light, with
patients being most sensitive to blue light than red light (Figure 1B). A key aspect of the
model is the inclusion of the 'guessing' parameter (α) for robustness. The 'guessing'
parameter is a measure of points that do not conform well with the logistic regression, i.e.

the fraction of outliers. The small 'guessing' parameter estimates (mode < 0.05) for most

eyes support the reliability of this test, with some notable exceptions where subjects seem to

be 'guessing' more than half of the time (Figure 1B). Figure 1C shows representative cases

with posterior predictions from our model alongside data measurements with the 'guessing'

rate presented on top of each panel. Typically, a subject's probability of correctly identifying

the light stimulus decreases as the light intensity (x_i) decreases, as described by the S-

shaped curve of the logistic function. Vertical lines show the threshold value (the light

intensity at which the probability of success equals 0.5). The sensitivity to color generally

decreases from Blue to Red, although the amount of the shift varies from patient to patient

and even from RE to LE (id =4 and id =28 for example). In samples with larger 'guessing',

subject's responses were seemingly random and measurements tended to fall outside the S-

shaped curve of the logistic regression (patient id = 16 and 28 RE). Although the difference

in threshold values between our estimates (solid line) and those made by the measurement

instrument (dotted line) are practically identical in most of the cases, there are also cases

where they differ significantly (id= 16). The dotted line is estimated from the

measurements for a particular sample (eye) at a particular color in isolation (no pooling),

whereas our estimates take into account all of the data, from all conditions and from all

samples (hierarchical partial pooling). In hierarchical partial pooling, estimates are more

skeptical of measurements that fall outside the group means. For example, the instrument

estimated threshold (dotted line, no pooling) would indicate that id = 16/RE has much

higher threshold for Blue light than for Red light, which is physiologically unlikely as the

response to Blue stimuli includes contributions from both rods and cones, whereas

responses to Red stimuli is solely driven by red cones. The hierarchical partial pooling

estimate is instead largely informed by the overall color effect (β_{clr}), which places Blue at

lower thresholds than Red, and patient and eye biases (β_{pat}, β_{eye}). The large 'guessing' is

also an important factor here, as some of the observations are seemingly random. As these

estimates are not strongly supported by the data, compatibility intervals encompass a wider

range of values. These estimates are nonetheless useful as they narrow the reasonable and

credible range of values. Similarly, using the available information, we are able to make

inferences even if data is entirely missing for a particular color (id = 3 Blue, id = 30 Green

for example), or in cases where no correct responses were obtained within the tested range

(id = 6 LE). Overall, we believe our model is able to capture and evaluate the uncertainty in

measurements, giving us estimates with reliable confidence intervals reflective of the

subject's true ability to detect light.

Analysis and characterization of pupil responses in eyes with advanced RP

Pupillometry measurements were analyzed using a multilevel (hierarchical) multivariate

model as illustrated in Figure 2A. Measurements consists of time series of the pupil

diameter changes to different stimulus (stimulus is applied at t=200 ms). For each

stimulations condition (rod, cone1, cone2, and mela) there are 5 repeats. While repeated

measurements seem to reliably reproduce pupil diameter changes for rod, cone1, and cone2

stimulations, this was not the case for mela stimulation. In most cases, the first repeat

elicited a large reduction in pupil diameter, while repeats 2-4 exhibited a more attenuated

response. We therefore separated the response to mela stimulation to mela1 and mela2 in

order to differentiate between the first and the remaining repeats. Although the data is a

time series, we focused on three key regions: before, peak, and after light stimulation, to

simplify the analysis. We extracted the pupil diameters from these regions for each

measurement and estimated their values. Figure 2B shows the posterior distribution of some

key parameters. We used a multivariate distribution as before, peak, and after pupil diameter

values tended to be highly correlated. We allow every eye and light stimulation condition to

vary while estimating the bias for each subject and the effect of light stimulation. Most of

the variance in the data is explained by patient-to-patient variation ($\beta_{pat} > \beta_{eye}, \beta_{eyeXclr}$),

although substantial LE/RE differences were observed in some cases. Figure 2C shows the

expected typical response ($\beta_0 + \beta_{clr}$). Note that peak is not defined for mela1 and mela2, as

we do not expect, or observe, a sharp peak for these. Rod, cone1, cone2, and mela1

measurements have a baseline (before) of about 5.5 (Rod: 5.6mm, Cone1/Cone2: 5.2) mm.

There is very little or no pupil change to Rod stimulation as is expected in RP patients.

Rod/Cone (cone1, cone2) stimulation induce a small change, with pupils transiently

constricting to about 4.3-4.5 mm, and immediately reverting back to baseline values. On the

other hand, there is a large constriction of the pupil for mela1 (3.2 mm) which only partially

reverts (3.5 mm), resulting in very small pupil diameter change for mela2. Figure 2D shows

the estimated correlation coefficients for the different light stimulation conditions (rod,

cone1, cone2, mela1, and mela2). Before, peak, and after values had a high correlation for

rod, cone1, cone2, and mela2 stimulations. This trend is disrupted for the correlation

between before and after measurements for mela1, which show most of the posterior

distribution with negative correlation values. This indicates that, pupil diameter changes in

mela1 are larger when pupil diameter is larger before the light stimulus.

Figure 2E shows some of the data highlighting typical and noteworthy cases. Model

predictions (posterior predictions = $\beta_0 + \beta_{clr} + \beta_{pat} + \beta_{eye} + \beta_{eyeXclr}$) reasonably describe

the data. Figure 2E id = 30 shows a typical response, with an absent rod response, a small

peak response for cone1 and cone2 stimuli, a large mela1 response (repeat #1) and a very

small mela2 response (repeat#2-5). In most cases the repeats of mela2 were largely

identical, however there were a few cases where melanopsin response seem to gradually

decrease. For example, figure 2E id =7 shows an example where baseline values for mela2

gradually decreased (repeat 1 (mela1) > repeat 2,3 > repeat 4,5). Figure 2E id = 11 shows

an example where cone response varies greatly. Also there appears to be a rapid oscillation

of the pupil following mela2 stimulation. Figure 2E id = 16 shows an example where all

responses were absent except for mela1. Figure 2E id = 39 is similar to id = 16 however

there is a much more robust response for mela1, and there is a large difference between

LE/RE.

FST and Pupil response is largely uncorrelated to VA and retinal thickness

Having analyzed FST and pupil response separately, we next focused our attention on how

these correlate to each other. Figure 3 shows the pair plot for some representative features.

We select sex, age, VA(logMAR), total retinal thickness within macular area (2mm from

central fovea), FST (White), pupil diameter before stimulation, and pupil change to cone1,

mela1, and mela2 stimulation for this analysis. We used the threshold for white light

stimulation for FST measurements as representative data for FST measurements, as

correlation between FST threshold for different stimulus color (Red, Green, White, Blue)

was extremely high ($\rho > 0.9$, data not shown) and using any color would have resulted in

practically identical results. Note the wide distribution of values for total retinal thickness,

FST, and pupil diameters within the same VA group (Figure 3 blue rectangle), indicating

that individuals within the same VA group may exhibit a wide range of visual functions and

retinal thickness. The correlation analysis shows that VA is not significantly correlated to

retinal thickness, or pupil responses, but moderately correlated to FST ((Figure 3 orange

rectangle, $\rho_{FST/VA} = 0.42$). Since central macular cone area is generally considered last to

degenerate in eyes with RP, we initially expected that macular retinal thickness to reasonably

correlate with measures of visual function (FST, or Pupil response). However, we only found

a moderate correlation with pupil response ($Pupil_{cone1}$) in these eyes with low vision.

Retinal thickness also seemed to be correlated to subject's sex, with female subjects

exhibiting a slightly thicker retina.

Figure 4 and 5 are per patient/eye summaries of collected data highlighting some

characteristic cases. As we noted above, FST and pupil response does not correlate with VA

or retinal thickness. The eye in Figure 4 shows that the retinal thickness and structure of the

layer are relatively well preserved, with clearly defined ONL in fovea with visual acuity of

0.01. However, FST was poor and with a rather high guessing rate. We further looked into

the Goldman visual field (VF) test to find that the eye had a very constricted central vision.

In contrast, Figure 5 shows the eye with very thin retina in the macular area with visual

acuity of HM. The eye however exhibited a relatively good FST with a clear shift between

Blue and Red lights indicating rod functionality. Pupil response to rod stimuli was weakly

present, supporting the presence of rod function in addition to cone function, which was

rare in our RP patients. Goldman VF test show the presence of peripheral vision. Together

these results indicate that both cone and rod function were preserved in spite of the very

thin central retina.

Discussion

Birch et al. performed an FST study in inherited retinal disease(IRD)patients with visual

acuity of 0.25 or better and reported that in some cases the threshold difference between

blue and red stimuli was >10 dB.[22] The decrease in the difference between Blue and Red

thresholds is generally thought to reflect the more advanced retinal degeneration status.[6–

8]Although our study cohort consisted of patients with more advanced retinal degeneration

than that of Birch et al., we were nonetheless able to detect differences between Blue and

Red FST (18 eyes (21%) (0.01/CF: 6 eyes, HM: 6 eyes, LP): 5 eyes, NLP: 1 eye).

Furthermore, we noted that eyes that have low threshold values tended to have a larger shift

between Blue and Red light than eyes that have high threshold values (compare for example

Figure1C id=43 and 3, note that higher threshold values indicate lower probability to

respond to the light stimulus). To further understand the relationship between the shift in

the threshold from Blue light, we plotted Green, White, and Red threshold values against

Blue threshold values (Figure 7A). While Green and White values largely mirror Blue

threshold values in all regions, red threshold values show a larger deviation from Blue values

at lower thresholds. This is consistent with the notion that reduced sensitivity to Blue light

represent advanced rod degeneration. Thus, while FST measurements for White or Green

colors are highly correlated, Blue and Red FST values provide important diagnostic

information in late-stage RP. Moreover, our data indicates that threshold values for White,

Green, and Red light can reasonably be predicted from the Blue light stimulus alone, or vice

versa.

OCT measurements, such as retinal thickness are widely and commonly used to diagnose

retinal health, with well-organized thicker retinas generally presumed to correlate with more

visual function. Since central cones are often preserved in RP, we measured retinal thickness

within the macular area. However, we found no correlation between retinal thickness and

indicators of visual function such as VA (logMAR), FST, or pupil response in our cohort of

advanced retinal degeneration with visual acuity of 0.01 or lower (Figure 6). In order to

further explore the relationship between OCT measurements and visual function we
stratified the data into four groups according to subject's visual acuity (0.01/CF, HM, LP,
and NLP). Unexpectedly, we found a reversal of correlation between retinal thickness and
FST values (Figure 6). Patients with thicker retina tended to have higher FST values (lower
sensitivity for light perception) in 0.01/FC and HM whereas this trend is reversed in LP and
NLP. Moreover, while Figure 6 shows the analysis using the average of 4 points for the entire
retinal thickness (2,000 μ m from the fovea), the same trend was observed at each of the
quadrant position individually (temporal, nasal, superior, inferior) and for different layers
(photoreceptor layer, INL, GCL), indicating that this is a global effect possibly affecting the
entire macula. Although human visual function largely depends on central vision, example
cases shown in Figure 4 and 5 suggest that in the eyes with advanced degeneration,
remaining central visual function may be complemented by peripheral vision, possibly
explaining the apparent discrepancy between the retinal thickness around the macula and
overall visual function. It is of note that the FST and pupillometry measurements seemed
sensitive enough to suggest the presence of remaining rod and cone function in the
peripheral vision in the eye with "HM" (Figure 5). These results highlight that, together,
FST, pupillometry and OCT measurements may help understand the pathological status of
retinal degeneration, allowing for a more graded and comprehensive assessment than the
conventional VA categories of FC, HM, LP, and NLP in ultra-low vision patients.
395
With the possibility of clinical application of novel treatments for visual restoration on the
horizon, such as gene therapy, cell-based treatments, optogenetics, and prosthetic devices, a
more detailed assessment of patient's visual potency and the state of the retina is urgently
needed. FST is very sensitive and has been reportedly used to detect subtle changes before
and after treatment in a clinical study of gene therapy for LCA patients[8,23,24]. Indeed, in the
current study, we obtained positive responses (defined as threshold below +15.0dB) from
almost half of the eyes with "no light perception" highlighting the sensitivity of this approach.
In the pupillary test, the two cone stimuli resulted in practically identical results and seem to
reflect the remaining photoreceptor pathway and RGC activities. Positive FST and cone
pupillary tests indicate a functional retinal circuitry even in the eyes with poor central vision.
In these eyes, gene therapy or cell-based approaches may help enhance visual responses. On
the other hand, eyes that only have a pupillary response to melanopsin stimuli, may better
benefit from direct stimulation of RGCs such as artificial eye or optogenetic treatment rather
than cell therapy. Lastly, eyes that completely lack FST and pupil responses may benefit from
cortical interventions (Figure 7).
These emerging therapies for visual restoration bring hope for patients with severe visual

impairments. The current study highlights the importance of developing diagnostic tools for
ultra-low vision patients, bringing these treatments one step closer to clinical application.
414
415
416   Acknowledgments
417   We thank Dr. Yozo Miyake and Dr. Kaoru Fujinami for invaluable advice and all members of
418   the Kobe City Eye Hospital for their ongoing support.
419
Author Contributions
T. Maeda, M.Y and N.M conceived the study and performed the examinations. M.M,T.
Matsuyama and M.Y analyzed and interpreted the examination data. T. Matsuyama
conducted statistically analysis. M.Y, T. Matsuyama , and M.M. wrote the manuscript, which
was approved by all authors prior to submission. A.M,Y.H,S.T, N.M and D.S, provided
helpful guidance and suggestions. Y.K and M.M directed all work.
426
427

References

1. Hartong DT, Berson EL, Dryja TP. Retinitis pigmentosa. Lancet 2006;368:1795–1809.

2. Sahel J-A, Boulanger-Scemama E, Pagot C, et al. Partial recovery of visual function in a

blind patient after optogenetic therapy. Nature Medicine 2021;27:1223–1229.

3. Assawachananont J, Mandai M, Okamoto S, et al. Transplantation of embryonic and

induced pluripotent stem cell-derived 3D retinal sheets into retinal degenerative mice. Stem

Cell Reports 2014;2:662–674.

4. Mandai M, Fujii M, Hashiguchi T, et al. iPSC-Derived Retina Transplants Improve Vision

in rd1 End-Stage Retinal-Degeneration Mice. Stem Cell Reports 2017;8:69–83.

5. kurimoto   yasuo. iPS cell-derived neural retinal sheet transplantation for retinitis

pigmentosa.

6. Roman AJ, Schwartz SB, Aleman TS, et al. Quantifying rod photoreceptor-mediated vision

in retinal degenerations: dark-adapted thresholds as outcome measures. Experimental Eye

Research 2005;80:259–272.

7. Roman AJ, Cideciyan AV, Aleman TS, Jacobson SG. Full-field stimulus testing (FST) to

quantify visual perception in severely blind candidates for treatment trials. Physiol Meas

2007;28:N51-56.

8. Roman AJ, Cideciyan AV, Wu V, et al. Full-field stimulus testing: Role in the clinic and as

an outcome measure in clinical trials of severe childhood retinal disease. Progress in Retinal

and Eye Research 2022;87:101000.

9. Anon. Analysis of retinal function using chromatic pupillography in retinitis pigmentosa

and the relationship to electrically evoked phosphene thresholds - Kelbsch - 2017 - Acta

Ophthalmologica   -   Wiley   Online   Library.   Available   at:

https://onlinelibrary.wiley.com/doi/10.1111/aos.13259 [Accessed March 29, 2022].

10. Park JC, Moura AL, Raza AS, et al. Toward a Clinical Protocol for Assessing Rod, Cone,

and Melanopsin Contributions to the Human Pupil Response. Investigative Ophthalmology

& Visual Science 2011;52:6624–6635.

11. Kardon R, Anderson SC, Damarjian TG, et al. Chromatic Pupillometry in Patients with

Retinitis Pigmentosa. Ophthalmology 2011;118:376–381.

12. Lisowska J, Lisowski L, Kelbsch C, et al. Development of a Chromatic Pupillography

Protocol for the First Gene Therapy Trial in Patients With CNGA3-Linked Achromatopsia.

Investigative Ophthalmology & Visual Science 2017;58:1274–1282.

13. Rukmini AV, Milea D, Gooley JJ. Chromatic Pupillometry Methods for Assessing

Photoreceptor Health in Retinal and Optic Nerve Diseases. Frontiers in Neurology 2019;10.

Available at: https://www.frontiersin.org/article/10.3389/fneur.2019.00076 [Accessed April

26, 2022].

14. Lorenz B, Strohmayr E, Zahn S, et al. Chromatic Pupillography and RPE65 Deficiency.

Investigative Ophthalmology & Visual Science 2011;52:3891.

15. Hattar S, Liao H-W, Takao M, et al. Melanopsin-Containing Retinal Ganglion Cells:

Architecture, Projections, and Intrinsic Photosensitivity. Science 2002;295:1065–1070.

16. Morin LP, Allen CN. The circadian visual system, 2005. Brain Research Reviews

2006;51:1–60.

17. Güler AD, Ecker JL, Lall GS, et al. Melanopsin cells are the principal conduits for rod–

cone input to non-image-forming vision. Nature 2008;453:102–105.

18. Klein DC, Moore RY, Reppert SM, Klein CS on NNI of CH and HDDC. Suprachiasmatic

Nucleus: The Mind's Clock. Oxford University Press; 1991.

19. Czeisler CA, Shanahan TL, Klerman EB, et al. Suppression of Melatonin Secretion in

Some Blind Patients by Exposure to Bright Light. New England Journal of Medicine

1995;332:6–11.

20. Morin LP. The circadian visual system. Brain Research Reviews 1994;19:102–127.

21. Ln J, Me G, Gb K, Rw M. Levodopa may improve vision loss in recent-onset, nonarteritic

anterior ischemic optic neuropathy. Ophthalmology 2000;107. Available at:

https://pubmed.ncbi.nlm.nih.gov/10711891/ [Accessed April 4, 2022].

22. Birch DG, Cheng P, Duncan JL, et al. The RUSH2A Study: Best-Corrected Visual Acuity,

Full-Field Electroretinography Amplitudes, and Full-Field Stimulus Thresholds at Baseline.

Translational Vision Science & Technology 2020;9:9.

23. Russell S, Bennett J, Wellman JA, et al. Efficacy and safety of voretigene neparvovec

(AAV2-hRPE65v2) in patients with RPE65-mediated inherited retinal dystrophy: a

randomised, controlled, open-label, phase 3 trial. Lancet 2017;390:849–860.

24. Jacobson SG, Cideciyan AV, Ratnakaram R, et al. Gene Therapy for Leber Congenital

Amaurosis caused by RPE65 mutations: Safety and Efficacy in Fifteen Children and Adults

Followed up to Three Years. Arch Ophthalmol 2012;130:9–24.

Figure legends

Fig. 1

Analysis of FST

A. FST Model. Patient's responses y_i (correct or wrong) to a light stimulus were analyzed

with a logistic regression model. In a typical FST experiment, the intensity of the light

stimulus (x_i, measured in dB) is gradually decreased as the subject pushes a bottom if they

can perceive the light. We used multilevel (hierarchical) modeling, to estimate the effect of

different light stimulus color (β_{clr}), patient (β_{pat}), and patient eye (β_{eye}) on the probability

of correctly detecting the light stimulus. In order to allow higher freedom for estimates of

each eye to vary at different light color we added an interaction term for patient eye and

light color ($\beta_{eyeXclr}$). Each parameter was constrained so that $\Sigma\beta_k = 0$ (sum to zero

constrain) and the effect of each parameter is shown as deviation from the overall mean

(β_0). The probability of seeing the light stimulus is given by the logistic model and a

'guessing' parameter (α), which is estimated for each eye. This parameter accommodates

data points that do not conform well with the logistic regression.

B. Posterior estimates of FST model. Estimates for patient (β_{pat}) vary widely whereas β_{eye}

and $\beta_{eyeXclr}$ variation is relatively smaller. Most of the patients have similar estimates for

LE and RE although some patients seem to have a significant LE/RE difference. There was

a clear effect for the light color, with subjects being more sensitive to blue light than red

light (Red < Green, White < Blue). Although most of the subjects had a relatively small

'guessing' estimate (α) with the mode and 95% interval below 0.1 (=10% guessing), there

were notable exceptions with very large guessing values (mode > 0.2).

C. Examples of raw data and model estimates. Points show the raw data for a series of

experiments for a patient. Note that points are shown slightly offset from their true values (0

or 1) to avoid RE and LE points overlapping. Model estimates are shown with a solid curved

line (mean) and ribbon (95% intervals). 'Guessing' estimates are shown as violin plots on

top of the FST results. The vertical lines show the FST threshold values (dB at which

probability of success is 0.5). The solid line represents the mean threshold value of our

estimates, whereas the dotted line shows the threshold estimated by the measurement

apparatus (threshold estimated individually). Color indicates LE (orange) and RE (blue).

Fig. 2

Analysis of chromatic pupillometry

A. Pupillometry model. Before, peak, and after were defined as the pupil diameter values

before light stimulation (0 – 200 ms), peak value following light stimulation (200-2000 ms,

defined only for rod, cone1, and cone2 stimuli), and value of the relatively flat region

following light stimulation (3000 – 6000 ms) respectively. These pupil diameter

measurements (x_{before}, x_{peak}, x_{after}) were modeled as a multivariate distribution, taking

into account their inter-correlation. We used the t-distribution for robustness since some

values may have been influenced by measurement artifacts (see Methods for details). Note

that de degrees of freedom ν is also a parameter that is estimated from the data. Similar to

the FST analysis, we incorporated patient (β_{pat}), eye (β_{eye}), color (β_{clr}), and an interaction

term for eye and color ($\beta_{eyeXclr}$) to allow estimation of overall trends while allowing for

flexibility to account specific trends for each eye/color combination if there is strong

evidence. Parameters were constrained so that $\Sigma\beta_k = 0$ (sum to zero constrain) and the

effect of each parameter is shown as deviation from the overall mean (β_0). The correlation

between measurements was estimated for each stimulation condition (ρ_{clr}). B. Posterior

estimates of PM model. Estimates for effect off *patient, eye, eyeXclr* are shown per patient

(rows). Note that for each patient id there are three estimates for before, peak, and after

(shown as different shapes). C. Posterior distribution of predicted typical response ($\beta_0 +$

β_{clr}). D. Posterior estimates of correlation coefficient for each light stimulus. E. Examples of

raw data and model estimates. Light blue traces show the individual pupillometry

measurements. Purple dots overlayed on the blue traces (id =16 and 39) show data points

that were excluded from the analysis, as they consist of regions with constant values (sd = 0)

and therefor likely represent regions where pupil detection failed. Rod, cone1, and cone2

measurements consist of 5 traces (repeats), whereas the 5 traces of mela light stimulus were

separated to mela1 (repeat 1) and mela2 (repeats 2-5), as we noticed that there was a large

difference between mela1 and mela2. The black and gray shaded area show the estimated

mean and 95% confidence interval for before, peak, and after pupil diameter.

Fig. 3

Pairwise relationship of dataset. We plotted (from left to right and top to bottom) sex, age,

VA(logMAR), total retinal thickness (μm), FST (db, white light stimulation), pupil diameter

before light stimulation ($Pupil_{before}$, mm), pupil change to cone1 stimuli ($Pupil_{cone1}$, mm),

pupil change to mela1 stimuli ($Pupil_{mela1}, mm$), pupil change to mela2 stimuli ($Pupil_{mela2}$,

mm). The diagonal show distribution of each variable (bar chart for categorical data and

histogram for continuous data). Lower triangular panels show the pairwise relationship of

features. The blue rectangle highlights the relationship of VA(logMAR) to other

measurements. Upper triangular panels show the estimated correlation coefficient,

estimated from a multivariate model, with density plots showing the distribution of

estimated correlation coefficients. Red indicates negative correlation ($\rho = -1$) and blue

indicates positive correlation ($\rho = 1$). The mean (black point) and 95% confidence interval

(black bar) are indicated on the bottom of the density plot. Numbers on top indicate the

mean, and the dotted line denotes the zero line. The orange rectangle highlights the

correlation coefficients of features with VA.

Fig. 4

Representative case of an eye with a thick retina and poor FST. Vertical and horizontal OCT

scans show ONL of the fovea is preserved and the structure of retina is appears to be well

preserved. The average retinal thickness is near the macula was 242µm, which is equivalent

to that of normal subject. The eye had been diagnosed as HM (center of middle panel). All

of the pupillay responses (right of lower panel) except for rod response were clearly present.

On the other hand, FST (right of middle panel) was relatively poor with threshold values

over 0dB. The result of Goldman VF test (left of lower panel) shows a very constricted VF

remained in the center.

Fig.5

Representative case with thin retina but good visual acuity and FST sensitivity. The average

of retinal thickness was 184µm and OCT images (upper panel) reveal photoreceptor cell

loss and lack of fovea. VA (center of middle panel) was CF, the highest category of this

study, and the FST sensitivities were similar to that of normal eyes. The response to

chromatic pupillometry (left of lower panel) were also good (even rod response were

confirmed which was rare in our study cohort) and GoldmanVF test (right of lower panel)

shows a relatively large residual field of view in the periphery.

Fig. 6

A. Relationship between FST values for Blue, White, Green and Red light. The Blue FST

value was plotted against White, Green, and Red FST values. The dotted diagonal line has

intercept = 0 and slope = 1. Points above the dotted line indicate higher FST value than

Blue light and points below the dotted line indicate lower FST. The solid line shows the

mean and the shaded area shows the 95% compatibility interval of quadratic regression for

each of the colors. B. Relationship between retinal thickness and FST (white), stratified by

four visual acuity groups. The upper left panel shows the distribution of retinal thickness

and the lower right panel shows the distribution of FST for white light. The lower left panel

show the distribution of the data (points) overlayed with estimated correlation (density

plot), with horizontal error bars showing the SD of retinal thickness measurement (average

of four points), and vertical error bars showing the estimated error (SD) from the FST

analysis. The upper right panel shows the distribution of posterior estimates for the

correlation between FST and retinal thickness (error bars show 95% confidence interval).

Fig. 7

FST and pupillometry can be used, complementary to VA and OCT observations, to more

precisely estimate pathological stage. As photoreceptors degenerate FST threshold

increases. We use Red – Blue FST > 10bB as a reference point, indicating advanced rod loss.

According to our analysis in figure 6A, this is equivalent to a blue FST < -10dB. The loss of

photoreceptor cells in central macula as well as peripheral area leads to >15dB FST test

results and the loss of pupillary cone response. Further degeneration and loss of retinal

inner cells results in the decrease in pupillary melanopsin responses.

Table 1

Clinical characteristic of the patients

A

$$\mu_i = \beta_0 + \beta_{pat,i} + \beta_{eye,i} + \beta_{clr,i} + \beta_{eyeXclr,i} + \beta x_i$$

$$\Sigma\beta_k = 0 \quad k = \{pat, eye, clr, eyeXclr\}$$

$$\theta_i = \alpha_{eye,i}\frac{1}{2} + (1 - \alpha_{eye,i})logistic(\mu_i)$$

'guessing'   logistic regression

$$y_i \sim Bernoulli(\theta_i)$$
$$y_i = \begin{cases} 1 & \text{if correct} \\ 0 & \text{if wrong} \end{cases}$$

B **Estimate of effect of patient(β_{pat}), eye(β_{eye}), eyeXclr($\beta_{eyeXclr}$)**

points: mode, lines: 95% intervals

eye ● LE ● RE

Estimate of color effect (β_{clr})
points: mode, lines: 95% intervals

Red −1.9 **−1.6** −1.3
White −0.2 **0.1** 0.3
Green −0.5 **−0.3** 0
Blue 1.5 **1.8** 2.1

Estimate of 'guessing' (α_{eye})
points: mode, lines: 95% intervals

C **FST data posterior predictive per patient (raw data vs model predictions)**

id = 4   eye LE RE
id = 43   eye LE RE
id = 28   eye LE RE
id = 30   eye LE RE

id = 1   eye LE RE
id = 6   eye LE RE
id = 3   eye LE RE
id = 16   eye LE RE

A

$$\vec{\mu}_i = \vec{\beta_0} + \vec{\beta_{pat,i}} + \vec{\beta_{eye,i}} +$$
$$\vec{\beta_{clr,i}} + \vec{\beta_{eyeXclr,i}}$$

$$\Sigma_{color} = diag(\vec{\sigma}) \times \Omega_{clr} \times diag(\vec{\sigma})$$

$$\Sigma\beta_k = 0$$
$$k = pat, eye, clr, eyeXclr$$

$$\vec{x_i} = t_\nu\left(\vec{\mu}_i, \Sigma_{clr}\right)$$

$$\vec{\mu}_i = \left[\mu_{before,i}, \mu_{peak,i}, \mu_{after,i}\right]$$
$$\vec{\sigma} = \left[\sigma_{before}, \sigma_{peak}, \sigma_{after}\right]$$
$$\vec{x_i} = \left[x_{before,i}, x_{preak,i}, x_{after,i}\right]$$

B Estimated effect of predictors

patient(β_pat), eye(β_eye), eyeXstim(β_eyeXstim)
points: mode, lines: 95% intervals, oragne:LE, blue:RE

before ● peak ▲ after ■

β_{pat} β_{eye} $\beta_{eyeXrod}$ $\beta_{eye,cone1}$ $\beta_{eye,cone2}$ $\beta_{eye,mela1}$ $\beta_{eye,mela2}$

patient id

pupil diameter (mm)

C Estimated typical pupil diameter response (β_0+β_clr)

before peak after

pupil diameter (mm)

D correlation (Ω_clr)

after/before after/peak peak/before

correlation

E Pupillometry data posterior predictive check

id = 30

before — peak — after

rod cone1 cone2 mela1 mela2

pupul diameter (mm)

LE

RE

time (ms)

id = 7

id = 11

id = 16

id = 39

OCT

**OCT
(macula)**

VA

**FST
(guessing)**

FST

thickness (µm)

count

'guessing' rate

probability of success

visual acuity: CF, HM, SL, NLP

dB

Blue, White, Green, Red

superior
temporal
nasal
inferior

photo | inl | gc

Chromatic Pupillometry

rod   cone1   cone2   mela1   mela2

pupul diameter (mm)

time

A

Blue vs Green|White|Red FST

B

Pair plot: FST and retinal thickness per VA group

Retina Pathology

	Progressive Loss of Photoreceptors	Complete Loss of Photoreceptors	Loss of Inner Retinal cells	Complete loss of retina function

Visual Acuity — low vision   **VA > 0.01**   ultra-low vision   → No light perception

FST — Red-Blue > 10dB ~ FST(Blue) <-10 dB   → FST > 15dB

Pupil Response — Decreased Rod Response   No Cone response   Decreased Mela response   No Mela2 response   → No pupil response

Table 1 Clinical charactaristic of the patients

	84 eyes of 43 patients
Patient age (yr)	59.3 ± 9.96
Sex (female:male)	17 : 26
BCVA	
0.01	13 (15.5%)
CF	1 (1.2%)
HM	23 (27.4%)
LP	40 (47.6%)
NLP	7 (8.3%)

Values are presented as number (%) or mean ± standard deviation.