

Health services for Aboriginal and Torres Strait Islander children in remote Australia: a scoping review

PJ Dossetor^{1*}, JM Freeman^{2*}, K Thorburn³, J Oscar⁴, M Carter⁵, HE Jeffery², D Harley^{1,6}, EJ Elliott^{2,7}, ALC Martiniuk^{2, 8, 9&}

¹ Clinical Medical School, College of Medicine, Biology & Environment, Australian National University, Canberra, ACT, Australia

² University of Sydney, Faculty of Medicine and Health, Sydney, Australia

³ Nulungu Research Institute, University of Notre Dame, Broome, Australia

⁴ Marninwarntikura Women's Resource Centre, Fitzroy Crossing, Australia

⁵ Nindilingarri Cultural Health Services, Fitzroy Crossing, Australia

⁶ Queensland Centre for Intellectual and Developmental Disability, Mater Research Institute-UQ, The University of Queensland, Brisbane, QLD, Australia

⁷ The Sydney Children's Hospital Network (Westmead), Kids Research, Westmead, Australia

⁸ Dalla Lana School of Public Health, University of Toronto, Toronto, Canada

⁹ George Institute for Global Health, Sydney, Australia

& Corresponding Author

E-mail: Alexandra.Martiniuk@sydney.edu.au

* These authors contributed equally to this work

Abstract

In Australia, there is a significant gap between health outcomes in Indigenous and non-Indigenous children, which may relate to inequity in health service provision, particularly in remote areas. The aim was to conduct a scoping review to identify publications in the academic and grey literature and describe 1) Existing health services for Indigenous children in remote Australia and service use, 2) Workforce challenges in remote settings, 3) Characteristics of an effective health service, and 4) Models of care and solutions. Electronic databases of medical/health literature were searched (Jan 1990 to May 2021). Grey literature was identified through investigation of websites, including of local, state and national health departments. Identified papers (n=1775) were screened and duplicates removed. Information was extracted and summarised from 116 papers that met review inclusion criteria (70 from electronic medical databases and 45 from the grey literature). This review identified that existing services struggle to meet demand. Barriers to effective child health service delivery in remote Australia include availability of trained staff, limited services, and difficult access. Aboriginal and Community Controlled Health Organisations are effective and should receive increased support including increased training and remuneration for Aboriginal Health Workers. Continuous quality assessment of existing and future programs will improve quality; as will measures that reflect aboriginal ways of knowing and being, that go beyond traditional Key Performance Indicators. Gold standard models for service delivery have community leadership and collaboration. Increased resources with a focus on primary prevention and health promotion are essential.

“The failure to close the gaps in Aboriginal and Torres Strait Islander health inequality, and other measures of social and economic disadvantage, cannot be justified by more rhetoric or data in another report. For us, the harrowing failure to close the gap is felt through sorry business, the countless funerals of family and friends, the hospital visits and the coronial inquiries that we continue to painfully endure. So many of our losses were and are preventable – that is the failure and pain we carry. A sensible way of doing business is long overdue as, apart from small gains, the attempts to close the gaps in Aboriginal and Torres Strait Islander life expectancy, health and education have failed.” June Oscar [1]

1 **Introduction**

2 The estimated 727,500 Aboriginal and Torres Strait Islander people, herein called Indigenous
3 Australians, comprise 3.3% of the total Australian population [2]. One fifth of Indigenous Australians
4 live in remote and very remote settings [2]. Indigenous children living in remote Australia experience
5 a greater burden of disease than children living in metropolitan settings [3-9]. Colonisation, forcible
6 separation and forced assimilation, violence and loss of culture and land, underlying intergenerational
7 trauma, discrimination, exclusion and racism which deeply impact health and the social determinants
8 of health including education, employment and housing, and access to health care.

9
10 Indicators of child health and life expectancy include the frequency of low birth weight (< 2500g) and
11 infant mortality [7, 10, 11]. In 2010, Indigenous babies were twice as likely (12%) as non-Indigenous
12 babies (6%) to have a low birth weight (LBW). The Indigenous infant mortality rate is 1.8 times
13 higher for non-Indigenous than Indigenous infants (5.1 versus 2.9 infant deaths per 1000 live births,
14 respectively) [12].

15
16 Indigenous Australians are hospitalised at 2.6 times the rate of non-Indigenous Australians [13]. Rates
17 of hospitalisation and emergency department presentation for Indigenous children living in remote
18 areas are high, often for potentially preventable skin, respiratory and gastrointestinal infections [8, 14,
19 15]. One in five Aboriginal and Torres Strait Islander children live with disability (22%), including
20 sensory (12%), cognitive (9%), physical (5%) and psychosocial disability (4%) [16].

21
22 Health is a human right so Indigenous children living in remote areas need priority access to high
23 quality health care [17, 18] to improve current health, decrease the risk of chronic disease and
24 increase life expectancy [19]. Social determinants contribute powerfully to Indigenous health
25 inequality and include poor nutrition, housing shortage, limited primary health care access, low
26 income and limited educational attainment [20, 21].

27
28 In response to the 2009 ‘Closing the Gap’ strategy each Australian State and Territory developed
29 plans and policies to address gaps between Indigenous and non-Indigenous health indicators and

30 improve the effectiveness of health service delivery to children in remote settings [22, 23]. For
31 example, the Queensland Government aimed to close the health gaps by 2033 by addressing risk
32 factors, increasing primary prevention strategies and access to multidisciplinary health services, and
33 providing nurturing safe environments for children [22].

34

35 The 2021 ‘Close the Gap’ report found mixed results [24]. There was a 17% reduction in avoidable
36 deaths between 2006 and 2018 [24], however this rate of decline is slowing. Indigenous Australians
37 currently die from avoidable causes at three times the rate of non-Indigenous Australians [24].
38 Mortality rates for Australian Indigenous children have not changed since 2005. While infant
39 mortality has declined, child injury deaths have remained the same. The number of Indigenous
40 children in out of home care has increased since 2013 [25].

41

42 This scoping review aimed to identify publications in the academic and grey literature and describe 1)
43 Existing health services and service use for Indigenous children in remote Australia, 2) Workforce
44 challenges in remote settings, 3) Characteristics of an effective health service, and 4) Effective models
45 of care.

46

47 **Methods**

48 ***Rationale for a scoping review***

49 To address our aims, we employed a scoping review methodology. This is because we sought to find
50 and assess the scope of the literature on the topic; as well as to describe and synthesise existing
51 evidence, sources, locations, breadth, definitions, and knowledge gaps on this topic.

52

53 ***Ethical Approval***

54 This study being a review, used only previously published and publicly accessible data. Review by an
55 ethical board or consent from participants was therefore not appropriate and not sought.

56

57 ***Search of Electronic Databases***

58 Electronic databases (MEDLINE, Cumulative Index for Nursing and Allied Health Literature
59 (CINAHL), Psychological Information Abstracts Services (PsycINFO), Web of Knowledge, Excerpta
60 Medica Database (EMBASE), Educational Resources Information Centre (ERIC), and Scopus) were
61 searched for relevant publications.

62

63 ***MeSH headings and key words***

64 The MeSH Headings used were health services; community health services; primary health care;
65 family practice; child health services; adolescent medicine; health services, Indigenous; rural health;
66 rural health services; rural population; rural; child; paediatrics; Oceanic ancestry group; and Australia.

67 The key words searched were health services; community health services; primary health care; family
68 practice; adolescent medicine; Indigenous health; Indigenous health services; rural; remote; rural
69 health services; child; infant; paediatrics; paediatrics; Indigen*; Aborigin*; Oceanic ancestry group;
70 allied health; Australia. The MEDLINE strategy is shown in Box 1 and was adapted to all other
71 databases.
72

73 **Box 1**

MEDLINE
1990 until May 28, 2021

health services.mp. or Health Services/ OR
community health services.mp. or Community Health Services/ OR
primary health care.mp. or Primary Health Care/ OR
family practice.mp. or Family Practice/
AND
Adolescent Medicine/ or Adolescent/ or Adolescent Health Services/ OR
Child Health Services/ or child*.mp. or Child/ OR
pediatrics.mp. or exp Pediatrics/
AND
rural.mp. or Rural Health/ or Hospitals, Rural/ or Rural Population/ or Rural Health Services/ OR
remote.mp.
AND
oceanic ancestry group.mp. or Oceanic Ancestry Group/ OR
indigen*.mp. OR
aborigin*.mp.
AND
health services indigenous.mp. or Health Services, Indigenous/ OR
australia.mp. or Australia/

limited to
yr = 1990 until May 28, 2021
Region = Australia

74

75

76 ***Websites and Databases for Grey Literature search***

77 The following websites and databases were searched: Informit Indigenous Collection (IIC); Lowitja
78 Institute, Australian Institute of Aboriginal and Torres Strait Islander Studies (AIATSIS) and
79 Australian Aboriginal Health Info-net; Websites of the Federal and State Governments Health
80 departments and associated agencies; Parliamentary Hearings and Senate committees, Commonwealth
81 and State; Aboriginal medical services; regional health services; Australian Human Rights
82 Commission; National Health and Medical Research Council (NHMRC) of Australia; Royal
83 Australasian College of Physicians; Royal College of Paediatrics and Child Health (UK); Australian
84 Bureau of Statistics, Australian Institute of Health and Welfare (AIHW), Research Institutes focussing
85 on Indigenous health e.g. Menzies, Centre for Aboriginal Economic Policy Research at Australian
86 National University, Telethon Institute for Child Health Research (Western Australia), Centre for
87 Remote Health, and Australian Primary Health Care Research Institute (APHCRI).

88

89 ***Inclusion criteria***

90 Medical publications and grey literature published January 1990-May 2021 describing existing rural
91 and remote health services, health service use, or needs in relation to Indigenous Australian children

92 (0-18 years) were included. Grey literature was only included if it provided original data or insights
93 on health service delivery to Indigenous children living in remote areas.

94

95 ***Exclusion Criteria***

96 Papers were excluded if they were not published in English, were outside the review time frame, or
97 focused on: health services in foreign countries or large metropolitan areas; adult, dental, oral, or
98 sexual health; education of children or health professionals; social or cultural primary outcomes;
99 attraction and retention of health professionals; substance abuse and misuse; childcare centres; the
100 judicial system; or health policy.

101

102 ***Article Selection and Review Process***

103 Two authors reviewed the title, abstract and, when relevant, full text of all publications identified for
104 eligibility. A third author resolved disagreements. Reference lists of publications were reviewed for
105 additional relevant citations. Data from all included papers were extracted and summarised (Appendix
106 Tables 1a and 1b). The Preferred Reporting Items for Systematic reviews and Meta-Analyses
107 (PRISMA) extension for scoping reviews was followed during the process of researching and writing
108 this review paper; with attention to the 20 recommended reporting items [26].

109

110 **Definitions**

111 ***Health Systems and Services***

112 ‘Health services’ were defined as any primary, secondary, or tertiary child health services, including
113 paediatric specialists, remote nursing clinics, allied health professionals, hospital inpatient and
114 emergency departments, patient retrieval through the Royal Flying Doctor Service, fly-in-fly-out
115 (FIFO) services, paediatric outreach services, multidisciplinary teams, public health, tele-paediatrics
116 and videoconferencing or ‘telehealth’ systems.

117

118 We used the World Health Organization (WHO) definition of ‘health systems’ to include all
119 organisations, people, and actions whose primary intent is to promote health [27].

120

121 ***Rural and Remote Locations***

122 The classification scale used to define rural and remote varies in studies, however the Rural, Remote
123 and Metropolitan Areas classification (RRMA) and the Accessibility Remoteness Index of Australia
124 (ARIA) scales were commonly used [28, 29].

125

126

127

128

129 **Results**

130 **Included publications**

131 We identified 1775 publications in seven databases, removed 406 duplicates, and reviewed 1369
132 abstracts. Of these, 1152 (84%) were excluded. Following full text review of the remaining 217
133 papers, 70 (32%) were included. In addition, 45 reports from the grey literature satisfied inclusion
134 criteria (Figure 1).

135 *Please place Figure 1 about here: PRISMA flowchart of peer-reviewed and grey literature*

136
137 Results of this review are presented under subheadings to address the stated aims of the review.

138

139 **1) Existing health services for Indigenous children in remote Australia and service use**

140 ***Children's Use of Services***

141 Indigenous adults are more likely to use health services than non-Indigenous adults [30, 31]. In
142 contrast, Indigenous children are less likely than non-Indigenous children to have used a health
143 service in the previous 12 months (mean 2.5 v 3.1, $p < 0.001$), or use maternal and child health services
144 (OR=0.35, 95% CI: 0.24-0.49), general practitioners (OR=0.45, 95% CI: 0.35-0.64) or paediatricians
145 (OR=0.52, 95% CI: 0.35-0.77) [32]. However, Indigenous children are more likely to be hospitalised
146 than non-Indigenous children (17 versus 9.9%, $p = 0.01$) [32].

147

148 Social structures and health outcomes interact [32-34]. Proportionally, more Indigenous than non-
149 Indigenous infants live remotely or very remotely, but it is not known whether health outcomes are
150 worse for these Indigenous infants because of lack of access to services or the impact of lower levels
151 of parent education, employment and private health insurance, younger maternal age, and increased
152 rates of single parenting [32]. The causal relationships between health service use and outcomes
153 cannot be assessed using cross-sectional data and some datasets do not incorporate Aboriginal-
154 specific health services [35].

155

156 When Indigenous children attend health services, the service must have culturally appropriate,
157 consistent assessment and reporting [36]. For instance, one medical record audit that aimed to
158 understand the quality of developmental monitoring in a remote Australian Aboriginal health service
159 identified that only 60-80% of attending children (N=151) received a developmental check [37].

160 Recommendations included implementing systems-wide, formal assessment and recording of
161 development in children, improved training for staff undertaking assessments, and use of culturally
162 appropriate assessment tools [37].

163

164 ***Challenges to Provision of remote health services***

165 Health professionals who work in remote communities understand the barriers to provision of health

166 services, but limited evidence is published [38-40]. Identified challenges to delivering effective and
167 equitable health services to Indigenous children in remote areas include:

- 168 • poor access (distance between services and communities, dirt roads, lack of private transport,
169 limited patient accommodation, lack of outreach services) [22, 41, 42]
- 170 • poor communication and infrastructure (lack of internet/telephone access, public transport,
171 sewerage, water supply, electricity) [43, 44]
- 172 • lack of skilled health professionals (difficulty in recruitment and retention and high staff
173 turnover, limited accommodation) [45-47]
- 174 • limited Aboriginal and Torres Strait Islander health workforce (lack of cultural competency) [48,
175 49]
- 176 • environmental factors (high temperatures, flooding, risks associated with travel) [50]
- 177 • health service factors (inflexible health service structures, poor communication and coordination
178 between services, resource constraints e.g., inadequate staff/equipment) [51-54]
- 179 • economic factors (increased cost of transport, food, accommodation, high unemployment) [55-
180 57]
- 181 • issues of cultural safety (lack of interpreters, poor clinician-patient communication; lack of
182 Indigenous workforce; different perceptions of health, illness and medicine; historical association
183 of hospitals with death and mistreatment of Indigenous people and forced removal of children)
184 [58, 59]
- 185 • failure of health services to engage and codesign with consumers including young people [60,
186 61]

187

188 *Access to Effective Health Services*

189 Accessing health services for remote-living families often means travelling long distances to reach
190 health care, relying on the Royal Flying Doctors Service, or waiting for outreach health services to
191 visit remote communities [62-64]. For instance, families must travel long distances to access allied
192 health services for children in remote settings, notably paediatric speech pathologists [65, 66]. Access
193 to speech therapy is particularly important because of high rates of otitis media, hearing problems and
194 poor language skills in remote Indigenous children [67, 68], which impact development, education
195 and intergenerational transfer of cultural knowledge. In rural NSW and Victoria, over 30% of
196 residents live beyond the ‘critical maximum distance’ of 50km from an allied health professional,
197 beyond which patients are less likely to travel. As a result, most (98.6%) allied health treatments in
198 these regions are not delivered at the ideal frequency of at least one session per week [65].

199

200 Extremely isolated communities often depend on the Royal Flying Doctor Service (RFDS) for basic
201 primary health care services and emergency responses [69, 70]. Remote communities rely on

202 aeromedical transport to bring in health professionals and evacuate children for medical care [71, 72].
203 Between 2003-2005, 6.5% of people in a very remote east Arnhem Land community were evacuated
204 by air, with one evacuation every 2.2 days. Evacuation rates were higher during the monsoon; 47%
205 occurred after hours; and the median wait time for the plane was 3 hours (1-21 hours). Respiratory
206 disease (21%), gastroenteritis (14%) and injury or poisoning (11%) accounted for 46% of all
207 aeromedical evacuations [71]. Children (37.7%) were over-represented in evacuations [71]. Reliance
208 on the RFDS highlights the need for support and funding for local staff and services to meet
209 prevention and primary health care (PHC) demands [71]. Many people transported by the RFDS
210 would also benefit from access to multidisciplinary care (53/78, 68%) or shared specialist care (41/78,
211 53%) provided locally [73].

212
213 Multiple examples show that outreach services improve access to healthcare and health outcomes in
214 remote settings [74-76]. In one study, interviews with health practitioners, outreach specialists,
215 regional health administrators and patients in remote Northern Territory (NT) communities, found
216 that outreach specialist services increased gynaecological consultations 4-fold over a three-year
217 period (1996-1999) [77]. Equivalent increases in consultations were not observed for specialties
218 without outreach services [39]. Other studies show that specialist outreach services increased access
219 to elective and urgent surgery [40] and are cost-effective [78, 79].

220
221 Andersen's health care utilisation model incorporates the dependence of health services on multiple
222 factors such as service availability (including out of hours), travel/transportation, costs, language and
223 cultural barriers [38, 39]. Although we know that access to health services is challenging for people
224 living in remote communities, we don't have standardised measures suitable for documenting or
225 comparing access to quality health services in such communities [80, 81]. Furthermore, the validity of
226 existing indicators of access to care, such as the Ambulatory Care Sensitive Conditions (SCSH) score,
227 is debated for remote areas [82].

228
229 Other measures of health service availability that assist our understanding of health service access
230 may not be relevant in remote settings. For example, although the population-based distribution of
231 intensive care units (ICU) correlated with population distribution in one study, accessibility varied
232 geographically; the median distance to the nearest ICU was 161.7km in Western Australia and 7.6km
233 in the ACT [83]. However, there are few data on the population-based availability of health services
234 or the relationship between access to primary health care services and health outcomes in regional and
235 remote Australia [4, 84, 85].

236
237 Even if appropriate indicators do become available to measure access to health services and ensure
238 their quality, it would be difficult to compare services across Australian States and Territories because

239 each jurisdiction manages its own health budget and policies [86]. It is also difficult to compare health
240 system performance within jurisdictions, apart from reviewing health indicators monitored by the
241 Coalition of Australian Governments (CoAG), e.g., Close the Gap targets. Life expectancy, mortality
242 rates, morbidity from chronic diseases and lifestyle factors (obesity, smoking, poor nutrition, physical
243 activity) are used as measures of population health but are less relevant to children [87]. Also,
244 Indigenous child health outcomes do not solely reflect health system functioning, but are impacted by
245 social and environmental factors, case-mix and public health campaigns [88-90].

246

247 ***Social and cultural factors impact service use***

248 Social and cultural factors influence service use by Indigenous people [52, 91]. Some health
249 professionals have ethnocentric attitudes and lack understanding of Indigenous culture and this deters
250 service use [38, 92]. Disrespectful or inappropriate communication and racism also impact
251 engagement [38, 93, 94]. Employment of Indigenous people in health-related occupations increased
252 from 96 to 173 per 10,000 between 1996 and 2016 [95]. Although 54% of the full-time equivalent
253 workforce in Commonwealth-funded Indigenous primary healthcare are Indigenous people, they are
254 three times less likely than non-Indigenous people to be part of the national health workforce [95].
255 Appropriate training in cultural competence of non-Indigenous staff and employment of more
256 Indigenous staff would increase availability of culturally appropriate health services [38].

257

258 ***Funding Complexity***

259 Local, State, and national government, privately funded organisations and NGOs all fund remote
260 health services. One specific national program to increase access to specialist medical care in remote
261 Australia is the Medical Specialist Outreach Assistance Program (MSOAP) which was established in
262 2000 [96]. Access to the funding requires a competitive application process [96]. In some jurisdictions
263 including the NT this funding is accessed directly from the Australian Department of Health, whereas
264 in Queensland (*Check-up*) and WA (*Rural Health West*), non-government entities receive the
265 MSOAP funding and report back to the Australian Government [96]. It is the responsibility of the
266 MSOAP to ensure that the services they fund are linked with existing services.

267

268 Overall expenditure for Indigenous health services and per capita expenditure on public hospital
269 services is higher for Indigenous than non-Indigenous Australians [95]. In 2015-16 the average
270 expenditure per Indigenous Australian was approximately \$8,494 (130% the amount for non-
271 Indigenous Australians \$6,657), with almost half (\$4,436) of this expenditure going to hospital
272 services [95]. In Western Australia from mid-2015 to mid-2017 preventable hospitalisations were 3.8
273 times higher in Indigenous Australians (91 per 1,000 compared to 24 per 1,000 non-Indigenous
274 people in age-standardised groups) [95].

275

276 **2) Workforce challenges in remote settings**

277 ***High Pressure on Workforce***

278 Rural and remote areas often struggle to retain staff [97]. Aboriginal Medical Services (AMS) and
279 specialist services in rural and remote Australia report increased workforce stress, compared with
280 metropolitan services [67, 98]. In rural and remote settings there is a high demand for local services,
281 but physicians report fewer child health services in rural/remote compared with urban settings for
282 instance for audiology (11.1% versus 0%), ENT (33.3% versus 3.9%) and hearing aid provision
283 (37.7% versus 1.9%) [67]. Children in rural and remote settings experience longer wait times for
284 audiologists than urban children (18.3% versus 1.9% waited over the recommended 3 months) [67]. A
285 review of child health services in a remote WA town with a population of approximately 3500
286 identified monthly outreach visits from a general paediatrician and three hospital-based medical
287 officers, only one of whom lived permanently in the town [99].

288

289 **Steps to alleviate pressure on rural health workforce**

290 Attracting and retaining skilled clinicians is crucial to decreasing workforce pressures [69, 99-101].
291 There are numerous workforce initiatives to help alleviate pressures on the rural health workforce,
292 many of which rely on encouraging doctors to work in rural settings [102-105]. Up-skilling of
293 paediatric nurses and Aboriginal Health Workers aims to alleviate workforce shortages [106].
294 Many children in remote communities have complex, chronic health problems and would benefit from
295 multidisciplinary teams that enable coordination and communication between clinicians, address
296 service delivery gaps and prevent duplication [102-105]. Regular meetings that facilitate
297 communication between health professionals, disciplines and organisations are recommended to
298 establish clear roles and responsibilities and decrease workforce stress [107, 108]. Employment of
299 administrative staff to communicate with and on behalf of communities alleviates time pressures
300 experienced by clinicians and improves coordination of services [104]. Additionally, improved
301 infrastructure (e.g. telehealth, internet and IT services) would support workforce in remote locations
302 [105].

303

304 ***Optimal staffing for remote health services***

305 We found no national recommendations for optimal staffing levels for health professionals in remote
306 Australia, other than for nurses [9]. However, some regional planning documents prescribe optimal
307 clinician to population ratios [38, 109]. For example, the recommended ratios for Central Australia
308 are 1 Aboriginal Health Worker per 100 Aboriginal people, 1 community nurse per 250 people, and 1
309 doctor per 600 people [110]. Based on these figures, communities with a stable population of 250
310 should have a health service located within the community and access to on-call services and
311 communities of between 100-250 people should have a clinic staffed by two health professionals,
312 either senior Aboriginal Health Workers or registered nurses [9]. The World Health Organisation

313 (WHO) recommends 4.45 skilled health workers (physicians/nurses/midwives) per 1000 population
314 [111]. Although these targets were established in 1997 it continues to be a challenge to meet targets
315 due to difficulties with workforce retention and resources [9, 112].

316

317 Between 2013 and 2018, the Australian health workforce increased by 5.5%, yet for all registered
318 health professions, the number of employed full-time equivalent clinicians working decreased with
319 remoteness [113]. In 2020 there were more than 104,000 registered medical doctors in Australia but
320 fewer than 1,500 of these worked in remote and very remote areas [114].

321

322 **3) Characteristics of an effective health service**

323 ***Measuring the Effectiveness of a Health Service***

324 The effectiveness of a health service can be indicated by its use (the number of presentations); the
325 system performance; and/or community health outcomes [115, 116]. Specific measurements of
326 improved effectiveness of a health service include timeliness of care, increased staff recruitment and
327 retention, decreased wait times, reduced readmissions, increased referrals, improved cost-
328 effectiveness, improved safety, improved access to records, decreased feelings of isolation by
329 clinicians, increased equity, community participation and satisfaction with care, and decreased
330 adverse outcomes including suicide [117]. However, these metrics are seldom available in remote
331 health settings [118]. There are some data that enable determination of the reach and effectiveness of
332 health services. For example, in 2018 national immunisation rates in 5-year-olds were higher for
333 Indigenous (97%) than non-indigenous (95%) children [95].

334

335 A comprehensive review of the features of effective primary health care models in rural and remote
336 Australia highlighted the importance of supportive policy, positive relationships with State and
337 Territory governments, and community commitment [117]. Fundamental requirements for effective
338 and sustainable health services include good governance and management, community involvement
339 and leadership, adequate financing, infrastructure and ample workforce supply [117]. Many authors
340 highlight the importance of evaluating health services to ensure optimal health services [3, 106, 115,
341 117, 119-123]. However, the evaluation method must be carefully considered. [107] For example,
342 Key Performance Indicators (KPI) can be used to quantitatively assess health care services. However
343 these are not ideal on their own because their narrow focus can overlook social theories and
344 Indigenous concepts of health. KPIs alone are likely to fail to accurately reflect the contribution of a
345 service to health [124].

346

347 Several studies discuss the value of continuous quality improvement (CQI) for assessing the
348 sustainability and effectiveness of remote health services [121-123]. Previous research indicates that
349 CQI is effective for assessing primary health care of chronic conditions in remote Australia [108, 121,

350 122]. When applied to regional services, CQI resulted in large increases in the numbers of patients
351 accessing services (from 12 to 4000 patients from 1995-2009) [123].

352

353 ***Limitations to the effectiveness of health services***

354 Remoteness, inadequate medical and health workforce, and poor coordination negatively impact
355 health service effectiveness [9, 125]. In some very remote settings Community Health staff, whose
356 primary role is in preventative health care, are forced to deliver acute medical care [69]. There is an
357 increase in the proportion of children receiving nursing (rather than doctor) consultations as
358 remoteness increases [69].

359

360 Aboriginal Medical Services (AMS) are important providers of culturally appropriate services for
361 remote Indigenous children. However, in WA (2004) fewer than 5% of doctors, including specialists,
362 practice in remote and very remote areas and only 5.8 full time equivalent (FTE) doctors work in
363 culturally sound organisations such as AMSs [69].

364

365 Individuals in the remote health workforce are often forced to manage extremely complex disorders
366 with few resources and limited options for specialist referral [102]. Paediatricians are often required
367 to meet a greater demand than their capacity and resources allow, due to limited primary care
368 services. Thus, time is allocated to children with the greatest urgency and acuity, restricting the time
369 available for primary and preventive health care [126]. These circumstances are experienced in many
370 remote communities and may contribute to high staff turnover [99].

371

372 One qualitative study of the effectiveness of a remote health system in northern Australia found six
373 themes that contributed to a 'very chaotic system'. The system was: 'very ad hoc', 'swallowed by
374 acute' needs, 'going under', 'a flux', had a 'them and us' mentality and was 'a huge barrier' to quality
375 health care. [127]. Evaluation of a system for maternal and infant health in remote parts of
376 Australia's 'Top End' found a mismatch between staff numbers and skills and the volume and
377 severity of disease in patients using the service [125]. Other key concerns included insufficient or
378 absent Aboriginal leadership and inadequate coordination between remote and tertiary services [125].

379

380 ***Difficulties with communication, coordination, collaboration***

381 Many services have difficulty communicating with patients and other health professionals, due to a
382 lack of cultural knowledge and skills and technological barriers and struggle to ensure cooperation,
383 collaboration and coordination between different organisations and remote communities. [38, 102]
384 This is aggravated by challenges in case planning and organising referrals [102]. It is recommended
385 that there should be community paediatricians (or other health professionals or staff) who commit
386 solely to advocating for and co-ordinating outreach services to alleviate this burden from

387 organisations and other clinicians [126]. It is estimated that one day of administration and liaison is
388 required for each day of clinical work in remote settings [99, 102, 126].

389

390 Few online resources are designed to assist in the co-ordination of health services. In the NT, the
391 Department of Health’s online Remote Atlas contains a section specifically for SONT (Specialist
392 Outreach NT). Protocols to facilitate efficient and co-ordinated health service delivery to remote
393 communities, including SharePoint access and an online calendar with schedules of all specialist and
394 other health services, would improve services [128].

395

396 ***What core health services should be available***

397 In 2014, Thomas et al aimed to determine what core primary health services should be available in
398 remote communities, using a survey and Delphi process with healthcare providers and other key
399 stakeholders, 4 (10%) of whom were Indigenous community members or consumers [129]. They
400 identified eight essential primary health care services: care of the sick and injured, mental health,
401 maternal/child health, allied health, sexual/reproductive health, rehabilitation, oral/dental health, and
402 public health/illness prevention. They also identified seven functions required for primary health care
403 support: good management, governance and leadership, coordination, health infrastructure, quality
404 systems, data systems, professional development, and community participation. Further, these authors
405 found that equity in rural and remote areas was improved by prioritising service coordination, having
406 diverse strategies, and addressing the difficulty in recruiting and retaining clinicians [129].

407

408 **4) Models of care and solutions**

409 Numerous recommendations have been made for models of care to improve Indigenous health
410 services, but the challenge remains to implement these [9, 115, 130-133]. Healthcare delivery can be
411 improved with national policy frameworks for maternal and child health [134, 135], child and
412 adolescent mental health services [136], and child nutritional supplementation [107, 137].

413

414 The ‘National Aboriginal and Torres Strait Islander Health Plan 2021–2031’ [138] is the guiding
415 document for Indigenous health policy, programs and initiatives for the decade. It claims to change
416 the way governments work with Indigenous people reflecting their priorities, and to provide a holistic
417 perspective. The framework asks health care delivery to be place based, person centred and, culturally
418 safe and responsive [138].

419

420 The National Aboriginal and Torres Strait Islander Early Childhood Strategy refocuses investment
421 and policy on Indigenous children, so that they can be the future Elders and Custodians of Country
422 [130]. It has five goals, that Indigenous children are: 1) born healthy and strong, 2) supported to thrive
423 in the early years, 3) supported to connect to culture, country, and language, 4) in safe nurturing

424 homes and strong families, 5) with their families, active partners in building a better service system
425 [130].

426

427 Indigenous led organisations are leading the way in guiding policy, building leadership and building
428 healing [138]. One example is ‘SNAICC – National Voice for Our Children’, which works to embed
429 Indigenous priorities and best practice in national child policy [138].

430

431 The Australian Productivity Commission found that service delivery in remote Indigenous
432 communities where there are multiple providers is rarely competitive or enabling of user choice.
433 [139]. They recommended that decision making should be decentralised enabling local staff and
434 communities to plan, engage and implement services [139]. Similar recommendations were made at a
435 State level when the Queensland Productivity Commission, in 2017, reported on service availability
436 in remote Indigenous communities [140]. The Queensland Government spends \$1.2 billion a year
437 (\$29,000 per person) on services for remote Indigenous groups. Despite this, Indigenous outcomes are
438 poor and there is reliance on external funding. The report found the current system ‘is broken’
439 because it undermines outcomes and fosters passive dependence [140]. They recommended that
440 overhaul of the system must come from Indigenous communities and cited three key reforms
441 required: 1) structural reform to transfer accountability and decision-making to communities; 2)
442 service delivery reform to better focus on the needs of individuals and communities; 3) economic
443 reform to facilitate economic participation and community development [140].

444

445 At a regional level, an example of a holistic model of care for remote Australia is provided by the
446 *Kimberley Aboriginal Health Performance Forum (KAHPF)* [9] which began as the Kimberley
447 Aboriginal Health Plan Steering Committee in 1998 and published the *Kimberley Regional*
448 *Aboriginal Health Plan* in 1999 [38]. Its members include the Kimberley Aboriginal Medical
449 Services, Western Australian Community Health Services (Kimberley), and State government
450 representatives. Key recommendations of KAHPF address social determinants [9, 38] and include :
451 better coordination of health service delivery; interagency collaboration; use of innovative health
452 promotion programmes targeting specific groups; increasing the Aboriginal health workforce;
453 increasing prevention and health promotion; use of culturally appropriate, locally relevant resources;
454 allied health supports for children in classrooms; a School Entry Check for early identification of
455 health and developmental problems; screening for common childhood problems including anaemia
456 and growth faltering; better access to specialists; use of KAHPF diagnostic and treatment protocols
457 and clinical guidelines; use of the Kimberley standard drug list; clear referral pathways; long term
458 funding for successful programs; and ongoing training of health professionals [9]. This model
459 provides a gold standard for future remote health services and highlights the complexity of providing
460 a thorough, effective service.

461

462 Over the last decade, Aboriginal Community Controlled Health Services have contributed to
463 significant gains in maternal and child health, reduced smoking and alcohol misuse, and reduced
464 morbidity from cardiovascular, kidney and respiratory diseases [138]. In 2021, more than 97% of
465 Indigenous children aged 5 were fully immunised, a rate 2% higher than in non-Indigenous children
466 [12]. Implementation of programs to augment Aboriginal Community Controlled Health Services,
467 particularly with a focus on health promotion and early intervention, would improve child health
468 outcomes [104, 105, 141].

469

470 Improving access to primary, secondary and tertiary health services is crucial to improve child health
471 [39, 142, 143]. In remote communities, patient access is in part limited by lack of transport and
472 accommodation options [104, 144]. Increasing public transport to isolated communities or
473 implementing other service delivery models (e.g. outreach or telemedicine) may increase equity of
474 access and service use [145]. Mobile services e.g., for ear screening and surveillance may be a
475 valuable and cost-effective [78]. Paediatric outreach services to rural and remote areas benefit
476 children and their families in part by reducing their need to travel to services. Such services also
477 increase health professionals' cultural understanding and engagement in communities [40, 120, 146,
478 147].

479

480 **Best Practice: Moving towards health equity for Aboriginal children**

481 *Models of care*

482 Best practice models of care for child and adolescent health include provision of prevention programs,
483 multidisciplinary assessment, and early and secondary intervention [137]. A holistic approach to
484 health engages families in identifying problems and informing solutions for child health, growth and
485 development [102]. Continuity of care in rural and remote settings is essential, including transition
486 from child to adult health services. [102] Investment in local, trusted, Indigenous people as
487 'navigators' of language and cultural barriers improves care [148].

488

489 Meeting optimal child healthcare recommendations and providing access to services in remote
490 communities underpins best practice service models [102]. Adjustments required to improve current
491 services include increasing Indigenous leadership and involvement in service delivery and prevention
492 programs, improving access to services, dismantling cultural barriers to use existing services, using a
493 multidisciplinary approach for diagnosis and management, increasing IT capabilities, introducing
494 workforce initiatives, and integrating mother and child health services.

495

496 *Community engagement*

497 Indigenous community leadership (e.g. through Aboriginal Community-Controlled Organisations) is
498 essential for identifying service needs and how they should be delivered [24, 38, 104] and forming
499 collaborative partnerships with external organisations to develop services [104, 123, 149, 150].
500 Ongoing program evaluation is important to capture feedback from Indigenous families using
501 services, measure health outcomes, and highlight contemporary priorities for child health [145].
502 Enhancing access to Aboriginal Health Worker training and increasing numbers of graduates would
503 help dismantle cultural and language barriers that limit service use and effectiveness [141]. Protocols
504 should be generated for individual agencies, outlining best local practice policies for child health and
505 providing guidance to ensure delivery of culturally appropriate services [134, 145].

506

507 Improved youth engagement is integral to promote use of adolescent services. The Derby Aboriginal
508 Health Service talked to young Aboriginal people about barriers and enablers to accessing healthcare
509 [151] and found that engagement with youth, fostered by trained staff members skilled in caring for
510 young people, and continuity of staff were key.

511

512 ***Improving communication***

513 Communication challenges between health professionals and agencies, communities and health
514 departments could be improved with better infrastructure, including videoconferencing, and shared
515 electronic medical information systems [104, 145]. Telehealth is valuable for specialist medical
516 consultations in remote settings and enables education and capacity-building of local health
517 professionals [152, 153]. In isolated areas videoconferencing saves patients time and money and
518 improve access to high quality health services. Videoconferencing is dependent on technology,
519 including secure internet coverage, which is inadequate in many remote settings [152-155]. Allied
520 healthcare is delivered via telehealth in rural areas and may help fill service gaps in remote locations
521 and improve Indigenous child health [156]. Children receiving therapy engage more with school and
522 benefits could be maximised by targeting therapy to certain year levels. Barriers include hardware and
523 software breakdowns in remote settings and lack of local capacity to fix IT problems.

524

525 ***Access to services***

526 The importance of well-maintained and uncrowded housing for Indigenous child health and safety in
527 remote settings is well known. Less frequently discussed however, is the lack of accommodation for
528 health workers and its impact on the duration of stay in remote communities [157]. Increasing the
529 availability of accommodation for health professionals in remote communities would decrease travel
530 time, increase face-to-face clinical time, facilitate multidisciplinary care, and reduce wait times [145].

531

532 Improving public and private transport options would also improve access to services [145]. Provision
533 of social support and accommodation for children and families travelling to health services may
534 increase the likelihood of subsequent visits and improve health outcomes [104].

535

536 *New services*

537 When new health services are developed, they require adequate infrastructure and staffing to avoid
538 overload. Collaboration with Indigenous-controlled health organisations to improve public and
539 environmental health (including preventative health strategies) and address the social determinants of
540 health is imperative [6, 104, 134]. Ensuring that new services undergo evaluation of their processes
541 and outcomes and undertake continuous quality improvement is crucial.

542

543 **Discussion**

544 In this review we highlight the challenges to provision of child health services in remote Indigenous
545 communities, including access costs, and workforce. We summarise strategies to improve healthcare
546 for Indigenous children, including development and implantation of plans in partnership with
547 community leaders. We highlight the need for culturally informed services and trained Indigenous
548 healthcare workers in remote communities. Government, NGO, specialist outreach, and virtual care
549 services all play important roles. However, coordination between primary, secondary, and tertiary
550 services is essential to prevent duplication and maximise communication.

551

552 Aboriginal and Torres Strait Islander cultures and people thrive in Australia despite the ongoing
553 legacy of dislocation and violence stemming from colonisation. The impact of the Stolen Generation,
554 born between 1910 and 1970 when one in three Indigenous children were taken from their families
555 under Australian Government policy, persists [138]. We acknowledge the strength and resilience
556 shown by Indigenous people in the face of historical trauma and the ongoing disadvantage and
557 disparity between health outcomes in Indigenous and non-Indigenous children, especially in remote
558 settings.

559

560 Ideally local community members receive training and remain or return to their community to provide
561 healthcare. Unfortunately, this is not the current reality for most remote communities so attracting and
562 retaining healthcare professionals to remote locations requires financial, educational, and collegiate
563 support and training in cultural competency. Increasing the availability of accommodation for
564 healthcare workers in remote settings is crucial. It is unquestionable that increasing the availability of
565 accommodation for local community members is also crucial. There must also be pathways for
566 training Indigenous people and use of hybrid services using telehealth. Gold standards should be
567 established for staff numbers in remote Australia, supported by national policy frameworks. Audits
568 which assess whether positions are filled, with a focus on staff consistency and continuity, will inform

569 health services for remote Indigenous communities. Aboriginal Community Controlled Health
570 Services are a particular strength, as is the provision of holistic, wrap-around services in remote
571 Indigenous communities [139].

572
573 Holistic services which embrace physical, mental (social/emotional), cultural and spiritual health and
574 focus on the social determinants of health will likely have the greatest impact in remote settings.
575 Removing barriers to adequate nutrition, good housing, educational supports, and preventative
576 programs including vaccination are key elements. Promoting maternal health literacy, pre-conception
577 and pregnancy care, parenting programs, and early infancy and childhood programs is paramount.
578 Further strengthening the ability of Indigenous people to share their own knowledge regarding the
579 importance of the developmental origins of health and disease is likely to support most effective
580 health promotion message and policies for the first 1000 days of a child's life [158].

581
582 The average annual cost of healthcare for an Indigenous Australian person (\$8,949) is 1.3 times that
583 for non-Indigenous Australians (\$6,657) and was highest for Indigenous Australians living in remote
584 and very remote areas (\$9,005) in 2015–16 [95], reflecting the increased morbidity among Indigenous
585 populations, expenses incurred by remote service delivery, and incentives for health professionals to
586 work in remote settings [104]. However, there is a disconnect between funding and implementation of
587 services, particularly relating to the social determinants of health [104]. Although funding for
588 Indigenous child health services has increased over time, implementation costs have increased in
589 keeping with inflation and demand [159]. In tertiary health care, the case-mix method used to allocate
590 hospital funding uses benchmarks based on national averages, cost weights, and length of stay [89].
591 This method is criticised as further disadvantaging Indigenous children, who present more frequently
592 to health services and more often have complex illnesses, comorbidities and prolonged hospital stays
593 [89]. Savings and improved child health could result from addressing social determinants of health
594 and improving maternal-child supports, primary health care, and disease prevention in remote
595 communities [145]. Continuous quality assessment of existing and future programs is crucial to
596 indicate needs and inefficiencies and inform funding allocation.

597
598 Additional support for Aboriginal Community Controlled Health Organisations and increased funding
599 for training and employment for Aboriginal Health Workers will be pivotal for improving patient
600 engagement with and cultural safety within health services. Expenditure on appropriate, well-
601 resourced services for maternal and child health in remote Australia will have a strong impact on child
602 and future adult well-being [145, 160].

Acknowledgements

We acknowledge the contribution of Prof Kirsty Douglas and Prof Kathryn Glass to an early draft of manuscript.

Funding

Philippa Dossetor was supported by a part-time PhD scholarship through the Australian National University Medical School and the College of Biology, Medicine and the Environment, a University of Sydney Poche Institute Scholarship and by a part-time Avant DiT scholarship. Joseph Freeman is supported by the Curtin PhD Scholarship for Clinical Research. Alexandra Martiniuk was funded by an NHMRC TRIP (Translating Research into Practice) Fellowship (2016–2018) and an NHMRC Investigator Grant (2021-current). Elizabeth Elliott is supported by a Medical Research Futures Fund Next Generation Fellowship (#1135959).

References

1. Oscar J, Lowitja Institute. Close the gap campaign report 2021. 2020.
2. Australian Bureau of Statistics. Census of Population and Housing: characteristics of Aboriginal and Torres Strait Islander Australians, 2016. Canberra: 2019.
3. Bailie RS, Si D, Dowden MC, Connors CM, O'Donoghue L, Liddle HE, et al. Delivery of child health services in Indigenous communities: implications for the federal government's emergency intervention in the Northern Territory. *Med J Aust*. 2008;188(10):615-8. Epub 2008/05/20. doi: 10.5694/j.1326-5377.2008.tb01806.x. PubMed PMID: 18484941.
4. Thomas S. Kids Pitstop: one stop community health shop for Indigenous families in the West Pilbara. *Rural Health: the place to be*; 17-20 May; Cairns 2009.
5. Pulver LJ, Haswell MR, Ring I, Waldon J, Clark W, Valorie Whetung, et al. Indigenous Health - Australia, Canada, Aotearoa New Zealand and the United States - Laying claim to a future that embraces health for us all. *World Health Report - Financing for Universal Health Coverage Background Paper*. 2010;No 33 1-79.
6. Australian Institute of Health and Welfare. Access to health services for Aboriginal and Torres Strait Islander people. Canberra: AIHW, 2011.
7. Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islander health services report, 2010-11: OATSIH services reporting - key results. Canberra: AIHW, 2012.
8. Bar-Zeev SJ, Kruske SG, Barclay LM, Bar-Zeev NH, Carapetis JR, Kildea SV. Use of health services by remote dwelling Aboriginal infants in tropical northern Australia: a retrospective cohort study. *BMC Pediatrics*. 2012;12:19. doi: 10.1186/1471-2431-12-19.
9. Lewis J. Kimberley Aboriginal Primary Health Plan 2012-2015. Broome: Kimberley Aboriginal Health Planning Forum, 2013.
10. Li Z. Australia's Mothers and Babies 2010. Canberra: Australian Institute of Health and Welfare; 2012.
11. National Health and Medical Research Council. A Healthy Start to Life for all Australians. NHMRC, 2002.
12. Australian Institute of Health Welfare. Australia's Mothers and Babies 2018: In Brief. Canberra: AIHW, 2020 176054681X.
13. Australian Institute of Health and Welfare. Admitted patient care 2017–18: Australian hospital statistics. Canberra: AIHW, 2019.
14. Dossetor PJ, Fitzpatrick EFM, Glass K, Douglas K, Watkins R, Oscar J, et al. Emergency Department Presentations by Children in Remote Australia: A Population-based Study. *Global Pediatric Health*. 2021;8. doi: 10.1177/2333794X21991006.
15. Dossetor PJ, Martiniuk ALC, Fitzpatrick JP, Oscar J, Carter M, Watkins R, et al. Pediatric hospital admissions in Indigenous children: a population-based study in remote Australia. *BMC Pediatrics*. 2017;17(1):195. doi: 10.1186/s12887-017-0947-0.
16. Gilchrist DJ, Perks B. Challenges of Delivering Regional and Remote Human Services and Supports. *The Palgrave Encyclopedia of Urban and Regional Futures*: Springer; 2022. p. 1-6.
17. Zwi K, Joshua P, Moran P, Casacelli M. Prioritising health services for high risk and disadvantaged children. *Journal of Paediatrics and Child Health*. 2011;47:23.
18. Adams E, Tongs J. Starting a perinatal and infant mental health service at Winnunga Nimmityjah. *Australasian Psychiatry*. 2011;19:S20-S2.
19. Freemantle C, Officer K, Mcaullay D, Anderson I. Australian Indigenous Health- Within an International Context. Darwin: Cooperative Research Centre for Aboriginal Health, 2007.

20. Carson B, Dunbar, T., Chenall, R.D., Bailie, R. Social determinants of Indigenous health Menzies School of Health Research: Allen and Unwin; 2007.
21. Kang M, Robards F, Luscombe G, Sancu L, Usherwood T. The relationship between having a regular general practitioner (GP) and the experience of healthcare barriers: a cross-sectional study among young people in NSW, Australia, with oversampling from marginalised groups. *BMC family practice*. 2020;21(1):1-9.
22. Queensland Health. Making Tracks toward closing the gap in health outcomes for Indigenous Queenslanders by 2033 Implementation Plan 2009-10 to 2011-12. Brisbane: Department of Health, 2010.
23. COAG Reform Council. Indigenous reform 2011-12: comparing performance across Australia. Sydney: COAG Reform Council, 2013.
24. Lowitja Institute. Close the gap campaign report 2021. 2021.
25. Australian Institute of Health Welfare. Australia's children. Canberra: AIHW, 2020 Contract No.: CWS 69.
26. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Annals of internal medicine*. 2018;169(7):467-73.
27. World Health Organization. The world health report 2000: health systems: improving performance: World Health Organization; 2000.
28. Australian Institute of Health and Welfare. Rural, regional and remote health: a guide to remoteness classifications. AIHW, 2004.
29. Wakerman J. Defining remote health. *Australian Journal of Rural Health*. 2004;12(5):210-4. doi: 10.1111/j.1440-1854.2004.00607.x.
30. Whelan S, Wright DJ. Health Services Use and Lifestyle Choices of Indigenous and non-Indigenous Australians. *Social Science & Medicine*. 2013.
31. Crouchley K, Carlse N. Aboriginal Health Profile: Kimberley Health Region. WA: Department of Health, 2012.
32. Ou L, Chen J, Hillman K, Eastwood J. The comparison of health status and health services utilisation between Indigenous and non-Indigenous infants in Australia. *Australian and New Zealand Journal of Public Health*. 2010;34(1):50-6.
33. Chandradasa M, Basu S. Collaborative networking between regional child mental health, paediatric and educational services in Gippsland, Australia: An online survey. *The Australian Journal of Rural Health*. 2019.
34. Achat HM, Thomas P, Close GR, Moerkerken LR, Harris MF. General health care service utilisation: where, when and by whom in a socioeconomically disadvantaged population. *Australian Journal of Primary Health*. 2010;16(2):132-40.
35. Barr M, Duncan J, Dally K. A systematic review of services to DHH children in rural and remote regions. *The Journal of Deaf Studies and Deaf Education*. 2018;23(2):118-30.
36. Morgan K. QT babies: Neonatal care in the Queensland tropics. *Journal of Paediatrics and Child Health*. 2012.
37. D'Aprano A, Silburn S, Johnston V, Bailie R, Mensah F, Oberklaid F, et al. Challenges in monitoring the development of young children in remote Aboriginal health services: clinical audit findings and recommendations for improving practice. *Rural Remote Health*. 2016;16(3):3852.
38. Atkinson D, Bridge C, Gray D. Kimberley Regional Aboriginal Health Plan: Aboriginal Health in the Kimberley: current circumstances and future directions. Broome: Kimberley Aboriginal Medical Services Council, 1999.
39. Gruen RL, Bailie RS, d'Abbs PH, O'Rourke IC, O'Brien MM, Verma N. Improving access to specialist care for remote Aboriginal communities: evaluation of a specialist

- outreach service. *Med J Aust.* 2001;174(10):507-11. Epub 2001/06/23. doi: 10.5694/j.1326-5377.2001.tb143400.x. PubMed PMID: 11419770.
40. Gruen RL, Bailie RS, Wang Z, Heard S, O'Rourke IC. Specialist outreach to isolated and disadvantaged communities: a population-based study. *Lancet.* 2006;368(9530):130-8. doi: 10.1016/S0140-6736(06)68812-0.
 41. Rural Health West. *Specialist Services in Rural Western Australia: Gap and Equity* Perth: RHW, 2013.
 42. The Senate Select Committee on Regional and Remote Communities. *Indigenous Australians, Incarceration and the Criminal Justice System.* Canberra: Senate, 2010.
 43. Phillips JH, Wigger C, Beissbarth J, McCallum GB, Leach A, Morris PS. Can mobile phone multimedia messages and text messages improve clinic attendance for Aboriginal children with chronic otitis media? A randomised controlled trial. *Journal of Paediatrics and Child Health.* 2014;50(5):362-7.
 44. Breen C, Altman L, Ging J, Deverell M, Woolfenden S, Zurynski Y. Significant reductions in tertiary hospital encounters and less travel for families after implementation of Paediatric Care Coordination in Australia. *BMC health services research.* 2018;18(1):1-10.
 45. Health Workforce Australia. *National Rural and Remote Workforce Innovation and Reform Strategy.* Adelaide: HWA, 2013.
 46. Health Workforce Australia. *Health Workforce 2025 – Doctors, Nurses and Midwives – Volume 2.* Adelaide: HWA, 2012.
 47. Australian Institute of Health and Welfare. *Expenditure on Health for Aboriginal and Torres Strait Islander People 2008-09: An Analysis by Remoteness and Disease.* AIHW, 2011.
 48. Office of Aboriginal and Torres Strait Islander Health. *Improving Health in Aboriginal and Torres Strait Islander Mothers, Babies and Young Children - A Literature Review.* 2005.
 49. Office for Aboriginal Torres Strait Islander Health. *Better health care: studies in the successful delivery of primary health care services for Aboriginal and Torres Strait Islander Australians.* Canberra: Commonwealth Department of Health and Ageing, 2001.
 50. WA Country Health Service. *Kimberley Health Profile.* Perth: WACHS, 2018.
 51. Nancarrow SA, Roots A, Grace S, Saberi V. Models of care involving district hospitals: a rapid review to inform the Australian rural and remote context. *Australian Health Review.* 2015;39(5):494-507.
 52. Menzies School of Health Research. *Effective Integration of Services for Children and Families: Making it Happen.* Darwin: 2011.
 53. Wakerman J, Humphreys J, Wells R, Kuipers P, Entwistle P, Jones J. A systematic review of primary health care delivery models in rural and remote Australia, 1993-2006. 2006.
 54. Schofield D, Fuller J, Wagner S, Friis L, Tyrell B. Multidisciplinary management of complex care. *Australian Journal of Rural Health.* 2009;17(1):45-8.
 55. Allen, Clarke. *Evaluation of the child health check initiative and the expanding health service delivery initiative: summary report.* Department of Health and Ageing Canberra; 2011.
 56. Standing Committee on Family and Community Affairs. *Standing Committee on Family and Community Affairs. Health is life: Inquiry into indigenous health.* Canberra: 2000.
 57. Standing community on Health and Ageing. *The Blame Game: Report on the inquiry into health funding.* Canberra: Senate, 2006.

58. McDonald L, Bailie R, Morris P, Rumbold A, Paterson B. Interventions to prevent growth faltering in remote Indigenous communities. Australian Primary Health Care Research Institute (APHCRI), The Australian National University; 2006.
59. Shaw G. An ethnography of changes in child rearing over time in the Ngaanyatjarra Lands: Implications for policy development on health in early childhood. CAEPR Seminar Series Canberra 2013.
60. Veit FC, Sanci LA, Coffey CM, Young DY, Bowes G. Barriers to effective primary health care for adolescents. *Medical Journal of Australia*. 1996;165(3):131-3.
61. Hussain R, Tait K. Parental perceptions of information needs and service provision for children with developmental disabilities in rural Australia. *Disability and Rehabilitation*. 2015;37(18):1609-16.
62. Gardiner F, Gale L, Ransom A, Laverty M. Looking Ahead: responding to the health needs of country Australians in 2028-the centenary year of the RFDS. Canberra, Australia: The Royal Flying Doctor Service. 2018.
63. Royal Flying Doctor Service. Annual Report 2011-12. 2012.
64. Medlin L, editor Indigenous Respiratory Outreach Care (IROC): Enhancing Respiratory Health In Rural And Remote Aboriginal And Torres Strait Islander Communities. *Respirology*; 2014.
65. Verdon S, Wilson L, Smith-Tamaray M, McAllister L. An investigation of equity of rural speech-language pathology services for children: a geographic perspective. *Int J Speech Lang Pathol*. 2011;13(3):239-50. doi: 10.3109/17549507.2011.573865.
66. O'Callaghan AM, McAllister L, Wilson L. Consumers' proposed solutions to barriers to access of rural and remote speech pathology services. *Advances in Speech Language Pathology*. 2005;7(2):58-64.
67. Gunasekera H, Morris PS, Daniels J, Couzos S, Craig JC. Otitis media in Aboriginal children: the discordance between burden of illness and access to services in rural/remote and urban Australia. *Journal of Paediatrics and Child Health*. 2009;45(7-8):425-30.
68. Jones DM, McAllister L, Lyle DM. Rural and remote speech-language pathology service inequities: An Australian human rights dilemma. *International journal of speech-language pathology*. 2018;20(1):98-101.
69. Zubrick SR, Lawrence DM, Silburn SR, Blair E, Milroy H, Wilkes T, et al. The Western Australian Aboriginal Child Health Survey: The Health of Aboriginal Children and Young People. Perth: Telethon Institute for Child Health Research, 2004.
70. Margolis SA. Is Fly in/Fly out (FIFO) a viable interim solution to address remote medical workforce shortages? *Rural and Remote Health*. 2012;12(4):1-6.
71. Peiris D, Wirtanen C, Hall J. Aeromedical evacuations from an east Arnhem Land community 2003-2005: the impact on a primary health care centre. *Aust J Rural Health*. 2006;14(6):270-4. doi: 10.1111/j.1440-1584.2006.00828.x.
72. Francis A, Didsbury M, Lim WH, Kim S, White S, Craig JC, et al. The impact of socioeconomic status and geographic remoteness on access to pre-emptive kidney transplantation and transplant outcomes among children. *Pediatric Nephrology*. 2016;31(6):1011-9.
73. Garne DL, Perkins DA, Boreland FT, Lyle DM. Frequent users of the Royal Flying Doctor Service primary clinic and aeromedical services in remote New South Wales: a quality study. *Med J Aust*. 2009;191(11-12):602-4. doi: 10.5694/j.1326-5377.2009.tb03344.x.
74. Smith AC, Armfield NR, White MM, Williams ML, Koh T, Hurley T, et al. Clinical services and professional support: a review of mobile telepaediatric services in Queensland. *Studies in Health Technology and Informatics*. 2010;161:149-58.

75. Allen O. Anthill and other injuries: a case for mobile allied health teams to remote Australia. *Australian Journal of Rural Health*. 1996;4(1):33-42.
76. Dossetor DR, Nunn KP, Fairley M, Eggleton D. A child and adolescent psychiatric outreach service for rural New South Wales: a telemedicine pilot study. *J Paediatr Child Health*. 1999;35(6):525-9. doi: 10.1046/j.1440-1754.1999.00410.x.
77. A L, R B. Aboriginal Maternal and Child Project Strengths and Needs Analysis. Perth: Aboriginal Health Council of Western Australia, 2010.
78. Nguyen KH, Smith AC, Armfield NR, Bensink M, Scuffham PA. Cost-Effectiveness Analysis of a Mobile Ear Screening and Surveillance Service versus an Outreach Screening, Surveillance and Surgical Service for Indigenous Children in Australia. *PLoS One*. 2015;10(9). doi: 10.1371/journal.pone.0138369.
79. Nguyen H, Zarnowiecki D, Segal L, Gent D, Silver B, Boffa J. Feasibility of implementing infant home visiting in a Central Australian Aboriginal community. *Prevention Science*. 2018;19(7):966-76.
80. Vos T, Carter R, Barendregt J, Mihalopoulos C, Veerman L, Magnus A, et al. Assessing cost-effectiveness in prevention: ACE-prevention September 2010 final report: University of Queensland; 2010.
81. Australian institute of Health and Welfare. Aboriginal and Torres Strait Islander Health Performance Framework. Perth: 2013.
82. Ansari Z, Laditka JN, Laditka SB. Access to health care and hospitalization for ambulatory care sensitive conditions. *Med Care Res Rev*. 2006;63(6):719-41. doi: 10.1177/1077558706293637.
83. Flabouris A, Hart GK, Nicholls A. Accessibility of the Australian population to an ICU, and of ICUs to each other. *Critical care and resuscitation : journal of the Australasian Academy of Critical Care Medicine*. 2012;14(3):177-84.
84. WA Department of Health. 2005 - 2015 WA Clinical services framework
85. Mitchinson C, Strobel N, McAullay D, McAuley K, Bailie R, Edmond KM. Anemia in disadvantaged children aged under five years; quality of care in primary practice. *BMC pediatrics*. 2019;19(1):1-11.
86. Department of Health and Ageing. National Strategic Framework for Rural and Remote Health. 2011.
87. Prime Minister's Report. Closing the Gap: The Prime Minister's Report Canberra: 2012.
88. Hoy WE. Closing the gap by 2030: aspiration versus reality in Indigenous health. *Medical Journal of Australia*. 2009;190(10):542-4.
89. Ruben AR, Fisher DA. The casemix system of hospital funding can further disadvantage Aboriginal children. *Medical Journal of Australia*. 1998;169(SUPPL.):S6-S10.
90. Russell-Weisz D, Hindle D. High length-of-stay outliers under casemix funding of a remote rural community with a high proportion of aboriginal patients. *Aust Health Rev*. 2000;23(2):47-61. doi: 10.1071/ah000047.
91. Edmond KM, McAuley K, McAullay D, Matthews V, Strobel N, Marriott R, et al. Quality of social and emotional wellbeing services for families of young Indigenous children attending primary care centers; a cross sectional analysis. *BMC health services research*. 2018;18(1):1-11.
92. Edmond KM, Tung S, McAuley K, Strobel N, McAullay D. Improving developmental care in primary practice for disadvantaged children. *Archives of Disease in Childhood*. 2019;104(4):372-80.
93. HealthInfoNet AI, Harford-Mills M, MacRae A, Drew N. Summary of Aboriginal and Torres Strait Islander health status 2018. 2019.

94. Australian Institute of Health and Welfare. Healthy for Life-Aboriginal Community Controlled Health Services: Report Card. AIHW Canberra; 2013.
95. Australian Institute of Health and Welfare. Aboriginal and Torres Strait Islander Health Performance Framework 2020 report: Western Australia. Canberra: AIHW, 2020.
96. Department of Health and Human Services. Medical Specialist Outreach Assistance Program (MSOAP) Guidelines. 2010.
97. Lenthall S, Wakerman J, Opie T, Dunn S, Macleod M, Dollard M, et al. Nursing workforce in very remote Australia, characteristics and key issues. *The Australian Journal of Rural Health*. 2011;19(1):32-7.
98. O'Kane A, Tsey K. Towards a needs based mental health resource allocation and service development in rural and remote Australia. *Australas Psychiatry*. 2004;12(4):390-5. doi: 10.1080/j.1440-1665.2004.02133.x.
99. Dossetor PJ, Thorburn K, Oscar J, Carter M, Fitzpatrick J, Bower C, et al. Review of Aboriginal child health services in remote Western Australia identifies challenges and informs solutions. *BMC health services research*. 2019;19(1):758. doi: 10.1186/s12913-019-4605-0.
100. Williams ML, Smith AC. Paediatric outreach services. *Journal of Paediatrics and Child Health*. 2004;40(9-10):501-3. doi: 10.1111/j.1440-1754.2004.00450.x.
101. Geelhoed GC, Geelhoed EA. Positive impact of increased number of emergency consultants. *Archives of disease in childhood*. 2008;93(1):62-4.
102. Grant S. Child Malnutrition / Failure to Thrive Project 2008: Final Report. Darwin, NT.: Department of Health and Families, Northern Territory Government, 2008.
103. Fitzpatrick JP, Elliott EJ, Latimer J, Carter M, Oscar J, Ferreira M, et al. The Lililwan Project: study protocol for a population-based active case ascertainment study of the prevalence of fetal alcohol spectrum disorders (FASD) in remote Australian Aboriginal communities. *BMJ Open*. 2012;2(3). doi: 10.1136/bmjopen-2012-000968.
104. Griew R, Tilton E, Cox N, Thomas D. The link between primary health care and health outcomes for Aboriginal and Torres Strait Islander Australians. Canberra: OATSIH, 2008.
105. Humphreys J, Wakerman J. Primary health care in rural and remote Australia: achieving equity of access and outcomes through national reform: A discussion paper. Canberra: National Health and Hospitals Reform Commission, 2008.
106. Price E, Considine G. Kimberley Primary Health Care Sustainability Study 2008 - 2030. Kimberley Division of General Practice Ltd, 2008.
107. Wakerman J. Innovative rural and remote primary health care models: what do we know and what are the research priorities? *Aust J Rural Health*. 2009;17(1):21-6. doi: 10.1111/j.1440-1584.2008.01032.x.
108. Gardner K, Bailie R, Si D, O'Donoghue L, Kennedy C, Liddle H, et al. Reorienting primary health care for addressing chronic conditions in remote Australia and the South Pacific: review of evidence and lessons from an innovative quality improvement process. *Aust J Rural Health*. 2011;19(3):111-7. doi: 10.1111/j.1440-1584.2010.01181.x.
109. Lewis J. Kimberley Aboriginal Primary Health Plan 2012-15. Broome: Kimberley Aboriginal Health Planning Forum, 2013.
110. Wakerman J, Bennett, M., Healy, V., & Warchivker, I. . Kimberley Regional Aboriginal Health Plan. Northern Territory Australia: Menzies School of Health Research. 1997.
111. Organization WH. Health workforce requirements for universal health coverage and the sustainable development goals.(human resources for health observer, 17). 2016.
112. Health Workforce Australia. Growing Our Future: the Aboriginal and Torres Strait Islander Health Worker Project Final Report. Adelaide: HWA, 2011.

113. Australian Institute of Health and Welfare. Health workforce. Canberra: AIHW, 2020.
114. Department of Health. Medical doctors and specialists in Australia. Canberra: DoH, 2020.
115. McCalman J, Tsey K, Clifford A, Earles W, Shakeshaft A, Bainbridge R. Applying what works: a systematic search of the transfer and implementation of promising Indigenous Australian health services and programs. *BMC Public Health*. 2012;12(1):600.
116. World Health Organization. Toolkit on monitoring health systems strengthening: Service Delivery. 2008.
117. Wakerman J, Humphreys JS, Wells R, Kuipers P, Jones JA, Entwistle P, et al. Features of effective primary health care models in rural and remote Australia: a case-study analysis. *Medical Journal of Australia*. 2009;191(2):88-91. doi: 10.5694/j.1326-5377.2009.tb02700.x.
118. Sanchez L, Carney S, Estermann A, Sparrow K, Turner D. An evaluation of the benefits of swimming pools for the hearing and ear health status of young Indigenous Australians: A whole-of-population study across multiple remote Indigenous communities. Adelaide: Flinders University. 2012.
119. Bailie RS, Togni SJ, Si D, Robinson G, d'Abbs PH. Preventive medical care in remote Aboriginal communities in the Northern Territory: a follow-up study of the impact of clinical guidelines, computerised recall and reminder systems, and audit and feedback. *BMC health services research*. 2003;3(1):15. doi: 10.1186/1472-6963-3-15.
120. Wakerman J, Humphreys JS, Wells R, Kuipers P, Entwistle P, Jones J. Primary health care delivery models in rural and remote Australia: a systematic review. *BMC health services research*. 2008;8:276. doi: 10.1186/1472-6963-8-276. PubMed PMID: 19114003.
121. Bailie R, Si D, Connors C, Weeramanthri T, Clark L, Dowden M, et al. Study protocol: Audit and Best Practice for Chronic Disease Extension (ABCDE) Project. *BMC health services research*. 2008;8(1):184. doi: 10.1186/1472-6963-8-184.
122. Bailie RS, Si D, O Donoghue L, Dowden M. Indigenous health: effective and sustainable health services through continuous quality improvement. *Medical Journal of Australia*. 2007;186(10):525.
123. Hayman N. Strategies to improve indigenous access for urban and regional populations to health services. *Heart Lung Circulation*. 2010;19(5-6):367-71. doi: 10.1016/j.hlc.2010.02.014.
124. McKeown S. Evaluating aboriginal primary health care services using national key performance indicators: What has happened to the social theories of aboriginal health? *Internal Medicine Journal*. 2011;41:37.
125. Barclay L, Kruske S, Bar-Zeev S, Steenkamp M, Josif C, Narjic CW, et al. Improving Aboriginal maternal and infant health services in the 'Top End' of Australia; synthesis of the findings of a health services research program aimed at engaging stakeholders, developing research capacity and embedding change. *BMC health services research*. 2014;14(1):241. doi: 10.1186/1472-6963-14-241.
126. Henderson R. Review of community paediatrics, the Central Australian Remote Health Service, Alices Springs. Prince Margaret Hospital for Children, Perth WA, 2008.
127. Josif CM, Kruske S, Kildea SV, Barclay LM. The quality of health services provided to remote dwelling aboriginal infants in the top end of northern Australia following health system changes: a qualitative analysis. *BMC Pediatrics*. 2017;17(1):93. doi: 10.1186/s12887-017-0849-1.
128. Northern Territory Department of Health. Specialist and Clinical Services Outreach PPHC Remote. Darwin: Department of Health, 2022 Contract No.: July.

129. Thomas SL, Wakerman J, Humphreys JS. What core primary health care services should be available to Australians living in rural and remote communities? *BMC Family Practice*. 2014;15:143. doi: 10.1186/1471-2296-15-143.
130. National Indigenous Australians Agency. National Aboriginal and Torres Strait Islander Early Childhood Strategy. Canberra: NIAA, 2021.
131. Strategy Policy Planning. The Best Opportunities in Life: Northern Territory Child and Adolescent health and Wellbeing Strategic Plan 2018-2028. Darwin: Department of Health, 2018.
132. Moss B, Harper H, Silburn S. Strengthening Aboriginal Child Development in Central Australia through a Universal Preschool Readiness Program. *Australasian Journal of Early Childhood*. 2015;40(4):13-20. doi: 10.1177/183693911504000403.
133. Healing Foundation, Emerging Minds. Improving the social and emotional wellbeing of Aboriginal and Torres Strait Islander children. 2020.
134. Australian Medical Association. Indigenous Health Report Card – "Ending the Cycle of Vulnerability: The Health of Indigenous Children". 2008.
135. Johns S. Early childhood service development and intersectoral collaboration in rural Australia. *Australian Journal of Primary Health*. 2010;16(1):40-6.
136. Seidler ZE, Rice SM, Dhillon HM, Cotton SM, Telford NR, McEachran J, et al. Patterns of youth mental health service use and discontinuation: Population data from Australia's headspace model of care. *Psychiatric services*. 2020;71(11):1104-13.
137. Trevena LJ. Problem-based learning in public health workforce training: a discussion of educational principles and evidence. 2007 Jan-Feb. Report No.: Contract No.: 1-2.
138. Department of Health. National Aboriginal and Torres Strait Islander Health Plan 2021–2031. Canberra: Australian Government, 2021.
139. Productivity Commission. Introducing competition and informed user choice into human services: Reforms to human services-inquiry report. 2018.
140. Queensland Productivity Commission. Service delivery in remote and discrete Aboriginal and Torres Strait Islander communities. 2018.
141. Couzos S. Practical measures that improve human rights - towards health equity for Aboriginal children. *Health Promotion Journal of Australia : Official Journal of Australian Association of Health Promotion Professionals*. 2004;15(3):186-92.
142. Australian Indigenous HealthInfoNet. Overview of Aboriginal and Torres Strait Islander health status, 2020. 2021.
143. Sheperd C, Zubrick S. Human capabilities and child development: Some inferences using the 2008 NATSISS. Perspectives on the 2008 National Aboriginal and Torres Strait Islander Social Survey; ANU 2011.
144. DoH Epidemiology Branch. Child Population Profile, Health and Wellbeing Surveillance System, 2009 Kimberley health region. Perth: Department of Health, 2010.
145. Larson A, Bradley R. Aboriginal maternal and child project: strengths and needs analysis. Geraldton, WA: Combined Universities Centre for Rural Health and Aboriginal Health Council of Western Australia, 2010.
146. Cord-Udy N. The Medical Specialist Outreach Assistance Programme in South Australia at 2 years. *Australasian Psychiatry*. 2004;12(2):161-5.
147. Humphreys JS. Key considerations in delivering appropriate and accessible health care for rural and remote populations: discussant overview. *Australian Journal of Rural Health*. 2009;17(1):34-8. doi: 10.1111/j.1440-1584.2008.01034.x.
148. Mitchell AG, Belton S, Johnston V, Ralph AP. Transition to adult care for Aboriginal children with rheumatic fever: a review informed by a focussed ethnography in northern Australia. *Australian Journal of Primary Health*. 2018;24(1):9-13. doi: 10.1071/PY17069.

149. Elliott EJ, Latimer J, Oscar J, Fitzpatrick J, Carter M. The Lililwan Collaboration: Inquiry into Fetal Alcohol Spectrum Disorders (FASD). 2012 978-0-9873014-0-6.
150. Jeffery V, Ervin K. Responding to rural health needs through community participation: addressing the concerns of children and young adults. *Australian Journal of Primary Health*. 2011;17(2):125-30. doi: 10.1071/PY10050.
151. Warwick S, Atkinson D, Kitaura T, LeLievre M, Marley JV. Young Aboriginal People's Perspective on Access to Health Care in Remote Australia: Hearing Their Voices. *Prog Community Health Partnersh*. 2019;13(2):171-81. doi: 10.1353/cpr.2019.0017.
152. Smith AC. Telepaediatrics in Queensland. *Telemedicine and Child Health*. 2005:25-39.
153. Moffatt JJ, Eley DS. The reported benefits of telehealth for rural Australians. *Australian Health Review*. 2010;34(3):276-81. doi: 10.1071/AH09794.
154. Buckley D, Weisser S. Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit. *Australian and New Zealand Journal of Public Health*. 2012;36(5):478-82. doi: 10.1111/j.1753-6405.2012.00915.x.
155. Australian Government. *Regional Telecommunications Review*. Canberra: 2021.
156. Langbecker DH, Caffery L, Taylor M, Theodoros D, Smith AC. Impact of school-based allied health therapy via telehealth on children's speech and language, class participation and educational outcomes. *Journal of Telemedicine and Telecare*. 2019;25(9):559-65. doi: 10.1177/1357633X19875848.
157. Australian Government. *Remote Housing Review*. Canberra: 2017.
158. Sivertsen N, Anikeeva O, Deverix J, Grant J. Aboriginal and Torres Strait Islander family access to continuity of health care services in the first 1000days of life: a systematic review of the literature. *BMC health services research*. 2020;20(1):829.
159. Australian Institute of Health and Welfare. *Health expenditure Australia 2010–11*. Canberra: AIHW, 2012.
160. Steenkamp M, Bar-Zeev S, Rumbold A, Barclay L, Kildea S. Pragmatic indicators for remote Aboriginal maternal and infant health care: why it matters and where to start. *Australian and New Zealand Journal of Public Health*. 2010;34 Suppl 1(s1):S5-8. doi: 10.1111/j.1753-6405.2010.00545.x.

APPENDIX

Table 1a: Peer-reviewed literature

Author, Date, Title	Location, Setting	Objective(s), Study type	Key findings	Analysis/Gaps
Achat, 2010 [1] General health care service utilisation: Where, when and by whom in a socioeconomically disadvantaged population	Disadvantaged population, health care use.	Examine the utilisation of and preferences related to health care services by residents of a disadvantaged area and to identify factors associated with levels of current and future use. Survey.	Disadvantaged communities make only moderate use of specialists and emergency departments, and little routine use of other primary health services.	Effect of length of visit and level of engagement not included in dataset.
Adams, 2011 [2] Starting a perinatal and infant mental health service at Winnunga Nimmityjah	Canberra, Australia, mental health service.	Describe how the contexts of social and emotional wellbeing in Aboriginal and Torres Strait Islander communities impacts on perinatal and infant mental health. Description of a health service.	The process of starting the clinic and some clinical themes are described.	Shows the need for perinatal and infant mental health services at AHC.
Allen, 1996 [3] Anthill and other injuries: a case for mobile allied health teams to remote Australia	Far North Queensland, Australia, mobile allied health teams	Assess a mobile allied health service visiting the remote western shires of the Peninsula and Torres Strait health region. Description of a health service.	There is demand for allied health services; mobile remote outreach services are cost effective.	Multidisciplinary outreach teams are cost-effective.
Ansari, 2006 [4] Access to health care and hospitalization for ambulatory care sensitive conditions	Victoria, Australia	Examine the external validity of the indicator Hospitalization for Ambulatory Care Sensitive Conditions (ACSH). Validation of tool.	Hospitalization for ACSH is an accepted indicator of access to health care and avoidable morbidity. Results support the ACSH indicator. When rural residence was considered, the covariate measuring access was not significant. However, rural residence also may contribute importantly to access. Results suggest both the complexity of the meaning of access and the desirability of further research to validate the ACSH indicator.	The use of ACSH as an indicator of health services access and avoidable morbidity

<p>Bailie, 2003 [5] Preventive medical care in remote Aboriginal communities in the Northern Territory: a follow-up study of the impact of clinical guidelines, computerised recall and reminder systems, and audit and feedback</p>	<p>Northern Territory, Remote Aboriginal Communities</p>	<p>Describe the impact on trends in delivery of preventive medical services following a multifaceted intervention.</p> <p>Post-intervention evaluation.</p>	<p>Clinical guidelines, computerised recall and reminder systems, and audit and feedback improved service outcomes.</p>	<p>Initial improvements were “not fully sustained over the three-year period” showing that the solution is not simply providing but maintaining an evidence-base service.</p>
<p>Bailie, 2008 [6] Delivery of child health services in Indigenous communities: implications for the federal government's emergency intervention in the Northern Territory</p>	<p>NT, Australia, Aboriginal communities</p>	<p>Describe delivery of child health services in Australian Aboriginal communities, identifies gaps in services required to improve the health of Aboriginal children.</p> <p>Cross-sectional baseline audit.</p>	<p>Ineffective systems for follow-up mean that screening children for disease and adverse social circumstances will result in little or no benefit.</p>	<p>Shows existing services are not enough and that there is a great need for long term, sustainable services</p>
<p>Bar-Zeev, 2012 [7] Use of health services by remote dwelling Aboriginal infants in tropical northern Australia: a retrospective cohort study</p>	<p>NT, Australia, Aboriginal communities</p>	<p>Describe health service utilisation patterns at the primary and referral level by remote dwelling Aboriginal infants from northern Australia.</p> <p>Retrospective cohort study.</p>	<p>Hospitalisation rate is high and admissions commence early in life, visits to the remote primary health centre are frequent. Half of all presentations are for new problems.</p>	
<p>Barclay, 2014 [8] Improving Aboriginal maternal and infant health services in the 'Top End' of Australia; synthesis of the findings of a health services research program aimed at engaging stakeholders, developing research capacity and embedding change</p>	<p>Northern Territory, Australia, remote maternity services</p>	<p>Present a five-year collaborative program between stakeholders and researchers that led to sustainable improvements in the maternity services for remote-dwelling Aboriginal women and their infants in the Top End (TE) of Australia.</p> <p>Service audit.</p>	<p>System-wide problems were identified, causing poor quality of infant services: unacceptable standards of infant care and parent support; no relationship between volume and acuity of presentations and staff numbers or required skills for providing care for infants; and an 'outpatient' model of care.</p> <p>Insufficient/absent Aboriginal leadership and inadequate coordination between remote and tertiary services.</p>	<p>Strength: worked closely with NT Health throughout: NT Health requested some parts of the study and so were invested in findings.</p>

Barr, 2018 [9] A systematic review of services to DHH children in rural and remote regions	Australia-wide, services for hearing impaired	Document service provision to children with hearing loss in regional, rural and remote Australia. Systematic review.	Reduced quality and frequency of service in rural and remote areas.	Small number of relevant studies.
Breen, 2018 [10] Significant reductions in tertiary hospital encounters and less travel for families after implementation of Paediatric Care Coordination in Australia	Australia-wide, care coordination	Evaluate the impacts of Care Coordination on tertiary hospital service use and family outcomes. Cohort study.	The Care Coordination service has clear benefits for the tertiary paediatric hospital network and for families. Au\$4.9 million was saved over 2 years due to prevented hospital encounters.	Took place in very well-funded tertiary network so less relevant for rural/remote; however rural/remote often travel to tertiary centres and this program shows reduction in this travel burden.
Buckley, 2012 [11] Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit	Regional NSW, acute mental health videoconferencing	Determine if adding video link to the existing phone connection would decrease transfer to the central mental health unit. Cohort study.	The ability for the psychiatrist or senior mental health clinician to see mental health patients via videoconferencing was associated with a reduced probability of the patient being transferred. This satisfies the preference of patients to remain in their community and access mental health services.	Videoconferencing allows patients to remain in their communities whilst accessing mental health services and reduces the rate of transferrals
Chandradasa, 2019 [12] Collaborative networking between regional child mental health, paediatric and educational services in Gippsland, Australia: An online survey	Regional Victoria, Australia, survey	Survey of health professionals on utility of collaborative networking between regional child mental health, paediatric and educational services in Gippsland, Australia. Survey.	Survey found that monthly meetings between CAMHS, Department of Education, paediatricians was beneficial.	Survey of existing regular interdisciplinary meeting; positive review may be expected from those already organising this meeting. Patient/family survey to determine what matters for them would strengthen study.
Cord-Udy, 2004 [13] The Medical Specialist Outreach Assistance Programme in South Australia at 2 years	Remote SA, Australia, psychiatry outreach	Describes the progress and achievements of the psychiatric services developed by the Medical Specialist Outreach Assistance Programme in South Australia over the last 2 years. Service audit.	During this period there has been an expansion of visiting psychiatric services to South Australia's rural and remote communities. There is increased access to Child and Indigenous services.	Psych services – the benefits of specialist outreach

<p>Couzos, 2004 [14] Practical measures that improve human rights - towards health equity for Aboriginal children</p>	<p>Australia-wide</p>	<p>Detail the requirements, in terms of health policy frameworks and preventative programs, for Australian Aboriginal maternal and child health.</p> <p>Commentary.</p>	<p>Recommendations include the need for the adoption of a policy framework for maternal and child health, concomitant national performance indicators, nutritional supplementation programs, multifaceted measures to enhance the immunisation coverage of Aboriginal children and preventive child health assessments.</p>	<p>Improve health policy and programs; improve the AHS and the child and mental health framework and KPIs</p>
<p>D'Aprano, 2016 [15] Challenges in monitoring the development of young children in remote Aboriginal health services: clinical audit findings and recommendations for improving practice</p>	<p>NT, Australia, remote Aboriginal Health Centres</p>	<p>Describe developmental monitoring practice in two remote Australian Aboriginal primary healthcare services and identify gaps in the delivery of developmental monitoring services.</p> <p>Service audit.</p>	<p>Developmental checks were more likely among children who attended services more regularly. Inconsistent documentation. Need for a systems-wide approach to the delivery and recording of developmental monitoring services. This will require routine training of remote Aboriginal health workers and remote area nurses in developmental monitoring practice including the use of a culturally appropriate, structured developmental screening measure.</p>	<p>Retrospective audit with difficulty differentiating inadequate care from poor documentation; authors note that poor documentation will in itself impact continuity of care.</p>
<p>Dossetor, 1999 [16] A child and adolescent psychiatric outreach service for rural New South Wales: a telemedicine pilot study</p>	<p>Rural NSW, Australia, psychiatry telehealth outreach</p>	<p>Examines the feasibility of a tertiary outreach service in child and adolescent psychiatry to two rural health centres in New South Wales, Australia.</p> <p>Prospective cohort study.</p>	<p>Telepsychiatry provides access to a flexible, effective tertiary service for those with special, complex needs, including the disadvantaged or isolated. It makes a valuable, economic contribution to supporting and educating rural health professionals, thereby enriching rural mental health services.</p>	
<p>Dossetor, 2017 [17] Paediatric hospital admissions in Indigenous children: a population-based study in remote Australia</p>	<p>Kimberley Region, WA, Australia, rural hospital</p>	<p>Analysis of hospital admissions of a predominantly Aboriginal cohort of children in the remote Fitzroy Valley in Western Australia during the first 7 years of life.</p> <p>Descriptive, quantitative.</p>	<p>In the Fitzroy Valley 70% of children were hospitalised at least once before age 7 years and over one third of admissions were in infants. Infections were the most common reason for admission in all age groups but comorbidities were common. Many hospitalizations were feasibly preventable.</p>	<p>Misses children transferred directly to larger hospitals.</p>

Dossetor, 2019 [18] Review of Aboriginal child health services in remote Western Australia identifies challenges and informs solutions	Kimberley region, Australia, health services	Identify and map child health services in the very remote Fitzroy Valley, West Kimberley, and document barriers to effective service delivery. Service audit.	There is no document providing a comprehensive overview of child health services in Fitzroy Valley. The need to attend to acute illness acts to deprioritise crucial primary and preventative health care. Critical shortage of Aboriginal Health Workers.	Strength: produced in collaboration with local community.
Dossetor, 2021 [19] Emergency department use in a rural Australian setting: are the factors prompting attendance appropriate?	Australia, a rural hospital	Examine if the reasons for attendance at rural and remote ED are appropriate. Survey.	The results highlighted the importance of the rural hospital ED as an additional and alternate service to existing primary care, particularly outside business hours. ED was utilised for clinically non-emergent conditions, which was justified given availability to services.	Highlights the importance of services additional to PHC services in remote areas.
Edmond, 2018 [20] Quality of social and emotional wellbeing services for families of young Indigenous children attending primary care centers; a cross sectional analysis	Australia-wide, PHC	Assess delivery of social and emotional wellbeing services to the families of young (3-11 months) and older (12-59 months) Indigenous children attending primary care centres. Cross-sectional study.	The families of young Indigenous children appear to receive priority for social and emotional wellbeing care in Australian primary care centres, however many Indigenous families are not receiving services.	Auditors were clinic staff with standardised training however may be increased inter-rater variability. Participation was voluntary a possible selection bias.
Edmond, 2019 [21] Improving developmental care in primary practice for disadvantaged children	Australia-wide, PHC	Assess if sustained participation in continuous quality improvement (CQI) activities could improve delivery of 'basic developmental care' to disadvantaged children in primary care settings. Cross-sectional study.	Sustained CQI can improve basic developmental care in primary care settings. However, many disadvantaged children are not receiving services.	No interviews with patients/families; missed patients who are not presenting for care at all.
Flabouris, 2012 [22] Accessibility of the Australian population to an ICU, and of ICUs to each other	Australia-wide, ICUs	Uses a geographic information system to illustrate the geospatial relationship of the Australian population to intensive care resources. Spatial survey.	The distribution of Australian ICUs and the Australian population was similar. However, accessibility varied by state/territory.	

Francis, 2016 [23] The impact of socioeconomic status and geographic remoteness on access to pre-emptive kidney transplantation and transplant outcomes among children	Australia-wide, nephrology	Determine whether access to pre-emptive transplantation and transplant outcomes differ according to SES and geographic remoteness in Australia. Cohort study.	In Australia, children from regional or remote regions are much less likely to receive pre-emptive kidney transplantation. Strategies such as improved access to nephrology services through expanding the scope of outreach clinics, and support for regional paediatricians to promote early referral may ameliorate this inequity.	Subgroup analyses may be underpowered; used a robust database.
Garne, 2009 [24] Frequent users of the Royal Flying Doctor Service primary clinic and aeromedical services in remote New South Wales: a quality study	Rural/remote NSW, Australia, retrieval/outreach	Examine activity patterns of the Royal Flying Doctor Service of Australia (RFDS) in far western New South Wales and to determine whether frequent use of RFDS services, particularly emergency evacuations, is a useful indicator of patients who may benefit from care planning and review. Audit.	Conclusions: Simple, practical clinical review systems can help health care organisations in rural and remote communities to achieve better outcomes by identifying patients who may benefit from planned care.	
Geelhoed, 2008 [25] Positive impact of increased number of emergency consultants	Australia, Paediatric emergency department	Determine the impact of increased paediatric emergency department consultants. Cohort study.	The provision of additional consultant medical staff in a paediatric ED coincided with a decrease in the percentage of children admitted, complaints to the department and average waiting times, and was cost effective.	Increasing use of ED services decreased the proportion of kids admitted to hospital and waiting times and was cost effective.
Gruen, 2006 [26] Specialist outreach to isolated and disadvantaged communities: a population-based study	Northern Australia, 3 remote Indigenous communities	Assess the effects of outreach clinics on access, referral patterns, and care outcomes in remote communities in Australia. Cohort study.	Specialist outreach visits to remote Indigenous communities in Australia improve access to specialist consultations and procedures without increasing elective referrals or demands for hospital inpatient services.	Not paediatric-focused

<p>Gunasekera, 2009 [27] Otitis media in Aboriginal children: the discordance between burden of illness and access to services in rural/remote and urban Australia</p>	<p>All Australian Aboriginal medical services</p>	<p>Compare burden of otitis media (OM) managed by Aboriginal Medical Service (AMS) practitioners and the availability of specialist ear health services in rural/remote versus urban Australian settings.</p> <p>Survey of AMS practitioners/services.</p>	<p>Rural/Remote AMS practitioners manage a greater OM burden than urban AMS practitioners. One in five rural/remote Aboriginal children wait longer than recommended for audiology testing.</p>	<p>Shortage of health services for the burden of disease.</p>
<p>Hayman, 2010 [28] Strategies to improve indigenous access for urban and regional populations to health services</p>	<p>Queensland, PHC</p>	<p>Describe strategies to improve indigenous health services access.</p> <p>Commentary.</p>	<p>Community consultation and participation were the main ingredients to improving indigenous access to the service. All primary health care services working in Aboriginal and Torres Strait Islander health settings should have access to funded continuous quality improvement activities.</p>	<p>Not solely paediatric-focused. New data.</p>
<p>Henderson, 2008 [29] Review of community paediatrics, the Central Australian Remote Health Service, Alice Springs</p>	<p>Central Australia, outreach paediatrics</p>	<p>Review outreach community paediatric service operating out of Central Australian Outreach Remote Health Service.</p> <p>Service audit.</p>	<p>Wide-ranging findings and recommendations relating to relationship of hospital and community-based paediatric services.</p>	
<p>Humphreys, 2009 [30] Key considerations in delivering appropriate and accessible health care for rural and remote populations: discussant overview</p>	<p>Australia-wide</p>	<p>Provide an overview of papers discussing optimal service delivery models for rural and remote Australia.</p> <p>Narrative discussion.</p>	<p>Systemic change is required to ensure equitable access to health care services in small rural and remote communities.</p>	<p>Requirements for equitable access to health care should be based on:</p> <ul style="list-style-type: none"> - overarching health goals - agreed service requirements - recognition of how rural and remote health contexts impact upon health service provision - the constraints limiting health service responses

<p>Hussain, 2015 [31] Parental perceptions of information needs and service provision for children with developmental disabilities in rural Australia</p>	<p>Australia-wide, consumer advocacy</p>	<p>Give voice to parents regarding challenges caring for children with developmental disabilities, faced by them due to their location in rural regions.</p> <p>Survey, interviews.</p>	<p>Feasible options within resource constraints to improve support include focused in-service training to narrow the information gap, improve provider-client interaction around attitudinal issues, and uptake of tele-health services. Setting up of parent-professional support groups as well peer support groups using digital technologies will help reduce the sense of isolation for rural carers and minimise impediments related to travelling long distances.</p>	<p>Interviews had no indigenous participants.</p>
<p>Jeffery, 2011 [32] Responding to rural health needs through community participation: addressing the concerns of children and young adults</p>	<p>Rural Vic, Australia, health service</p>	<p>Detail a needs analysis of a small rural health service in 2008 to identify gaps in service delivery and duplication of services. This exercise was intended to inform strategic direction but the result was consumer and community consultation and outcomes that far exceeded everyone's expectations.</p> <p>Participatory Rural Appraisal.</p>	<p>The project described resulted in targeted, purposeful action regarding community engagement, and the findings and outcomes are reflective of this.</p>	<p>Partnerships between HS and communities</p>
<p>Johns, 2010 [33] Early childhood service development and intersectoral collaboration in rural Australia</p>	<p>Tasmania, Australia, in three small communities</p>	<p>Examine community-based intersectoral collaborations involving government and non-government organisations from the health and allied health, education and community service sectors.</p> <p>Qualitative study.</p>	<p>Three groups of factors influence collaborations: social capital, leadership and environmental. Internal and external leaders are needed.</p>	<p>Infant service development and function depends on: Effective models of early childhood development require strong local and external leadership; harness the skills, knowledge and resources of internal and external leaders; the ability to this empowers them to take control/ownership of health & wellbeing</p>

Jones, 2018 [34] Rural and remote speech-language pathology service inequities: An Australian human rights dilemma	Australia-wide, review	Examine lack of health services, specifically speech-language pathology with a human rights approach. Literature review. Narrative discussion.	NA	Narrative synthesis of literature
Josif, 2017 [35] The quality of health services provided to remote dwelling aboriginal infants in the top end of northern Australia following health system changes: A qualitative analysis	Top End, Australia, Aboriginal communities	Investigate service quality, from the clinicians' perspective and as observed and recorded by the researcher, in two large Aboriginal communities in the Top End of northern Australia following health system changes. Observational study.	A range of negative factors persisted beyond the examined health system change: ineffective service delivery, inadequate staffing, culturally unsafe practices. The six themes identified in the data: 'very adhoc', 'swallowed by acute', 'going under', 'a flux', 'a huge barrier' and 'them and us' illustrate how these factors continue, and when combined portray a 'very chaotic system'.	A few hours of observation in two communities brings into question the saturation and generalisability of identified themes.
Kang, 2020 [36] The relationship between having a regular general practitioner (GP) and the experience of healthcare barriers: a cross-sectional study among young people in NSW, Australia, with oversampling from marginalised groups	NSW, Australia, PHC	Explore the association between having a regular GP and experience of healthcare barriers and attitudes to health system navigation among young people in New South Wales (NSW), Australia. 'Over-sampling' of ATSI and rural/remote groups. Cross-sectional study.	Having a regular GP was associated with having more positive attitudes to health system navigation. Those with regular GP were less likely impacted by cost, other structural barriers, feeling judged, and not knowing which service to go to.	Oversampled ATSI and rural/remote; Cross-sectional design though strengthened by qualitative component.
Langbecker, 2019 [37] Impact of school-based allied health therapy via telehealth on children's speech and language, class participation and educational outcomes	QLD, Australia, schools	Examine the impact of school-based allied health therapy via telehealth on children's speech and language, class participation and educational outcomes. Observational study.	Delivering allied health therapy by videoconference may enable children to better engage with schooling. Differences in improvements by grade suggest improvements may be maximised by targeting therapy at certain year levels. This model may be useful in areas with limited access to allied health services.	Didn't compare with telehealth with standard in-person service. Didn't note if participants were ATSI, or how program could work for ATSI students

<p>Larson, 2010 [38] Aboriginal maternal and child project: strengths and needs analysis</p>	<p>WA, Australia, Epidemiological</p>	<p>Strengths and needs analysis of aboriginal maternal and child health.</p> <p>Strengths and Needs analysis.</p>	<p>Aboriginal community-controlled health services provide a value maternal health service and have the potential to contribute to effective child health care. Greater investment in the basic infrastructure necessary to deliver services – the space, funding and partnerships with other health services – is directly translated into improved quality of service provision.</p>	<p>Recommendations for models of care, communication & infrastructure, access, workforce, integration, and referral for ATSI mothers and babies.</p>
<p>Lenthall S, 2011 [39] Nursing workforce in very remote Australia, characteristics and key issues</p>	<p>Remote Australia, nursing workforce</p>	<p>Describe the nursing workforce in very remote Australia, characteristics, and key issues.</p> <p>Descriptive and narrative discussion.</p>	<p>The workforce is ageing, the numbers of nurses per population has fallen and the numbers of midwives and child health nurses have dropped significantly over the last 15 years. As many of these nurses work in Indigenous communities, if these trends continue it is likely to have a negative effect on 'closing the gap' in Indigenous health outcomes.</p>	<p>Decrease in nursing workforce particularly remote, paediatric nurses</p>
<p>Margolis, 2012 [40] Is Fly in/Fly out (FIFO) a viable interim solution to address remote medical workforce shortages?</p>	<p>Remote Australia</p>	<p>Describe Fly In Fly Out (FIFO) viability, anecdotal synthesis.</p> <p>Narrative discussion.</p>	<p>Effective FIFO needs commitment from the sponsoring organisation for short, balanced, flexible, family friendly rosters. A positive organizational structure with effective communication between management and front-line staff is also required. FIFO can capitalise on mid-career doctors who are experienced in remote work but have moved to cities.</p>	<p>Advocates for the FIFO model – to address rural health service workforce shortages.</p>
<p>McCalman, 2012 [41] Applying what works: a systematic search of the transfer and implementation of promising Indigenous Australian health services and programs</p>	<p>Australia-wide</p>	<p>Evaluate transfer and implementation of Indigenous health services.</p> <p>Systematic review.</p>	<p>Few studies focus on the process of transfer and implementation or the effectiveness of the service or program in the new setting.</p>	<p>Once implemented, services aren't being assessed on their effectiveness. This requires active partnerships between services and researchers.</p>

McKeown, 2011 [42] Evaluating aboriginal primary health care services using national key performance indicators: What has happened to the social theories of aboriginal health?	Australia-wide, primary health care (PHC)	Examine whether KPIs focused on process and outcome of clinical service delivery are appropriate to Aboriginal-controlled PHCs? Commentary.	Evaluation frameworks for Aboriginal PHC services need to encompass wider PHC frameworks rather than a narrower biomedical approach to improving health.	Suggests that using the KPI to evaluate Aboriginal PHC services won't provide an accurate reflection of their contribution to improving health.
Medlin, 2014 [43] Indigenous respiratory outreach care (IROC): Enhancing respiratory health in rural and remote aboriginal and Torres strait islander communities	QLD, Australia, specialist outreach	Assess the Indigenous Respiratory Outreach (nurse-led) Care (IROC) (adults and paediatrics) in rural/remote ATSI. Program evaluation Cohort study.	IROC started in 2011, now 12 community sites. Early data shows communities like it and it has picked up previously undetected paediatric chronic lung disease. IROC provides a successful model for achieving this goal that is readily transferrable to other regions in Australia.	Preliminary data
Mitchell, 2018 [44] Transition to adult care for Aboriginal children with rheumatic fever: a review informed by a focussed ethnography in northern Australia	Northern Territory, Australia, remote communities	Describe transition to adult care for children with chronic conditions and consider applicability to the care of children. Narrative review, ethnographic study of four remote communities.	Recommended tailoring of transition care to engage and value local navigators who can address language and cultural barriers to provide a sustainable alternative to transition coordinators in mainstream programs. This has potential to improve care without further burdening overstretched clinical resources.	Explicitly took account of power imbalances. Assessed an evidenced program in new setting
Mitchinson, 2019 [45] Anaemia in disadvantaged children aged under five years; quality of care in primary practice	Australia-wide, PHC	Assess quality of care of anaemic disadvantaged children. Cross-sectional study.	The burden of anaemia and quality of care for disadvantaged Indigenous children was concerning across all remote and urban locations assessed in this study.	Self-selected study group may limit generalisability; Only 70% children screened for anaemia, remainder may be more severe (differential error)

Moffatt, 2010 [46] The reported benefits of telehealth for rural Australians	Australia-wide	Identify reported benefits attributed to telehealth for people living and professionals working in rural and remote areas of Australia. Narrative review.	Telehealth may decrease urban-rural health disparities and improve workforce retention. Benefits of telehealth for patients: cheaper, convenient, better access, better quality; for clinicians: educational opportunities, experiential learning, networking, collaboration.	
Morgan, 2012 [47] QT babies: Neonatal care in the Queensland tropics	Far North QLD, Australia	Review the requirements of this diverse and geographically isolated population and how they are met by the team at The Townsville Hospital. Narrative discussion.	The challenge of providing neonatal care to this unique population continues to be successfully achieved by the dedicated extended multidisciplinary team at The Townsville Hospital.	Neonatal care is successfully achieved at the Townsville hospital ATSI, shows the success of a multidisciplinary team
Nancarrow, 2015 [48] Models of care involving district hospitals: a rapid review to inform the Australian rural and remote context	Australia-wide	Identify models of care that incorporate district hospitals and have relevance to the Australian rural and remote context. Service audit.	District hospitals are important symbolically and functionally for small towns. They assist in vertical integration of health services.	Study commissioned for policy development. Noted that literature focused on rural health systems more broadly, not the target of review: 'district hospitals'
Nguyen, 2015 [49] Cost-effectiveness analysis of a mobile ear screening and surveillance service versus an outreach screening, surveillance and surgical service for Indigenous children in Australia	Rural Queensland, Australia, ENT mobile outreach	Evaluate the cost-effectiveness of the community-based mobile telemedicine-enabled screening and surveillance (MTESS) service using a lifetime Markov model that compares two options: (i) the Deadly Ears Program alone (current practice involving an outreach ENT surgical service and screening program), and (ii) the Deadly Ears Program supplemented with the MTESS service. Economic analysis.	The MTESS service is cost-effective. It presents an opportunity to resolve major issues confronting Australia's health system such as the inequitable provision and access to quality healthcare for rural and remote communities, and for Indigenous Australians.	Modelling relied on nascent epidemiology of ear disease in Indigenous People.
Nguyen, 2018 [50] Feasibility of Implementing Infant Home Visiting in a Central Australian Aboriginal Community	Central Australia, home visits	Assess the feasibility of Implementing Infant Home Visiting in a Central Australian Aboriginal Community (as is/was done in Alice Springs since 2009). Descriptive, cohort study.	This study finds it could be done; needs outcome evaluation to test hypothesised benefits.	

O'Callaghan, 2005 [51] Consumers' proposed solutions to barriers to access of rural and remote speech pathology services	Remote NSW, Australia, paediatrics speech pathology	Investigate solutions to perceived barriers of consumers when accessing paediatric speech pathology services in rural and remote New South Wales (NSW). Survey design.	Consumer-based solutions to barriers to access to maximize the effectiveness of sparse rural and remote speech pathology services, by matching the beliefs and expectations of consumers with the characteristics of services provided.	Barriers to services – allied health, speech paths.
O'Kane, 2004 [52] Towards a needs based mental health resource allocation and service development in rural and remote Australia	Central Australia, Remote mental health services	Develop a transparent, needs- and evidence-based framework to guide resource allocation and mental health service development in rural and remote settings. Review of services.	Existing funding allocation was half that required for adequate, equitable level of care.	Demonstrates great need for mental services, particularly for children, adolescents and remote-dwelling Aboriginal people.
Ou, 2010 [53] The comparison of health status and health services utilisation between Indigenous and non-Indigenous infants in Australia	Australia-wide, LSAC analysis	Describe differences in health services utilisation and the associated risk factors between Indigenous and non-Indigenous infants at a national level in Australia. Analysis of a national longitudinal dataset.	Indigenous infants use less healthcare than non-Indigenous. Determining factors include mothers' characteristics and socio-economic status.	ATSI children are more likely to use a hospital outpatient clinic than: maternal & child health centre, help lines, maternal and child health nurse visits, GP, or paediatrician
Peiris, 2006 [54] Aeromedical evacuations from an east Arnhem Land community 2003-2005: The impact on a primary health care centre	Arnhem Land, Australia, PHC evacuations	Describe the profile and impact of aeromedical evacuations by Royal Flying Doctor Service in remote Indigenous communities. Service audit.	Aeromedical evacuations burden primary health centres. Meeting this burden competes with regular clinic duties. Clinic staff are regularly required to provide hospital-level acute care, often for several hours at a time.	High service usage but is expensive and could be decreased by improving availability of remote area health centres.

<p>Phillips, 2014 [55] Can mobile phone multimedia messages and text messages improve clinic attendance for Aboriginal children with chronic otitis media? A randomised controlled trial</p>	<p>Remote NT, Australia, community</p>	<p>Examine effect of phone multimedia messages (MMS) to families of Indigenous children with tympanic membrane perforation (TMP): (i) increase clinic attendance; (ii) improve ear health; and (iii) provide a culturally appropriate method of health promotion?</p> <p>Randomised-controlled trial.</p>	<p>No improvement in clinic attendance or ear health. But this study shows that MMS in local languages can be culturally appropriate.</p>	<p>Affected by social conditions: didn't own, or couldn't regularly access mobile phone</p>
<p>Ruben, 1998 [56] The case mix system of hospital funding can further disadvantage Aboriginal children</p>	<p>NT, Australia, funding model.</p>	<p>Examine effect of mixed hospital funding using national averages and national cost weights as benchmarks for length of stay and funding.</p> <p>Retrospective cohort study.</p>	<p>ATSI children had more comorbidities, and prolonged hospital stays. These results confirm clinical impressions about disease patterns and length of hospital stay in ATSI children, and highlight the problems of imposing a case mix classification system for a "typical" Australian population on a region with a high proportion of people of ATSI descent.</p>	<p>Explores problems associated with case mix system of hospital funding for complex patients.</p>
<p>Schofield, 2009 [57] Multidisciplinary management of complex care</p>	<p>Northern New South Wales, Australia, in community</p>	<p>Describe the advantages of multidisciplinary care in terms of patient outcomes, clinician satisfaction and system efficiency are considered with reference to an innovative rural multidisciplinary model that highlights how these positive outcomes can be achieved.</p> <p>Pre and post descriptive study.</p>	<p>Multidisciplinary care can provide better health outcomes and a more satisfying work environment for health professionals. Especially in rural areas where the only colleagues may be from other disciplines.</p>	

<p>Seidler, 2020 [58] Patterns of Youth Mental Health Service Use and Discontinuation: Population Data from Australia's Headspace Model of Care</p>	<p>Australia-wide, mental health services.</p>	<p>Explore sociodemographic predictors of attendance and discontinuation of mental health services in a large, population-based sample.</p> <p>Cohort study.</p>	<p>Sociodemographic factors were found to be associated with treatment discontinuation, and some young people followed a pathway in and out of mental health treatment. Further exploration is needed to determine the appropriate length and type of care for specific socio-demographic groups and how best to tailor treatment accordingly.</p>	<p>Used existing dataset (minimum data set; MDS).</p>
<p>Smith, 2010 [59] Clinical services and professional support: a review of mobile telepaediatric services in Queensland</p>	<p>Rural and remote QLD, Australia, health services</p>	<p>Review of telepaediatric service records to determine which clinical and educational services had been delivered through the mobile videoconference systems.</p> <p>Narrative review.</p>	<p>Our service model provided a streamlined method of delivering specialist health services to children and families living in rural and remote regions of Queensland.</p>	
<p>Thomas, 2014 [60] What core primary health care services should be available to Australians living in rural and remote communities?</p>	<p>Australia-wide</p>	<p>Describe what core primary health care services should be available to rural/remote Australians.</p> <p>Delphi survey of remote health experts including policy-makers, academics, clinicians and consumers.</p>	<p>This study identifies a basket of PHC services that consumers in rural and remote communities can expect to access. It provides rigorously derived evidence that will contribute to a more systematic approach to PHC service planning and availability and will assist policy makers in the allocation of scarce resources necessary to improve the health outcomes of residents of rural and remote areas.</p>	<p>Only 4 (10%) Delphi panellists were from community ('consumer representatives').</p>
<p>Veit, 1996 [61] Barriers to effective primary health care for adolescents</p>	<p>Victorian GPs</p>	<p>Describe general practitioners' (GPs) perceptions of barriers in the health care system that hinder provision of effective adolescent health care; and assess the training needs of GPs for a future education program in adolescent health.</p> <p>Survey of GPs.</p>	<p>Medicare cards should be automatically issued to adolescents from age 16 years to improve their access to health care. Revision of Medicare rebate system and clarification of HIC's investigative functions may improve GP's effectiveness.</p>	

Verdon, 2011 [62] An investigation of equity of rural speech-language pathology services for children: a geographic perspective	NSW and Victoria, Australia, speech pathologists	Assess speech pathology services in rural areas of New South Wales and Victoria/ Structured interviews with 74 speech-language pathologists.	Almost all (98.60%) localities are underserved. Fifty kilometres is the critical maximum distance past which consumers become unable or unwilling to travel to access weekly rural speech-language pathology services in rural New South Wales and Victoria.	Speech pathology may be not representative of wide access.
Wakeman, 2008 [63] Primary health care delivery models in rural and remote Australia: a systematic review	Australia-wide	Describe what health service models were reported to work, where they worked and why. Systematic review.	Sustainable models may address diseconomies of scale which result from large distances and small dispersed populations. The authors developed a conceptual framework for development of sustainable remote PHC models.	Systematic review of PHC models in remote Australia enabled development of a conceptual framework to help in the sustainability of these PHC models.
Wakeman, 2009 [64] Innovative rural and remote primary health care models: what do we know and what are the research priorities?	Australia-wide	Examine literature on 'innovative' primary health care models in rural and remote areas to identify areas where knowledge is lacking and describes future research priorities. Narrative synthesis.	Health services evaluation should focus on: optimal financing systems, the optimal range and mix of providers, supports for team practice, appropriate community participation mechanisms, improved health information systems and relevant performance indicators.	Need for evaluation of services to inform best practice.
Wakeman, 2009 [65] Features of effective primary health care models in rural and remote Australia: a case-study analysis	Australia-wide, primary health care	Describe the factors and processes that facilitate or inhibit implementation, sustainability, and generalisation of effective models of primary health care (PHC) service delivery in rural and remote Australia. Literature review, interviews, direct observation.	In Australia, establishment of a single national health system, operationalised at a regional level, would obviate much of the current inefficiency and poor coordination.	Highlights need for improved: <ul style="list-style-type: none"> - governance - management - community involvement

Ward, 2006 [66] Correlations of siblings' and mothers' utilisation of primary and hospital health care: A record linkage study in Western Australia	WA, Australia	Examine the relationships between the utilisation of health care by siblings and mothers over 1984-97. Observational study.	Strong relationship between siblings' use of GPs and a weaker but still significant association between the hospital admissions of siblings. Suggests the presence of intergenerational correlation of morbidity; that behavioural factors are not sufficiently explanatory.	Correlation between mothers and children, siblings for GP and Hospital service usage.
Warwick, 2019 [67] Young Aboriginal People's Perspective on Access to Health Care in Remote Australia: Hearing Their Voices	Kimberley region, WA, Australia, Aboriginal Health Service	Identify barriers and enablers of access for adolescent and young adult Aboriginal people to Derby Aboriginal Health Service (DAHS), a remote Western Australian Aboriginal Community Controlled Health Service (ACCHSs), to assist in improving access. Qualitative study.	Improving youth engagement seems to be central to increasing acceptability and, hence, use. This requires that staff able to engage with young people are recruited, trained, and retained.	Respondents were patients/consumers. Used local researchers to collect data however researchers were also health service staff which may pressure respondents to reply favourably and introduce bias into results.
Whelan, 2013 [68] Health Services Use and Lifestyle Choices of Indigenous and non-Indigenous Australians	Australia-wide	Analyse national data to examine whether Indigenous Australians make different lifestyle choices and health services use than non-Indigenous Australians. Observational database study.	Indigenous Australians are more likely to make poorer lifestyle choices but are more likely to use health services. These results are magnified for Indigenous Australians who live in remote areas.	Focuses on adults.
Williams, 2004 [69] Paediatric outreach services	Australia-wide	Describe Paediatric outreach services. Narrative review.	A combination of telehealth clinics and outreach visits may be most efficient for delivering paediatric specialty and subspecialty care to remote Australia.	Synthesis but nil new data.

<p>Zwi, 2011 [70] Prioritising health services for high risk and disadvantaged children</p>	<p>Australia wide</p>	<p>Document prioritisation strategies for health services for vulnerable children, map current prioritisation methods in an urban area health service, and analyse the views of service directors, clinicians, and intake workers regarding the ethics and practicalities of prioritising care.</p> <p>Narrative review.</p>	<p>Health staff favour prioritisation strategies for vulnerable children. Prioritisation strategies can burden staff and so require extra resources.</p>	
--	-----------------------	--	--	--

Table 1b: Source of grey literature

Source	Overview	Reports used
A Country Health Service: Kimberley Health	Western Australia rural health organisation	- Aboriginal Health Profile: Kimberley Health Region, 2012 [71] - Kimberley Health Profile 2018 [72] - Child Population Profile, Health and Wellbeing Surveillance System, 2009: Kimberley health region [73]
Australian Indigenous Health Infonet	Aims to make knowledge available to health organisations	- Overview of Aboriginal and Torres Strait Islander health status, 2020 [74] - Summary of Aboriginal and Torres Strait Islander health status 2018 [75]
Australian Institute of Health and Welfare	Reports relating to health workforce stats, expenditure, Aboriginal community-controlled health services, ATSI health performance framework reports for each state and territory. Majority national level data analysis, some state level comparative	- Expenditure on health for Aboriginal and Torres Strait Islander people 2008-09: an analysis by remoteness and disease [76] - Access to health services for Aboriginal and Torres Strait Islander people [77] - Aboriginal and Torres Strait Islander health services report, 2010-11: OATSIH services reporting [78] - Healthy for Life—Aboriginal Community Controlled Health Services: report card [79] - Aboriginal and Torres Strait Islander Health Performance Framework 2013: Western Australia [80] - Health workforce 2020 [81]
Australian Medical Association	Peak body for medical professionals in Australia	- Ending the Cycle of Vulnerability: The Health of Indigenous Children [82]
Australian Primary Health		- A systematic review of primary health care delivery models in rural and remote Australia 1993 -2006 [83] - Interventions to prevent growth faltering in remote Indigenous communities [84]
Boab Health	Includes the Kimberley Division of General Practice	- Kimberley Primary Health Care Sustainability Study 2008 – 2030 [85]
Centre for Aboriginal Economic Policy Research	Think tank based at Australian National University	- Human capabilities and child development: Some inferences using the 2008 NATSISS [86] - An ethnography of changes in child rearing over time in the Ngaanyatjarra Lands: Implications for policy development on health in early childhood [87]
Commonwealth Department of Families, Housing, Community Services and Indigenous Affairs	National government department	- Indigenous reform 2011-12: comparing performance across Australia [88]
Commonwealth Department of Health and Ageing	National government department	- Primary health care in rural and remote Australia: achieving equity of access and outcomes through national reform: A discussion paper [89] - Evaluation of the Child Health Check Initiative and the Expanding Health Service Delivery Initiative: Summary Report [90]
Commonwealth House of Representatives Parliamentary Committees	Indigenous health reported in across several Parliamentary Committee Inquiries	- Health is life: inquiry into Indigenous health [91] - Inquiry into Health Funding Report: The Blame Game [92]
Commonwealth Senate Committee	Senate Select Committee on Regional and Remote Indigenous Communities	- Indigenous Australians, Incarceration and the Criminal Justice System [93]

Health Workforce Australia	National research and policy development relating to the development of a sustainable health workforce across Australia	<ul style="list-style-type: none"> - National Rural and Remote Workforce Innovation and Reform Strategy [94] - Health Workforce 2025 – Doctors, Nurses and Midwives – Volume 2 [95] - Growing Our Future: the Aboriginal and Torres Strait Islander Health Worker Project Final Report [96]
KAMSC – Kimberley Aboriginal Medical Services Council	Aboriginal controlled coordination group	<ul style="list-style-type: none"> - Regional Aboriginal Health Plan: Aboriginal Health in the Kimberley: current circumstances and future directions [97] - Kimberley Aboriginal Primary Health Plan 2012-2015 [98]
Lowitja Institute	Aboriginal organisation promoting research and knowledge translation	<ul style="list-style-type: none"> - Assessing Cost-Effectiveness in Prevention (ACE-Prevention): Final Report. [99] - Australian Indigenous Health - Within an International Context [100] - An evaluation of the benefits of swimming pools for the hearing and ear health status of young Indigenous Australians: a whole-of-population study across multiple remote Indigenous communities [101] - Leadership and legacy through crises: keeping our mob safe Close the Gap campaign report 2021 [102]
Menzies School of Health Research	Research organisation in the Northern Territory	<ul style="list-style-type: none"> - Effective Integration of Services for Children and Families: Making it Happen [103]
National Rural Health Conference 2009	Run through the National Rural Health Alliance	<ul style="list-style-type: none"> - Pitstop: one stop community health shop for Indigenous families in the West Pilbara [104]
Northern Territory Department of Health and Community Services Digital library	Many policy documents relating to child health, and material for health staff to draw on	<ul style="list-style-type: none"> - Child Malnutrition / Failure to Thrive Project 2008: Final Report [105]
Office for Aboriginal and Torres Strait Islander Health	Produces information on performance of States and Territories vis-à-vis Closing the Gap targets, Clinical guidelines, and Health programs	<ul style="list-style-type: none"> - Improving Health in Aboriginal and Torres Strait Islander Mothers, Babies and Young Children - A Literature Review [106] - Evaluation of the Child Health Check Initiative and the Expanding Health Service Delivery Initiative: Summary Report [90] - The link between primary health care and health outcomes for Aboriginal and Torres Strait Islander Australians [107] - Better Health Care: Studies in the successful delivery of primary health care services for Aboriginal and Torres Strait Islander Australians [108]
Queensland Health	State health department	<ul style="list-style-type: none"> - Making tracks: Toward closing the gap in health outcomes for Indigenous Queenslanders by 2033 [109]
Royal Flying Doctor Service	Health charity specialist in medical retrieval and remote primary care	<ul style="list-style-type: none"> - Annual report 2011/12 - service provision at a local level, relevant to rural/remote [110] - Looking Ahead: Responding to the health needs of country Australians in 2028/ the centenary year of the RFDS [111]
Rural and Regional Health Australia	www.ruralhealthaustralia.gov.au/inter-net/rha/publishing.nsf/Content/NSFR_RH-HealthServices (National Strategic framework for rural and remote health)	<ul style="list-style-type: none"> - National Strategic framework for rural and remote health [112]
Rural health west	Contains factsheets relating to, for example, distribution of particular	<ul style="list-style-type: none"> - Specialist Services in Rural Western Australia: Gap and Equity [113]

	health professionals in WA, useful for planning at a state level	
Telethon Institute for Child Health Research	Research institute in Western Australia	- The Western Australian Aboriginal Child Health Survey: The Health of Aboriginal Children and Young People [114]
World Health Organisation	International standard setting	- Toolkit on monitoring health systems strengthening: Service Delivery [115]

References for tables in Appendix

1. Achat HM, Thomas P, Close GR, Moerkerken LR, Harris MF. General health care service utilisation: where, when and by whom in a socioeconomically disadvantaged population. *Australian Journal of Primary Health*. 2010;16(2):132-40.
2. Adams E, Tongs J. Starting a perinatal and infant mental health service at Winnunga Nimmityjah. *Australasian Psychiatry*. 2011;19:S20-S2.
3. Allen O. Anthill and other injuries: a case for mobile allied health teams to remote Australia. *Australian Journal of Rural Health*. 1996;4(1):33-42.
4. Ansari Z, Laditka JN, Laditka SB. Access to health care and hospitalization for ambulatory care sensitive conditions. *Med Care Res Rev*. 2006;63(6):719-41. doi: 10.1177/1077558706293637.
5. Bailie RS, Togni SJ, Si D, Robinson G, d'Abbs PH. Preventive medical care in remote Aboriginal communities in the Northern Territory: a follow-up study of the impact of clinical guidelines, computerised recall and reminder systems, and audit and feedback. *BMC health services research*. 2003;3(1):15. doi: 10.1186/1472-6963-3-15.
6. Bailie RS, Si D, Dowden MC, Connors CM, O'Donoghue L, Liddle HE, et al. Delivery of child health services in Indigenous communities: implications for the federal government's emergency intervention in the Northern Territory. *Med J Aust*. 2008;188(10):615-8. Epub 2008/05/20. doi: 10.5694/j.1326-5377.2008.tb01806.x. PubMed PMID: 18484941.
7. Bar-Zeev SJ, Kruske SG, Barclay LM, Bar-Zeev NH, Carapetis JR, Kildea SV. Use of health services by remote dwelling Aboriginal infants in tropical northern Australia: a retrospective cohort study. *BMC Pediatrics*. 2012;12:19. doi: 10.1186/1471-2431-12-19.
8. Barclay L, Kruske S, Bar-Zeev S, Steenkamp M, Josif C, Narjic CW, et al. Improving Aboriginal maternal and infant health services in the 'Top End' of Australia; synthesis of the findings of a health services research program aimed at engaging stakeholders, developing research capacity and embedding change. *BMC health services research*. 2014;14(1):241. doi: 10.1186/1472-6963-14-241.
9. Barr M, Duncan J, Dally K. A systematic review of services to DHH children in rural and remote regions. *The Journal of Deaf Studies and Deaf Education*. 2018;23(2):118-30.
10. Breen C, Altman L, Ging J, Deverell M, Woolfenden S, Zurynski Y. Significant reductions in tertiary hospital encounters and less travel for families after implementation of Paediatric Care Coordination in Australia. *BMC health services research*. 2018;18(1):1-10.
11. Buckley D, Weisser S. Videoconferencing could reduce the number of mental health patients transferred from outlying facilities to a regional mental health unit. *Australian and New Zealand Journal of Public Health*. 2012;36(5):478-82. doi: 10.1111/j.1753-6405.2012.00915.x.
12. Chandradasa M, Basu S. Collaborative networking between regional child mental health, paediatric and educational services in Gippsland, Australia: An online survey. *The Australian Journal of Rural Health*. 2019.
13. Cord-Udy N. The Medical Specialist Outreach Assistance Programme in South Australia at 2 years. *Australasian Psychiatry*. 2004;12(2):161-5.
14. Couzos S. Practical measures that improve human rights - towards health equity for Aboriginal children. *Health Promotion Journal of Australia : Official Journal of Australian Association of Health Promotion Professionals*. 2004;15(3):186-92.

15. D'Aprano A, Silburn S, Johnston V, Bailie R, Mensah F, Oberklaid F, et al. Challenges in monitoring the development of young children in remote Aboriginal health services: clinical audit findings and recommendations for improving practice. *Rural Remote Health*. 2016;16(3):3852.
16. Dossetor DR, Nunn KP, Fairley M, Eggleton D. A child and adolescent psychiatric outreach service for rural New South Wales: a telemedicine pilot study. *J Paediatr Child Health*. 1999;35(6):525-9. doi: 10.1046/j.1440-1754.1999.00410.x.
17. Dossetor PJ, Martiniuk ALC, Fitzpatrick JP, Oscar J, Carter M, Watkins R, et al. Pediatric hospital admissions in Indigenous children: a population-based study in remote Australia. *BMC Pediatrics*. 2017;17(1):195. doi: 10.1186/s12887-017-0947-0.
18. Dossetor PJ, Thorburn K, Oscar J, Carter M, Fitzpatrick J, Bower C, et al. Review of Aboriginal child health services in remote Western Australia identifies challenges and informs solutions. *BMC health services research*. 2019;19(1):758. doi: 10.1186/s12913-019-4605-0.
19. Dossetor PJ, Fitzpatrick EFM, Glass K, Douglas K, Watkins R, Oscar J, et al. Emergency Department Presentations by Children in Remote Australia: A Population-based Study. *Global Pediatric Health*. 2021;8. doi: 10.1177/2333794X21991006.
20. Edmond KM, McAuley K, McAullay D, Matthews V, Strobel N, Marriott R, et al. Quality of social and emotional wellbeing services for families of young Indigenous children attending primary care centers; a cross sectional analysis. *BMC health services research*. 2018;18(1):1-11.
21. Edmond KM, Tung S, McAuley K, Strobel N, McAullay D. Improving developmental care in primary practice for disadvantaged children. *Archives of Disease in Childhood*. 2019;104(4):372-80.
22. Flabouris A, Hart GK, Nicholls A. Accessibility of the Australian population to an ICU, and of ICUs to each other. *Critical care and resuscitation : journal of the Australasian Academy of Critical Care Medicine*. 2012;14(3):177-84.
23. Francis A, Didsbury M, Lim WH, Kim S, White S, Craig JC, et al. The impact of socioeconomic status and geographic remoteness on access to pre-emptive kidney transplantation and transplant outcomes among children. *Pediatric Nephrology*. 2016;31(6):1011-9.
24. Garne DL, Perkins DA, Boreland FT, Lyle DM. Frequent users of the Royal Flying Doctor Service primary clinic and aeromedical services in remote New South Wales: a quality study. *Med J Aust*. 2009;191(11-12):602-4. doi: 10.5694/j.1326-5377.2009.tb03344.x.
25. Geelhoed GC, Geelhoed EA. Positive impact of increased number of emergency consultants. *Archives of disease in childhood*. 2008;93(1):62-4.
26. Gruen RL, Bailie RS, Wang Z, Heard S, O'Rourke IC. Specialist outreach to isolated and disadvantaged communities: a population-based study. *Lancet*. 2006;368(9530):130-8. doi: 10.1016/S0140-6736(06)68812-0.
27. Gunasekera H, Morris PS, Daniels J, Couzos S, Craig JC. Otitis media in Aboriginal children: the discordance between burden of illness and access to services in rural/remote and urban Australia. *Journal of Paediatrics and Child Health*. 2009;45(7-8):425-30.
28. Hayman N. Strategies to improve indigenous access for urban and regional populations to health services. *Heart Lung Circulation*. 2010;19(5-6):367-71. doi: 10.1016/j.hlc.2010.02.014.
29. Henderson R. Review of community paediatrics, the Central Australian Remote Health Service, Alice Springs. Prince Margaret Hospital for Children, Perth WA, 2008.

30. Humphreys JS. Key considerations in delivering appropriate and accessible health care for rural and remote populations: discussant overview. *Australian Journal of Rural Health*. 2009;17(1):34-8. doi: 10.1111/j.1440-1584.2008.01034.x.
31. Hussain R, Tait K. Parental perceptions of information needs and service provision for children with developmental disabilities in rural Australia. *Disability and rehabilitation*. 2015;37(18):1609-16.
32. Jeffery V, Ervin K. Responding to rural health needs through community participation: addressing the concerns of children and young adults. *Australian Journal of Primary Health*. 2011;17(2):125-30. doi: 10.1071/PY10050.
33. Johns S. Early childhood service development and intersectoral collaboration in rural Australia. *Australian Journal of Primary Health*. 2010;16(1):40-6.
34. Jones DM, McAllister L, Lyle DM. Rural and remote speech-language pathology service inequities: An Australian human rights dilemma. *International journal of speech-language pathology*. 2018;20(1):98-101.
35. Josif CM, Kruske S, Kildea SV, Barclay LM. The quality of health services provided to remote dwelling aboriginal infants in the top end of northern Australia following health system changes: a qualitative analysis. *BMC Pediatrics*. 2017;17(1):93. doi: 10.1186/s12887-017-0849-1.
36. Kang M, Robards F, Luscombe G, Sancu L, Usherwood T. The relationship between having a regular general practitioner (GP) and the experience of healthcare barriers: a cross-sectional study among young people in NSW, Australia, with oversampling from marginalised groups. *BMC family practice*. 2020;21(1):1-9.
37. Langbecker DH, Caffery L, Taylor M, Theodoros D, Smith AC. Impact of school-based allied health therapy via telehealth on children's speech and language, class participation and educational outcomes. *Journal of Telemedicine and Telecare*. 2019;25(9):559-65. doi: 10.1177/1357633X19875848.
38. Larson A, Bradley R. Aboriginal maternal and child project: strengths and needs analysis. Geraldton, WA: Combined Universities Centre for Rural Health and Aboriginal Health Council of Western Australia, 2010.
39. Lenthall S, Wakerman J, Opie T, Dunn S, Macleod M, Dollard M, et al. Nursing workforce in very remote Australia, characteristics and key issues. *The Australian Journal of Rural Health*. 2011;19(1):32-7.
40. Margolis SA. Is Fly in/Fly out (FIFO) a viable interim solution to address remote medical workforce shortages? *Rural and Remote Health*. 2012;12(4):1-6.
41. McCalman J, Tsey K, Clifford A, Earles W, Shakeshaft A, Bainbridge R. Applying what works: a systematic search of the transfer and implementation of promising Indigenous Australian health services and programs. *BMC Public Health*. 2012;12(1):600.
42. McKeown S. Evaluating aboriginal primary health care services using national key performance indicators: What has happened to the social theories of aboriginal health? *Internal Medicine Journal*. 2011;41:37.
43. Medlin L, editor *Indigenous Respiratory Outreach Care (IROC): Enhancing Respiratory Health In Rural And Remote Aboriginal And Torres Strait Islander Communities*. *Respirology*; 2014.
44. Mitchell AG, Belton S, Johnston V, Ralph AP. Transition to adult care for Aboriginal children with rheumatic fever: a review informed by a focussed ethnography in northern Australia. *Australian Journal of Primary Health*. 2018;24(1):9-13. doi: 10.1071/PY17069.

45. Mitchinson C, Strobel N, McAullay D, McAuley K, Bailie R, Edmond KM. Anemia in disadvantaged children aged under five years; quality of care in primary practice. *BMC pediatrics*. 2019;19(1):1-11.
46. Moffatt JJ, Eley DS. The reported benefits of telehealth for rural Australians. *Australian Health Review*. 2010;34(3):276-81. doi: 10.1071/AH09794.
47. Morgan K. QT babies: Neonatal care in the Queensland tropics. *Journal of Paediatrics and Child Health*. 2012.
48. Nancarrow SA, Roots A, Grace S, Saberi V. Models of care involving district hospitals: a rapid review to inform the Australian rural and remote context. *Australian Health Review*. 2015;39(5):494-507.
49. Nguyen KH, Smith AC, Armfield NR, Bensink M, Scuffham PA. Cost-Effectiveness Analysis of a Mobile Ear Screening and Surveillance Service versus an Outreach Screening, Surveillance and Surgical Service for Indigenous Children in Australia. *PLoS One*. 2015;10(9). doi: 10.1371/journal.pone.0138369.
50. Nguyen H, Zarnowiecki D, Segal L, Gent D, Silver B, Boffa J. Feasibility of implementing infant home visiting in a Central Australian Aboriginal community. *Prevention Science*. 2018;19(7):966-76.
51. O'Callaghan AM, McCallister L, Wilson L. Consumers' proposed solutions to barriers to access of rural and remote speech pathology services. *Advances in Speech Language Pathology*. 2005;7(2):58-64.
52. O'Kane A, Tsey K. Towards a needs based mental health resource allocation and service development in rural and remote Australia. *Australas Psychiatry*. 2004;12(4):390-5. doi: 10.1080/j.1440-1665.2004.02133.x.
53. Ou L, Chen J, Hillman K, Eastwood J. The comparison of health status and health services utilisation between Indigenous and non-Indigenous infants in Australia. *Australian and New Zealand Journal of Public Health*. 2010;34(1):50-6.
54. Peiris D, Wirtanen C, Hall J. Aeromedical evacuations from an east Arnhem Land community 2003-2005: the impact on a primary health care centre. *Aust J Rural Health*. 2006;14(6):270-4. doi: 10.1111/j.1440-1584.2006.00828.x.
55. Phillips JH, Wigger C, Beissbarth J, McCallum GB, Leach A, Morris PS. Can mobile phone multimedia messages and text messages improve clinic attendance for Aboriginal children with chronic otitis media? A randomised controlled trial. *Journal of Paediatrics and Child Health*. 2014;50(5):362-7.
56. Ruben AR, Fisher DA. The casemix system of hospital funding can further disadvantage Aboriginal children. *Medical Journal of Australia*. 1998;169(SUPPL.):S6-S10.
57. Schofield D, Fuller J, Wagner S, Friis L, Tyrell B. Multidisciplinary management of complex care. *Australian Journal of Rural Health*. 2009;17(1):45-8.
58. Seidler ZE, Rice SM, Dhillon HM, Cotton SM, Telford NR, McEachran J, et al. Patterns of youth mental health service use and discontinuation: Population data from Australia's headspace model of care. *Psychiatric services*. 2020;71(11):1104-13.
59. Smith AC, Armfield NR, White MM, Williams ML, Koh T, Hurley T, et al. Clinical services and professional support: a review of mobile telepaediatric services in Queensland. *Studies in Health Technology and Informatics*. 2010;161:149-58.

60. Thomas SL, Wakerman J, Humphreys JS. What core primary health care services should be available to Australians living in rural and remote communities? *BMC Family Practice*. 2014;15:143. doi: 10.1186/1471-2296-15-143.
61. Veit FC, Sancu LA, Coffey CM, Young DY, Bowes G. Barriers to effective primary health care for adolescents. *Medical Journal of Australia*. 1996;165(3):131-3.
62. Verdon S, Wilson L, Smith-Tamaray M, McAllister L. An investigation of equity of rural speech-language pathology services for children: a geographic perspective. *Int J Speech Lang Pathol*. 2011;13(3):239-50. doi: 10.3109/17549507.2011.573865.
63. Wakerman J, Humphreys JS, Wells R, Kuipers P, Entwistle P, Jones J. Primary health care delivery models in rural and remote Australia: a systematic review. *BMC health services research*. 2008;8:276. doi: 10.1186/1472-6963-8-276. PubMed PMID: 19114003.
64. Wakerman J. Innovative rural and remote primary health care models: what do we know and what are the research priorities? *Aust J Rural Health*. 2009;17(1):21-6. doi: 10.1111/j.1440-1584.2008.01032.x.
65. Wakerman J, Humphreys JS, Wells R, Kuipers P, Jones JA, Entwistle P, et al. Features of effective primary health care models in rural and remote Australia: a case-study analysis. *Medical Journal of Australia*. 2009;191(2):88-91. doi: 10.5694/j.1326-5377.2009.tb02700.x.
66. Ward AM, de Klerk N, Pritchard D, Firth M, Holman CD. Correlations of siblings' and mothers' utilisation of primary and hospital health care: a record linkage study in Western Australia. *Soc Sci Med*. 2006;62(6):1341-8. doi: 10.1016/j.socscimed.2005.08.027.
67. Warwick S, Atkinson D, Kitaura T, LeLievre M, Marley JV. Young Aboriginal People's Perspective on Access to Health Care in Remote Australia: Hearing Their Voices. *Prog Community Health Partnersh*. 2019;13(2):171-81. doi: 10.1353/cpr.2019.0017.
68. Whelan S, Wright DJ. Health Services Use and Lifestyle Choices of Indigenous and non-Indigenous Australians. *Social Science & Medicine*. 2013.
69. Williams ML, Smith AC. Paediatric outreach services. *Journal of Paediatrics and Child Health*. 2004;40(9-10):501-3. doi: 10.1111/j.1440-1754.2004.00450.x.
70. Zwi K, Joshua P, Moran P, Casacelli M. Prioritising health services for high risk and disadvantaged children. *Journal of Paediatrics and Child Health*. 2011;47:23.
71. Crouchley K, Carlose N. Aboriginal Health Profile: Kimberley Health Region. WA: Department of Health, 2012.
72. WA Country Health Service. Kimberley Health Profile. Perth: WACHS, 2018.
73. DoH Epidemiology Branch. Child Population Profile, Health and Wellbeing Surveillance System, 2009 Kimberley health region. Perth: Department of Health, 2010.
74. Australian Indigenous HealthInfoNet. Overview of Aboriginal and Torres Strait Islander health status, 2020. 2021.
75. HealthInfoNet AI, Harford-Mills M, MacRae A, Drew N. Summary of Aboriginal and Torres Strait Islander health status 2018. 2019.
76. Australian Institute of Health and Welfare. Expenditure on Health for Aboriginal and Torres Strait Islander People 2008-09: An Analysis by Remoteness and Disease. AIHW, 2011.
77. Australian institute of Health and Welfare. Access to health services for Aboriginal and Torres Strait Islander people. 2011.

78. Australian institute of Health and Welfare. Aboriginal and Torres Strait Islander health services report, 2010-11: OATSIH services reporting - key results. Canberra: 2013.
79. Australian Institute of Health and Welfare. Healthy for Life-Aboriginal Community Controlled Health Services: Report Card. AIHW Canberra; 2013.
80. Australian institute of Health and Welfare. Aboriginal and Torres Strait Islander Health Performance Framework. Perth: 2013.
81. Australian Institute of Health and Welfare. Health workforce. Canberra: AIHW, 2020.
82. Australian Medical Association. Indigenous Health Report Card – "Ending the Cycle of Vulnerability: The Health of Indigenous Children". 2008.
83. Wakerman J, Humphreys J, Wells R, Kuipers P, Entwistle P, Jones J. A systematic review of primary health care delivery models in rural and remote Australia, 1993-2006. 2006.
84. McDonald L, Bailie R, Morris P, Rumbold A, Paterson B. Interventions to prevent growth faltering in remote Indigenous communities. Australian Primary Health Care Research Institute (APHCRI), The Australian National University; 2006.
85. Price E, Considine G. Kimberley Primary Health Care Sustainability Study 2008 - 2030. Kimberley Division of General Practice Ltd, 2008.
86. Sheperd C, Zubrick S. Human capabilities and child development: Some inferences using the 2008 NATSISS. Perspectives on the 2008 National Aboriginal and Torres Strait Islander Social Survey; ANU 2011.
87. Shaw G. An ethnography of changes in child rearing over time in the Ngaanyatjarra Lands: Implications for policy development on health in early childhood. CAEPR Seminar Series Canberra 2013.
88. COAG Reform Council. Indigenous reform 2011-12: comparing performance across Australia. 2013.
89. Humphreys J, Wakerman J. Primary health care in rural and remote Australia: achieving equity of access and outcomes through national reform: A discussion paper. Canberra: National Health and Hospitals Reform Commission, 2008.
90. Allen, Clarke. Evaluation of the child health check initiative and the expanding health service delivery initiative: summary report. Department of Health and Ageing Canberra; 2011.
91. Standing Committee on Family and Community Affairs. Standing Committee on Family and Community Affairs. Health is life: Inquiry into indigenous health. Canberra: 2000.
92. Standing community on Health and Ageing. The Blame Game: Report on the inquiry into health funding. Canberra: Senate, 2006.
93. The Senate Select Committee on Regional and Remote Communities. Indigenous Australians, Incarceration and the Criminal Justice System. Canberra: Senate, 2010.
94. Health Workforce Australia. National Rural and Remote Workforce Innovation and Reform Strategy. Adelaide: HWA, 2013.
95. Health Workforce Australia. Health Workforce 2025 – Doctors, Nurses and Midwives – Volume 2. Adelaide: HWA, 2012.
96. Health Workforce Australia. Growing Our Future: the Aboriginal and Torres Strait Islander Health Worker Project Final Report. Adelaide: HWA, 2011.

97. Atkinson D, Bridge C, Gray D. Regional Aboriginal Health Plan: Aboriginal Health in the Kimberley: current circumstances and future directions. 2000.
98. Lewis J, editor Kimberley aboriginal primary health plan 2012-2015. Kimberley Aboriginal Health Planning Forum Broome; 2013.
99. Vos T, Carter R, Barendregt J, Mihalopoulos C, Veerman L, Magnus A, et al. Assessing cost-effectiveness in prevention: ACE-prevention September 2010 final report: University of Queensland; 2010.
100. Freemantle C, Officer K, Mcaullay D, Anderson I. Australian Indigenous Health-Within an International Context. Darwin: Cooperative Research Centre for Aboriginal Health, 2007.
101. Sanchez L, Carney S, Estermann A, Sparrow K, Turner D. An evaluation of the benefits of swimming pools for the hearing and ear health status of young Indigenous Australians: A whole-of-population study across multiple remote Indigenous communities. Adelaide: Flinders University. 2012.
102. Lowitja Institute. Close the gap campaign report 2021. 2021.
103. Menzies School of Health Research. Effective Integration of Services for Children and Families: Making it Happen. Darwin: 2011.
104. Thomas S. Kids Pitstop: one stop community health shop for Indigenous families in the West Pilbara. Rural Health: the place to be; 17-20 May; Cairns 2009.
105. Grant S. Child Malnutrition / Failure to Thrive Project 2008: Final Report. Darwin, NT.: Department of Health and Families, Northern Territory Government, 2008.
106. Office of Aboriginal and Torres Strait Islander Health. Improving Health in Aboriginal and Torres Strait Islander Mothers, Babies and Young Children - A Literature Review. 2005.
107. Griew R, Tilton E, Cox N, Thomas D. The link between primary health care and health outcomes for Aboriginal and Torres Strait Islander Australians. Report for the Office of Aboriginal and Torres Strait Islander Health Canberra: DoHA. 2008.
108. Office for Aboriginal Torres Strait Islander Health. Better health care: studies in the successful delivery of primary health care services for Aboriginal and Torres Strait Islander Australians. Canberra: Commonwealth Department of Health and Ageing, 2001.
109. Queensland Health. Making Tracks toward closing the gap in health outcomes for Indigenous Queenslanders by 2033 Implementation Plan 2009-10 to 2011-12. Brisbane: Department of Health, 2010.
110. Royal Flying Doctor Service. Annual Report 2011-12. 2012.
111. Gardiner F, Gale L, Ransom A, Laverty M. Looking Ahead: responding to the health needs of country Australians in 2028-the centenary year of the RFDS. Canberra, Australia: The Royal Flying Doctor Service. 2018.
112. Department of Health and Ageing. National Strategic Framework for Rural and Remote Health. 2011.
113. Rural Health West. Specialist Services in Rural Western Australia: Gap and Equity Perth: RHW, 2013.
114. Zubrick SR, Lawrence DM, Silburn SR, Blair E, Milroy H, Wilkes T, et al. The Western Australian Aboriginal Child Health Survey: The Health of Aboriginal Children and Young People. Perth: Telethon Institute for Child Health Research, 2004.

115. World Health Organization. Toolkit on monitoring health systems strengthening: Service Delivery. 2008.

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources




Figure 1: PRISMA flowchart of peer-reviewed and grey literature

Adapted from: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71.