Distinct origins and transmission pathways of *bla*KPC Enterobacterales across three U.S. states

Zena Lapp[1], Rany Octaria[2,3], Sean M. O'Malley[4], Tu Ngoc Nguyen[5], Hannah Wolford[6], Ryan Crawford[1], Christina Moore[3], Paula Snippes Vagnone[4], Diane Noel[5], Nadezhda Duffy[6], Ali Pirani[7], Linda S. Thomas[3], Brittany Pattee[4], Claire Pearson[5], Sandra N. Bulens[6], Sophie Hoffman[1], Marion Kainer[3], Melissa Anacker[4], James Meek[8], Isaac See[6], Allison Chan[3], Ruth Lynfield[4], Meghan Maloney[5], Mary K. Hayden[9,10], Evan Snitkin[7], Rachel B. Slayton[6]

[1] Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan, USA

[2] Department of Medicine, Division of Epidemiology, Vanderbilt University, Nashville, Tennessee, USA

[3] Tennessee Department of Health, Nashville, Tennessee, USA

[4] Minnesota Department of Health, USA

[5] Connecticut Department of Public Health, USA

[6] Centers for Disease Control and Prevention, USA

[7] Department of Microbiology and Immunology, University of Michigan, Ann Arbor, Michigan, USA

[8] Connecticut Emerging Infections Program, Yale School of Public Health, New Haven, CT

[9] Department of Medicine, Division of Infectious Diseases, Rush University Medical Center, Chicago, Illinois, USA

[10] Department of Pathology, Rush University Medical Center, Chicago, Illinois, USA

Keywords:

Transmission, carbapenem-resistant Enterobacterales (CRE), importation, horizontal gene

transfer (HGT), patient transfer

Running title:

CRE origin and transmission in 3 states

Corresponding author:

Evan Snitkin

esnitkin@med.umich.edu

1520D MSRB I, 1150 W. Medical Center Dr., Ann Arbor, MI, 48109-5680

 (734) 647-6472

Alternate corresponding author:

Rachel B. Slayton

via3@cdc.gov

1600 Clifton Rd NE, MS H16-3, Atlanta, GA 30329

404-636-4566

47   **Summary:**

48   Analysis of whole-genome sequences of bla_{KPC} Enterobacterales from three U.S. states revealed

49   that the underlying processes driving CRE burden in different regions can differ substantially,

50   and that within an individual region, different STs can spread via different pathways.

Abstract

Background

Carbapenem-resistant Enterobacterales (CRE) are among the most concerning antibiotic

resistance threats due to high rates of multidrug resistance, transmissibility in healthcare settings,

and high mortality rates. We evaluated the potential for regional genomic surveillance to track

bla_{KPC}-carrying CRE transmission across healthcare facilities in three U.S. states.

Methods

Clinical isolates were collected from Connecticut (CT; 2017-2018), Minnesota (MN; 2012-

2018), and Tennessee (TN; 2016-2017) through the U.S. Centers for Disease Control and

Prevention's Multi-site Gram-negative Surveillance Initiative and additional surveillance.

bla_{KPC}-containing *Enterobacter hormaechei* and *Klebsiella pneumoniae* isolates were whole-

genome sequenced, and case report data on patient comorbidities, healthcare utilization, and

interfacility patient transfer were extracted.

Findings

In CT, most CRE isolates showed evidence of importation from outside the state. In MN, cases

were primarily from clonal expansion of epidemic lineages ST258 *K. pneumoniae* and ST171 *E.

hormaechei*. In TN, ST258, ST307 *K. pneumoniae*, and ST114 *E. hormaechei* were most

common; the latter two showed evidence of local acquisition of bla_{KPC} followed by onward

dissemination. Patients from different facilities with genetically similar isolates were enriched in

shared facility exposures, consistent with patient transfer driving regional spread. Placing

genomic data in the context of statewide patient transfer networks in MN and TN revealed

largely non-overlapping facility networks mediating spread of ST258 versus other STs.

Conclusions

74   These findings demonstrate that the underlying processes driving CRE burden in different

75   regions can differ substantially, and that within a region, different STs can spread via different

76   pathways. Integrating genomic and epidemiological data, including interfacility patient transfers,

77   can provide insights to target interventions.

Introduction

Carbapenem-resistant Enterobacterales (CRE) accounted for an estimated 13,100 infections among hospitalized patients in the United States (U.S.) in 2017 [1], and likely colonized upwards of 100,000 additional patients [2]. Infections with CRE are considered an urgent public health threat as they can be difficult to treat due to their namesake resistance to carbapenems, as well as increasing rates of resistance to other last-resort antibiotics [3,4]. Of concern are CRE that contain a carbapenemase gene found on a plasmid or other mobile genetic element due to their high propensity to spread [1]. Due to the mobile nature of the genetic elements harboring these genes, carbapenem resistance can disseminate within and between different species via horizontal gene transfer (HGT). Once acquired by a strain sufficiently adapted to the healthcare environment, carbapenem-resistant organisms can spread through clonal dissemination [5]. Others have noted that different carbapenemases, which typically reside on distinct mobile elements, have varying tendencies for HGT vs. clonal spread. While HGT and clonal dissemination have been observed to play a role in CRE dynamics at both local and regional levels [5], the relative contributions of these processes in seeding and maintaining CRE burden across different geographic regions is still unclear.

Understanding the transmission pathways for CRE in hospitalized patients is critical for effective and efficient surveillance and intervention strategies. The CRE burden in some regions appears to be driven by the evolution or importation of lineages with epidemic potential, such as bla_{KPC}-carrying ST258 *Klebsiella pneumoniae* (e.g. California) [5] and ST171 *Enterobacter hormaechei* (e.g. Minnesota) [6] in the United States. Following initial regional emergence, these strains can be amplified via transmission within healthcare facilities and can spread between facilities via

the transfer of infected and colonized patients [5,7,8]. In regions with sustained transmission of

epidemic clones, effective regional control requires identification of locations where

transmission is occurring and impacting the burden in the region more broadly, as well as

monitoring the movement of CRE carriers between healthcare facilities [9,10]. In contrast, others

have observed a significant role for HGT of mobile carbapenemase elements in driving CRE

spread within individual healthcare facilities [11−13]. If HGT is a significant contributor to

overall regional CRE burden, effective prevention and control requires identification of

reservoirs of these resistance elements and an understanding of the propensity of different

strain/mobile element combinations to spread between patients or act as HGT donors.

Here, we evaluate the potential for regional genomic surveillance to inform the pathways leading

to CRE cases across the regional healthcare networks in three U.S. states involved in the U.S.

Centers for Disease Control and Prevention's (CDC) Multi-site Gram-negative Surveillance

Initiative (MuGSI). In particular, we focus on the most commonly observed CRE in the MuGSI

catchment areas, *K. pneumoniae* and *E. hormaechei* harboring the *K. pneumoniae*

carbapenemase (KPC). This builds off of the MuGSI aim to quantify the burden of certain

resistant gram-negative bacteria in the U.S. by supplementing preexisting data with whole-

genome sequencing of CRE isolates collected through MuGSI and supplemental state-wide

surveillance. By integrating genomic data from densely sampled CRE cases and healthcare

exposure data, and comparing local strains with global isolates, we gained insight into local

strain origins and regional transmission pathways.

Methods

This retrospective study was a collaboration between the Emerging Infections Program (EIP) and CDC Prevention Epicenters. The EIP is a collaboration between 10 state health departments and their partnering academic institutions and the Centers for Disease Control and Prevention and other federal agencies [14]. The Healthcare-Associated Infections Community Interface (HAIC) is one of the core components of the EIP and conducts active population- and laboratory-based surveillance for CRE [15]. Three EIP sites participated in this study (Connecticut, CT; Minnesota, MN; Tennessee, TN).

CRE surveillance

Each of the participating states conducted statewide public health surveillance for CRE. In addition, MN and TN conducted active surveillance for CRE through the HAIC MuGSI [16] in selected counties within the state and CT conducted active surveillance statewide. In both instances, carbapenem-resistant isolates were submitted by clinical laboratories to each state's respective State Public Health Laboratory (SPHL; see details below). Details about each state's CRE definition and MuGSI catchment area can be found in the Supplementary material. Multiple isolates can be obtained from the same patient if the new isolate was obtained at least 30 days after a previous case isolate was identified.

For this study, CRE are defined as any organism in the Enterobacterales order isolated from any clinical specimen and resistant to doripenem, meropenem, or imipenem (minimum inhibitory concentrations [MIC] of ≥ 4 µg/ml), ertapenem (MIC ≥ 2 µg/ml), or demonstrates production of a carbapenemase [16]. Specifically, isolates from residents of the surveillance area identified at the

clinical laboratory as CRE from *Escherichia coli*, *Enterobacter cloacae* complex species (i.e., *E.*

cloacae, *E. asburiae*, *E. bugandensis*, *E. hormaechei*, *E. kobei*, *E. ludwigii*, and *E.*

nimipressuralis), and *Klebsiella* species (i.e., *K. aerogenes*, *K. oxytoca*, and *K. pneumoniae*)

were included. All CRE isolates identified through the described public health surveillance

systems were characterized at the respective SPHL, which tested for carbapenemase genes (e.g.,

$bla_{KPC,}$ $bla_{NDM,}$ bla_{OXA-48}, bla_{VIM}, bla_{IMP}; see Supplemental Methods for more details). The

percentage of CRE tested for carbapenemases was 81.5% in CT, 90% in MN, and 78% in TN.

CRE considered for whole-genome sequencing were collected in CT from 2017-2018, in MN

from 2012-2018, and in TN from 2016-2017. For CT and MN, isolates from the entire state were

included. In TN, only isolates from TN Emergency Medical Services (EMS) Region 2-7, which

includes all TN counties except those in the Memphis-Delta and Northeast Tennessee Region,

were included. These were excluded due to significant healthcare utilization across state lines.

Table S1 provides information on the number of each type of healthcare facility in each state, as

well as the number of each type for which we have at least one whole-genome sequenced isolate.

<u>Isolate selection for whole-genome sequencing</u>

Over 90% of CRE isolates considered for whole-genome sequencing were bla_{KPC}-containing *K.*

pneumoniae or *E. hormaechei*. We elected to sequence this subset of isolates as the sample size

for the other carbapenemases and species would be too small to make any meaningful inferences.

<u>Isolate and genomic data processing</u>

DNA extraction and Illumina whole-genome sequencing were performed on all isolates. See

Supplemental Methods for state-specific details. We identified species and sequence types

[17,18], called single-nucleotide variants using species-specific reference genomes (*K.*

pneumoniae: KPNIH1, GenBank accession number CP008827.1, 5,394,056 base pairs; *E.*

hormaechei: MNCRE9, GenBank accession number JZDE00000000.1, 4,911,317 base pairs)

[19–25], built reference-based phylogenetic trees [24–28], and generated and annotated

assemblies [19,20,29,30]. Public CRE assemblies were downloaded from PATRIC [18,31], core

genome phylogenies were generated [24,32,33], and bla_{KPC} was identified [31,34]. See

Supplemental Methods for more details about the genomic data processing. Isolates from

BioProject numbers PRJNA272863 [6] and PRJNA873034 (new from this study) were used (see

Supplemental File S1 for isolate information).

<u>Epidemiologic data</u>

Epidemiologic data were obtained through medical record review and in TN supplemented by

additional data sources (see Appendix). Variables of interest collected for all isolates in this

study include state, treatment facility ID, patient ID, age, sex, hospitalizations in the previous

year, and culture source. In addition, MuGSI surveillance data included information on infection

type and underlying conditions, which was only obtained for selected non-MuGSI isolates. The

intra-facility transmission analysis investigating shared underlying conditions was limited to

samples for which we had data on these conditions; other analyses included all isolates (i.e.,

MuGSI and non-MuGSI). See Supplemental Methods for details on state-specific clinical case

data collection, facility identification, and matching facility names to Centers for Medicare &

Medicaid Services (CMS) facility IDs.

<u>Generation of aggregate patient transfer networks</u>

Aggregate patient transfer networks (i.e., data about patient flow across healthcare facilities [5])

for each state for the year 2017 were derived from CMS fee-for-service beneficiary claims data

linked to the CMS Minimum Data Set by Medicare beneficiary ID. The number of transfers

between two facilities includes transfers directly from one facility to another as well as transfers

with an intervening stay in the community of less than 365 days. See Supplemental Methods for

additional details.

<u>Public CRE isolates</u>

Public KPC-positive and KPC-negative *K. pneumoniae* and *E. hormaechei* isolates were

downloaded from the PATRIC database on 04/23/2021 [18] (see supplemental methods for more

details). These isolates were used to contextualize study isolates geographically and with respect

to KPC presence/absence. See below and supplementary methods for more details.

<u>Integrated genomic and epidemiologic data analysis</u>

To prevent overcounting of intra-facility transmission events, isolates were limited to the first

isolate per patient for each sequence type (ST)/facility combination. We then identified putative

groups of isolates related by geographic importation or HGT using maximum likelihood

ancestral reconstruction of core genome trees including study and public isolates. The discrete

trait used for ancestral reconstruction was geographic region or KPC presence/absence,

respectively. See supplementary methods for details. Next, we computed pairwise single

nucleotide variant (SNV) distances and patristic distances for study isolates from each sequence

alignment and classified isolates pairs as intra- or inter-facility pairs [35–37]. For most pairwise

SNV distance analyses, rather than choosing a specific pairwise SNV distance threshold, we

highlight various thresholds ($\leq$ 5, 10, 15) as a sensitivity analysis. A threshold of 10 was chosen

where needed based on a decrease in the fraction of intra-facility pairs at various SNV distances

(**Figure S4**). For each intra-facility patient pair, we determined whether they had at least one

shared comorbidity and used a Fisher's exact test [38] to compare sharing of comorbidities

between patients with isolates $\leq$10 SNVs and >10 SNVs. This was done based on the assumption

that patients with shared comorbidities would be provided care in more proximate inpatient

locations. For each inter-facility patient pair, we determined whether they had a shared

healthcare exposure in the past year (data collection method described in the Supplemental

Methods). Then we compared the extent of exposure for pairs of patients from different facilities

with isolates $\leq$10 SNVs and >10 SNVs using a Fisher's Exact test [38]. We also compared paths

of maximum patient flow computed from aggregate patient transfers and compared it to the

minimum pairwise SNV distance of isolate pairs from each facility pair using a Wilcox test

comparing facilities where the minimum pairwise SNV distance was $\leq$10 with those where it was

>10. An empirical p-value was also computed to compare the distribution of transmission

between ST171 and ST258 in MN. See Supplemental Methods for more details on these

analyses.

Data analysis and visualization

R v4.1.1 was used for all data analysis and visualization using the following packages: tidyverse

v1.3.1 [39], ggtree v2.5.2 [40,41], igraph v1.2.6 [42], regentrans v0.1 [37], tidygraph v1.2.0 [43],

ggraph v2.0.5 [44], exact2x2 v1.6.5 [38], cowplot v1.1.1 [45], readxl v1.3.1 [46], ape v5.5 [36].

The code used to perform all analyses can be found here: https://github.com/Snitkin-Lab-

Umich/eip-cre-transmission-ms.

<u>Human subjects research</u>

The study was reviewed and approved by the Tennessee Department of Health institutional

review board (IRB). See 45 CFR. part 46; 21 CFR part 56. The Minnesota and Connecticut

Department of Health IRB determined that the study was exempt from IRB review in accordance

with 45 CFR 46.101(b)(4)." The study was approved by the CDC IRB with a waiver of HIPAA

authorization under the Privacy Rule as per 45 CFR 46.512(i). The University of Michigan

Medical School IRB approved this protocol.

Results

We performed whole-genome sequencing on bla_{KPC}-positive *K. pneumoniae* and *E. hormaechei*

clinical isolates from CT, MN, and TN and included the first isolate of each unique

ST/facility/patient combination for downstream analyses (**Table 1**). For the purposes of

readability, we refer to this set of isolates as CRE throughout the manuscript. CRE were present

across numerous healthcare facilities in each state (CT: 22, MN: 60, TN: 40; **Table S1**; **Figure**

S1). The dominant STs across these three states were *K. pneumoniae* ST258 in all three states, *E.*

hormaechei ST171 in MN, and *E. hormaechei* ST114 and *K. pneumoniae* ST307 in TN.

Different extents of clonal dissemination after importation or HGT across states and STs

We were first interested in investigating the origins and distribution of bla_{KPC}-positive CRE STs across states. To do this we used previously sequenced *Enterobacter* spp. and *Klebsiella* spp. genomes deposited in the PATRIC database (**Figure S2**) to provide geographic and genetic context. We then classified phylogenetic clusters of isolates within each region as originating from either importation of a preexisting strain harboring bla_{KPC}-into the state or horizontal gene transfer of bla_{KPC} into a previously bla_{KPC}-negative strain within the state (**Figure 1**; **Figures S3-4**). We found that bla_{KPC}-positive *K. pneumoniae* ST258 is prevalent across all three states and, except for 3/66 isolates of unknown origin, exclusively originated from importation from outside the state. These importation events were followed by varying degrees of onward dissemination. This is consistent with *K. pneumoniae* ST258 being commonly disseminated across U.S. healthcare facilities and circulating lineages having a common ancestor dating back to the early 2000's [47]. Additionally, we found extensive dissemination of *E. hormaechei* ST171 in MN after an inferred single introduction into the state, consistent with previous reports of *E. hormaechei* ST171 transmission in MN following importation from North Dakota sometime prior to 2009 [6]. We also observed clonal dissemination of the emerging epidemic clones *K. pneumoniae* ST307 [48] and *E. hormaechei* ST114 [49] in TN following HGT of plasmids harboring bla_{KPC} into those respective strains as recently as 2016. In contrast, we found little evidence of dissemination after HGT in MN or CT, with many sporadic bla_{KPC} acquisitions found in one or a few patients.

Genomic analysis of regional clinical isolate collections allowed for the detection of local transmission in MN and TN, and lack thereof in CT

Having characterized the origin of circulating strains, we next sought to understand whether the

regional genomic surveillance of clinical isolates was dense enough to allow insight into

transmission pathways. To this end we investigated the extent to which regional phylogenetic

clusters that represented regional emergence events contained putative recent local transmission

events. We defined recent local transmission events as pairs of isolates with small pairwise SNV

distances from different patients and performed a sensitivity analysis with pairwise SNV distance

thresholds of ≤5, ≤10, and ≤15 (**Figure 2**; **Figures S5-6**) as well as with patristic distance

(**Figure S7**). Within an ST, smaller pairwise SNV distances indicate potential local transmission,

while larger pairwise SNV distances are not indicative of local transmission [50]. We captured

very few local transmission events in CT, suggesting that importation from other geographic

regions may be driving case load there. In contrast, we captured some local transmission events

in MN for *K. pneumoniae* ST258 and *E. hormaechei* ST171 and some local transmission of *K.*

pneumoniae ST258, *K. pneumoniae* ST307 and *E. hormaechei* ST114 in TN, consistent with our

finding that there was importation or HGT followed by dissemination of these STs. Closely

related inter-facility isolate pairs were most likely to occur between acute care hospitals (**Table**

S2).

To examine whether WGS of clinical isolate collections can shed light on which facilities had

transmission occur, we also investigated the relatedness of isolates within each facility (**Figure**

S8). We observed very little putative intra-facility transmission in CT. Conversely, we found that

facilities in MN tended to have closely related ST171 and ST258 isolates for all three pairwise

SNV distance thresholds, consistent with detection of intra-facility transmission. In TN, we saw

a similar trend, but for ST114 and ST307. Taken together, these findings suggest that clinical

isolate collections indeed capture recent transmission events, although the amount of

transmission appears different across regions, consistent with the phylogenetic analysis.

Patients with closely related intra-facility pairs do not share underlying conditions

Having detected putative intra-facility transmission in regional surveillance collections, we next

investigated whether these patients shared any underlying conditions, which may be indicative of

an outbreak related to a certain location or procedure in the hospital. However, we found that

patients residing in the same facility with closely related isolates did not consistently share

underlying conditions relative to patients with distantly related isolates (**Table S3**). This may be

due to extensive intermingling of patients with different chronic conditions in acute care

hospitals in the United States.

Aggregate- and individual-level patient transfer data support patient transfer mediating

regional spread

We next explored evidence for patient transfer mediating inter-facility spread of different CRE

STs using two patient sharing metrics: an aggregate metric based on CMS-derived patient

transfer frequencies between facilities and an individual metric based on patient-level shared

facility exposures (see Methods for details). A positive correlation between patient sharing and

isolate relatedness would suggest that the detection of closely related isolates in different

facilities is due to inter-facility spread via the transfer of colonized or infected patients. The

relationship between the aggregate and individual-level patient sharing metrics (aggregate,

Figure S9; and individual, **Table 2**, **Figure S10**), with isolate pair relatedness were generally

consistent for a given ST and state (**Figure S11**). For ST258 in all states ($p < 0.05$), facility pairs

with closely related isolate pairs tend to be ones with more patient sharing, consistent with our

knowledge that ST258 often traverses the healthcare network via patient transfer [5,7].

Moreover, we noted that in all three states, as compared to other STs, genetic linkages for ST258

were most strongly associated with shared facility exposures among linked patients. In MN we

also observe a significant association between genetic relatedness and shared exposures for

ST307, ST171 and ST114 (all p < 0.05), suggesting that patient transfer may be driving local

transmission in these cases as well. In TN we do not observe a significant association between

genetic linkage among non-ST258 isolates and shared exposures, due to a combination of small

numbers (e.g. ST171), and high frequencies of shared exposures among patients with isolates

with greater than 10 SNVs (e.g. ST114, ST307 and other STs).

Comparison of clonal spread across MN facilities reveals different patterns of transmission

across STs

As we observed evidence of substantial inter-facility transmission of both *E. hormaechei* ST171

and *K. pneumoniae* ST258 in MN, we next investigated the geographic spread of these STs

across the healthcare network. We found very different patterns of spread between these two

STs, with putative *K. pneumoniae* ST258 inter-facility transmission occurring sporadically

across the healthcare network (empirical p = 0.3), while putative *E. hormaechei* ST171

transmission was concentrated around a single facility we call Facility W (empirical p < 0.001;

Figure 3; **S12**). Further investigation of the connectedness of Facility W in the patient transfer

network revealed that it is highly connected to other facilities (**Figure S13**). Additionally, 10/24

(41.7%) Facility W patients with *E. hormaechei* ST171 had a shared prior healthcare exposure at

a specific other facility (Facility X). These findings indicate that, while both *E. hormaechei*

ST171 and *K. pneumoniae* ST258 may traverse the healthcare network via patient transfer, the

pattern by which these two STs disseminated across facilities, and the potential role of specific

facilities, is very different.

Distinct facility sub-networks in TN harbor different lineages

The examination of shared facility exposures in TN revealed an unexpected amount of shared

facility exposures among patients with *E. hormaechei* ST114 and *K. pneumoniae* ST307,

regardless of the precise genetic relatedness of their isolates (**Table 2**). To gain insight into this

pattern, we investigated what facilities patients with these STs had been to in the previous year

and found a cluster of seven facilities where patients with these STs, as well as STs categorized

as Other, had been (**Figure 4A**). Visualizing these seven facilities in the context of the aggregate

patient transfer network revealed that they are closely connected in the patient transfer network

(**Figure 4B**) and have more patient sharing among themselves than with other facilities (**Figure**

4C). Additionally, this facility cluster is distinct from a cluster containing predominately *K.*

pneumoniae ST258 (**Figure 4B**).

Discussion

As public health laboratories increase their capacity for genomic surveillance, it is critical to

understand if and how genomic data can be used to guide regional intervention efforts. Here, we

evaluated whether genomic analysis of clinical bla_{KPC}-positive CRE isolates from three U.S.

states could inform our understanding of the origins and transmission pathways of circulating

strains. We found that, in addition to detecting differences in strain composition across regions,

we could discern the relative contributions of importation, clonal dissemination and HGT to each

state's bla_{KPC}-*positive K. pneumoniae* and *E. hormaechei* case burden, as well as highlight

individual facilities and connected regional sub-networks where putative transmission occurred.

Notably, these insights did not rely on detailed clinical meta-data from affected patients,

indicating that other regions interested in identifying hotspots and pathways for regional spread

may be able to do so with only knowledge of recent healthcare exposures. These types of

insights can enable both targeting of regional infection prevention efforts, as well as prioritizing

surveillance to monitor transmission hotspots and track the reservoirs of circulating mobile

elements.

Despite focusing on the same two Enterobacterales species harboring the same carbapenemase

gene, we observed stark differences in the underlying drivers of each state's bla_{KPC}-positive CRE

burden. In CT, neither established epidemic lineages nor emergent CRE strains showed evidence

of onward transmission. Frequent importation of bla_{KPC}-positive CRE into CT is unsurprising,

given the proximity to other major CRE hotspots in the Northeast U.S. [51]. However, this still

leaves unanswered why imported bla_{KPC}-positive CRE have apparently not gained a foothold.

One reason may be that structural differences in regional healthcare networks influence local

propensity for bla_{KPC}-positive CRE amplification. For example, recent work has shown that post-

acute care settings such as long-term acute care hospitals and ventilator skilled nursing facilities

can act as initiators and amplifiers of regional CRE epidemics [52,53]. Thus, infection

prevention practices in these high-risk facilities, and their patient transfer connections with other

regional facilities may have large impacts on overall regional CRE spread. Regardless, our

observations suggest that focusing resources on testing admission cultures from out of state may

allow CT to continue having a low burden of these CRE strains.

Conversely, in MN, bla_{KPC}-positive CRE burden was driven by importation events of the major

epidemic lineages ST258 *K. pneumoniae* and ST171 *E. hormaechei*, followed by subsequent

dissemination within the region. However, there may be distinct reservoirs and/or transmission

pathways for ST258 and ST171 in MN as virtually all ST171 can be traced back to a single

importation event, while ST258 has entered the region numerous times. Furthermore,

examination of healthcare exposures for patients with genetically linked ST258 isolates was

consistent with patient transfer between facilities driving regional spread. In contrast, despite a

larger number of genetically linked ST171 isolate pairs, we observed a weaker enrichment in

shared patient exposures among genetically linked patients from different facilities. However,

despite a lack of individual-level shared healthcare exposures, we observed that genetically

linked isolates from different facilities were largely connected to isolates from a single hub

facility, Facility W. This facility is a hospital with many referrals that provides specialty care.

The lack of direct patient overlap or exposure to Facility W could indicate a large unobserved

pool of transmission intermediates at feeder healthcare facilities (e.g. Facility X), and/or

potentially an environmental reservoir within or proximate to Facility W. This suggests that MN

should focus their infection control efforts more on clonal dissemination rather than HGT. In

particular, Facility W could prioritize patients admitted from connected facilities, and state

epidemiologists could investigate facilities, such as Facility X, that are connected to Facility W

to identify other potential reservoirs and locations where intervention may have a broader

positive impact on regional CRE burden.

In contrast to MN where the bla_{KPC}-positive CRE burden was due to spread of two major

epidemic lineages, TN showed a much greater diversity in circulating strains. Most notably, in

addition to a significant burden of ST258, we observed large numbers of isolates from two more

recently emerged CRE lineages - ST307 *K. pneumoniae* and ST114 *E. hormaechei*. Although

their recent emergence limits genetic context provided by public databases, it is of note that the

most recent ancestor of clones disseminating in the region were bla_{KPC}-negative, consistent with

intra-region HGT initiating regional spread. In addition to putative regional HGT events in

ST307 and ST114, we also observed evidence of HGT events in less common CRE strains that

also seeded onward transmission. This pattern is in stark contrast with MN, where there were

very few cases of onward transmission detected after a non-epidemic strain acquired bla_{KPC}. It is

unclear whether these differences in the fate of novel plasmid acquisitions are due to the lack of

promiscuity of circulating plasmids, differences in the propensity of regional non-CRE strains to

maintain a bla_{KPC} plasmid, or some combination of the two. Prior work has suggested co-

adaptation of epidemic strains and their plasmids [54], which could point to the dominant

presence of ST171 and ST258 hindering onward dissemination of their bla_{KPC} plasmids, whereas

the relatively recent associations of dominant CRE strains and their plasmids in TN allows for

increased chance of productive HGT. This hypothesis is supported by the co-association of lower

frequency STs with ST307 and ST114 in a highly connected regional subnetwork in TN, as

compared to regional subnetworks where ST258 is dominant. Therefore, in TN we believe that

regional CRE prevention should focus on containing ST307 and ST114 as wel as continued

surveillance for new HGT events leading to broader clonal dissemination.

In addition to differences in drivers of CRE burden in different regions, we also observed some

similarities that are of note. In particular, despite significant differences across states in the

overall burden of ST258 and the number of observed genomic transmission linkages, we

consistently observed that patients with genetically linked ST258 isolates from different facilities

were strongly enriched for having histories of common healthcare exposures. This observation is

consistent with several previous studies showing evidence that ST258 spreads across regions via

the transfer of colonized and infected patients between healthcare facilities [5,55]. In contrast,

there were weaker, and sometimes no, association between shared facility exposures and

transmission for other STs, even in several cases where large numbers of genetically linked

patients were captured. One possible explanation for this is that non-ST258 sequence types have

reservoirs outside of hospitals that influence patterns of regional spread. An alternative

hypothesis is that the iceberg effect is much larger for non-ST258 STs, such that we are missing

more links in the transmission chain, and thereby failing to detect relevant exposures [2].

Our study has several limitations. First, we only included clinical isolates in our analysis and not

isolates that attempt to capture the burden of colonization, and we have an incomplete sampling

of isolates from TN. Both of these factors may lead to an underestimation of clonal

dissemination, that may be different for different STs depending on their virulence and

associated colonization-to-infection ratio (i.e. ST-specific iceberg effect). However, it is

noteworthy that we were still able to uncover patterns of clonal dissemination using these data,

suggesting that regional clinical isolate collection can provide actionable insights for regional

public health labs. Moreover, given that performing any kind of active surveillance at a regional

level is likely beyond the capacity of most local health departments, understanding how to

extract insights from clinical isolate collections, and potentially guide more targeted surveillance

efforts, is critical. While transmission may not have occurred at the facilities where clinical

isolates were collected, this nevertheless provides a location at which to initiate an investigation.

A second limitation is that the isolate collection periods were different for each state. This may

result in different levels of contextual information provided by public databases, through which

importation and HGT were inferred. However, our observations are consistent with prior studies

in that KPC is known to be stably associated with ST258, ST171 emerged in the upper Midwest

and ST307 and ST114 only recently became associated with KPC. Another limitation is that the

public isolates we used from PATRIC to inform the origin of locally circulating strains are not

equally representative of all geographic regions, which may underestimate the extent of

importation, as multiple importations from an unsampled reservoir may be merged. While the

analyses performed here would be enhanced by additional context, we note that inferences made

into importation using phylogenetic clustering are largely consistent with SNV-distance based

analyses, supporting the robustness of our conclusions. Furthermore, we only investigated HGT

of KPC rather than the entire bla_{KPC} plasmid, which could lead to an underestimate of the

number of HGT events. However, identification of plasmids with short-read data is fraught, and

we were still able to identify several putative HGT events using our simpler definition of HGT.

Furthermore, the aggregate patient transfer data used to understand the connectivity between

regional healthcare facilities was derived from Medicare and Medicaid patient transfers, which

may lead to biases in patient transfer connections. However, we note that, despite this, we

observed a strong relationship for many CRE strains, whereby closely related isolates tended to

come from pairs of facilities with high patient flow. We also observed differences between the

individual-level and aggregate-level patient transfer analyses, which may be due to smaller

sample sizes for these STs, leading to insufficient power to identify associations in the aggregate

patient transfer analysis.

In conclusion, our results suggest that, not only can the origins and transmission patterns of

bla_{KPC}-positive CRE be investigated using genomic surveillance of sufficiently comprehensive

regional clinical isolate collections, but also that these transmission dynamics can vary across

strains and regions. Importantly, the differences underlying the bla_{KPC}-positive CRE burden in

these three states would not have been discernible without genomic data, and in fact would be

strengthened further by the existence of larger genomic surveillance initiatives that could provide

more granular context for the origin of strains and their mobile elements to guide prevention

efforts. Similarly, our understanding of how bla_{KPC}-positive CRE proliferates once in a region

would be extremely limited without knowledge of patient healthcare exposures and transfer

networks, which enable a more nuanced understanding of where transmission is occurring, where

reservoirs might exist and where additional surveillance or intervention is required. Taken

together, these observations support the value of robust regional genomic surveillance for

antibiotic resistance threats and the need for analytic platforms capable of integrating genomic

and patient movement data to guide local and state infection prevention efforts.

Acknowledgements

Pamela Talley and Matt Estes from the Tennessee Department of Health. The MN-PHL

Microbiology and Sequencing & Bioinformatics laboratories for their work in identifying and

sequencing all of the MN isolates in the project. Gwendolyn Hughes and Christina Sancken from

CDC.

Funding

This work was supported by the CDC through the Emerging Infections Program cooperative

agreement (grant number CK17-1701). It was also supported in part by CDC Prevention

Epicenter Program Cooperative Agreement No. U54CK000481-S1. ZL received support

from the National Science Foundation Graduate Research Fellowship Program under Grant No.

DGE 1256260. Any opinions, findings, and conclusions or recommendations expressed in this

material are those of the authors and do not necessarily reflect the views of the National Science

Foundation.

Conflicts of interest

M.K.H. was a member of a clinical adjudication panel for an investigational SARS-CoV-2

vaccine developed by Sanofi for a year and a half, ending in July 2022.

Disclaimer: The findings and conclusions in this report are those of the authors and do not

necessarily represent the views of the Centers for Disease Control and Prevention. The use of

trade names is for identification only and does not imply endorsement by the Centers for Disease

Control and Prevention.

References

1. Centers for Disease Control and Prevention (U.S.). Antibiotic resistance threats in the

   United States, 2019. Centers for Disease Control and Prevention (U.S.), 2019. Available at:

   https://stacks.cdc.gov/view/cdc/82532. Accessed 11 June 2020.

2. Lee BY, Bartsch SM, Wong KF, et al. Tracking the spread of carbapenem-resistant

Enterobacteriaceae (CRE) through clinical cultures alone underestimates the spread of CRE

even more than anticipated. Infection Control & Hospital Epidemiology. **2019**; 40: 731–

734.

3. World Health Organization, editor. Antimicrobial resistance: global report on surveillance.

Geneva, Switzerland: World Health Organization, 2014.

4. Potter RF, D'Souza AW, Dantas G. The rapid spread of carbapenem-resistant

Enterobacteriaceae. Drug Resistance Updates **2016**; 29:30–46.

5. Han JH, Lapp Z, Bushman F, et al. Whole-genome sequencing to identify drivers of

carbapenem-resistant Klebsiella pneumoniae transmission within and between regional

long-term acute-care hospitals. Antimicrobial agents and chemotherapy **2019**; 63:e01622-

19.

6. Hargreaves ML, Shaw KM, Dobbins G, et al. Clonal Dissemination of Enterobacter cloacae

Harboring blaKPC-3 in the Upper Midwestern United States. Antimicrobial Agents and

Chemotherapy **2015**; 59:7723–7734.

7. Snitkin ES, Won S, Pirani A, et al. Integrated genomic and interfacility patient-transfer data

reveal the transmission pathways of multidrug-resistant Klebsiella pneumoniae in a regional

outbreak. Science Translational Medicine **2017**; 9:eaan0093.

8. Lapp Z, Crawford R, Miles-Jay A, et al. Regional Spread of blaNDM-1-Containing

Klebsiella pneumoniae ST147 in Post-Acute Care Facilities. Clinical Infectious Diseases

2021; Available at: https://doi.org/10.1093/cid/ciab457. Accessed 15 July 2021.

9. Slayton RB, Toth D, Lee BY, et al. Vital Signs: Estimated Effects of a Coordinated

Approach for Action to Reduce Antibiotic-Resistant Infections in Health Care Facilities —

United States. MMWR Morb Mortal Wkly Rep **2015**; 64:826–831.

10.   Toth DJA, Khader K, Slayton RB, et al. The Potential for Interventions in a Long-term

Acute Care Hospital to Reduce Transmission of Carbapenem-Resistant Enterobacteriaceae

in Affiliated Healthcare Facilities. Clin Infect Dis **2017**; 65:581–587.

11.   Mathers AJ, Cox HL, Kitchel B, et al. Molecular Dissection of an Outbreak of

Carbapenem-Resistant Enterobacteriaceae Reveals Intergenus KPC Carbapenemase

Transmission through a Promiscuous Plasmid. mBio **2011**; Available at:

https://journals.asm.org/doi/abs/10.1128/mBio.00204-11. Accessed 21 February 2022.

12.   Mathers AJ, Stoesser N, Sheppard AE, et al. Klebsiella pneumoniae Carbapenemase

(KPC)-Producing K. pneumoniae at a Single Institution: Insights into Endemicity from

Whole-Genome Sequencing. Antimicrob Agents Chemother **2015**; 59:1656–1663.

13.   León-Sampedro R, DelaFuente J, Díaz-Agero C, et al. Pervasive transmission of a

carbapenem resistance plasmid in the gut microbiota of hospitalized patients. Nat Microbiol

2021; 6:606–616.

14.   About EIP | DPEI | CDC. 2021. Available at: https://www.cdc.gov/ncezid/dpei/eip/eip-

about.html. Accessed 2 April 2022.

15.   Magill SS, Dumyati G, Ray SM, Fridkin SK. Evaluating Epidemiology and Improving

Surveillance of Infections Associated with Health Care, United States - Volume 21,

Number 9—September 2015 - Emerging Infectious Diseases journal - CDC. Available at:

https://wwwnc.cdc.gov/eid/article/21/9/15-0508_article. Accessed 16 February 2022.

16.   Guh AY, Bulens SN, Mu Y, et al. Epidemiology of Carbapenem-Resistant

Enterobacteriaceae in 7 US Communities, 2012-2013. JAMA **2015**; 314:1479–1487.

17.   Hunt M, Mather AE, Sánchez-Busó L, et al. ARIBA: rapid antimicrobial resistance

genotyping directly from sequencing reads. Microb Genom **2017**; 3. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5695208/. Accessed 26 March 2021.

18. Davis JJ, Wattam AR, Aziz RK, et al. The PATRIC Bioinformatics Resource Center:

expanding data and analysis capabilities. Nucleic Acids Res **2020**; 48:D606–D612.

19. Andrews S. s-andrews/FastQC. 2021. Available at: https://github.com/s-andrews/FastQC.

Accessed 10 March 2021.

20. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics **2014**; 30:2114–2120.

21. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.

Bioinformatics **2009**; 25:1754–1760.

22. broadinstitute/picard. Broad Institute, 2021. Available at:

https://github.com/broadinstitute/picard. Accessed 20 February 2021.

23. Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and SAMtools.

Bioinformatics **2009**; 25:2078–2079.

24. Croucher NJ, Page AJ, Connor TR, et al. Rapid phylogenetic analysis of large samples of

recombinant bacterial whole genome sequences using Gubbins. Nucleic Acids Res **2015**;

43:e15–e15.

25. Poplin R, Ruano-Rubio V, DePristo MA, et al. Scaling accurate genetic variant discovery to

tens of thousands of samples. bioRxiv **2018**; :201178.

26. Kurtz S, Phillippy A, Delcher AL, et al. Versatile and open software for comparing large

genomes. Genome Biology **2004**; 5:R12.

27. Arndt D, Grant JR, Marcu A, et al. PHASTER: a better, faster version of the PHAST phage

search tool. Nucleic Acids Res **2016**; 44:W16–W21.

28.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective

607   Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol

608   **2015**; 32:268–274.

609   29.   Bankevich A, Nurk S, Antipov D, et al. SPAdes: A New Genome Assembly Algorithm and

610   Its Applications to Single-Cell Sequencing. Journal of Computational Biology **2012**;

611   19:455–477.

612   30.   Walker BJ, Abeel T, Shea T, et al. Pilon: An Integrated Tool for Comprehensive Microbial

613   Variant Detection and Genome Assembly Improvement. PLOS ONE **2014**; 9:e112963.

614   31.   Brettin T, Davis JJ, Disz T, et al. RASTtk: A modular and extensible implementation of the

615   RAST algorithm for building custom annotation pipelines and annotating batches of

616   genomes. Sci Rep **2015**; 5:8365.

617   32.   Crawford RD, Snitkin ES. cognac: rapid generation of concatenated gene alignments for

618   phylogenetic inference from large, bacterial whole genome sequencing datasets. BMC

619   Bioinformatics **2021**; 22:70.

620   33.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective

621   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol **2015**;

622   32:268–274.

623   34.   Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-generation

624   sequencing data. Bioinformatics **2012**; 28:3150–3152.

625   35.   R Core Team. R: A Language and Environment for Statistical Computing. **2020**; Available

626   at: https://www.R-project.org/.

627   36.   Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary

628   analyses in R. Bioinformatics **2019**; 35:526–528.

37. Hoffman S, Lapp Z, Wang J, Snitkin ESY 2022. regentrans: a framework and R package

for using genomics to study regional pathogen transmission. Microbial Genomics 8:000747.

38. Fay MP. Confidence intervals that match Fisher's exact or Blaker's exact tests. Biostatistics

2010; 11:373–374.

39. Wickham H, Averick M, Bryan J, et al. Welcome to the Tidyverse. Journal of Open Source

Software **2019**; 4:1686.

40. Yu G, Smith DK, Zhu H, Guan Y, Lam TT-Y. ggtree: an r package for visualization and

annotation of phylogenetic trees with their covariates and other associated data. Methods in

Ecology and Evolution **2017**; 8:28–36.

41. Yu G, Lam TT-Y, Zhu H, Guan Y. Two Methods for Mapping and Visualizing Associated

Data on Phylogeny Using Ggtree. Mol Biol Evol **2018**; 35:3041–3043.

42. Csardi G, Nepusz T. The Igraph Software Package for Complex Network Research.

InterJournal **2005**; Complex Systems:1695.

43. Pedersen TL. tidygraph: A Tidy API for Graph Manipulation. 2020. Available at:

https://CRAN.R-project.org/package=tidygraph. Accessed 16 February 2022.

44. Pedersen TL, RStudio. ggraph: An Implementation of Grammar of Graphics for Graphs and

Networks. 2021. Available at: https://CRAN.R-project.org/package=ggraph. Accessed 16

November 2021.

45. Wilke CO. cowplot: Streamlined Plot Theme and Plot Annotations for 'ggplot2'. 2019.

Available at: https://CRAN.R-project.org/package=cowplot. Accessed 15 April 2020.

46. Wickham H, Bryan J, attribution) Rs (Copyright holder of all R code and all C code without

explicit copyright, et al. readxl: Read Excel Files. 2019. Available at: https://CRAN.R-

project.org/package=readxl. Accessed 16 February 2022.

47. Bowers JR, Kitchel B, Driebe EM, et al. Genomic Analysis of the Emergence and Rapid

Global Dissemination of the Clonal Group 258 Klebsiella pneumoniae Pandemic. PLoS

ONE **2015**; 10:e0133727.

48. Peirano G, Chen L, Kreiswirth BN, Pitout JDD. Emerging Antimicrobial-Resistant High-

Risk Klebsiella pneumoniae Clones ST307 and ST147. Antimicrobial Agents and

Chemotherapy **2020**; 64. Available at: http://aac.asm.org/content/64/10/e01148-20.

Accessed 26 February 2021.

49. Izdebski R, Baraniak A, Herda M, et al. MLST reveals potentially high-risk international

clones of Enterobacter cloacae*. Journal of Antimicrobial Chemotherapy **2015**; 70:48–56.

50. Worby CJ, Chang H-H, Hanage WP, Lipsitch M. The Distribution of Pairwise Genetic

Distances: A Tool for Investigating Disease Transmission. Genetics **2014**; 198:1395–1404.

51. Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant Enterobacteriaceae:

The Impact and Evolution of a Global Menace. J Infect Dis **2017**; 215:S28–S36.

52. Lin MY, Lyles-Banks RD, Lolans K, et al. The Importance of Long-term Acute Care

Hospitals in the Regional Epidemiology of Klebsiella pneumoniae Carbapenemase–

Producing Enterobacteriaceae. Clin Infect Dis **2013**; 57:1246–1252.

53. Pacilli M, Kerins JL, Clegg WJ, et al. Regional Emergence of Candida auris in Chicago and

Lessons Learned From Intensive Follow-up at 1 Ventilator-Capable Skilled Nursing

Facility. Clin Infect Dis **2020**; 71:e718–e725.

54. Harrison E, Brockhurst MA. Plasmid-mediated horizontal gene transfer is a coevolutionary

process. Trends in Microbiology **2012**; 20:262–267.

55. Snitkin ES, Won S, Pirani A, et al. Integrated genomic and interfacility patient-transfer data

reveal the transmission pathways of multidrug-resistant Klebsiella pneumoniae in a regional

outbreak. Science Translational Medicine **2017**; 9.

Figures and tables

Table 1: Overview of whole-genome sequenced isolates from each state.

State	Surveillance area	Time period	Species	Number of isolates
CT	Entire state	2017-2018	*E. hormaechei*	8
			K. pneumoniae	44
MN	Entire state[1]	2012-2018	*E. hormaechei*	95
			K. pneumoniae	103
TN	83 counties[2]	2016-2017	*E. hormaechei*	58
			K. pneumoniae	78

681   [1]The MuGSI catchment area is Hennepin and Ramsey counties.

682   [2]4 counties in the Memphis-Delta Emergency Medical Services (EMS) and 8 in Northeastern TN

683   were excluded due to extensive healthcare utilization across state lines; the MuGSI catchment

684   area is TN EMS Region 5, an 8-county region encompassing the metropolitan Nashville area, of

685   TN.

Table 2: Individual-level healthcare exposure data provides some insight into potential sites of inter-facility transmission.

State	ST	Shared facility exposures[1] (%)		Fisher's exact p
		Pairwise SNV distance ≤10	Pairwise SNV distance > 10	
CT	*K. pneumoniae* ST258	1/2 (50)	1/197 (1)	0.02
	K. pneumoniae ST307	0/1 (0)	0/2 (0)	1
	Other	0/3 (0)	0/4 (0)	1
MN	*E. hormaechei* ST114	3/9 (33)	1/43 (2)	0.02
	E. hormaechei ST171	23/430 (5)	62/1956 (3)	0.03
	K. pneumoniae ST258	10/62 (16)	23/1956 (1)	$2*10^{-8}$
	K. pneumoniae ST307	3/3 (100)	0/7 (0)	0.008
	Other	2/4 (50)	0/6 (0)	0.1

	E. hormaechei ST114	4/16 (25)	24/79 (30)	0.8
	E. hormaechei ST171	1/4 (25)	1/21 (5)	0.3
TN	*K. pneumoniae* ST258	7/21 (33)	7/369 (2)	$1*10^{-6}$
	K. pneumoniae ST307	5/17 (29)	30/104 (29)	1
	Other	7/13 (54)	4/18 (22)	0.1

689 [1]A shared facility exposure is when individuals with isolates from different facilities both spent

690 time in the same facility at some point in the last year.

Figure 1: Different extents of clonal dissemination after importation or horizontal gene transfer (HGT) across states and sequence types (STs). Clusters are separated by white lines between the colored bars, where the bar height of a given cluster is the cluster size. Clusters were defined as subclades that were monophyletic for a single state. Importation events were considered those with an edge transition from another state to the EIP state. HGT events were considered those with a KPC- to KPC+ transition regardless of the state of origin of the KPC- strain. See methods for more details. We observed several importation events with limited onward transmission in all states. We also see a large cluster of *E. hormaechei* ST171 in MN that arose from a putative importation event, indicating importation followed by sustained dissemination. Additionally, we observed two large clusters of *E. hormaechei* ST114 and *K. pneumoniae* ST307, respectively, in TN, indicating HGT events followed by dissemination.

Figure 2: Small pairwise single nucleotide variant (SNV) distances between isolates suggest

that local transmission was captured. Each point is an isolate pair. The y axis is log10-scaled

and points are jittered horizontally for visualization purposes. Varying extents of transmission

were captured for different states and STs. Putative local transmission events are those in the tail

of the distribution (lower pairwise SNV distances). SNV bins were chosen to highlight closely

related pairs of isolates at different SNV thresholds as a sensitivity analysis.

Figure 3: Different patterns of transmission across STs in MN. Facilities are oriented as they

are geographically in space, but longitude and latitude are anonymized. The grey box is an insert

of the left map. Each isolate was paired to their closest isolate in another facility with a

subsequent collection date. *E. hormaechei* ST171 inter-facility transmission is concentrated

around facility W (29/40 [72.5%] of inter-facility pairs with a pairwise SNV distance ≤10 are

connected to facility W), while *K. pneumoniae* ST258 transmission is more evenly distributed

between facilities (maximum 3/8 [37.5%] of inter-facility pairs with a pairwise SNV distance

≤10 are connected to any facility). SNV = single nucleotide variant.

Figure 4: Distinct facility sub-networks in TN harbor different lineages. (A) Number of

patient facility exposures in the prior year for each facility and ST. Each patient may be

represented more than once if they have more than one facility exposure. A cluster of facilities

identified using the complete linkage clustering method (dashed box) are common in patients

with ST114, ST307, and other isolates. (B) Patient transfer network of facilities with at least one

whole-genome sequenced isolate plotted using the Kamada-Kawai algorithm. The cluster of

facilities identified in the heatmap in panel A (Kp ST307/Ec ST114/Other facility cluster) are

also clustered in the network. (C) Number of patient transfers for facility pairs where 0, 1, or 2 of

the facilities are in the facility cluster identified in the heatmap in panel A. Facility pairs where

both facilities are in the cluster have more patient sharing than facility pairs where only 0 or 1 of

the facilities is in the identified cluster.

Supplemental information

Supplemental methods

<u>Catchment areas</u>

The Multi-site Gram-negative Surveillance Initiative (MuGSI) [16] CRE surveillance catchment

areas for each state participating in this study are: the entire state of CT; Hennepin and Ramsey

counties, encompassing the metropolitan Minneapolis-St. Paul area, of MN; and TN EMS

Region 5, an 8-county region encompassing the metropolitan Nashville area, of TN. The TN the

Memphis-Delta and Northeast Regions were excluded due to healthcare utilization across state

lines.

<u>CRE surveillance</u>

Connecticut

Connecticut did not participate in MuGSI surveillance during the study period, so all study

isolates from Connecticut came from statewide passive surveillance for CRE. CRE surveillance

of all Enterobacterales isolates collected from all invasive clinical sites, respiratory sources, and

urine began on Jan 1, 2017. All CRE isolates undergo antimicrobial susceptibility testing with a

custom Sensititre and disk diffusion panel, phenotypic detection of carbapenemase activity using

the mCIM test and PCR detection of carbapenemase genes (bla_{KPC}, bla_{NDM}, bla_{OXA-48}, bla_{VIM},

bla_{IMP}). Isolates carrying bla_{KPC} that were identified in 2017 and 2018 from this collection were

included.

Minnesota

MuGSI surveillance in Minnesota is conducted in Ramsey and Hennepin counties. MuGSI

isolates are obtained from sterile sites or urine and include *E. coli*, *Klebsiella* spp., and

Enterobacter spp. Non-MuGSI isolates are collected from a variety of culture sources (e.g.,

blood, urine, respiratory) and represent multiple Enterobacterales species. Isolate identification is

confirmed by the MALDI Biotyper CA System (Bruker Daltonics, Inc., Billerica, MA). All CRE

isolates are tested for carbapenemase production using the modified carbapenem inactivation

method (mCIM). Isolates demonstrating phenotypic carbapenemase production undergo

molecular testing (i.e., polymerase chain reaction [PCR]) for detection of carbapenemase genes

bla_{KPC}, bla_{NDM}, bla_{OXA-48}, bla_{VIM}, bla_{IMP}. All bla_{KPC} -positive *K. pneumoniae* and *E. cloacae*

complex isolates submitted to MN PHL from 2012-2018 were analyzed for this study.

Tennessee

EMS region 2-7 were selected because it provides a good representation of patient referrals;

captures most isolates in patient transfer networks covering the area; and includes both the

Knoxville area in which CRE emerged in 2016, and the Nashville area (the MuGSI catchment

area: Dickson, Cheatham, Robertson, Sumner, Wilson, Rutherford, Williamson and Davidson

counties) where CRE increased in 2017. CRE isolates underwent organism confirmation using

the MALDI-TOF bioMerieux. Isolates were tested for antimicrobial resistance with the Kirby

Bauer method. Those isolates that were resistant to at least one carbapenem were then tested by

the CarbaNP Method for carbapenemase production. Those that demonstrated carbapenemase

production from the CarbaNP underwent a PCR assay developed by the CDC to test for the

bla_{KPC} and bla_{NDM}. All bla_{KPC}-positive isolates collected during 2016–2017 and submitted to the

TN SPHL were included.

<u>Epidemiologic data</u>

MuGSI surveillance data were collected through medical record review by EIP surveillance

officers in each site. Additional details on data collection for isolates identified outside of the

MuGSI program are listed below for each site.

Connecticut

Clinical case details were abstracted from medical records by EIP surveillance officers with the

Connecticut Department of Public Health.

Minnesota

At the time of culture collection, surveillance staff completed CRF through electronic or in-

person medical chart review.

Tennessee

For all carbapenemase-producing Enterobacterales cases, a questionnaire is completed by the

healthcare facility of origin. For cases that did not have a completed questionnaire,

hospitalization details were gathered from the Statewide Hospital Discharge data (described

below).

<u>Facility identification</u>

Facility identification was performed through chart review at the reporting facility. State-specific

details are described below.

Connecticut

819 Chart review was performed at the reporting facility for all inpatient and hospital network events.

820 When additional referring facilities (including nursing homes, long term acute care hospitals,

821 etc.) were identified on medical record review, follow-up investigation was performed at the

822 additional facilities. Upon identification of outpatient isolates, patient details were collected from

823 ordering providers by phone call and/or faxed data collection instrument.

Minnesota

826 MN staff maintain a compendium with facility name and type according to CMS. The facility

827 name and type are recorded in the MN surveillance database for each isolate and verified using

828 this compendium. For MuGSI cases, chart review at the facility where the isolate was collected

829 was also performed by MN staff.

Tennessee

TN staff linked patient identifiers of each case in the National Electronic Disease Surveillance

System (NEDDS)-Base System (NBS) surveillance data to the TN Hospital Discharge Data

System (HDDS) to identify and confirm the healthcare facility where they had the first specimen

culture collected for each CP-CRE case in TN. Alternatively, if the patient was not hospitalized

in a TDH-licensed hospital and therefore, was not found in the HDDS, TN staff used

surveillance data from NBS and the afore-mentioned questionnaire to identify the healthcare

facility of origin. Using this approach, the healthcare facility of origin was identified for 139

(89%) of the 157 included isolates. TN staff also gathered information on healthcare exposures

within one year prior to specimen collection by linking the patient identifiers in the surveillance

data with the inpatient and outpatient HDDS dataset. Isolates with no healthcare exposure

information were not used for the patient sharing network analysis.

<u>Linkage/matching of facility names to CMS ID</u>

Linkage of facility names to CMS ID was performed at each respective SPHL.

Connecticut

Facilities were matched with CMS IDs using CMS Provider of Services Files available at:

https://www.cms.gov/Research-Statistics-Data-and-Systems/Downloadable-Public-Use-

Files/Provider-of-Services

Minnesota

In MN, CMS IDs were manually entered for each isolate using Excel.

Tennessee

In TN, the name healthcare facility of origin was linked with the registered facility name by

CMS ID using the SPEDIS and COMPGED commands in SAS 9.4 (SAS Institute, Cary, North

Carolina) to allow for inexact matches that accounted for minor spelling variations between the

facility names from TN's surveillance data and those from the CMS dataset. Due to the data-

sharing agreement between TDH and the University of Michigan, the facility names and CMS

IDs in the WGS dataset and patient transfer data from CMS were de-identified before sharing

with The University of Michigan.

<u>Generation of aggregate patient transfer networks</u>

Aggregate patient transfer networks for each state for the year 2017 were derived from Centers

for Medicare and Medicaid Services (CMS) fee for service beneficiary claims data linked to the

CMS Minimum Data Set (MDS) by Medicare beneficiary ID. The dataset comprises a network

of medical facilities, including but not limited to short-term acute care hospitals, skilled nursing

facilities and long-term acute care hospitals, but may not be representative of non-Medicare

beneficiaries. To create an annual transfer network, CDC staff linked each unique beneficiary's

healthcare utilization. This included CMS hospital claims and the MDS to ascertain a

beneficiary's presence in a nursing home, regardless of payer for that stay. Transfer networks

were comprised of the year of interest and looked back to the prior year to obtain all discharging

facilities where patients had an opportunity to acquire CRE.

The number of transfers between two facilities includes transfers directly from one facility to

another as well as transfers with an intervening stay in the community of less than 365 days. We

chose to use a 365-day time interval rather than a smaller time interval because this gives us

more resolution in comparing the extent of patient flow between facilities. Any facility pairs with

10 or fewer patient transfers were censored, as per data use agreements between CDC and CMS.

<u>DNA extraction</u>

Connecticut

Total DNA extractions performed by CT DPHL for WGS were either by the manual method or

automated on the QIAcube. Manual extractions from pure single colony Gram-negative bacterial

isolates by QIAgen DNeasy Blood and Tissue kit and follow-up quality control of all purified

extracts were conducted following PulseNet Protocols for Whole Genome Sequencing on the

Illumina MiSeq system. Automated extractions of bacterial cell pellets were performed using the

QIAcube protocol for Isolation of DNA from Gram-negative bacteria and QIAGEN DNeasy

Blood and Tissue kit. To obtain the cell pellets, colonies from overnight plates were resuspended

in 1x Phosphate buffered saline (PBS, Sigma) to an OD600 of 0.35-0.40. Two milliliter cell

suspension were centrifuged at 5,000 x g for 5 minutes at 2-8 °C and used for extractions.

Minnesota

DNA extraction of the isolates was performed in MN using either the QIAcube (Qiagen,

Germantown, MD) extractor and QIAmp DNA Mini extraction kit (Qiagen, Germantown, MD)

or the Roche MagnaPure LC and the TNA Isolation extraction kit.

Tennessee

DNA extraction of the isolates was performed in TN on a Thermo Fisher, KingFisher Flex auto-

extraction instrument, using an Applied Biosystems MagMAX DNA Multi-Sample Ultra

extraction kit.

<u>Whole-genome sequencing (WGS)</u>

The extracted DNA was prepared for WGS using Illumina Nextera XT library Prep kits with

Illumina, Nextera XT Index kits (MN, TN, CT) or with Illumina DNA Prep kit and Illumina CD

Indexes (CT) (Illumina, San Diego, CA). The Library preparation process was performed on a

Beckman Coulter Biomek Fxp instrument (Beckman Coulter, Brea, CA) (TN), on the Eppendorf

epMotion (CT) (Eppendorf, Enfield, CT), or manually (MN, CT). The prepared libraries were

sequenced on an Illumina MiSeq sequencing instrument (Illumina, San Diego, CA) using 2x250

(MN, TN, CT) or 2x150 (CT) paired end sequencing with V2 chemistry. The fastq files

generated were transferred to the University of Michigan for processing and analysis.

Genomic data processing

All genomic data processing and analysis was performed by members of Evan Snitkin's lab at

the University of Michigan.

Species/ST identification

MLST was assigned with Ariba v2.14.4 [17]. For PATRIC isolates mlst were extracted from the

isolate metadata [18].

Sequence alignments

Single-nucleotide variants were called in study sequences using SNPKIT

(https://github.com/Snitkin-Lab-Umich/snpkit). Quality of reads was assessed with FastQC

v0.11.9 [19]. Adapter sequences and low-quality bases were removed with Trimmomatic v0.36

[20]. Variants were identified by mapping filtered reads to the KPNIH1 reference genome

(GenBank accession number CP008827.1; 5,394,056 base pairs) for *K. pneumoniae* sequences

and the MNCRE9 reference genome (GenBank accession number JZDE00000000.1; 4,911,317

base pairs) for *E. hormaechei* sequences using bwa v0.7.17 [21], removing polymerase chain

reaction duplicates with Picard 2.21.7 [22], removing clipped alignments using Samclip 0.4.0,

and calling variants with SAMtools v1.11and bcftools [23]. Variants were filtered from raw

results using GATK's VariantFiltration v3.8 (QUAL, >100; MQ, >50; ≥10 reads supporting

variant; and FQ, <0.025) [25]. Consensus files generated during variant calling were

recombination filtered using Gubbins v3.0.0 [24]. The alleles at each position that passed

filtering were concatenated to generate a non-core variant alignment relative to the respective

reference genome. Alleles that did not pass filtering were considered unknown (denoted as N in

the alignment).

Reference-based phylogenies

A custom Python script was used to filter out (mask) single nucleotide variants in the whole-

genome alignment that were: (i) <5 base pairs (bp) in proximity to indels that were identified by

GATK HaplotypeCaller [25], (ii) in a recombinant region identified by Gubbins v3.0.0 [24], (iii)

in a phage region identified by the Phaster web tool [27] or (iv) they resided in tandem repeats of

length greater than 20bp as determined using the exact-tandem program in MUMmer v3.23 [26].

This whole-genome masked variant alignment was used to reconstruct a maximum likelihood

phylogeny with IQ-TREE v1.6.12 using the general time reversible model GTR+G and ultrafast

bootstrap with 1000 replicates (-bb 1000) [33].

Genome assemblies

Quality of reads was assessed with FastQC v0.11.9 [19]. Adapter sequences and low-quality

bases were removed with Trimmomatic v0.36 [20]. The clean reads were assembled with Spades

v3.14.1 (careful mode) [29]followed by Pilon v1.23 assembly correction [30]. Contigs smaller

than 500bp were discarded.

Public isolates

All assemblies from the species identified within the EIP dataset were downloaded from the

PATRIC database on 04/23/2021 [18]. These data were merged with assemblies from

PRJNA603790, PRJNA690239, PRJNA401340, and PRJNA415194 for a total of 74,367

assemblies. Genome annotations for the dataset were generated with RAST v1.035 [31].

Core genome phylogenies

A concatenated gene alignment of core genes was generated for the entire dataset with cognac

v1.0 [32]. Relevant isolates for understanding the population structure and geographic

distribution of the isolates collected as part of the EIP study were identified as any isolate within

25 substitutions or the nearest phylogenetic neighbor by alignment distance. This yielded a

subset of the 5,346 isolates, including study isolates and public isolates; and a second,

concatenated, core gene alignment was generated with cognac for these most relevant isolates,

returning the nucleotide alignment for more precise distance comparisons. To generate sequence

type specific alignments, these data were divided into relevant clonal groups. Any clonal group

with less than 10 members was merged into a group of its respective species. Outgroups were

identified as the isolate with the lowest substitution distance to an isolate assigned to that clonal

group or species, for clonal group or species alignments, respectively. A core, concatenated gene

alignment for each set of isolates corresponding to species/clonal group was generated using

cognac with the corresponding outgroup specified, returning the nucleotide alignment.

Recombinant positions were removed with Gubbins v3.0.0 [24]. For alignments with more than

1000 isolates included, the alignments were randomly downsampled into subsets of 250

sequences, including the outgroup in each alignment, and gubbins was run on the downsampled

alignment. This was performed in triplicate, and any position identified as recombinant in the

three runs was masked from the alignment. The recombination filtered alignments were then

input to IQ-TREE v1.6.12 [33] to generate maximum likelihood phylogenies.

bla$_{KPC}$ identification

The *bla*$_{KPC}$ gene was identified using the CD-HIT v4.7 [34] data generated with cognac. The

gene cluster containing coding sequences annotated by RAST v1.035 [31] as *bla*$_{KPC}$ was

identified. *bla*$_{KPC}$ status was determined for each genome as having representation in the *bla*$_{KPC}$

cluster.

<u>Integrated genomic and epidemiologic data analysis</u>

Identification of putative geographic importation and HGT events

For each core genome tree, maximum likelihood ancestral reconstruction of each study site

location (MN, TN, CT) and presence of *bla*$_{KPC}$ presence was performed individually in R v4.0.2

[35] using the ace() function in ape v5.5 [36]. Nodes with an ancestral reconstruction confidence

of < 0.875 were discarded. All transition events (for each geographic location and *bla*$_{KPC}$) were

then mapped onto the edges of the phylogeny. Each study isolate tip was traversed towards the

root of the tree until the first transition event was identified. This transition event was considered

a putative geographic importation if the transition was a geographic transition and a putative

HGT event if the transition was for *bla*$_{KPC}$. Study isolates with the same edge transition event

were considered part of a cluster. We consider the number of transition events for a given

geographic location/HGT event to be the number of identified clusters. Some study isolates were

unable to be classified using this method and are considered of unknown origin. These isolates

were clustered together if they appeared as a pure cluster on the phylogeny.

1002

Intra- and inter-facility transmission analysis

Pairwise single nucleotide variant (SNV) distances for isolates from each sequence alignment

were calculated in R v4.0.2 [35] using the dist.dna() function in ape v5.5 [36] with

pairwise.deletion = TRUE. These pairwise SNV distances were classified as intra- and inter-

facility pairs using the get_pair_types() function in regentrans v0.1 [37]. This function was also

used to identify closely related pairs within a given facility.

1009

1010 *Individual level patient healthcare exposure analysis*

1011 For each inter-facility patient pair, we determined whether they had a shared healthcare exposure

1012 (data collection method described above). Then we compared the extent of exposure for pairs of

1013 patients from different facilities with isolates ≤10 SNVs compared to >10 SNVs using a Fisher's

1014 Exact test [38].

1015

Aggregate patient transfer analysis

For each state, we subset the aggregate patient transfer network (described above) to only

include the facilities in our dataset or ones that are connected to one of the facilities in our

dataset by at least one patient transfer event over the course of a year. We then identified paths of

maximum patient flow between this subset of facilities using the get_patient_flow() function in

regentrans v0.1 [37]. Patient flow was then compared to the minimum pairwise SNV distance of

isolate pairs from each facility pair using a Wilcox test comparing facilities where the minimum

pairwise SNV distance was ≤10 with those where it was >10.

1024

Comparison of ST171 and ST258 transmission patterns in MN

Each isolate was paired to their closest isolate in another facility with a subsequent collection

date. To assess whether *E. cloacae* ST171 and *K. pneumoniae* ST258 in MN had more

transmission surrounding a central facility than expected by chance, we calculated empirical p-

values as follows. We define *n* as 2x the total number of closely related inter-facility pairs (i.e.

edges ≤10 SNVs different; see **Figure 4**), because each pair is connected to 2 facilities. We then

identified facilities that were connected by at least one closely related edge. Next, we randomly

sampled these facilities with replacement *n* times, counted the number of times each facility

appeared, found the proportion of total edges connected to that facility, and took the maximum.

We repeated this process 1000x and compare dthe maximum proportion from the sampled data

to the actual maximum proportion.

1036

1037

1038

 Supplemental table and figures

 Table S1: Number of each facility type by state and how many facilities were represented

 in whole-genome sequencing (in parentheses).

Facility type	CT	MN	TN
ACH	30 (19)	50 (23)	92 (31)
LTACH	3 (0)	2 (2)	10 (4)
Other	7 (0)	96 (6)	48 (1)
SNF	224 (0)	366 (9)	309 (0)
Unknown	0 (3)	0 (20)	0 (4)

 Table S2: Facility type pairs of inter-facility isolate pairs with a pairwise SNV distance ≤10.

State	ST	ACH-ACH	ACH-LTACH	ACH-SNF	Other
CT	*K. pneumoniae* ST258	2	0	0	0
CT	*K. pneumoniae* ST307	1	0	0	0
CT	Other	1	0	0	2
MN	*E. hormaechei* ST114	2	5	1	1
MN	*E. hormaechei* ST171	109	92	56	101

MN	*K. pneumoniae* ST258	9	1	6	42
MN	*K. pneumoniae* ST307	1	0	0	2
MN	Other	1	0	0	1
TN	*E. hormaechei* ST114	9	1	0	2
TN	*E. hormaechei* ST171	1	1	0	2
TN	*K. pneumoniae* ST258	13	6	0	0
TN	*K. pneumoniae* ST307	17	0	0	0
TN	Other	6	1	0	3

1047 Abbreviations: SNV=single nucleotide variant; ST=sequence type; ACH=acute care hospital;

1048 LTACH=long-term acute care hospital; SNF=skilled nursing facility

1049

1050 **Table S3: Patients with closely related isolates from the same facility with the same ST do**

1051 **not share comorbidities more than patients with distantly related isolates.**

State	ST	Shared comorbidity (%)		Fisher's exact p
		Pairwise SNV distance ≤10	Pairwise SNV distance > 10	
CT	*K. pneumoniae* ST258	1/1 (100)	5/10 (50)	1
MN	*E. hormaechei* ST114	1/1 (100)	0/2 (0)	0.3

	E. hormaechei ST171	6/86 (0.07)	17/229 (0.07)	1
	K. pneumoniae ST258	0/25 (0)	3/102 (0.03)	1
	E. hormaechei ST114	1/1 (100)	6/9 (0.67)	1
	E. hormaechei ST171	0/0	1/3 (50)	1
TN	*K. pneumoniae* ST258	2/5 (40)	5/11 (45)	1
	K. pneumoniae ST307	5/8 (0.625)	7/7 (100)	0.2

1052　Abbreviations: SNV=single nucleotide variant; ST=sequence type

1053

1054

1057

Figure S1: Distribution of species and sequence types across healthcare facilities with at least one identified CRE with *bla*KPC. X axis shows facilities. Grey is any other ST of the species corresponding to that faceted plot.

1061

1062

1063

1064 **Figure S2: Information about public isolates downloaded from the PATRIC database that**

1065 **were included for context in the investigation of importation and horizontal gene transfer.**

1066 (A) Distribution of year of isolation for isolates where this information was provided. (B)

1067 Distribution of location and KPC presence/absence.

1068

1069

1070 **Figure S3: Phylogenies used to identify importation and HGT events.**

1071

1072

1073 **Figure S4: HGT locations.** In-state means that the KPC- ancestor was in the same state,

1074 unknown means the KPC- ancestor may have been in-state or out-of-state.

1075

Figure S5: Fraction of intra-facility isolate pairs for pairwise SNV distances to help identify

pairwise SNV distance cutoffs. We expect there to be an enrichment in closely related intra-

facility pairs under the assumption that intra-facility transmission is more common than inter-

facility transmission. This plot helps us identify potential pairwise SNV distance thresholds by

looking for a SNV distance where there is a decrease in the fraction of intra-facility isolate pairs

(around 5 and 10 appear to be reasonable thresholds for a sensitivity analysis). SNV = single

nucleotide variant.

1083

1084

Figure S6: Pairwise single nucleotide variant (SNV) distances of all STs. SNV = single

nucleotide variant. Vertical line in each plot represents 10 SNVs, used as a cutoff for

determining what isolates are considered closely related (note that we also did sensitivity

analyses with ≤ 5, 10, and 15 SNVs).

1089

1090

Figure S7: Pairwise patristic distances concordant with pairwise SNV distance thresholds.

1092

1093

Figure S8: The relatedness of isolates within facilities varies. For a given facility, sometimes

the isolates are relatively closely related, while sometimes they are distantly related. SNV =

single nucleotide variant.

1097

1098

Figure S9: Aggregate-level patient transfer data provides insight into pathways of

transmission. Each point is a facility pair. Patient flow was calculated using the

get_patient_flow function in regentrans and is a metric of the extent of patient transfer between

facilities that considers patient transfer to intermediate facilities along the path between two

facilities in the patient transfer network. Both the y and x axis are shown on a log10 scale and

points are jittered for visualization purposes. Wilcox p-values were calculated by comparing

patient flow between facility pairs with at least one isolate pair ≤10 SNVs apart and facility pairs

with no isolate pairs ≤10 SNVs apart. For most sequence types, facility pairs with closely related

isolates tend to have higher patient flow between them.

1108

1109

Figure S10: Patients from different facilities with closely related isolates are sometimes linked by a shared facility exposure. Nodes are facilities, line width indicates number of isolate pairs ≤10 SNVs shared by the two facilities. SNV = single nucleotide variant.

1113

1114

1115

1116

1117

1118 **Figure S11: Differences between individual- and aggregate-level patient transfer analysis**

1119 **results.**

1120

1121

Figure S12: Comparison of actual and randomized distributions of facilities with closely related inter-facility isolate pairs. See supplemental material for details.

1124

Figure S13: Facility W is more connected in the MN healthcare patient transfer network than most other healthcare facilities. Facilities are weighted by the number of patient transfers, generating a directed graph. Facility W is shown as the red diamond. For closeness, degree, and strength the in- and out- values were summed (from and to a facility).

1125

1126

1127

1128

1129

1130

1131