

1 **Multimorbidity and its associated factors in Indonesia**
2 **through universal health coverage scheme: A cross-sectional study**
3 **based on national claims data**

4 Noor Afif Mahmudah¹, Mesfin Kassaye Tessma^{2*}, Yodi Mahendradhata³

5

6 ¹Family and Community Medicine Department, Faculty of Medicine, Public Health and
7 Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

8 ²Department of Learning, Informatics, Management and Ethics, Karolinska Institutet,
9 Stockholm, Sweden

10 ³Health Policy and Management Department, Faculty of Medicine, Public Health and
11 Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

12

13 *Corresponding author

14 Email: mesfin.kassaye.tessma@ki.se

15

16

17 **Abstract**

18 Multimorbidity has been increasing globally and is usually associated with higher health
19 care utilization and costs. Indonesia has been implementing Universal Health Coverage
20 (UHC) program since 2014. However, there is a limited study of the prevalence of
21 multimorbidity and its impact on health care utilization and costs through the UHC
22 scheme in Indonesia. This study aimed to determine the prevalence of multimorbidity and
23 its associated factors, particularly the health care utilization and costs of patients with
24 multimorbidity compared to patients with one chronic condition in the hospitals in

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

25 Indonesia based on the UHC scheme. The study was a comparative cross-sectional
26 design. The data was collected from the Social Security Agency for Health/Badan
27 Penyelenggara Jaminan Sosial Kesehatan (BPJS Health) National Sample Data 2015-
28 2016. All patients ≥ 60 years of age and have at least one chronic condition in the hospital
29 were included. Descriptive statistics, bivariate analysis, and multivariable regression
30 analysis were conducted to analyze the data. In a sample of 23,460 patients, the
31 prevalence of multimorbidity was 44.4% among patients with chronic conditions in the
32 hospital. We observed significant difference in gender, marital status, and membership
33 segmentations between patients with multimorbidity and one chronic condition ($p < 0.05$).
34 Health care utilization and costs were significantly higher in multimorbid patients
35 ($p < 0.001$). This positive association between multimorbidity and health care utilization
36 (OR: 1.70, 95% CI: 1.61-1.79) and health care costs (unstandardized coefficient 0.483,
37 95% CI: 0.443-0.524, $p < 0.001$) remains significant after adjusting for age and gender.
38 The analysis found that the prevalence of multimorbidity was high and positively
39 associated with higher health care utilization and costs through the UHC scheme in
40 Indonesia. Therefore, health policymakers and healthcare professionals need to consider
41 the burden of multimorbidity more when structuring health care.

42

43 **Keywords:** *Multimorbidity, health care utilization, health care costs, prevalence,*
44 *universal health coverage*

45

46 **Introduction**

47 Multimorbidity is a coexistence of two or more chronic conditions in a single individual
48 [1,2]. Compared to patients without a chronic condition and patients with only one

49 chronic condition, multimorbidity is usually associated with higher health care utilization
50 and cost, including more frequent hospital admission, longer hospital stays, and a need
51 for a larger number of different medical specialists during a year [3–5]. Currently, the
52 prevalence of multimorbidity is increasing both in high-income countries (HICs) and low
53 and middle-income countries (LMICs), which increases with age, including in Indonesia
54 [4,6].

55 Indonesia has been implementing Universal Health Coverage (UHC) program
56 since 2014. It is implemented by the Social Security Agency for Health (Badan
57 Penyelenggara Jaminan Sosial Kesehatan/BPJS Health) and covers 84.2% of the
58 population in 2020[7,8]. In 2018, BPJS Health reported that chronic conditions were the
59 highest claimed health expenditure with total of 42,171,533 cases and IDR
60 8,806,699,514,190 (equivalent to 638,239,743.66 USD) in outpatient care of hospitals
61 [9].

62 To reduce the high treatment and financial burden of patients with chronic
63 conditions, it is crucial to analyze the prevalence, associated factors, particularly the
64 health care utilization and costs in patients, both with multiple chronic conditions
65 (multimorbidity) and one chronic condition in Indonesia [3]. However, there is a limited
66 study of the prevalence of multimorbidity and its impact on health care utilization and
67 costs through the UHC scheme in Indonesia. This study aimed to determine the
68 prevalence of multimorbidity and its associated factors, particularly the health care
69 utilization and costs of patients with multimorbidity compared to patients with one
70 chronic condition in the hospitals in Indonesia based on the UHC scheme.

71

72 **Methods**

73 This study used a comparative cross-sectional design that determined the prevalence of
74 multimorbidity and compared the health care utilization and costs between patients with
75 multimorbidity and one chronic condition. We conducted and reported our study
76 following the STROBE guideline [10].

77 **Sample and data**

78 The data were collected from the BPJS Health National Sample Data from 2015 – 2016
79 in hospitals [11]. BPJS Health National Sample Data is an open-source database that
80 provides information about the representative of the members nationally at the individual
81 level. It comprises the characteristics of BPJS Health members, health care utilization,
82 diagnosis, and related costs. The BPJS Health National Sample Data were collected by
83 stratified random sampling method from the database of all members per December 31st,
84 2016. Each case has a different weight based on the stratification to reduce the bias
85 between sample and population [12].

86 In this study, we included all patients aged 60 years, and older who were
87 diagnosed with at least one chronic condition in hospital providers. This age group was
88 chosen considering the life expectancy and the high prevalence of multimorbidity from
89 a previous study in Indonesia [13]. The hospital setting was chosen because it was the
90 only setting that provided data with more than one diagnosis to classify the cases into a
91 patient with multimorbidity or a patient with one chronic condition. In this study, 19
92 chronic conditions were included based on the multimorbidity prevalence in Indonesia
93 and ICD-10 codes used in the previous study [14,15] (Table 1).

94

95

Table 1. List of 19 chronic conditions based on ICD-10 codes

No.	Chronic condition	ICD-10 codes
1.	Hypertension	I10 – I15
2.	Lipid metabolism disorders	E78
3.	Severe vision reduction	H17 – H18, H25 – H28, H31, H33, H34.1 – H34.2, H34.8 – H.34.9, H35 – H36, H40, H43, H47, H54
4.	Joint arthrosis	M15 – M19
5.	Diabetes mellitus	E10 – E14
6.	Chronic ischemic heart disease	I20, I25, I21
7.	Cardiac arrhythmias	I44 – I45, I46.0, I46.9, I47 – I48, I49.1 –I49.9
8.	Obesity	E66
9.	Hyperuricemia/gout	E79, M10
10.	Liver disease	K70, K71.3 – L71.5, K71.7, K72.1, K72.7, K72.9, K73 – K74, K76
11.	Depression	F32 – F33
12.	Asthma/COPD	J40 – J45, J47
13.	Atherosclerosis/PAOD	I65 – I66, I67.2, I70, I73.9
14.	Cerebral ischemia/chronic stroke	I60 – I64, I69, G45
15.	Cardiac insufficiency	I50
16.	Severe hearing loss	H90, H91.0, H91.1, H91.3, H91.8, H91.9
17.	Rheumatoid arthritis/chronic polyarthritis	M05–M06, M79.0
18.	Cancer	C00–C14, C15–C26, C30–C39, C40–C41, C43–C44, C45–C49, C50, C51–C58, C60–C63, C64–C68, C69– C72, C73–C75, C81–C96, C76–C80, C97, D00–D09, D37–D48

19. Tuberculosis A15 – A19

96

97 Twenty-three thousand four hundred sixty subjects in the database that meet the
98 inclusion criteria in the study were identified and included in this study (Fig 1). Subjects
99 were then classified into with one chronic disease and with multimorbidity, two or more
100 chronic diseases.

101

102 **Fig 1. Selection of study sample from BPJS Health National Sample Data**

103

104 **Variables**

105 **Sociodemographic and Illness characteristics**

106 The sociodemographic characteristics are determined by age group, gender, marital
107 status, region, and membership segmentation. Membership segmentation of BPJS Health
108 was classified into Premium Assistance Beneficiaries/*Penerima Bantuan Iuran* (PBI),
109 who receives the assistance of contribution to BPJS health from government; and Non-
110 Premium Assistance Beneficiaries/Non-*Penerima Bantuan Iuran* (Non-PBI), whose
111 premium to BPJS Health is paid by the employers with contribution from employees, or
112 themselves, or government (for the pensioners and veterans). Meanwhile, the illness
113 characteristics are the severity level of diseases and the group of chronic conditions. The
114 severity level of diseases ranged from level 0-3, with level 0 being the mildest, and level
115 3 being the most severe. Whereas the group of chronic conditions is based on the primary
116 diagnosis of patients: cardiovascular, respiratory, special senses, cancer, endocrine,
117 musculoskeletal, mental health [9,12,16].

118 **Outcome variables**

119 Health care utilization and costs were determined as the primary outcome variables.

120 Health care utilization is a description of the use of health care services by persons to
121 prevent and cure health problems [17]. This study includes the type of service (inpatient,
122 outpatient), type of hospital (type A, type B, type C, type D, others), and length of stay.

123 The health care costs were according to the mix case-based group tariff rates,
124 namely Indonesia Case-Based Groups (INA-CBGs), which follow the code of disease,
125 severity level of disease, type of hospitalized room, and the number of secondary
126 diagnosis[18,19]. The costing of INA-CBGs tariff rates was based on the overhead cost,
127 intermediate cost, and final cost. The overhead cost is the cost for general hospital
128 support, for example, the administration, maintenance, communication, cleaning service,
129 information system, and nutrition. The intermediate cost is the medical hospital support,
130 for instance, the pharmacy, radiology, laboratories, Intensive Care Unit (ICU), and
131 hemodialysis. Meanwhile, the final cost is the medical care cost, including the outpatient
132 and inpatient care [18,19]. The period for the cost for each individual is 2015 - 2016. The
133 prices of INA-CBGs costs has been set by Ministry of Health, Republic of Indonesia in
134 2013 [19,20].

135 **Statistical analysis**

136 Differences in patient characteristics between multimorbidity and one chronic condition
137 were examined using a chi-square or a Fischer's Exact test for categorical variables and
138 a t-test for continuous variables. However, Mann-Whitney was performed for ordinal
139 variables and in the case of a deviation from normality. For the ordinal variable level of
140 exposure, we employed extended Mantel-Haenzel Chi-square for linear trend.

141 Multivariable regression was employed to control potential confounders and
142 examine the relationship between the outcome and explanatory variables. Multiple linear
143 regression was used to estimate the relationship between the continuous outcome variable
144 (health care costs) and multimorbidity while controlling for other predictor or explanatory
145 variables. Log transformation was applied to health care costs due to its high skewness.
146 Multiple logistic regression was performed to examine the associations between the
147 multimorbidity and demographic and other relevant explanatory variables. Explanatory
148 variables were selected based on earlier research and univariable simple regression. The
149 crude association of each explanatory variable was determined to examine its relationship
150 with the outcome variable in univariate models.

151 Upon completing the univariate analyses, we selected variables for the multivariable
152 analyses. Any variable whose univariate test had a P-value <0.15 was considered a
153 candidate for the multivariable model along with variables of known practical
154 importance. Once the variables were identified, they were entered into a multivariable
155 model. Normal probability plots and residual plots were performed to assess violations
156 of assumptions, and a Cook's distance was examined to assess outlier problem. All
157 analyses were conducted using IBM SPSS Statistic version 25.

158

159 **Results**

160 **Sociodemographic and Illness characteristic of patients with** 161 **multimorbidity versus one chronic condition**

162 The total sample consisted of 23,460 patients with age equal to or more than 60 years and
163 one or more medically diagnosed chronic conditions in the hospital providers. Of those

164 individuals, 55.6% were patients with one chronic condition (n=13,034) and 44.4% were
 165 patients with multimorbidity (n=10,426). The mean (SD) age of the total sample was 70.6
 166 (± 8.1), with 70.6 (± 8.1) for the patients with one chronic condition and 70.8 (± 8.0) for
 167 the multimorbidity patients.

168 Descriptive statistics of the sociodemographic and Illness characteristics of
 169 multimorbidity versus one chronic condition patients are shown in Table 2. We observed
 170 a statistically significant difference in some sociodemographic characteristics: gender,
 171 marital status, and membership segmentation.

172

Table 2. Sociodemographic characteristics of population aged ≥ 60 years with one chronic condition and multimorbidity in hospitals

Characteristics n (%)	Total study sample (n = 23460)	One chronic condition (n = 13034)	Multimorbidity (n = 10426)	Sig
Gender:				*
Male	11760 (50.1)	6455 (49.5)	5305 (50.9)	
Female	11700 (49.9)	6579 (50.5)	5121 (49.1)	
Age group:				-
60 – 69	12240 (52.2)	6897 (52.9)	5343 (51.2)	
70 – 79	7692 (32.8)	4178 (32.1)	3514 (33.7)	
≥ 80	3528 (15.0)	1959 (15.0)	1569 (15.0)	
Marital status:				***
Married	13351 (56.9)	7539 (57.8)	5812 (55.7)	
Unmarried	10109 (43.1)	5495 (42.2)	4614 (44.3)	
Region:				-
Java Bali	13052 (55.6)	7309 (56.1)	5743 (55.1)	
Others	10408 (44.4)	5752 (43.9)	4683 (44.9)	

Membership				***
segmentation:				
PBI	6907 (29.4)	3671 (28.2)	3236 (31.0)	
Non-PBI	16553 (70.6)	9363 (71.8)	7190 (69.0)	

***p-value <0.001 **p-value<0.01 *p-value<0.05 -non-significant.

PBI: contribution beneficiaries

173

174 A significant difference was not observed in the age and region of patients with
 175 one chronic condition and multimorbidity. However, there was an increase in odds ratio
 176 among the age groups in multimorbidity (age group 60-69 as reference, OR=0.967, CI
 177 95%:0.897-1.043 in the age group of 70-79, OR=1.05, CI 95%: 0.969-1.138 in age group
 178 ≥ 80).

179 Table 3 demonstrated illness characteristics. For the severity level of Illness,
 180 patients with one chronic condition had a higher proportion of level 0 compared to
 181 multimorbid patients. However, at levels 1-3, the proportion was higher in multimorbid
 182 patients compared to patients with one chronic condition. Besides, for the prevalence of
 183 chronic conditions, the three most frequent groups of chronic conditions were
 184 cardiovascular, respiratory, and endocrine. In contrast, the least frequent was mental
 185 health.

Table 3. Illness characteristics of population aged ≥ 60 years with one chronic condition and multimorbidity in hospitals

Characteristics n (%)	Total study sample (n = 23460)	One chronic condition (n = 13034)	Multimorbidity (n = 10426)	Sig
Severity level of Illness:				***
Level 0	13174 (56.2)	8067 (61.9)	5107 (49.0)	
Level 1	6276 (26.8)	2968 (22.8)	3308 (31.7)	

Level 2	231 (10.8)	1247 (9.6)	1284 (12.3)
Level 3	1479 (6.3)	752 (5.8)	727 (7.0)
Group of chronic conditions:			***
Cardiovascular	6546 (50.4)	2109 (43.8)	4436 (54.3)
Respiratory	2057 (15.8)	811 (39.4)	1246 (15.2)
Endocrine	1718 (13.2)	669 (13.9)	1049 (12.8)
Special senses	1551 (11)	546 (11.4)	1005 (12.3)
Cancer	580 (4.5)	367 (7.6)	213 (2.6)
Musculoskeletal	539 (4.1)	302 (6.3)	237 (2.9)
Mental health	8 (0.1)	6 (0.1)	2 (0)

***p-value <0.001 **p-value<0.01 *p-value<0.05 -non-significant.

186

187 **Health care utilization and costs of patients with** 188 **multimorbidity versus one chronic condition**

189 As Table 4 shows, the proportion of patients admitted to the hospital (inpatient) was
190 higher among patients with multimorbidity than patients with one chronic condition. The
191 length of stay also was higher in patients with multimorbidity than patients with one
192 chronic condition. However, the proportion of hospital type in patients with one chronic
193 condition was similar to patients with multimorbidity in which the majority utilize the
194 type C hospitals. The mean length of stay in days was 5.05 and 5.25 for patients with one
195 chronic condition and patients with multimorbidity, respectively.

Table 4. Health care utilization and costs in population aged ≥ 60 years with one chronic condition and multimorbidity in hospitals, 2015-2016

Health care utilization	Total study sample	One chronic condition	Multimorbidity	Sig
n (%)	(n = 23460)	(n = 13034)	(n = 10426)	

Type of service:				***
Inpatient	10286 (43.8)	4967 (38.1)	5319 (51.0)	
Outpatient	13174 (56.2)	8067 (61.9)	5107 (49.0)	
Type of hospital:				***
Type A	1209 (5.2)	770 (5.9)	439 (4.2)	
Type B	8331 (35.5)	4681 (35.9)	3650 (35.0)	
Type C	10333 (44.0)	5741 (44.0)	4592 (44.0)	
Type D	2597 (11.1)	132 (10.1)	1276 (12.1)	
Others	990 (4.2)	521 (4.0)	469 (4.5)	
Length of stay, days	4 (4)	4 (4)	4 (5)	-
median (IQR)				
Health care costs, IDR	307 450.00	190 900.00	2 160 050.00	***
median (IQR)	(3 552 200)	(3 034 175)	(4 117 800)	

***p-value <0.001 **p-value<0.01 *p-value<0.05 -non-significant. IDR: Indonesian Rupiah,

IQR: interquartile

196

197 For the health care costs, multimorbid patients also had higher costs than patients
 198 with one chronic condition (Table 4). The average health care costs in patients with one
 199 chronic condition was IDR 2,122,912.19 per year (equal to USD 140.47). While the
 200 average health care costs in patients with multimorbidity was IDR 2,884,995.96 per year
 201 (equal to USD 190.90).

202 **Underlying factors associated with health care utilization and** 203 **costs of patients with multimorbidity**

204 In the multiple logistic regression model on health care utilization, multimorbidity was
 205 positively associated with inpatient care. The odds ratio (OR) in multimorbidity for
 206 inpatient care remains statistically significant, and the association positive when corrected

207 for gender and age. Thus, adjusting for gender and age, the inpatient costs is higher for
208 multimorbid patients. Table 5 shows no remarkable difference in the unadjusted and
209 adjusted OR.

Table 5. Results of multiple logistic regression model on the dependent variable of inpatient health care utilization (yes/no) in population aged ≥ 60 years

Model	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
Multimorbidity	1.69 (1.61-1.78)	1.70 (1.61-1.79)
Age group	1.47 (1.42-1.53)	1.48 (1.43-1.53)
Gender	0.92 (0.88-0.97)	0.92 (0.87-0.97)

OR: Odds ratio, CI: Confidence intervals

210

211 In addition, a multiple linear regression was performed to determine if the
212 observed variations in health care costs were either attenuated or eliminated after
213 adjusting for demographic confounding variables. The dependent variable was the log
214 health care costs. After adjustment, statistically significant differences were observed in
215 the variables multimorbidity, gender, and age group. Multimorbidity remained
216 statistically significant after adjusting for gender and age group with higher Beta values.
217 Thus, multimorbidity was observed to be a significant positive predictor of log health
218 care costs, regression coefficient (B) = .48 (95% CI = .44, .52, $P < 0.001$). Female gender
219 was negatively related to log health care costs, $B = -0.07$ (95% CI = $-.11, -.03$, $P < 0.001$).
220 Age group was positively related to the dependent variable, $B = .30$ (95%CI= .27, .33,
221 $P < 0.001$).

222 The coefficient of multimorbidity indicated that for patient with multimorbidity,
223 it was expected that the log costs increase by an average of 0.48. The coefficient of age
224 group indicated that for increasing age group, the log costs are expected to increase by an

225 average of 0.30 The coefficient of gender indicated that in female gender compared to
 226 male, the log costs expected to reduce by an average of 0.07. Hence, multimorbidity,
 227 gender, and age group were significant predictors of log health care costs (Table 6).

Table 6. Results of multiple linear regression model for the dependent variable log health care costs, in population aged ≥ 60 years

Predictor	Unstandardized		Standardized	t	Sig.	95% Confidence	
	Coefficients		Coefficients			Interval for B	
	B	Std. Error	Beta			Lower Bound	Upper Bound
Multimorbidity	.483	.021	.148	23.094	***	.443	.524
Gender	-.070	.021	.134	20.970	***	-.112	-.030
Age group	.298	.014	-.022	-3.394	***	.270	.326

***p-value <0.001

228

229

230 Discussion

231 Principal findings

232 Prevalence of multimorbidity in Indonesia based on Indonesia UHC

233 program

234 In the hospitals, half of the patients aged 60 years and above had at least one
 235 chronic condition, in which 44% had multimorbidity. This prevalence of multimorbidity
 236 was lower than the previous study in Indonesia by Hussain et al. [14], which stated that
 237 half of the individuals over 60 years in Indonesia had multimorbidity. This difference is

238 possible due to the limited population in the hospitals in the current study rather than in
239 all health care settings.

240 The sociodemographic characteristics which significantly different were gender,
241 marital status, and membership segmentation. In contrast to majority studies, the males
242 had a higher prevalence of multimorbidity than females in this study [5,14]. However,
243 this reverse trend was also observed in other study [21]. The finding for marital status
244 was in line with previous studies [22,23]. This proportion might be due to the high number
245 of people that married among the population aged 60 years and above [24]. Additionally,
246 the membership segmentation that reflected the socioeconomic situation of the patients
247 was also in line with other studies [14,25]. A previous study in Indonesia [14] also showed
248 that higher socioeconomic status is more vulnerable to multimorbidity as the
249 epidemiological transition in LMICs. The wealthier stratum has a higher risk for Non-
250 Communicable Diseases in LMICs with higher levels of health risk such as high
251 cholesterol, waist-hip ratio, and reduced physical activity [22].

252 On the other hand, the mean of age, proportion of age group and region were not
253 significantly different in multimorbid patients compared to the patients with one chronic
254 condition. It might be due to the current epidemiological transition in Indonesia, which
255 leads to chronic conditions in all ages and across the geographic area [26]. However, the
256 prevalence of multimorbidity increased with age, which aligned with numerous studies
257 [5,13,27].

258 The illness characteristics of severity level and group of chronic conditions were
259 significantly different. In alignment with previous studies [22,28], the severity level of
260 Illness showed that the multimorbidity had a higher proportion in more severe conditions.
261 The negative effect on multimorbidity in elderly age groups has been reported from high-

262 and low-income settings [4,27]. The prevalence of a group of chronic conditions confirms
263 some of the findings from a previous study in Indonesia [14], including that the most
264 frequent chronic conditions group was cardiovascular.

265

266 **Health care utilization and costs of patients with multimorbidity and** 267 **with one chronic condition**

268 The results showed that health care utilization and costs were higher in patients
269 with multimorbidity than patients with one chronic condition. The inpatient hospital
270 admission was higher in multimorbid patients. Moreover, the length of stay was also
271 longer compared to patients with one chronic condition. These findings confirm that
272 multimorbidity is related to higher health care utilization [4,5,29,30]. A cross-sectional
273 study in LMICs showed that visits to doctors in secondary care and hospitalization
274 increased with multimorbidity [31].

275 The utilization of different types of hospital was also significantly different
276 between patients with multimorbidity and one chronic condition. Nevertheless, the
277 proportion of hospital type in patients with one chronic condition was similar to patients
278 with multimorbidity in which the majority utilize the type C hospitals. This proportion is
279 possibly due to the tiered referral system of BPJS Health from primary care to type D or
280 C hospitals, then to type B, and type A as top referral [16].

281 Health care costs were higher in patients with multimorbidity than with one
282 chronic condition, which is similar to other studies [27,30,32]. For instance, a recent
283 systematic review study, which included studies from the United States, Europe, Canada,
284 stated that there was a positive correlation between multimorbidity and health care costs

285 [27]. Glynn et al. also reported that health care costs were significantly increased among
286 multimorbid patients [27].

287

288 **Underlying factors associated with health care utilization and costs of** 289 **patients with multimorbidity**

290 This study found a positive association between the utilization of inpatient health
291 services and multimorbidity. This association remains positive and increased to 70% after
292 adjusting for gender and age. Age group and gender also significantly associated with
293 health care utilization. In contrast, the female gender had negative association to the type
294 of services.

295 Those findings were in line with several previous studies [27,32,33], both in HICs
296 and LMICs. Studies in high-income countries found that hospital admission was
297 significantly increased with an increase in the number of chronic conditions after
298 adjusting for age, gender, and free medical care eligibility [27,32,34]. A study in low-
299 middle income countries found that the increased number of chronic conditions was
300 significantly associated with higher health care utilization, including hospitalization and
301 duration of stay in hospital [35].

302 Multimorbidity also had a positive association with health care costs. The result
303 showed that multimorbidity was a positive predictor of health costs in alignment with
304 other research findings [35]. The increased costs may be related to the associated costs of
305 caring for individuals with multimorbidity problems due to increased medical
306 expenditures. Thus, multimorbidity also incurs significant social and economic burdens
307 due to complex health and welfare demands and higher health costs.

308 Increasing age group also was shown to be a positive predictor. Consistent with
309 the previous study which have found that this increase in health care costs is reasonable
310 considering the ageing population with multimorbidity [36]. However, female gender was
311 a negative predictor. A recent study also reported that the combination of diseases and
312 intensity of multimorbidity differed in older men and women [36]. Further, several
313 studies also found that the number of chronic conditions was associated with higher total
314 health care costs after controlling for age and gender, severity of illness, and hospital
315 characteristics [37–39].

316

317 **Strength and limitation**

318 To the best to our knowledge, this study is the first to determine the multimorbidity and
319 its associated factors in the Indonesian population with an age equal to or more than 60
320 years using data from BPJS Health National sample data. A major strength of the current
321 study was the large stratified randomly selected representative sample of the Indonesian
322 population through BPJS Health. Thus, minimizing selection bias and enables for larger
323 generalization of results. Nevertheless, there are some study limitations.

324 The study was cross-sectional, which did not allow to determine any causal
325 interferences. Our findings may have underestimated the prevalence of multimorbidity
326 since the scope was limited to hospital settings and based on the claims data. Nonetheless,
327 the diagnosis of chronic conditions was ensured following the ICD-10 diagnosis rather
328 than self-report. **The current study** may also have underestimated the health care
329 utilization and costs of multimorbidity. The data only provided the total health care costs
330 based on INA-CBGs rather than in specific detail of each unit of costs. This limited data
331 made the study unable to examine the health care costs of multimorbidity. Lastly, the

332 administrative claims also did not record more details in economic and social factors such
333 as income and education levels that might independently influence the health care
334 utilization and costs.

335 **Implications to Health Care System**

336 The prevalence of multimorbidity among patients with chronic conditions and the
337 positive association between multimorbidity with higher health care utilization and costs
338 in BPJS Health should urgently be addressed by health care professionals and policy
339 makers in Indonesia.

340 Multimorbidity is a complex disease, and its management needs to be improved
341 [5,40]. However, the current administration has been focused on single conditions
342 approach rather than multiple conditions [5,40,41]. Therefore, cooperation, coordination,
343 and communication between different levels of health care and specialties need to be
344 increased. It is also crucial to develop an integrated care pathway and guideline to manage
345 the multimorbidity. Further, cost-effective management of multimorbidity and their risk
346 factors should also be improved [5].

347 Due to the high burden of multimorbidity in BPJS Health, the health policymakers
348 and health care professionals should promote a healthy lifestyle to prevent the risk factors.
349 BPJS Health should also expand the coverage to preventive measures than focus only on
350 curative to avoid the high burden of health financing, particularly multimorbidity in the
351 future [16,26].

352 The findings showed the association and impact of multimorbidity in health care
353 utilization and costs, particularly in Indonesia through UHC scheme. However, further
354 research is needed to understand the pattern of chronic conditions in multimorbidity.
355 Cluster analysis by grouping similar diseases may improve the knowledge of how

356 multimorbidity as complex disease impact health care utilization and costs. A study with
357 more comprehensive data from different health care levels, including sociodemographic
358 of respondents, could provide further context for multimorbidity research.

359

360 **Conclusions**

361 This study is a step forward to examine the impact of multimorbidity on health care
362 utilization and costs in Indonesia through UHC scheme. The study found a 44.4%
363 prevalence of multimorbidity among patients with chronic conditions in hospitals.
364 Multimorbidity also had a positive association with higher health care utilization and
365 costs compared to patients with one chronic condition. Therefore, health policymakers
366 and health care professionals need to take the burden of multimorbidity more into
367 consideration when structuring health care.

368 Further research could be directed toward the pattern of chronic conditions in
369 multimorbidity with more comprehensive data from different levels of health care.

370

371 **Acknowledgement**

372 The authors express sincere gratitude to BPJS Health for providing access to the data.

373

374 **References**

- 375 1. Mercer S, Furler J, Moffat K, Fischbacher-Smith D, Sanci LA, Organization WH,
376 et al. Multimorbidity. 2016.
- 377 2. The Academy of Medical Sciences. Multimorbidity a priority of global health
378 research. London, UK: The Academy of Medical Sciences; 2018.

- 379 3. Hopman P, Heins MJ, Rijken M, Schellevis FG. Health care utilization of patients
380 with multiple chronic diseases in The Netherlands: Differences and underlying
381 factors. *Eur J Intern Med*. 2015;26(3):190–6.
- 382 4. Marengoni A, Angleman S, Melis R, Mangialasche F, Karp A, Garmen A, et al.
383 Aging with multimorbidity: A systematic review of the literature. *Ageing Res Rev*.
384 2011;10(4):430–9.
- 385 5. Navickas R, Petric VK, Feigl AB, Seychell M. Multimorbidity: What Do We
386 Know? What Should We Do? *J Comorbidity*. 2016;6(1):4–11.
- 387 6. Singh K, Patel SA, Biswas S, Shivashankar R, Kondal D, Ajay VS, et al.
388 Multimorbidity in South Asian adults: prevalence, risk factors and mortality. *J*
389 *Public Health*. 2019;41(1):80–9.
- 390 7. BPJS Kesehatan. BPJS Kesehatan [Internet]. 2020 [cited 2020 Feb 24]. Available
391 from: <https://bpjs-kesehatan.go.id/bpjs/>
- 392 8. Ministry of Health R of I. Indonesia Health Profile 2018. Jakarta, Indonesia; 2019.
- 393 9. BPJS Kesehatan. Program Management and Financial Report of National Health
394 Insurance and Social 2018. Jakarta, Indonesia; 2018.
- 395 10. Erik von Elm, Douglas G Altman, Matthias Egger, Stuart J Pocock and PCG& JPV,
396 Initiative for the S. Policy and practice. Strength Report Obs Stud Epidemiol
397 STROBE Statement Guidel Report Obs Stud. 2007;85(November):867–72.
- 398 11. Hemant Kulkarni SA. Choosing Statistical and Epidemiologic Methods [Internet].
399 June 2010. 2010. Available from: <https://www.openepi.com/Search/Choosing.htm>
- 400 12. Ariawan I, Sartono B, Jaya C, Mawardi J. Report of Health BPJS Sample Data.
401 Jakarta, Indonesia: BPJS Kesehatan; 2019.
- 402 13. Hussain MA, Mamun A Al, Reid C, Huxley RR. Prevalence, Awareness,
403 Treatment and Control of Hypertension in Indonesian Adults Aged ≥ 40 Years:
404 Findings from the Indonesia Family Life Survey (IFLS). *PLoS ONE*. 2016;11(8).
- 405 14. Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in
406 Indonesian adults: An exploratory study using national survey data. *BMJ Open*.
407 2015;5(12):e009810.
- 408 15. Koller D, Schön G, Schäfer I, Glaeske G, van den Bussche H, Hansen H.
409 Multimorbidity and long-term care dependency—a five-year follow-up. *BMC*
410 *Geriatr*. 2014;14(1):70.
- 411 16. Mahendradhata Y, Trisnantoro L, Listyadewi S, Soewondo P, Marthias T. The
412 Republic of Indonesia Health System Review, Health Systems in Transition. Vol.
413 7. India: WHO Regional Office for South-East Asia; 2017.

- 414 17. Carrasquillo O. Health Care Utilization. In: Gellman MD, Turner JR, editors.
415 Encyclopedia of Behavioral Medicine. New York, NY: Springer; 2013. p. 909–10.
- 416 18. Ministry of Health R of I. Regulation of Ministry of Health, Indonesia, no. 27 year
417 2014, about INA CBGs System. Ministry of Health Indonesia; 2014.
- 418 19. Satibi S, Andayani TM, Endarti D, Suwantara IPT, Wintariani NP, Agustini NPD.
419 Comparison of Real Cost Versus the Indonesian Case Base Groups (INA-CBGs)
420 Tariff Rates Among Patients of High-Incidence Cancers Under the National
421 Health Insurance Scheme. *Asian Pac J Cancer Prev APJCP*. 2019;20(1):117.
- 422 20. Ministry of Health R of I. Regulation of Ministry of Health, Republic of Indonesia,
423 no. 69 year 2013 about The Healthcare Tariff in Unversal Health Coverage
424 Program. Indonesia; 2013.
- 425 21. Nguyen H, Manolova G, Daskalopoulou C, Vitoratou S, Prince M, Prina AM.
426 Prevalence of multimorbidity in community settings: A systematic review and
427 meta-analysis of observational studies. *J Comorbidity*. 2019
428 Jan;9:2235042X1987093.
- 429 22. Arokiasamy P, Uttamacharya U, Jain K, Biritwum RB, Yawson AE, Wu F, et al.
430 The impact of multimorbidity on adult physical and mental health in low- and
431 middle-income countries: what does the study on global ageing and adult health
432 (SAGE) reveal? *BMC Med*. 2015;13(1):178.
- 433 23. Prazeres F, Santiago L. Prevalence of multimorbidity in the adult population
434 attending primary care in Portugal: a cross-sectional study. *BMJ Open* [Internet].
435 2015 Sep 25;5(9). Available from:
436 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4593139/>
- 437 24. Ritonga DR. Statistic Data of Marriage in Indonesia. 2015.
- 438 25. Vellakkal S, Subramanian S V, Millett C, Basu S, Stuckler D, Ebrahim S.
439 Socioeconomic Inequalities in Non-Communicable Diseases Prevalence in India:
440 Disparities between Self-Reported Diagnoses and Standardized Measures. *PLOS*
441 *ONE*. 2013;8(7):e68219.
- 442 26. Moeloek NF. Indonesia national health policy in the transition of disease burden
443 and health insurance coverage. *Med J Indones*. 2017;26(1):3–6.
- 444 27. Glynn LG, Valderas JM, Healy P, Burke E, Newell J, Gillespie P, et al. The
445 prevalence of multimorbidity in primary care and its effect on health care
446 utilization and cost. *Fam Pract*. 2011;28(5):516–23.
- 447 28. Pati S, Swain S, Knottnerus JA, Metsemakers JFM, van den Akker M. Health
448 related quality of life in multimorbidity: a primary-care based study from Odisha,
449 India. *Health Qual Life Outcomes*. 2019;17(1):116.

- 450 29. Buja A, Rivera M, De Battisti E, Corti MC, Avossa F, Schievano E, et al.
451 Multimorbidity and Hospital Admissions in High-Need, High-Cost Elderly
452 Patients. *J Aging Health*. 2018;089826431881709.
- 453 30. Frølich A, Ghith N, Schiøtz M, Jacobsen R, Stockmarr A. Multimorbidity,
454 healthcare utilization and socioeconomic status: A register-based study in
455 Denmark. Kwon YD, editor. *PLOS ONE*. 2019;14(8):e0214183.
- 456 31. Lee JT, Hamid F, Pati S, Atun R, Millett C. Impact of Noncommunicable Disease
457 Multimorbidity on Healthcare Utilisation and Out-Of-Pocket Expenditures in
458 Middle-Income Countries: Cross Sectional Analysis. *PLoS ONE*. 2015;10(7).
- 459 32. Bähler C, Huber CA, Brüngger B, Reich O. Multimorbidity, health care utilization
460 and costs in an elderly community-dwelling population: a claims data based
461 observational study. *BMC Health Serv Res*. 2015;15(1):23.
- 462 33. van Oostrom SH, Picavet HSJ, de Bruin SR, Stirbu I, Korevaar JC, Schellevis FG,
463 et al. Multimorbidity of chronic diseases and health care utilization in general
464 practice. *BMC Fam Pract*. 2014 Dec;15(1):61.
- 465 35. Huntley AL, Johnson R, Purdy S, Valderas JM, Salisbury C. Measures of
466 Multimorbidity and Morbidity Burden for Use in Primary Care and Community
467 Settings: A Systematic Review and Guide. *Ann Fam Med*. 2012;10(2):134–41.
- 468 36. Hu RH, Hsiao FY, Chen LJ, Huang PT, Hsu WWY. Increasing age- and gender-
469 specific burden and complexity of multimorbidity in Taiwan, 2003-2013: a cross-
470 sectional study based on nationwide claims data. *BMJ Open*. 2019;9(6):e028333.
- 471 37. Kuo RN, Lai MS. The influence of socio-economic status and multimorbidity
472 patterns on healthcare costs: a six-year follow-up under a universal healthcare
473 system. *Int J Equity Health*. 2013;12(1):69.
- 474 38. Nagl A, Witte J, Hodek JM, Greiner W. Relationship between multimorbidity and
475 direct healthcare costs in an advanced elderly population. Results of the PRISCUS
476 trial. *Z Gerontol Geriatr*. 2012;45(2):146–54.
- 477 39. Friedman B, Jiang HJ, Elixhauser A, Segal A. Hospital Inpatient Costs for Adults
478 with Multiple Chronic Conditions. *Med Care Res Rev*. 2006;63(3):327–46.
- 479 40. Valderas JM. Multimorbidity, not a health condition or complexity by another
480 name. *Eur J Gen Pract*. 2015;21(4):213–4.
- 481 41. Pefoyo AJK, Bronskill SE, Gruneir A, Calzavara A, Thavorn K, Petrosyan Y, et
482 al. The increasing burden and complexity of multimorbidity. *BMC Public Health*.
483 2015;15(415):1–11.

484

Total samples from BPJS Health National Sample Data 2015 - 2016

N = 1697452

(male= 874480 and female= 822970)



Total samples aged 60 years and above in the hospitals

N = 44868

(male= 23016 and female = 21852)



Total samples with one or more chronic conditions

N = 23460

(male= 11760 and female = 11700)

Figure