1　Wastewater surveillance of human influenza, metapneumovirus, parainfluenza, respiratory
2　syncytial virus (RSV), rhinovirus, and seasonal coronaviruses during the COVID-19 pandemic
3
4

5　Alexandria B. Boehm[1*], Bridgette Hughes[2], Dorothea Duong[2], Vikram Chan-Herur[2], Anna
6　Buchman[2], Marlene K. Wolfe[3], Bradley J. White[2]
7
8　　1. Department of Civil & Environmental Engineering, School of Engineering and Doerr
9　　　 School of Sustainability, Stanford University, 473 Via Ortega, Stanford, CA, USA 94305
10　　2. Verily Life Sciences LLC, South San Francisco, CA, USA, 94080
11　　3. Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory
12　　　 University, 1518 Clifton Rd, Atlanta, GA, USA, 30322
13
14
15
16　* Author to whom correspondence should be addressed. Email: aboehm@stanford.edu, Tel:
17　650-724-9128
18
19

20
21

22 **Abstract**
23
Background: Respiratory disease is a major cause of morbidity and mortality; however, current
surveillance for circulating respiratory viruses is passive and biased. Seasonal circulation of
respiratory viruses changed dramatically during the COVID-19 pandemic. More active methods
for understanding respiratory disease dynamics are needed to better inform public health
response and to guide clinical decision making. Wastewater-based epidemiology has been used
to understand COVID-19, influenza A, and RSV infection rates at a community level, but has not
been used to investigate other respiratory viruses.
31
Methods: We measured concentrations of influenza A and B, RSV A and B, human
parainfluenza (1-4), rhinovirus, seasonal human coronaviruses, and human metapneumovirus
RNA in wastewater solids three times per week for 17 months spanning the COVID-19
pandemic at a wastewater treatment plant in California, USA. Novel probe-based assays were
developed and validated for non-influenza viral targets. We compared viral concentrations to
positivity rates for viral infections from clinical specimens submitted to sentinel laboratories.
38
Findings: We detected RNA from all target viruses in wastewater solids. Human rhinovirus and
seasonal coronaviruses were found at highest concentrations. Concentrations of viruses
correlated significantly and positively with positivity rates of associated viral diseases from
sentinel laboratories. Measurements from wastewater indicated limited circulation of RSV A and
influenza B, and human coronavirus OC43 dominated the seasonal human coronavirus
infections while human parainfluenza 1 and 4A dominated among parainfluenza infections.
45
46 Interpretation: Wastewater-based epidemiology can be used to obtain information on circulation
47 of respiratory viruses at a community level without the need to test many individuals because a
48 single sample of wastewater represents the entire contributing community. Results from
49 wastewater can be available within 24 hours of sample collection, allowing real time information
50 to inform public health response, clinical decision making, and individual behavior modifications.

Introduction

Acute respiratory illnesses (ARI) are defined as symptomatic infections of the lower or upper respiratory tract, and may also result in systemic symptoms and secondary infections. They are consistently the top cause of illness and mortality in children under 5 globally after the neonatal period, and represent a large burden of infectious disease[1]. ARI etiology can be bacterial, fungal, or viral; lower respiratory infections most commonly lead to severe conditions like pneumonia and bronchiolitis, while upper respiratory infections are largely self limiting but can result in secondary infections. Viral ARI are most commonly caused by SARS-CoV-2, seasonal human coronavirus, human parainfluenza, human metapneumovirus, influenza, respiratory syncytial virus, rhinovirus, adenoviruses, human bocaviruses, and non-rhinovirus enteroviruses[2]. In the United States, viral ARI causes nearly 400,000 child hospitalizations at a cost of $1 billion annually[3].

Respiratory disease surveillance in the United States is passive and draws from many sources, often relying on institutions to identify specific diseases through key words in clinical records, testing of clinical specimens, and searching for keywords on death certificates. The passive respiratory disease surveillance system is voluntary and biased towards identifying infections in individuals with co-morbidities and/or where symptoms are severe and patients require clinical care[4]. Since many ARIs are self-limiting, diagnostic testing of most individuals with respiratory illness is uncommon. Lack of robust, unbiased data on disease occurrence and resultant lack of knowledge on occurrence and trends in circulating respiratory diseases limits public health response, institutional awareness that can guide clinical decision making regarding testing and treatment, and efforts to understand disease epidemiology including disease interference and co-infection[5].

Prior to the COVID-19 pandemic, viral respiratory infections largely circulated seasonally with predictable dynamics[2]. However, seasonal occurrence of respiratory viruses has changed dramatically during the COVID-19 pandemic[6], presumably due to both imposed and voluntary behavior changes in response to the pandemic, such as masking and stay-at-home orders[7]. It is uncertain if and when respiratory disease dynamics will return to those observed pre-pandemic. More active approaches to disease surveillance are needed to better understand and respond to atypical respiratory disease dynamics.

Wastewater-based epidemiology (WBE) uses the information from wastewater to gain insight into infectious disease occurrence in contributing communities. Wastewater represents a composite biological sample containing bodily excretions including urine, feces, sputum, and mucus. Excretions of infected individuals can contain markers of infectious disease (including live and dead organisms, proteins and nucleic acids) and methods have been developed to detect and quantify these in the wastewater matrix. Benefits of WBE are that it does not require that infected individuals receive medical care or testing or even experience symptoms to be represented in the community-level data source.

The utility of WBE has been demonstrated for understanding community circulation of poliovirus[8], *Salmonella*[9], hepatitis A[10], and enteroviruses[11], but its use has not widely been a part of disease surveillance outside of polio in endemic regions. The application of WBE to respiratory diseases is relatively new and was prompted by the COVID-19 pandemic, during which it was widely shown that concentrations of SARS-CoV-2 RNA in wastewater correlate to reported COVID-19 incidence rates during periods of widely accessible clinical testing[12]. Thereafter, Hughes et al.[13] and Wolfe et al.[14] showed RSV and influenza A RNA

107 concentrations, respectively, in wastewater solids were associated with disease occurrence in
108 associated sewersheds.
109
Here we test the utility of WBE for a suite of viral respiratory diseases. We developed and
validated novel hydrolysis probe-based RT-PCR assays that target respiratory viral genomes
and then applied the assays to wastewater solids collected three times per week at a
wastewater treatment plant over 17 months during the COVID-19 pandemic. We measured
concentrations of respiratory syncytial virus (RSV) A and B, influenza A and B,
metapneumovirus, parainfluenza (1-4), seasonal human coronaviruses (229E, OC43, NL63,
and HKU-1), and rhinovirus RNA. We chose to focus on these respiratory viruses as passive
clinical surveillance occurs for these diseases in the region where the sampling was conducted
allowing us to compare wastewater to clinical surveillance data. Additionally, these viruses
represent important and common viral etiologies of respiratory infections[15].
120
121 **Methods**
122
RT-PCR assays. We used published hydrolysis probe-based RT-PCR assays for Influenza A
(IAV) and influenza B (IBV)[16]. We designed novel RT-PCR primers and internal hydrolysis
probes for rhinovirus (hereafter, "HRV" targeting rhinovirus A, B and C together); human
parainfluenza (HPIV) 1, 2, 3, 4A, 4B; human coronavirus (HCoV) 229E, NL63, OC43, HKU-1;
human metapneumovirus (HMPV); and Respiratory Syncytial Virus (RSV) A and B. To develop
the assays, viral genome sequences were downloaded from NCBI between between February
and June 2022 and aligned to identify conserved regions. Assays were developed *in silico* using
Primer3Plus (https://primer3plus.com/). Parameters used in assay development (e.g. sequence
length, GC content, and melt temperatures) are provided in Table S1. Primers and probes were
then screened for specificity *in silico*, and *in vitro* against virus panels, intact respiratory viruses,
synthetic viral genomic RNA, or cDNA sequences (see Supplementary Material (SM) including
Table S2).
135
136 **Wastewater analyses.** A wastewater treatment plant that serves 75% (1,500,000 people) of
137 Santa Clara County, California (San José-Santa Clara Regional Wastewater Facility, RWF) was
138 included in the study. Further descriptions of RWF can be found in elsewhere[12] .
139
Samples were collected daily for a COVID-19 WBE effort starting in November 2020[12], and a
subset of those samples (three samples per week, 216 samples total) are used in this study.
The samples were chosen to span a 17 month period (2/2/21 - 6/21/22, month/day/year format)
that included implementation and easing of indoor mask mandates, changes in COVID-19
vaccine availability, public health promotion of hand hygiene and mask wearing, and periods of
both high and low COVID-19 incidence.
146
Fifty mL of settled solids were collected using sterile technique in clean bottles from the primary
clarifier. Twenty-four hour composite samples were collected by combining grab samples from
the sludge line every six hours[12]. Samples were stored at 4°C, transported to the lab, and
processed within six hours. Solids were then dewatered[12] and frozen at -80°C for 4 - 60 weeks.
Samples were thawed overnight and then RNA was obtained from the dewatered solids
following previously published protocols[12] (see SM). RNA was obtained from 10 replicate
sample aliquots. Each replicate RNA extract from each sample was subsequently processed
immediately to measure viral RNA concentrations using digital RT-PCR with multiplexed assays
for IAV, IBV, HMPV, total HPIV (HPIV 1, HPIV 2, HPIV 3, HPIV 4A, and HPIV 4B), total HCoV
(HCoV 229E, HCoV NL63, HCoV OC43, HCoV HKU-1), RSV A, RSV B, HRV, and pepper mild
mottle virus (PMMoV) (see SI). PMMoV is highly abundant in human stool and wastewater

globally[17] and is used as an internal recovery and fecal strength control[18] . Nine samples
spanning the duration of the time series were selected to measure each HCoV and HPIV
individually (Table S3). Each 96-well PCR plate of wastewater samples included PCR positive
controls for each target assayed on the plate in 1 well, PCR negative no template controls in two
wells, and extraction negative controls (consisting of water and lysis buffer) in two wells. PCR
positive controls consisted of viral gRNA or gene blocks (Table S2). Results from replicate
wells were merged for analysis. In order for a sample to be recorded as positive, it had to have
at least 3 positive droplets.
166
Concentrations of RNA targets were converted to concentrations per dry weight of solids in units
of copies (cp)/g dry weight using dimensional analysis. The total error is reported as standard
deviations and includes the errors associated with the Poisson distribution and the variability
among the 10 replicates. Three positive droplets across 10 merged wells corresponds to a
concentration between ~500-1000 cp/g; the range in values is a result of the range in the
equivalent mass of dry solids added to the wells.
173
174 **Wastewater SARS-CoV-2 N gene.** We obtained SARS-CoV-2 N gene concentrations at the
175 POTW from a regional monitoring program[12]. These data were used in a supplementary manner
176 to provide insight into the progress of the COVID-19 pandemic (see SM).
177
Clinical data. California Sentinel Clinical Laboratories (hereafter, sentinel laboratories) test
clinical specimens for influenza A (IAV) and influenza B (IBV), respiratory syncytial virus (RSV),
parainfluenza viruses (HPIV) (1-4), human coronaviruses (HCoV) (229E, NL63, OC43, and
HKU-1), human metapneumovirus (HMPV), and rhinovirus (HRV). Sentinel laboratories do not
differentiate rhinovirus from other enteroviruses. Specimens tested by the laboratories may
represent inpatient or outpatient samples of people receiving medical care. Positivity rates were
calculated using data from all state sentinel laboratories and are reported by MMWR (Morbidity
and Mortality Weekly Report) week and assigned to the first day of MMWR week. Positivity
rates were also aggregated at the county-level for comparison (see SM).
187
Statistical analysis. Kendall's tau was used to test the null hypotheses (1) clinical specimen
positivity rates are not associated with viral concentrations in wastewater solids, and (2) viral
RNA concentrations were not associated with each other. Because case data are aggregated
by MMWR week, we used median wastewater measurements from the same MMWR week to
match clinical and wastewater data. To test correlations between wastewater viral RNA
concentrations, wastewater data were smoothed by calculating a rolling three-point median
(hereafter "smoothed wastewater concentrations"). A total of 50 Kendall's tests of association
were carried out so a p value of 0.001 was used to identify tau values significantly different from
0 based on Bonferroni's correction. To compare viral RNA concentrations visually on a similar
scale, we standardized their concentrations by subtracting the minimum and dividing by the
range.
199
200 **Results**
201
QA/QC. Results are reported as suggested in the EMMI guidelines[19]. All extraction and PCR
negative and positive controls performed as expected (negative and positive, respectively) with
the exception of the positive control failure for an HCoV run including samples before 2/26/22;
HCoV results prior to 2/26/22 were therefore omitted from the analysis. Median PMMoV
concentrations across the samples were 1.6×10^9 copies/g, similar to measurements previously
reported for the plant[12]. In addition, PMMoV levels were stable across samples (interquartile

range = 0.7x10^9 cp/g) suggesting consistent fecal strength and RNA extraction efficiency. See
SM for additional QA/QC details.
210
211 **Novel assay sensitivity and specificity.** *In silico* analysis indicated no cross reactivity of the
212 novel RT-PCR probe-based assays (Table 1) with sequences deposited in NCBI. The novel
213 assays for HPIV, HMPV, HRV, and the HCoV were tested *in vitro* against panels of non-target
214 viral gRNA as well as target gRNA (Table S2). No cross reactivity was observed.
215
Respiratory virus RNA in wastewater solids. RNA from all viruses were detected in
wastewater samples (Figure 1). Concentrations varied from non-detect to over 10^6 copies/g. In
order of highest to lowest, median concentrations were 35000 cp/g for HCoV, 8500 cp/g for
HRV, 2500 cp/g for HPIV, 1700 cp/g for RSV B, 760 cp/g for HMPV, and non-detect for IAV,
IAB, and RSV A (Table 2). While median IAV, IBV, and RSV A were non-detect, viral RNA from
these viruses was detected in 28%, 6% and 14% of the samples, respectively. For context,
median concentrations for SARS-CoV-2 was 48000 cp/g. In general, concentrations were
lowest at the beginning of the study (February 2021), and increased until January 2022 at which
time viral RNA for all viruses showed a steep drop in concentration. Coincidentally, January
2022 marked the onset of the Omicron BA.1 wave in the region which caused the highest
COVID-19 reported incidence rates to date. After the steep drop off, viral RNA concentrations
began to rise until the end of the time series with the exception of RSV B for which we did not
observe a rebound. HCoV rebounded first when all other viral RNA concentrations, including
those of SARS-CoV-2, were still at relatively low concentrations (Figure 2).
230
In order to assess which HCoV were circulating during the study, we measured concentrations
of OC43, HKU-1, 229E, and NL63 RNA in nine samples spanning the time series (Figure 3).
Nearly 100% of the total HCoV RNA was OC43 RNA in seven of the nine samples, all collected
prior to April 2022. For the two samples collected thereafter, the contribution of OC43 RNA
declined to 20-30% and contributions of 229E and HKU-1 increased from <10% to greater than
50% and 20-30%, respectively.
237
238 Similarly, we measured RNA concentrations of each HPIV (1, 2, 3, 4A, and 4B) in samples
239 spanning the time series. RNA from all four HPIV were present; HPIV 3 dominated in most
240 samples (40-90% of total HPIVs), although each HPIV 2 and 4B also dominated in one of the
241 nine tested samples. HPIV 1 and 4A contributed the least to total HPIV present (median < 10%,
242 Figure 3).
243
Associations between viral RNA and confirmed case positivity rates. We tested whether
state-aggregated positivity rates for respiratory viral infections was associated with the
wastewater concentrations of viral RNA. Sentinel laboratories tested a median of 5918 (range
1809-14813) clinical specimens per week for influenza (IAV and IAB), 2419 (619-10907) for
RSV, 1084 (574 - 1984) for HPIV, HMPV, and HRV, and 931 (469-1833) for HCoV. The
associations between wastewater RNA concentrations and positivity rates were significant and
positive for all viruses except for IBV. Kendall's tau between positivity rates and wastewater
concentration were 0.36 (p<10^{-5}) for HRV, 0.45 (p<10^{-5}) for IAV, 0.52 (p<10^{-9}) for HPIV, 0.57
(p<10^{-10}) for RSV (using RSV B wastewater concentrations as RSV A was mostly non-detect),
0.32 (p=0.00018) for HMPV, 0.32 (p=0.00011) for HCoV, and -0.010 (p=0.92) for IBV.
Wastewater concentrations and positivity rates aggregated across viruses were positively
correlated (tau = 0.50, p <10^{-15}) (Figure S1). Results for county-aggregated positivity rates are
reported in the SM (Figure S2).
257

258 State-aggregated positivity rates for individual HCoV infections are similar to observations of the
259 relative occurrence of their RNA in wastewater solids (Figure 3). OC43 had the highest positivity
260 relative to other HCoV, in agreement with the wastewater measurements. Wastewater indicates
261 that 229E dominated at the end of the study (spring 2022), but this is not reflected in the clinical
262 data.
263
264 Positivity rates for individual HPIV suggests HPIV 3 dominated HPIV infections throughout the
265 study period; HPIV 2 and 4 also circulated with lower positivity rates. HPIV 1 was rarely
266 detected in clinical specimens. This is in agreement with the wastewater data that indicated
267 HPIV 3 dominated with HPIV 2 and 4B also present at relatively high levels, and HPIV 1 rarely
268 detected (Figure 3).
269
Associations between wastewater viral RNA concentrations. Viral RNA concentrations
were associated with each other, as well SARS-CoV-2 N gene concentrations available from a
regional monitoring program at the POTW (Table S4). Three-day smoothed wastewater
concentrations of HPIV, HRV, HMPV, IAV, RSV B, and SARS-CoV-2 RNA were significantly
correlated with each other (p<0.001) with tau from 0.18 (IAV and HRV) to 0.61 (HRV and
HMPV). Given the low occurrence of IBV and RSV A RNA, it is not surprising that they were not
correlated with other viral RNA concentrations. HCoV RNA concentrations were not correlated
to concentrations of other viral RNA targets.
278
279 **Ethics.** This study was reviewed by State of California Health and Human Services Agency
280 Committee for the Protection of Human Subjects and determined to be Exempt.
281
282 **Role of funding source.** CDC Foundation played no role in study design, execution, or result
283 interpretation.
284
285 **Discussion**
286
We detected RNA from all tested respiratory viruses in wastewater solids, including human
rhinovirus, parainfluenza, metapneumovirus, influenza, respiratory syncytial virus, and seasonal
human coronaviruses. Additionally, we detected four types of seasonal human coronaviruses
including OC43, 229E, NL63, and HKU-1, and five different parainfluenza viruses (1, 2, 3, 4A,
and 4B). Non-influenza respiratory viral RNA targets were measured using novel assays; these
assays were found to be specific to the intended viral genomic targets when tested against a
range of viruses. Previous studies have documented human rhinovirus[11,20], RSV[13], and
influenza[14] RNA in wastewater, but to our knowledge, no study to date has reported
concentrations of other respiratory viruses.
296
RSV A RNA was rarely detected in wastewater, yet RSV B concentrations were >10,000 cp/g at
times, and RSV B wastewater trends mirrored RSV clinical surveillance data. This suggests
limited local co-circulation of the two RSV groups. Changes in dominant RSV strains have been
documented to occur at continental[21] and community scales[22] with resultant changes in
population immunity[23]. Sequencing of SARS-CoV-2 in wastewater has been successful at
identifying variants[24], and sequencing of RSV genomes from wastewater may yield additional
insight into strain circulation and their potential susceptibility to different pharmaceutical
interventions.
305
The association between respiratory viral RNA in wastewater solids and their respective virus
clinical specimen positivity rates as available through a passive disease surveillance system
suggests that wastewater data are representative of respiratory disease occurrence in the

contributing population. This is despite limitations with passive disease surveillance, and the fact
that data were aggregated across the entire state which obfuscates disease occurrence
patterns at more local spatial scales. For example, state-aggregated HCoV case positivity
suggests OC43 infections dominated HCoV infections, yet wastewater solids measurements
suggest 229E dominated locally in spring of 2022. Similarly, based on wastewater
measurements, HPIV 1 and 4 may have contributed to more HPIV infections locally than on
average across the state.
316
There are uncertainties associated with understanding disease occurrence in a community
through both clinical surveillance and wastewater testing. Clinical surveillance data is biased
towards those receiving medical care and not all specimens are tested for all viruses, making it
difficult to estimate disease occurrence from positivity rate data. Positivity rates may not be
associated with disease occurrence in the same manner for all considered viruses. For
wastewater, there is limited information on concentrations of viral RNA excreted from infected
individuals into wastewater despite evidence showing that shedding does occur for viruses
included in this study[25,26]. Given the current state of knowledge, it is difficult to know how inputs
of viral RNA to the wastewater system differ between individuals infected with different viruses
and with different disease severities.
327
Despite this, the association between viral RNA in wastewater solids and case positivity holds
when data from the measured viruses are aggregated, suggesting that even across
viruses,concentrations in wastewater solids are related to their different rates of disease
occurrence. HCoV and HRV concentrations were highest across our study period, suggesting
seasonal human coronavirus and human rhinovirus infections were most common. Conversely,
IBV and RSV A concentrations were lowest suggesting limited circulation of those viruses.
Future work is needed to document time varying patterns of respiratory viral RNA inputs from
infected individuals to the wastewater system to enable modeling of the number of infected
individuals in a sewershed from wastewater viral RNA concentrations[27] and to enable modeling
of epidemiological parameters like the reproductive number from wastewater data[28].
338
Viral RNA concentrations in wastewater solids, including those of SARS-CoV-2, generally
followed similar trends over time, with the exception of HCoV. Evident in the trends is a notable
drop-off in concentrations of all viruses after the Omicron BA.1 surge in January 2022. The
drop-off may suggest that disease mitigation measures practiced by the community in response
to the Omicron surge (isolation due to illness, working remotely, wearing masks) reduced the
spread of all respiratory viruses. Local indoor mask mandates ended shortly after the Omicron
surge, and this change was followed by an increase in concentrations of all respiratory viruses
in wastewater (Figure 3). HCoV, specifically OC43, was the first virus to re-appear in
wastewater after the Omicron surge. Non-SARS-CoV-2 HCoV are highly transmissible with R0
values higher, on average, than influenza and rhinovirus[29], but lower than RSV and similar to
HPIV; there is limited R0 data for HMPV[15]. It is also possible that infection with Omicron
reduced susceptibility to other respiratory viruses and contributed to the decrease in all viral
activity; antibody cross immunity has been suggested to control temporal patterns of some
respiratory viruses[5,30] .
353
Data generated from wastewater can be available within 24 hours of sample collection, and
samples represent the entire contributing population - even those with asymptomatic infections
- thereby overcoming various biases and the inherent reporting delays of the passive
surveillance system. Wastewater surveillance for multiple respiratory viruses that commonly
circulate can provide information at a community scale on the most likely causes of ARI.
Although testing of clinical specimens is often not done for ARI, the viruses included in this

panel have differing levels of likely severity and associated prevention measures, treatments,
and complications. Therefore, more information about the abundances of circulating viruses can
inform local clinical decision making including the prescription of drugs and testing, availability of
community testing resources, and vaccination or behavior change campaigns. This information
can also be communicated to the public, like a weather report, so the community and
particularly vulnerable individuals can make informed decisions about behaviors that may put
them at risk.
367

Acknowledgements. This work is supported by a gift from the CDC Foundation. We
acknowledge Payal Sarkar at RWF for overseeing sample collection. We acknowledge Erin
Murray, Monica Sun, Tomás Leon, and Alexander Yu at California Department of Public Health
for their assistance. This study was performed on the ancestral and unceded lands of the
Muwekma Ohlone people. We pay our respects to them and their Elders, past and present, and
are grateful for the opportunity to live and work here.

374
375　**Declaration of interests.** BH, DD, VCH, ABu, and BW are employees of Verily Life Sciences,
376　LLC.
377
378　**Data sharing statement.** Wastewater data are available publicly at the Stanford Digital
379　Repository (https://doi.org/10.25740/vx726fw9373). State-aggregated surveillance data is
380　available upon request from California Department of Health.

389
Figure 1. State-aggregated positivity rate reported as a percent from sentinel laboratories (top panel) and
viral RNA in wastewater solids for each virus reported in copies per gram dry weight (bottom panel),
exception is the top right panel which shows SARS-CoV-2 wastewater results for context. RSV is
respiratory syncytial virus (A is shown in green and B in grey/black, RSV A scale is 1/10th that of RSV B
and, like the RSV B axis, is scaled by 10^3), HCoV is the sum of all four seasonal human coronaviruses
(OC43, HKU-1, 229E, and NL63), HPIV is human parainfluenza viruses (1-4), HRV is human rhinovirus A,
B and C, HMPV is human metapneumovirus, and IAV and IBV are influenza A and B viruses,
respectively. Gray symbols represent measurements, error bars are 68% confidence intervals
representing the total error around 10 replicates. The black line is the 3-point rolling median of the
measurements. For HCoV there are three measurements located off scale (1.1×10^6, 3.6×10^5, and 5.1×10^5
cp/g on 4/9/21, 6/24/21, and 3/3/22, respectively). For HPIV there is one measurement located off scale
(3.6×10^4 cp/g on 4/9/21). For HRV there is one measurement located off scale (3.6×10^5 cp/g on 4/14/22).
For IAV there are 3 measurements located off scale (2.1×10^4, 7.5×10^4, 5.1×10^5 and 1.5×10^4 cp/g on
9/30/21, 10/31/21, 3/3/22, and 4/14/22, respectively). Dates are in month/day/year format.

404
405

406
407 Figure 2. Standardized concentrations of viral RNA for all viruses except for RSV A and IBV which were
408 rarely detected. Shaded background indicates periods of time that indoor masking mandates were in
409 effect locally. Dates of COVID-19 vaccine availability are provided at top. Dates are in month/day/year
410 format. Standardized concentrations were calculated by subtracting the minimum and dividing by the
411 range. SC2 is SARS-CoV-2.
412
413
414
415

416
417
418

419
Figure 3. Data from wastewater solids on the (A) total concentration of HCoV RNA (shown as gray filled
areas) and individual HCoVs (reported as fractions of total) and (B) total concentration of HPIV RNA
(shown as gray filled areas) and individual HPIVs (reported as fractions of total). State-aggregated
percent positivity for tests for (C) individual HCoVs and (D) individual HPIVs. Note that percent positivity
for "unknown HCoV" and "unknown HPIV" from the sentinel surveillance system are not shown in panels
C and D. Dates are in month/day/year format.
426

Target	Primer/Probe	Sequence
HRV	Forward	GCCYGCGTGGCKGCC
	Reverse	GAAACACGGACACCCAAAG
	Probe	TCCTCCGGCCCCTGAATG
HPIV 1	Forward	AAGTTCAGTACAAAGCGGGA
	Reverse	GCTCARTAGGGGTTCTCCTA
	Probe	AGCAAAGCAGAGATCTCACACA
HPIV 2	Forward	AATACAACAGGGCARTGGG
	Reverse	GATAAAATAGCGTGAGGACTGC
	Probe	TCCTGTATATGGTGGTCTCATAAATGG
HPIV 3	Forward	TGTGGTGACCAACAGATCAA
	Reverse	CCCTCCAAAGAATCGTCCTG
	Probe	TCCAATGAAAACACTGATCCCAGA
HPIV 4A	Forward	TGAACGGTTGCATTCAGG
	Reverse	TTGACTGGTTGCACCTAATTCT
	Probe	CTGGCAATCTCAACATAGACCATG
HPIV 4B	Forward	GGAGAACTTTGAAACCACCTCTAA
	Reverse	TCTCCTTTAACTACCCTATCTTTGC
	Probe	ACCCCCATAAGGCAAGAAGC
HCoV 229E	Forward	GGATGACATCATGAAGGCAG
	Reverse	TACCCGTTTTCGCTGACTTT

	Probe	TTCCTGAGGCTTGTCAAAACCT
HCoV NL63	Forward	GAAGCGTGTTCCTACCAGAG
	Reverse	TGGCATCAACACCATTCTGA
	Probe	CAGTGCTTTGGTCCTCGTGA
HCoV OC43	Forward	GTCTTTTACTCCTGGTAAGCAATC
	Reverse	GGGTACAACATTCCCTCCTG
	Probe	CCGATCAGTCCGACCAGTTTAG
HCoV HKU-1	Forward	CCTGGTACGATTTTGCCTCA
	Reverse	ATTGGGTCCACGTGATTGAG
	Probe	AGGCTCAGGAAGGTCTGCTT
HMPV	Forward	ACTTTATTGGAGAAGGAGCAGG
	Reverse	GGGTAATGRTGATCAAGRTCA
	Probe	AYTGGATGGCMAGAACAGCA
IAV	Forward	CAAGACCAATCYTGTCACCTCTGA C
	Reverse	GCATTYTGGACAAAVCGTCTACG
	Probe	TGCAGTCCTCGCTCACTGGGCACG
IBV	Forward	TCCTCAAYTCACTCTTCGAGCG
	Reverse	CGGTGCTCTTGACCAAATTGG
	Probe	CCAATTCGAGCAGCTGAAACTGCGGTG
RSV A	Forward	AGAGGTGGCAGTAGAGTTGA

	Reverse	CTCCACAACTTGTTCCATTTCTG	
	Probe	ATGGTGCAGGGCAAGTGATG	
RSV B	Forward	TGACACTCCCAATTATGATGTGC	
	Reverse	CCTGTGAATTTATGATTTGCATCTTCAG	
	Probe	ACACCTAAACAAACTATGTGGTATGC	

Table 1. Primer and probe sequences for detection of respiratory viral RNA used in this study. Additional details of the target region of the genomes is the Supplementary Material. Primers and probes were purchased from IDT. All probes contained a fluorescent molecule and quenchers (5′ FAM and/or HEX/ZEN/3′ IBFQ). FAM, 6-fluorescein amidite; HEX, hexachloro-fluorescein; ZEN, a proprietary internal quencher from IDT; IBFQ, Iowa Black FQ. The Supplementary Material indicates whether HEX or FAM molecules were used.

Statistic	HPIV	HCoV	HMPV	HRV	RSV A	RSV B	IAV	IBV	SARS-CoV-2
median	3.5×10^3	3.5×10^4	7.6×10^2	8.5×10^3	0	1.7×10^3	0	0	4.8×10^4
25th %ile	1.4×10^3	1.7×10^4	0	4.6×10^3	0	0	0	0	2.5×10^4
75th %ile	6.3×10^3	5.6×10^4	2.0×10^3	1.7×10^4	0	5.4×10^3	6.6×10^2	0	1.3×10^5
# of ND	21	0	95	18	186	89	156	204	0
N	216	207	216	216	216	216	216	216	216

Table 2. Summary statistics of measurements of viral RNA in wastewater solids. Units of concentration is copies per gram dry weight of wastewater solids. RSV is respiratory syncytial virus, HCoV is the sum of all four non-SARS human coronaviruses (OC43, HKU-1, 229E, and NL63), HPIV is human parainfluenza viruses (1-4), HRV is human rhinovirus A, B and C, HMPV is human metapneumovirus, and IAV and IBV are influenza A and B viruses, respectively. SARS-CoV-2 data are included for context. # ND is number of samples for which the viral RNA was not detected. N is number of measurements made.

References

Wang H, Bhutta ZA, Coates MM, *et al.* Global, regional, national, and selected subnational levels of stillbirths, neonatal, infant, and under-5 mortality, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet* 2016; **388**: 1725–74.

Moriyama M, Hugentobler WJ, Iwasaki A. Seasonality of Respiratory Viral Infections. *Annu Rev Virol* 2020; **7**: 83–101.

Henrickson KJ. Advances in the laboratory diagnosis of viral respiratory disease. *The Pediatric Infectious Disease Journal* 2004; **23**. https://journals.lww.com/pidj/Fulltext/2004/01001/Advances_in_the_laboratory_diagnosis_of_viral.2.aspx.

Killerby ME, Biggs HM, Haynes A, *et al.* Human coronavirus circulation in the United States 2014–2017. *Journal of Clinical Virology* 2018; **101**: 52–6.

Nickbakhsh S, Mair C, Matthews L, *et al.* Virus–virus interactions impact the population dynamics of influenza and the common cold. *Proceedings of the National Academy of Sciences* 2019; **116**: 27142–50.

Kuitunen I, Artama M, Haapanen M, Renko M. Respiratory virus circulation in children after relaxation of COVID-19 restrictions in fall 2021—A nationwide register study in Finland. *Journal of Medical Virology* 2022; **94**: 4528–32.

Liu P, Xu M, Lu L, *et al.* The changing pattern of common respiratory and enteric viruses among outpatient children in Shanghai, China: Two years of the COVID-19 pandemic. *Journal of Medical Virology* 2022; **94**: 4696–703.

Brouwer AF, Eisenberg JNS, Pomeroy CD, *et al.* Epidemiology of the silent polio outbreak in Rahat, Israel, based on modeling of environmental surveillance data. *Proc Natl Acad Sci USA* 2018; **115**: E10625.

Diemert S, Yan T. Clinically Unreported Salmonellosis Outbreak Detected via Comparative Genomic Analysis of Municipal Wastewater Salmonella Isolates. *Appl Environ Microbiol* 2019; **85**: e00139-19.

McCall C, Wu H, O'Brien E, Xagoraraki I. Assessment of enteric viruses during a hepatitis outbreak in Detroit MI using wastewater surveillance and metagenomic analysis. *Journal of Applied Microbiology* 2021; **131**: 1539–54.

Brinkman NE, Shay FG, Keely SP. Retrospective Surveillance of Wastewater To Examine Seasonal Dynamics of Enterovirus Infections. *mSphere* 2017; **2**: e00099-17.

Wolfe MK, Topol A, Knudson A, *et al.* High-Frequency, High-Throughput Quantification of SARS-CoV-2 RNA in Wastewater Settled Solids at Eight Publicly Owned Treatment Works in Northern California Shows Strong Association with COVID-19 Incidence. *mSystems* 2021; **0**: e00829-21.

Hughes B, Duong D, White BJ, *et al.* Respiratory Syncytial Virus (RSV) RNA in Wastewater Settled Solids Reflects RSV Clinical Positivity Rates. *Environ Sci Technol Lett* 2022; **9**: 173–8.

Wolfe MK, Duong D, Bakker KM, *et al.* Wastewater-Based Detection of Two Influenza Outbreaks. *Environ Sci Technol Lett* 2022; **9**: 687–92.

Leung NHL. Transmissibility and transmission of respiratory viruses. *Nature Reviews Microbiology* 2021; **19**: 528–45.

United States Center for Disease Control. Research Use Only CDC Flu SC2 Multiplex Assay Primers and Probes. 2022. https://www.cdc.gov/coronavirus/2019-ncov/lab/multiplex-primer-probes.html (accessed Aug 18, 2022).

Symonds EM, Nguyen KH, Harwood VJ, Breitbart M. Pepper mild mottle virus: A plant

pathogen with a greater purpose in (waste)water treatment development and public health management. *Water Research* 2018; **144**: 1–12.

McClary-Gutierrez JS, Aanderud ZT, Al-faliti M, *et al.* Standardizing data reporting in the research community to enhance the utility of open data for SARS-CoV-2 wastewater surveillance. *Environ Sci: Water Res Technol* 2021; **7**: 1545–51.

Borchardt MA, Boehm AB, Salit M, Spencer SK, Wigginton KR, Noble RT. The Environmental Microbiology Minimum Information (EMMI) Guidelines: qPCR and dPCR Quality and Reporting for Environmental Microbiology. *Environ Sci Technol* 2021; **55**: 10210–23.

Blomqvist S, Savolainen-Kopra C, Paananen A, Hovi T, Roivainen M. Molecular characterization of human rhinovirus field strains isolated during surveillance of enteroviruses. Journal of General Virology,. 2009; **90**: 1371–81.

Cantú-Flores K, Rivera-Alfaro G, Muñoz-Escalante JC, Noyola DE. Global distribution of respiratory syncytial virus A and B infections: a systematic review. *null* 2022; : 1–12.

Peret TCT, Hall CB, Schnabel KC, Golub JA, Anderson LJ. Circulation patterns of genetically distinct group A and B strains of human respiratory syncytial virus in a community. Journal of General Virology,. 1998; **79**: 2221–9.

Melero JA, Moore ML. Influence of Respiratory Syncytial Virus Strain Differences on Pathogenesis and Immunity. In: Anderson LJ, Graham BS, eds. Challenges and Opportunities for Respiratory Syncytial Virus Vaccines. Berlin, Heidelberg: Springer Berlin Heidelberg, 2013: 59–82.

Crits-Christoph A, Kantor RS, Olm MR, *et al.* Genome Sequencing of Sewage Detects Regionally Prevalent SARS-CoV-2 Variants. *mBio*; **12**: e02703-20.

Branche AR, Walsh EE, Formica MA, Falsey AR. Detection of Respiratory Viruses in Sputum from Adults by Use of Automated Multiplex PCR. *Journal of Clinical Microbiology* 2014; **52**: 3590–6.

Minodier L, Charrel RN, Ceccaldi P-E, *et al.* Prevalence of gastrointestinal symptoms in patients with influenza, clinical significance, and pathophysiology of human influenza viruses in faecal samples: what do we know? *Virology Journal* 2015; **12**: 215.

Soller J, Jennings W, Schoen M, *et al.* Modeling infection from SARS-CoV-2 wastewater concentrations: promise, limitations, and future directions. *Journal of Water and Health* 2022; **20**: 1197–211.

Huisman JS, Scire J, Caduff L, *et al.* Wastewater-based estimation of the effective reproductive number of SARS-CoV-2. *Environmental Health Perspectives* 2022; **In press**.

Spencer JA, Shutt DP, Moser SK, *et al.* Distinguishing viruses responsible for influenza-like illness. *Journal of Theoretical Biology* 2022; **545**: 111145.

Bhattacharyya S, Gesteland PH, Korgenski K, Bjørnstad ON, Adler FR. Cross-immunity between strains explains the dynamical pattern of paramyxoviruses. *Proceedings of the National Academy of Sciences* 2015; **112**: 13396–400.