Infliximab for Treatment of Adults Hospitalized with Moderate or Severe Covid-19

Jane A. O'Halloran, Eyal Kedar, Kevin J. Anstrom, Matthew W. McCarthy, Emily R. Ko, Patricia

Segura Nunez, Cynthia Boucher, P. Brian Smith, Reynold A. Panettieri, Jr, Sabina Mendivil

Tuchia de Tai, Martin Maillo, Akram Khan, Alfredo J. Mena Lora, Matthias Salathe, Gerardo

Capo, Daniel Rodríguez Gonzalez, Thomas F. Patterson, Christopher Palma, Horacio Ariza,

Maria Patelli Lima, Anne M. Lachiewicz, John Blamoun, Esteban C. Nannini, Eduardo Sprinz,

Analia Mykietiuk, Radica Alicic, Adriana M. Rauseo, Cameron R. Wolfe, Britta Witting, Daniel K.

Benjamin, Jr, Steven E. McNulty, Pearl Zakroysky, Susan Halabi, Sandra Butler, Jane Atkinson,

Stacey J. Adam, Richard Melsheimer, Soju Chang, Lisa LaVange, Michael Proschan, Samuel

A. Bozzette, William G. Powderly for the ACTIV-1 IM study group members*

*Author degrees and affiliations are listed at the end of the article.

Running title: Infliximab for hospitalized Covid-19

Word count: 2699

Address for correspondence: William G. Powderly, MD, Division of Infectious Diseases,

Department of Medicine, Washington University School of Medicine, St. Louis, MO 63110.

Email: Wpowderly@wustl.edu.

ABSTRACT

Background: Immune dysregulation contributes to poorer outcomes in severe Covid-19. Immunomodulators targeting various pathways have improved outcomes. We investigated whether infliximab provides benefit over standard of care.

Methods: We conducted a master protocol investigating immunomodulators for potential benefit in treatment of participants hospitalized with Covid-19 pneumonia. We report results for infliximab (single dose infusion) versus shared placebo both with standard of care. Primary outcome was time to recovery by day 29 (28 days after randomization). Key secondary endpoints included 14-day clinical status and 28-day mortality.

Results: A total of 1033 participants received study drug (517 infliximab, 516 placebo). Mean age was 54.8 years, 60.3% were male, 48.6% Hispanic or Latino, and 14% Black. No statistically significant difference in the primary endpoint was seen with infliximab compared with placebo (recovery rate ratio 1.13, 95% CI 0.99–1.29; p=0.063). Median (IQR) time to recovery was 8 days (7, 9) for infliximab and 9 days (8, 10) for placebo. Participants assigned to infliximab were more likely to have an improved clinical status at day 14 (OR 1.32, 95% CI 1.05–1.66). Twenty-eight-day mortality was 10.1% with infliximab versus 14.5% with placebo, with 41% lower odds of dying in those receiving infliximab (OR 0.59, 95% CI 0.39–0.90). No differences in risk of serious adverse events including secondary infections.

Conclusions: Infliximab did not demonstrate statistically significant improvement in time to recovery. It was associated with improved 14-day clinical status and substantial reduction in 28-day mortality compared with standard of care.

Trial registration: ClinicalTrials.gov (NCT04593940).

Keywords: infliximab, immune modulators, COVID-19, master protocol, shared placebo, TNF alpha inhibitors

INTRODUCTION

Immune dysregulation induced by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) is a major cause of morbidity and mortality.[1] While treatments directly targeting SARS-CoV-2 have shown significant impact in earlier stages of Covid-19,[2] immunoodulating agents provide benefit in hypoxia observed in later disease stages.[3-6]

Tumor necrosis factor alpha (TNF) is a pro-inflammatory cytokine that plays a role in nearly all acute inflammatory reactions.[7] In patients hospitalized for Covid-19, increased TNF levels are associated with more severe disease and death.[8] Inhibition of TNF reduces disease severity in mouse models of other respiratory viruses.[9] Infliximab, a TNF inhibitor that binds both soluble and transmembrane forms of TNF, is approved and commonly used to treat autoimmune diseases, including inflammatory bowel disease and rheumatoid arthritis. Data on infliximab use in Covid-19 treatment is limited.[10-12] However, patients who developed Covid-19 while on TNF inhibitors for other indications did not experience adverse outcomes.[13,14]

In April 2020, the National Institutes of Health (NIH) launched a public-private partnership, Accelerating Covid-19 Therapeutic Interventions and Vaccines (ACTIV), to develop a coordinated research response to Covid-19. ACTIV-1 IM was a master protocol designed to evaluate immunomodulatory agents in hospitalized patients with moderate/severe Covid-19. Infliximab was one of the agents included in ACTIV-1 IM based on its mechanism of action and efficacy and safety profile in inflammatory disorders. We report the results of a randomized, double-blind, placebo-controlled evaluation of infliximab compared with placebo in addition to standard of care.

METHODS

Study design

The ACTIV-1 IM master protocol was developed to allow for parallel investigation of efficacy and safety of multiple immunomodulators compared with a shared placebo, with standard of care

given as background therapy in both arms. Remdesivir (Gilead Sciences, Foster City, CA) was

provided as standard of care and given to eligible participants.

Eligibility

Eligibility criteria are outlined in the **Supplementary Appendix**. Briefly, adults ≥18 years with

confirmed moderate/severe SARS-CoV-2 infection admitted to hospital were eligible.

Radiological evidence of pulmonary involvement or oxygen saturation ≤94% on room air or

supplemental oxygen, mechanical ventilation, or extracorporeal membrane oxygenation

(ECMO) was required. Exclusion criteria included aspartate aminotransferase or alanine

aminotransferase >10 times the upper limit of normal, estimated glomerular filtration rate <30

mL/min, history of New York Heart Association class III/IV congestive heart failure, neutropenia,

lymphopenia, targeted immune therapies for any indication in the last four weeks or five drug

half-lives, or evidence of untreated tuberculosis or other untreated infections. Participants were

excluded from the infliximab substudy if they had a history of hepatosplenic T-cell lymphoma or

other lymphoma within five years before screening, history of or current diagnosis of multiple

sclerosis or other significant demyelinating condition.

Eligible participants were randomized in a two-stage process (**Supplementary**

Appendix). Between October 16, 2020, and December 31, 2021, 1061 participants were

randomized at 69 sites across five countries. Infliximab was administered on day 1 as a single-

dose intravenous infusion of 5 mg/kg over at least 2 hours. All participants received local

standard of care.

Procedures

Participants' clinical status was captured daily through day 29 if hospitalized. For discharged

participants, clinical status was assessed on days 8, 11, 15, and 29 in-person or by telephone if

in-person assessment was not possible. Day 60 follow-up was conducted by telephone

(**Supplementary Appendix**).

Outcomes and statistical analysis

The primary outcome was time to recovery evaluated up to day 29 (for clinical status on day

28). Recovery was defined as the first day on which participants attained category 6, 7, or 8 on

the 8-point ordinal scale (OS) defined in **Supplementary Appendix**. The primary efficacy

analysis was based on the Fine-Gray model with stratification by region and baseline disease

severity.[15]

Key secondary outcomes were mortality and clinical status assessed by 8-point OS at

days 14 and 28. Logistic and ordinal logistic regression models were used to estimate treatment

effects for mortality and 8-point OS endpoints. A multiple imputation approach was used to

account for the small amount of missing data for key secondary endpoints (**Statistical Analysis**

Plan [SAP]). The gatekeeping approach for controlling Type I error for the primary endpoint,

day 14 clinical status, and day 28 mortality is described in the SAP, including relevant p-value

cutoffs (**Table S1**).

Safety assessments included a composite endpoint of death, serious adverse events

(SAEs), or grade 3 (severe) and 4 (potentially life-threatening) AEs occurring through day 60.

Secondary infections as AEs of special interest through day 60 and discontinuation or

temporary suspension of trial-product administration for any reason were also assessed.

All efficacy and safety analyses reported are based on the modified intention-to-treat

(mITT) population consisting of all randomized participants who received at least one dose of

assigned study drug (infliximab or shared placebo), limiting shared placebo participants to those

eligible for infliximab.

RESULTS

Participants

Of 1061 participants who underwent randomization in this substudy, 531 were assigned to the

infliximab group and 530 to the shared placebo group (**Figure S1**). Ultimately, 517 participants

in the infliximab group and 516 in the shared placebo group received at least one dose of

assigned treatment and constitute the mITT population. Consistent with the protocol-specified

mITT definition, four participants were excluded post-randomization because they received

incorrect study drug. At baseline, 578 (56.0%) participants had moderate disease (52.1% and

3.9% OS 4 and 5) and 455 (44.0%) had severe disease (10.7% and 33.3% OS 2 and 3). Study

discontinuation by day 29 occurred in 4.8% in the infliximab group and 4.5% in the shared

placebo group. Mean age was 54.8 years and 60% were male. Overall, 652 (63.1%) were

White, 145 (14.0%) Black, 27 (2.6%) Asian, and 9 (0.9%) American Indian or Alaska Native; 502

(48.6%) were Hispanic or Latino. Overall, 975 (94.4%) participants received remdesivir, while

950 (92.0%) received corticosteroids (**Table 1**). Characteristics by region are shown in **Table**

S2, S3.

Primary outcome

Median time to recovery for infliximab was 1 day shorter than with shared placebo, but the

difference did not reach statistical significance (median 8 vs. 9 days; recovery rate ratio [RRR]

1.13, 95% confidence interval [CI] 0.99–1.29; p=0.063) (**Table 2, Figure S2**). Across OS

subgroups, the interaction p-value was 0.36 indicating no difference. Median time to recovery

among participants receiving mechanical ventilation/ECMO at enrollment (OS 2) was 23 days

for infliximab and >28 for shared placebo (RRR 1.117, 95% CI 0.612–2.039). For those on

noninvasive ventilation or high-flow oxygen (OS 3), median time to recovery was 11 days for

infliximab and 13 for shared placebo (RRR 1.326, 95% CI 1.039–1.693). Among those

hospitalized requiring supplemental oxygen (OS 4) and those hospitalized not requiring oxygen

(OS 5), median time to recovery was 6 versus 7 days (RRR 1.090, 95% CI 0.923–1.286) and 5

versus 4 days (RRR 0.767, 95% CI 0.404–1.455) (**Table 2**).

Key secondary outcomes

Mortality

Mortality at day 28 was 10.1% for infliximab and 14.5% for shared placebo (odds ratio [OR] for

death 0.59, 95% CI 0.39–0.90), resulting in 41% lower adjusted odds of dying (**Table 2, Figures**

1, 2A). When 28-day mortality was examined by OS, the interaction p-value was 0.31 indicating

no difference across subgroups. No difference in mortality was observed in the most severe

disease (OS 2) (OR 1.11, 95% CI 0.45–2.72) (**Figure 2C**). However, mortality decreased in

those who received infliximab compared with placebo for OS 3 (OR 0.52, 95% CI 0.29–0.91)

and OS 4/5 (OR 0.47, 95% CI 0.20–1.13). (**Figure 2D, E**) No deaths occurred in the OS 5

group. Day 14 mortality was 5.6% for infliximab and 8.1% for shared placebo (OR 0.63, 95% CI

0.36–1.08). Additional analysis revealed 60-day mortality rates of 12.6% in the infliximab group

and 16.5% in the shared placebo group, with a 32% reduction of odds of mortality observed (OR

0.68, 95% CI 0.46–0.999). The vast majority of those who died between day 29 and 60 were

intubated at day 28. In subgroup analyses, although point estimates indicated benefit for

infliximab in both c-reactive protein (CRP) subgroups, there was a trend for a stronger mortality

reduction at 28 days in participants with CRP >75 mg/L at baseline compared with CRP ≤75

mg/L.

Clinical status

The odds of improvement in clinical status at day 14 and day 28, as assessed by the OS, were

greater with infliximab compared with shared placebo (OR for improvement 1.32, 95% CI 1.05–

1.66) and (1.45 95% CI 1.14–1.85) **Tables 2 & S4, Figure S3**. For day 14, the interaction p-

value for subgroup analysis by OS was 0.90. On day 14, clinical status was improved with

infliximab compared with shared placebo for those at OS 2, 3, and 4 at randomization with the

strongest effect in OS 3 (OR 1.48, 95% CI 1.01–2.17). No difference was observed for OS 5. All

intention-to-treat data are presented in **Tables S5 and S6**.

Safety Assessments

No difference was observed in the composite safety endpoint at day 60 (32.9% vs. 33.7%;

hazard ratio [HR] 0.96, 95% CI 0.78–1.19). The number of participants with one or more grade 3

or 4 AEs was similar, (infliximab 146 [28.2%], shared placebo 131 [25.4%], risk difference 2.9;

95% CI -2.6 to 8.3) (**Table 3**). SAEs occurred in 125 (24.2%) participants in the infliximab group,

with seven events in six participants (1.2%), assessed by investigators as infliximab-related.

SAEs occurred in 130 (25.2%) participants in the control group, and the events were attributed

to trial product in seven of these participants (1.4%). One participant experienced grade 1 (mild)

infusion reaction in the infliximab group.

The percentage of participants who had any secondary infection was similar at day 60

with infliximab and shared placebo (79 [15.3%] vs. 72 [14.0%]) (**Table 3**). The most common

secondary infections were bacterial pneumonia, bloodstream and urinary tract infections (**Table

3**). All secondary infections were adjudicated by an independent safety officer.

DISCUSSION

This randomized, double-blind, placebo-controlled trial evaluated the addition of infliximab to

standard of care in hospitalized participants with moderate-to-severe Covid-19 pneumonia. The

trial displayed a strong, but not statistically significant, improvement in the primary endpoint of

time to recovery. We did, however, observe substantial improvements for key secondary

endpoints of 28-day mortality and 14-day clinical status. Infliximab was associated with a 41%

lower adjusted odds of death at 28 days in participants hospitalized with Covid-19. The mortality

benefit was observed across age groups, sexes, and races/ethnicities. Analysis demonstrated

that this mortality benefit was maintained to the completion of study at 60 days. Although some

additional deaths occurred after day 28 in both groups, these were mainly in participants

intubated at day 28. Sub-analysis by OS showed infliximab, when added to standard of care,

reduced mortality across a spectrum of disease severity with hospitalized participants requiring

supplemental oxygen and those on high-flow oxygen devices benefiting. In contrast, we

observed no benefit in those on mechanical ventilation or ECMO or those not requiring

supplemental oxygen. In another subgroup analysis, a stronger mortality benefit was observed

with the addition of infliximab to standard of care in participants with CRP >75mg/L at baseline

compared with those with CRP ≤75 mg/L.

Identification of anti-inflammatory agents to prevent or reverse dysregulated immune

cascades characteristic of severe Covid-19 has are an important area of investigation. The

addition of dexamethasone was shown to improve survival in patients requiring supplemental

oxygen and became embedded in standard of care.[3] However, the high morbidity and mortality

of Covid-19 and heterogeneity of therapeutic responses suggested the need for additional

immunomodulators. The JAK1/JAK2 inhibitor, baricitinib, and IL-6 antibody, tocilizumab, have

shown benefit for patients particularly in the setting of progressive respiratory failure.[4,6,16,17]

Results from these trials prompted the NIH guidelines panel to recommend tocilizumab or

baricitinib as a second immunomodulator in addition to dexamethasone for patients with

progressive respiratory failure and evidence of systemic inflammation.[18]

The infliximab data reported here, and a parallel report of the study of abatacept from

ACTIV-1 IM, adds to our knowledge by demonstrating mortality benefit for two separate

immunomodulators with unique and different mechanisms of action. Taken together with prior

studies, we now have substantial evidence that additional immunomodulation added to

corticosteroids reduces mortality and improves clinical outcomes in hospitalized patients with

moderate/severe Covid-19. The fact that this occurs with a variety of immunomodulatory agents,

all with different targets, is particularly interesting and establishing a better understanding of this

synergy is warranted at a mechanistic level.

Infliximab, in particular, and TNF blockade, in general, may be beneficial in Covid-19

management where elevated cytokine levels are seen in severe disease. TNF activates a wide

variety of immune cells and aberrant TNF-signaling is a central feature of cytokine release

syndrome.[19] One concern with TNF blockade in the setting of any viral illness is secondary

infection. Encouragingly, there were no new or negative safety signals observed during this trial,

including no differences between the infliximab and placebo groups in secondary infections.

These observations are consistent with infliximab's known safety profile having been used in the

treatment of inflammatory diseases for over twenty years. Likewise, the simplicity of a single

infusion and the global availability of infliximab could potentially increase the arsenal of

therapies available for treatment of moderate/severe Covid-19 in settings where current

guideline recommended therapies are not widely available.

With each completed trial we gain greater understanding of Covid-19, although

numerous clinical questions remain. In particular, there is lack of clarity around the optimal

management of patients requiring low-flow oxygen when first hospitalized. Subgroup analyses

presented here begin to address this. We show an apparent mortality benefit for

immunomodulators in both moderate and severe illness. Our results for participants with

moderate disease show improvement independent of inflammatory markers or clinical factors,

suggesting treatment with infliximab early in the disease process could provide benefit. In

contrast, our data do not support the addition of infliximab to dexamethasone in patients already

requiring mechanical ventilation. Additionally, subgroup analysis suggested a greater benefit in

both time to recovery and mortality in participants with CRP >75mg/L, suggesting patients with

evidence of more severe systemic inflammation at presentation to hospital may benefit more

from the addition of a second agent, such as infliximab, to dexamethasone. Future studies

should examine if a biomarker-driven approach facilitates early identification of people at-risk for

progression who could benefit from additional immunomodulation.

A limitation of this study is that the primary endpoint did not reach statistical significance.

Therefore, based on the pre-defined gatekeeping approach, the key secondary endpoints of 28-

day mortality and day 14 clinical status are not considered statistically significant although very

clinically relevant. A major challenge for this and other studies during the pandemic was the

appropriate selection of a primary endpoint in the setting of rapidly changing clinical scenarios.[20]

While a mortality primary endpoint is sometimes considered definitive and lacking in subjectivity,

it can require large participant numbers and potentially result in prolonged study recruitment and

delayed results. In addition, mortality does not encompass other patient-centered outcomes

which are a greater focus in endpoints such as time to recovery or clinical status. These issues

represent some factors that led to the selection of the primary endpoint in this study. Secondly,

this study was performed in the pre-omicron era. However, despite changes in the predominant

circulating variant over time and decreased disease severity overall, we continue to see patients

admitted to hospital with respiratory failure and evidence of immune dysregulation. Finally, while

there is a theoretical concern that a shared control group could negatively impact multiple

treatment evaluations in a platform study such as this, an examination of baseline

characteristics does not indicate this issue for ACTIV-1 IM. Integrity of the comparative analyses

was preserved through inclusion of only those participants in the shared placebo group who

were eligible to receive infliximab, and through the requirement that placebo participants were

shared only among agents active in the master protocol at the same time. More importantly, the

use of a shared control group has significant practical and ethical benefits.[21]

This report from ACTIV-1 IM shows that infliximab added to standard of care was

associated with clinically meaningful improvements in a number of key secondary outcomes,

including a 4% absolute reduction in 28-day mortality. As SARS-CoV-2 moves from being

pandemic to endemic, and while new variants continue to emerge, ongoing morbidity and

mortality from this disease is likely. Expanding our treatment toolbox and developing optimized

treatment strategies remains paramount.

Author Information

Jane A. O'Halloran, MD, PhD, Eyal Kedar, MD, Kevin J. Anstrom, PhD, Matthew W. McCarthy,

MD, Emily Ko, MD, PhD, Patricia Segura Nunez, MD, Cynthia Boucher, MS, P. Brian Smith,

MD, MPH, MHS, Reynold A. Panettieri, Jr, MD, Sabina Mendivil Tuchia de Tai, MD, Martin

Maillo, MD, Akram Khan, MD, Alfredo J. Mena Lora, MD, Matthias Salathe, MD, Gerardo Capo,

MD, Daniel Rodríguez Gonzalez, MD, Thomas F Patterson, MD, Christopher Palma, MD, ScM,

Horacio Ariza, MD, Maria Patelli Lima, MD, Anne M. Lachiewicz, MD, John Blamoun, MD,

Esteban C. Nannini, MD, Eduardo Sprinz, MD, Analia Mykietiuk, MD, Radica Alicic, Adriana M.

Rauseo, MD, Cameron R. Wolfe, MBBS, MPH, Britta Witting, Daniel K. Benjamin, Jr, MD, MPH,

PhD, Steven E. McNulty, MS, Pearl Zakroysky, S. Halabi, PhD, Sandra Butler, PhD, Jane

Atkinson, DDS, Stacey J. Adam, PhD, Richard Melsheimer, Soju Chang, MD, MPH, Lisa

LaVange, PhD, Michael Proschan, PhD, Samuel A. Bozzette, MD, PhD, William G. Powderly,

MD

Washington University St. Louis, St Louis, MO (J.A.O., A.R.A., W.G.P.); Weill Cornell Medicine

in New York (M.W.M., B.W.), University of North Carolina, Chapel Hill, NC (K.J.A., L.L., A.M.L.);

St. Lawrence Health, Potsdam, NY. (E.K.); Duke University Medical Center, Durham, NC (E.K.,

C.R.W.), Hospital Nacional Hipolito Unanue, Lima, Peru (P.S.N); National Center for Advancing

Translational Sciences (C.B., S.C., J.A., S.A.B.); Duke Clinical Research Institute (DCRI)

(D.K.B., P.B.S., P.Z., S.M., S.H.), Robert Wood Johnson Medical School, New Brunswick, NJ

(R.A.P.); Hospital Central de la Fuerza Aerea del Peru, Lima, Peru (S.M.T.T.); Sanatorio

Diagnostico, Santa Fe, Argentina (M.M.); Oregon Health and Science University, Portland, OR

(A.K.); University of Illinois at Chicago, Chicago, IL (A.M.L.); University of Kansas Medical

Acknowledgements

We thank the members of the ACTIV-1 Study Team (see Supplementary Appendix) for their many contributions in conducting the trial, the members of the data and safety monitoring board, William Blackwelder, PhD (chair) (University of Maryland School of Medicine, Baltimore,MD), Timothy G. Buchman, PhD, MD (Emory University School of Medicine, Atlanta, GA), Wilbur H. Chen, MD, MS (University of Maryland School of Medicine, Center for Vaccine Development and Global Health, Baltimore, MD), Lawrence H. Moulton, PhD (Johns Hopkins University, Bloomberg School of Public Health, Baltimore, MD), David M. Parenti, MD (The George Washington University School of Medicine, Washington, DC), Carol O. Tacket, MD (University of Maryland School of Medicine, Center for Vaccine Development and Global Health, Baltimore, MD), William Checkley, MD, PhD (Johns Hopkins University, Baltimore, MD), and Beatriz Grinsztejn, MD (Instituto Nacional de Infectologia Evandro Chagas-Fiocruz, Rio de Janeiro, Brazil) for their oversight; the Community Advisory Board, Larisa Caicedo, Ashish Cowlagi,

Anna Davis, Lincoln Larmond, Doug Lindsey, Bob Pearson and the patients and their families

for their altruism in participating in this trial.

Funding

This project has been funded with federal funds from the U.S. Department of Health and Human

Services, Office of the Assistant Secretary for Preparedness and Response, Biomedical

Advanced Research and Development Authority, under contract number

HHSO100201400002I/75A50120F33002. The findings and conclusions in this publication are

those of the authors and do not necessarily represent the views of the Department of Health

and Human Services or its components. Janssen provided infliximab for use in this trial but did

not provide any financial support. Gilead Sciences provided remdesivir for use in this trial but

did not provide any financial support. Employees of Janssen and Gilead Sciences participated

in discussions about protocol development and in weekly protocol team calls. The final trial

protocol was developed by the protocol chair, Dr. William Powderly, the IND sponsor Dr. Daniel

K. Benjamin, Jr, and a protocol development committee including representatives from the

National Center for Advancing Translational Sciences (NCATS). NCATS had a collaborative

role in the trial design, management, interpretation of the data and the preparation of the

manuscript along with the protocol chair, the study statisticians and the study writing group.

Infrastructure support and resources for research reported in this publication were provided in

part by NCATS of the National Institutes of Health under award number(s):UL1TR002345 to

Washington University in St. Louis; the Duke University – Vanderbilt University Medical Center

Trial Innovation Center (U24TR001608) (NCATS, Trial Innovation Network).

Data Sharing

A data sharing statement provided by the authors is available with this article.

REFERENCES

1. Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. Nat Rev Immunol 2020;20:355-62.

2. Gottlieb RL, Vaca CE, Paredes R, et al. Early Remdesivir to Prevent Progression to Severe Covid-19 in Outpatients. N Engl J Med 2022;386:305-15.

3. Group RC, Horby P, Lim WS, et al. Dexamethasone in Hospitalized Patients with Covid-19. N Engl J Med 2021;384:693-704.

4. Group RC. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. Lancet 2021;397:1637-45.

5. Gordon AC, Angus DC, Derde LPG. Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. Reply. N Engl J Med 2021;385:1147-9.

6. Marconi VC, Ramanan AV, de Bono S, et al. Efficacy and safety of baricitinib for the treatment of hospitalised adults with COVID-19 (COV-BARRIER): a randomised, double-blind, parallel-group, placebo-controlled phase 3 trial. Lancet Respir Med 2021;9:1407-18.

7. Feldmann M, Maini RN, Woody JN, et al. Trials of anti-tumour necrosis factor therapy for COVID-19 are urgently needed. Lancet 2020;395:1407-9.

8. Del Valle DM, Kim-Schulze S, Huang HH, et al. An inflammatory cytokine signature predicts COVID-19 severity and survival. Nat Med 2020;26:1636-43.

9. Hussell T, Pennycook A, Openshaw PJ. Inhibition of tumor necrosis factor reduces the severity of virus-specific lung immunopathology. Eur J Immunol 2001;31:2566-73.

10. Hachem H, Godara A, Schroeder C, et al. Rapid and sustained decline in CXCL-10 (IP-10) annotates clinical outcomes following TNFalpha-antagonist therapy in hospitalized patients with severe and critical COVID-19 respiratory failure. J Clin Transl Sci 2021;5:e146.

11. Fisher BA, Veenith T, Slade D, et al. Namilumab or infliximab compared with standard of

care in hospitalised patients with COVID-19 (CATALYST): a randomised, multicentre,

multi-arm, multistage, open-label, adaptive, phase 2, proof-of-concept trial. Lancet

Respir Med 2022;10:255-66.

12. Dolinger MT, Person H, Smith R, et al. Pediatric Crohn Disease and Multisystem

Inflammatory Syndrome in Children (MIS-C) and COVID-19 Treated With Infliximab. J

Pediatr Gastroenterol Nutr 2020;71:153-5.

13. Brenner EJ, Ungaro RC, Gearry RB, et al. Corticosteroids, But Not TNF Antagonists, Are

Associated With Adverse COVID-19 Outcomes in Patients With Inflammatory Bowel

Diseases: Results From an International Registry. Gastroenterology 2020;159:481-91

e3.

14. Melong Pianta Taleng CM, Lauper K, Gilbert B, et al. Incidence of COVID-19 in patients

treated with infliximab compared with patients treated with rituximab. RMD Open 2021;7.

15. Fine JP, Gray RJ. A Proportional Hazards Model for the Subdistribution of a Competing

Risk. J Am Stat Assoc 1999;94:496-509.

16. Kalil AC, Patterson TF, Mehta AK, et al. Baricitinib plus Remdesivir for Hospitalized

Adults with Covid-19. N Engl J Med 2021;384:795-807.

17. Investigators R-C, Gordon AC, Mouncey PR, et al. Interleukin-6 Receptor Antagonists in

Critically Ill Patients with Covid-19. N Engl J Med 2021;384:1491-502.

18. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19)

Treatment Guidelines. National Institutes of Health. Available at

https://www.covid19treatmentguidelines.nih.gov/. Accessed [8/21/2022].

19. Guo Y, Hu K, Li Y, et al. Targeting TNF-alpha for COVID-19: Recent Advanced and

Controversies. Front Public Health 2022;10:833967.

20. Douin DJ, Siegel L, Grandits G, et al. Evaluating Primary Endpoints for COVID-19

Therapeutic Trials to Assess Recovery. Am J Respir Crit Care Med 2022.

21.   LaVange L, Adam SJ, Currier JS, et al. Accelerating COVID-19 Therapeutic

Interventions and Vaccines (ACTIV): Designing Master Protocols for Evaluation of

Candidate COVID-19 Therapeutics. Ann Intern Med 2021;174:1293-300.

Figure Legends

Figure 1. Forest plot of 28-day mortality by subgroup (modified intent-to-treat population)

Figure 2. Kaplan-Meier curves of cumulative incidence of overall mortality at (A) day 28, (B) day

60, and by illness severity by ordinal scale for (C) mechanical ventilation or ECMO, (D) high-

flow oxygen devices or non-invasive ventilation, and (E) low-flow supplemental oxygen.

Table 1. Demographics and baseline characteristics (modified intent-to-treat)

Demographics and baseline characteristics	Infliximab (N=517)	Shared Placebo (N=516)	Total (N=1033)
Age, mean (SD), yrs	54.7 (14.94)	54.9 (14.65)	54.8 (14.79)
Male sex, no. (%)	325 (62.9%)	298 (57.8%)	623 (60.3%)
Race, no. (%)			
White	319 (61.7%)	333 (64.5%)	652 (63.1%)
Black or African American	77 (14.9%)	68 (13.2%)	145 (14.0%)
American Indian or Alaska Native	5 (1.0%)	4 (0.8%)	9 (0.9%)
Asian	11 (2.1%)	16 (3.1%)	27 (2.6%)
Native Hawaiian or Other Pacific Islander	1 (0.2%)	0 (0.0%)	1 (0.1%)
Multiracial	2 (0.4%)	2 (0.4%)	4 (0.4%)
Other	76 (14.7%)	71 (13.8%)	147 (14.2%)
Unknown	26 (5.0%)	22 (4.3%)	48 (4.6%)
Hispanic or Latino ethnicity, no. (%)	252 (48.7%)	250 (48.4%)	502 (48.6%)
Disease severity at baseline[*], no. (%)			
Severe disease	228 (44.1%)	227 (44.0%)	455 (44.0%)
Moderate disease	289 (55.9%)	289 (56.0%)	578 (56.0%)
Clinical status (8-point ordinal scale) at baseline, no. (%)			
1. Death	0	0	0

2. Hospitalized, on invasive ventilation or ECMO	58 (11.2%)	53 (10.3%)	111 (10.7%)
3. Hospitalized, on non-invasive ventilation or high flow oxygen devices	170 (32.9%)	174 (33.7%)	344 (33.3%)
4. Hospitalized, requiring supplemental oxygen	268 (51.8%)	270 (52.3%)	538 (52.1%)
5. Hospitalized, not requiring supplement oxygen, requiring ongoing medical care	21 (4.1%)	19 (3.7%)	40 (3.9%)
6. Hospitalized, not requiring supplement oxygen, not requiring ongoing medical care	0	0	0
7. Not hospitalized, limitations on activity and/or requiring home oxygen	0	0	0
8. Not hospitalized, no limitations on activities	0	0	0
Geographic region, no. (%)			
Argentina	56 (10.8%)	59 (11.4%)	115 (11.1%)
Brazil	54 (10.4%)	54 (10.5%)	108 (10.5%)
Mexico	15 (2.9%)	16 (3.1%)	31 (3.0%)
Peru	62 (12.0%)	56 (10.9%)	118 (11.4%)
USA - Northeast	119 (23.0%)	118 (22.9%)	237 (22.9%)
USA - Midwest	81 (15.7%)	77 (14.9%)	158 (15.3%)
USA - South	93 (18.0%)	94 (18.2%)	187 (18.1%)
USA - West	37 (7.2%)	42 (8.1%)	79 (7.6%)
Days from symptom onset, mean (SD)	-9.9 (4.41)	-9.9 (5.61)	-9.9 (5.04)
BMI, mean (SD), kg/m^2	32.1 (7.97)	32.7 (8.11)	32.4 (8.04)

Comorbidities, no. (%)

Hypertension	207 (40.0%)	209 (40.5%)	416 (40.3%)
Obesity (BMI ≥30 kg/m^2)	268 / 507 (52.9%)	299 / 502 (59.6%)	567 / 1009 (56.2%)
Diabetes mellitus	138 (26.7%)	144 (27.9%)	282 (27.3%)
Coronary artery disease	38 (7.4%)	27 (5.2%)	65 (6.3%)
History of heart failure	16 (3.1%)	14 (2.7%)	30 (2.9%)
History of cancer	34 (6.6%)	34 (6.6%)	68 (6.6%)
Asthma	35 (6.8%)	53 (10.3%)	88 (8.5%)
Chronic obstructive pulmonary disease	24 (4.6%)	26 (5.0%)	50 (4.8%)
Tuberculosis	3 (0.6%)	4 (0.8%)	7 (0.7%)
HIV/AIDS	2 (0.4%)	4 (0.8%)	6 (0.6%)
Severe liver disease	3 (0.6%)	2 (0.4%)	5 (0.5%)
Severe kidney disease	3 (0.6%)	7 (1.4%)	10 (1.0%)

Concomitant medication, no. (%)

Remdesivir (Day 1 - Day 5)	486 (94.0%)	489 (94.8%)	975 (94.4%)
Corticosteroids (Day 1 - Day 5)	468 (90.5%)	482 (93.4%)	950 (92.0%)
Tocilizumab (any time post randomization)	7 (1.4%)	12 (2.3%)	19 (1.8%)
Baricitinib (any time post randomization)	5 (1.0%)	11 (2.1%)	16 (1.5%)

415 BMI indicates body mass index; ECMO, extracorporeal membrane oxygenation; SD, standard deviation; USA, United States of America.
416 *Disease severity at baseline calculated as moderate disease = ordinal 5 + ordinal 4, severe disease = ordinal 3 + ordinal 2
417

418   **Table 2. Primary and Key Secondary Endpoints (Modified Intent-to-Treat Population)**

	Infliximab	Shared Placebo	Infliximab versus Shared Placebo	
	Proportion of Participants		Estimate (95% CI)	p value
Recovery through Day 28[*]				
Overall	416/517 (80.5%)	398/516 (77.1%)	1.130 (0.993, 1.286)	0.0631
Baseline 8-point ordinal scale				
2. Hospitalized, on invasive ventilation or ECMO	24/58 (41.4%)	20/53 (37.7%)	1.117 (0.612, 2.039)	
3. Hospitalized, on non-invasive ventilation or high flow oxygen devices	124/170 (72.9%)	111/174 (63.8%)	1.326 (1.039, 1.693)	
4. Hospitalized, requiring supplemental oxygen	247/268 (92.2%)	248/270 (91.9%)	1.090 (0.923, 1.286)	
5. Hospitalized, not requiring supplement oxygen, requiring ongoing medical care	21/21 (100.0%)	19/19 (100.0%)	0.767 (0.404, 1.455)	
Baseline C-reactive protein				
≤ 75 mg/L	187/216 (86.6%)	182/210 (86.7%)	1.018 (0.829, 1.250)	
> 75 mg/L	171/228 (75.0%)	147/209 (70.3%)	1.203 (0.974, 1.486)	
Mortality at Day 28[†]				
Overall	52/517 (10.1%)	75/516 (14.5%)	0.593 (0.390, 0.904)	
Baseline 8-point ordinal scale				
2. Hospitalized, on invasive ventilation or ECMO	18/58 (31.0%)	16/53 (30.2%)	1.106 (0.449, 2.723)	
3. Hospitalized, on non-invasive ventilation or high	26/170 (15.3%)	43/174 (24.7%)	0.516 (0.291, 0.914)	

flow oxygen devices			
4 or 5: Hospitalized, with or without supplemental oxygen, requiring ongoing medical care[‡]	8/289 (2.8%)	16/289 (5.5%)	0.470 (0.195, 1.131)
Baseline C-reactive protein			
≤ 75 mg/L	19/216 (8.8%)	18/210 (8.6%)	0.788 (0.372, 1.668)
> 75 mg/L	28/228 (12.3%)	42/209 (20.1%)	0.551 (0.304, 0.996)
Clinical status at Day 14 (N)[§]			
Overall	502	501	1.318 (1.047, 1.659)
Baseline 8-point ordinal scale			
2. Hospitalized, on invasive ventilation or ECMO	58	53	1.284 (0.655, 2.519)
3. Hospitalized, on non-invasive ventilation or high flow oxygen devices	165	169	1.477 (1.008, 2.165)
4. Hospitalized, requiring supplemental oxygen	258	261	1.260 (0.914, 1.737)
5. Hospitalized, not requiring supplement oxygen, requiring ongoing medical care	21	18	0.972 (0.367, 1.090)
Mortality at Day 60	65/517 (12.6%)	85/516 (16.5%)	0.679 (0.461, 0.999)

419 *Time to recovery calculated as recovery rate ratio using stratified Fine-Gray model. A number greater than 1 favors Infliximab.
420 [†]Mortality at 28 days calculated as odds of dying using logistic regression. A number less than 1 favors Infliximab.
421 [‡]No deaths were reported for baseline 8-point ordinal scale 5. Baseline 8-point ordinal scale values 4 and 5 were combined to address modeling issues.
422 [§]Clinical status at 14 days calculated as proportional odds model by ordinal logistic regression. A number greater than 1 favors Infliximab.
423 CI indicates confidence interval; ECMO, extracorporeal membrane oxygenation.
424
425

Table 3. Safety composite and adverse events through day 60

Safety Composite and Adverse Events	Infliximab (N=517)	Shared Placebo (N=516)	Total (N=1033)	Risk Difference (95% CI)
Safety Composite*	170 (32.9%)	174 (33.7%)	344 (33.3%)	-0.8 (-6.6, 4.9)
SAE	125 (24.2%)	130 (25.2%)	255 (24.7%)	-1.0 (-6.3, 4.2)
Grade 3 or 4 AE	146 (28.2%)	131 (25.4%)	277 (26.8%)	2.9 (-2.6, 8.3)
Grade 4 AE	63 (12.2%)	64 (12.4%)	127 (12.3%)	-0.2 (-4.3, 3.8)
Grade 3 AE	124 (24.0%)	97 (18.8%)	221 (21.4%)	5.2 (0.2, 10.2)
Secondary Infections				
Any Secondary Infection/Superinfection	79 (15.3%)	72 (14.0%)	151 (14.6%)	
Confirmed	24 (4.6%)	26 (5.0%)	50 (4.8%)	
Probable	55 (10.6%)	46 (8.9%)	101 (9.8%)	
Any Bacterial	71 (13.7%)	55 (10.7%)	126 (12.2%)	
Bacterial Pneumonia	49 (9.5%)	36 (7.0%)	85 (8.2%)	
Bloodstream infections	17 (3.3%)	15 (2.9%)	32 (3.1%)	
Urinary tract infections	15 (2.9%)	16 (3.1%)	31 (3.0%)	
Other bacterial infections	6 (1.2%)	2 (0.4%)	8 (0.8%)	
Tuberculosis	1 (0.2%)	0 (0.0%)	1 (0.1%)	
Any Fungal	14 (2.7%)	23 (4.5%)	37 (3.6%)	

Oral/oropharyngeal candidiasis	5 (1.0%)	8 (1.6%)	13 (1.3%)
Invasive candidiasis	4 (0.8%)	5 (1.0%)	9 (0.9%)
Other fungi infections	3 (0.6%)	8 (1.6%)	11 (1.1%)
Mold Infection (Aspergillus species, mucormycosis and other)	2 (0.4%)	2 (0.4%)	4 (0.4%)
Any Viral	2 (0.4%)	2 (0.4%)	4 (0.4%)

427   *The safety composite endpoint includes any of the following events: deaths, SAEs and Grade 3 or 4 AEs through Day 60.
428   AE indicates adverse event: CI, confidence interval; SAE, serious adverse event.
429
430

Figure 1.

Forest Plot of 28-Day Mortality by Subgroup
Modified Intent-to-Treat Population

434　**Figure 2A.**

436   **Figure 2B.**

439 **Figure 2C.**

442 **Figure 2D.**

445 **Figure 2E.**

446